



**HAL**  
open science

## Complete mitogenome of the giant invasive hammerhead flatworm *Bipalium kewense*

Romain Gastineau, Jean-Lou Justine, Claude Lemieux, Monique Turmel,  
Andrzej Witkowska

### ► To cite this version:

Romain Gastineau, Jean-Lou Justine, Claude Lemieux, Monique Turmel, Andrzej Witkowska. Complete mitogenome of the giant invasive hammerhead flatworm *Bipalium kewense*. *Mitochondrial DNA Part B Resources*, 2019, 4 (1), pp.1343-1344. 10.1080/23802359.2019.1596768 . hal-02090399


**HAL Id: hal-02090399**

**<https://hal.science/hal-02090399>**

Submitted on 21 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

# Complete mitogenome of the giant invasive hammerhead flatworm *Bipalium kewense*

Romain Gastineau, Jean-Lou Justine, Claude Lemieux, Monique Turmel & Andrzej Witkowski

To cite this article: Romain Gastineau, Jean-Lou Justine, Claude Lemieux, Monique Turmel & Andrzej Witkowski (2019) Complete mitogenome of the giant invasive hammerhead flatworm *Bipalium kewense*, Mitochondrial DNA Part B, 4:1, 1343-1344, DOI: [10.1080/23802359.2019.1596768](https://doi.org/10.1080/23802359.2019.1596768)

To link to this article: <https://doi.org/10.1080/23802359.2019.1596768>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 02 Apr 2019.



Submit your article to this journal [↗](#)



View Crossmark data [↗](#)

## Complete mitogenome of the giant invasive hammerhead flatworm *Bipalium kewense*

Romain Gastineau<sup>a</sup>, Jean-Lou Justine<sup>b</sup>, Claude Lemieux<sup>c</sup>, Monique Turmel<sup>c</sup> and Andrzej Witkowski<sup>a</sup>

<sup>a</sup>Palaeoceanology Unit, Faculty of Geosciences, and Natural Sciences Research and Educational Centre, University of Szczecin, Szczecin, Poland; <sup>b</sup>Institut Systématique Évolution Biodiversité (ISYEB) Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Paris, France; <sup>c</sup>Département de biochimie, de microbiologie et de bio-informatique, Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec, Canada

### ABSTRACT

We determined the sequence of the complete mitogenome of *Bipalium kewense*, an invasive land flatworm with worldwide distribution. All 37 genes encoded on this 15,666-bp genome reside on the same DNA strand. Phylogenomic analyses of mitochondrial proteins from taxa of the order Tricladida associated *B. kewense* with another invasive species of land planarians (*Obama* sp.).

### ARTICLE HISTORY

Received 11 February 2019  
Accepted 13 March 2019

### KEYWORDS


*Bipalium*; flatworm;  
invasive; Tricladida;  
Geoplanidae

The platyhelminth *Bipalium kewense* (family Geoplanidae) is an invasive land flatworm originating from southeastern Asia (Winsor 1983; Justine et al. 2018). Like other Bipaliinae, it features a hammerhead but is exceptional by its giant size (up to 30 cm). Remarkably, *B. kewense* and *Bipalium adventitium* were shown to produce tetrodotoxin, making these species the first known terrestrial invertebrates to possess this neurotoxin (Stokes et al. 2014). The current worldwide distribution of *B. kewense* is believed to be the result of man-mediated transportation of plants (Winsor 1983; Justine et al. 2018). Given that this invasive species preys on earthworms in cultivated areas (Justine et al. 2018), its dispersal has potentially devastating ecological impact. Recently, analysis of *cox1* from 22 *B. kewense* specimens collected from Metropolitan France, other European countries, and overseas French territories on three continents revealed 100% sequence identity for this mitochondrial gene (Justine et al. 2018), suggesting that worldwide dispersal of a single invasive clone occurred through asexual reproduction.

We sequenced the complete mitogenome of *B. kewense*. DNA was obtained from a specimen that was previously sampled for *cox1* analysis; this specimen collected in Ustaritz (Pyrénées-Atlantiques, France) is registered as MNHN JL184A

in the collections of the Muséum National d'Histoire Naturelle (Paris, France). Sequencing was performed by the Beijing Genomic Institute (Shenzhen, China) on the BGISEQ-500 platform. A total of 60 million paired-end reads were assembled using SPAdes 3.12.0 (Bankevich et al. 2012) and a single mitogenome contig was recovered. Gene identification was carried out using MITOS (Bernt et al. 2013).

The *B. kewense* mitogenome (GenBank: MK455837) is 15,666 bp long and encodes 13 proteins, two rRNAs, and 22 tRNAs. All genes are located on the same DNA strand, with the protein- and rRNA- coding genes ordered as follows: *cox1*, *ND6*, *ND5*, *cox3*, *atp6*, *ND1*, *cox2*, *ND3*, *ND2*, *rrnS*, *rrnL*, *cob*, *ND4L*, and *ND4*. We inferred a maximum likelihood tree using the mitogenome-encoded proteins of *B. kewense* and the few Tricladida taxa whose complete mitogenomes were available (Figure 1). The sequences of the proteins shared by all taxa were concatenated and analyzed with RAXML version 8.2.12 (Stamatakis 2014) using the MtArt model of sequence evolution (Abascal et al. 2007) and the Polycladida *Prosthlostomum siphuncululus* (KT363736) as outgroup. *B. kewense* was found to strongly associate with the only other Geoplanidae taxon in the tree (*Obama* sp., most probably *O. nungara*), a strain from South America which is also an invasive land flatworm.



**Figure 1.** Maximum Likelihood tree obtained on concatenated amino-acid sequences of all mitochondrial protein-coding genes from *Bipalium kewense* and other flatworms, using the MtArt model of evolution and after 100 bootstrap replications.

### Disclosure statement

The authors report no conflict of interest. The authors alone are responsible for the content and writing of this article.

### References

- Abascal F, Posada D, Zardoya R. 2007. MtArt: a new model of amino acid replacement for Arthropoda. *Mol Biol Evol.* 24:1–5.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19:455–477.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol.* 69:313–319.
- Justine J-L, Winsor L, Gey D, Gros P, Thévenot J. 2018. Giant worms chez moi! Hammerhead flatworms (Platyhelminthes, Geoplanidae, *Bipalium* spp., *Diversibipalium* spp.) in metropolitan France and overseas French territories. *PeerJ.* 6:e4672.
- Stamatakis A. 2014. RAxML Version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 30:1312–1313.
- Stokes AN, Ducey PK, Neuman-Lee L, Hanifin CT, French SS, Pfreder ME, Brodie ED, Brodie ED. 2014. Confirmation and distribution of tetrodotoxin for the first time in terrestrial invertebrates: two terrestrial flatworm species (*Bipalium adventitium* and *Bipalium kewense*). *PLoS One.* 9:e100718.
- Winsor L. 1983. A revision of the Cosmopolitan land planarian *Bipalium kewense* Moseley, 1878 (Turbellaria: Tricladida: Terricola). *Zoo J Linn Soc.* 79:61–100.