



A conserved fungal hub protein involved in adhesion and drug resistance in the human pathogen *Candida albicans*

Hélène Martin-Yken, Tina Bedekovic, Alexandra Brand, Mathias M. Richard, Sadri Znaidi, Christophe d'Enfert, Etienne Dague

► To cite this version:

Hélène Martin-Yken, Tina Bedekovic, Alexandra Brand, Mathias M. Richard, Sadri Znaidi, et al.. A conserved fungal hub protein involved in adhesion and drug resistance in the human pathogen *Candida albicans*. *The Cell Surface*, 2018, 4, pp.10-19. 10.1016/j.tcs.2018.10.002 . hal-02091662

HAL Id: hal-02091662

<https://hal.science/hal-02091662>

Submitted on 4 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A conserved fungal hub protein involved in adhesion and drug
resistance in the human pathogen *Candida albicans*.

Hélène Martin-Yken^{*1,2}, Tina Bedeković³, Alexandra Brand³, Mathias L. Richard⁴, Sadri
Znaidi^{5,6} Christophe d'Enfert and Etienne Dague².

1 INSA, UPS, INP, ISAE, LAAS, Université de Toulouse, Toulouse, France;
UMR792 Ingénierie des Systèmes Biologiques et des Procédés, INRA, Toulouse,
France; UMR5504, CNRS, Toulouse, France.

2 LAAS CNRS UPR 8001, Université de Toulouse, Toulouse, France.

3 MRC Centre for Medical Mycology, School of Medicine, Medical Sciences &
Nutrition, University of Aberdeen, Aberdeen, United Kingdom.

4 INRA, UMR1319 Micalis, AgroParisTech, Jouy-en-Josas, France.

5 Institut Pasteur de Tunis, Laboratoire de Microbiologie Moléculaire, Vaccinologie et
Développement Biotechnologique, 13 Place Pasteur, Tunis-Belvédère, Tunisia

6 Institut Pasteur, INRA, Unité Biologie et Pathogénicité Fongiques, 25 rue du
Docteur Roux, Paris, France.

*For correspondance: helene.martin@insa-toulouse.fr

Declarations of interest: none.

ABSTRACT

Drug resistance and cellular adhesion are two key elements of the worldwide dissemination and prevalence of the human fungal pathogen *Candida albicans*. Smi1 belongs to a family of hub proteins conserved among the fungal kingdom, whose functions in cellular signaling affect morphogenesis, cell wall synthesis and stress resistance. The data presented here indicate that *C. albicans SMII* is a functional homolog of *Saccharomyces cerevisiae KNR4* and is involved in the regulation of the fungal cell wall synthesis. Expression of *SMII* in the *S. cerevisiae knr4Δ* null mutant rescued its sensitivity to Caspofungin and to heat stress. Deletion of *SMII* in *C. albicans* resulted in sensitivity to the cell-wall-perturbing compounds Calcofluor White and Caspofungin. Analysis of wild-type and mutant cells by Atomic Force Microscopy showed that the Young's Modulus (stiffness) of the cell wall was significantly reduced upon deletion of *SMII*, while cell surface adhesion measured by Single Force Spectroscopy showed that the surface expression of adhesive molecules was also reduced in the mutant. Over-expression of *SMII gene*, on the contrary, drastically increased the cell surface adhesion phenotype vs the control strain. Finally, Smi1 cellular localization as cytoplasmic patches and concentrated spots at the sites of new cell wall synthesis is consistent with its role in the control of cell wall synthesis and appears conserved with the ones of its counterparts in *S. cerevisiae* and *Neurospora crassa* respectively for yeast cells and hyphae of *C. albicans*.

INTRODUCTION

Fungal infections are responsible of the death of about 1.5 million people per year worldwide. Hemiascomycetous yeasts of the genus *Candida* are the second most numerous agents of fungal infections worldwide, with a prominent contribution by *Candida albicans*, which causes over 400,000 cases of life-threatening systemic infections and 200,000 deaths per year (Brown et al., 2012). Only four classes of antifungal (AF) drugs are available for patient treatment and the emergence of resistance is becoming a serious concern. The fungistatic group of Azoles and the more recently-developed fungicidal group of Echinocandins constitute the two major classes of AF used to treat patients. Azoles block the biosynthesis of ergosterol - an essential sterol for fungal cell membranes - by targeting the cytochrome P450 14- α demethylase enzyme, Erg11, which catalyzes the conversion of lanosterol to ergosterol, thereby affecting membrane integrity and inhibiting fungal growth (Kathiravan et al., 2012; Vanden Bossche et al., 1995). On the other hand, Echinocandins, a class of compounds developed between 2001 and 2006, target the catalytic sub-unit of the β -1,3 glucan synthase protein complex, Fks1 (Odds et al., 2003). Studies of the molecular mechanisms of resistance to these two classes of AF compounds (recently reviewed in (Scorzoni et al., 2017)) have revealed that there are three major mechanisms leading to resistance: overexpression of multidrug efflux pump-encoding genes, notably CDR1, CDR2 and MDR1 (Sanglard and Odds, 2002; Sanglard et al., 1995), point mutations in the target proteins (ex: Erg11, Fks1) and alteration in the expression of proteins involved in sensitivity to the drug (ex: Erg3, Erg11). In addition, the formation of fungal biofilms can also be considered as a form of antifungal resistance due to the ability of the biofilm extracellular matrix (ECM) to bind and entrap antifungal compounds, particularly Azoles and Amphotericin B (Desai et al., 2014; Taff et al., 2013; Vedyappan et al., 2010; Zarnowski et al., 2014).

In this context, researchers have sought alternative antifungal targets and/or ways to improve the fungicidal effect of existing AFs. Such approaches notably involve targeting chaperones such as Hsp90 or components of stress signaling pathways, since these targets are more likely to simultaneously affect resistance to different classes of AF, morphogenesis mechanisms, cellular fitness and adaptation to changing environments. Key studies have been conducted in these areas by, for example, Brown and colleagues (Brown et al., 2010) and

Cowen and coworkers (Singh et al., 2009). Works of this latter group established the complex connections between Pkc1, Hsp90 and calcineurin suggesting interesting new strategies to treat fungal infections (LaFayette et al., 2010). However, these cellular targets suffer from a major drawback in that they are conserved in mammalian host cells, which make achieving fungal specificity a real challenge. Factors that regulate the pathogen's cell wall therefore remain a strong target for new, fungus-specific, therapeutic approaches.

Here we describe the role of Smi1, a *C. albicans* protein homologous to the *Saccharomyces cerevisiae* hub protein Knr4 which interacts physically with both the Slr2 MAP kinase and calcineurin, thus connecting the two primary signaling pathways involved in cell wall maintenance during stress: the cell wall integrity pathway (CWI) and the calcineurin pathway (Dagkessamanskaia et al., 2010; Martin-Yken et al., 2016). Knr4 is required for resistance to cell wall stress induced by elevated temperature or the presence of antifungal compounds, including Caspofungin (Lesage et al., 2004; Markovich et al., 2004). Knr4 also plays a role in filamentous and pseudohyphal growth, mucin secretion and agar invasion (Birkaya et al., 2009). Similarly, GS1 protein, the homolog of Knr4 and Smi1 in the model filamentous fungi *Neurospora crassa*, is also involved in the control of morphogenesis, Caspofungin sensitivity and new cell wall synthesis notably β -glucans (Enderlin and Selitrennikoff, 1994; Resheat-Eini et al., 2008; Seiler and Plamann, 2003).

The *Candida albicans* genome encodes 2 homologs of *KNR4*: *SMI1* and *SMI1B*. Previous studies have shown that deletion of *SMI1* affects cell wall β -glucan synthesis, biofilm formation and biofilm extracellular matrix production, as well as biofilm-associated resistance to Fluconazole (Nett et al., 2011). Global transcriptomic studies indicate that *SMI1* expression is induced in hyphal and planktonic cells by Cyr1 and is also biofilm-induced, while it is repressed by the Hap43 regulatory protein and Caspofungin (Liu et al., 2005). Much less is known about *SMI1B*, although it appears to be the closest homolog of *S. cerevisiae* *KNR4* according to phylogeny.

In this paper we further characterize the function of *SMI1* in *C. albicans*. We provide evidence that Smi1 is a functional homolog of *S. cerevisiae* protein Knr4, that its accurate expression is critical for the regulation of fungal cell wall integrity and biophysical properties, and that the cellular localization of Smi1-GFP in yeast and hyphal cells is consistent with the ones observed for its counterparts, *i.e.* Knr4 in *S. cerevisiae* yeasts and GS1 in the hyphae of *N. crassa*.

Material and Methods

Strains and growth media

The *C. albicans* and *S. cerevisiae* strains used in this study are listed in Table 1.

Depending on experimental conditions, yeast strains were grown in YPD (1 % (W/V) yeast extract, 2 % (W/V) peptone, and 1 % (W/V) dextrose), YP (1 % (W/V) yeast extract, 2 % (W/V) peptone) supplemented with 10 % Fetal Bovine Serum (FBS), or SD (synthetic dextrose, 0.67 % (W/V) yeast nitrogen base (YNB; Difco) with 2 % glucose) supplemented as necessary with arginine, histidine or uridine (20 mg/l). Agar (2 %) was used for growth on solid medium. *Escherichia coli* strains TOP10 (Invitrogen) or DH5 α were used for DNA cloning and maintenance of the plasmid constructs.

Plasmid construction and generation of epitope-tagged or mutant strains:

C. albicans *SMI1* gene was PCR amplified with primers: Sense SMI1 New start and Antisense SMI1 (sequences in Supplementary Material Table S1). The PCR product was then cloned in the *S. cerevisiae* expression vector YEplac195 *PGK/CYC1* between *PGK* promotor *P_{PGK1}* and *CYC1* terminator sequences (personal gift of Dr J.L. Parrou, based on YEplac195 (Gietz and Sugino, 1988)), thus yealding p*SMI1* plasmid. Plasmid p*KNR4* expressing the *S. cerevisiae* *KNR4* gene with its own promotor and terminator on a multicopy vector has been described previously (Martin et al., 1999).

S. cerevisiae cells were transformed according to the lithium acetate method (Gietz and Woods, 2006). *C. albicans* cells were transformed using the lithium-acetate protocol of Walther and Wendland (Walther and Wendland, 2003), followed by selection of transformants for uridine, arginine or histidine prototrophy when using the *URA3*, *ARG4* or the *HIS1* markers, respectively.

Construction of *C. albicans* *smi1* Δ/Δ and *smi1B* Δ/Δ knock-out mutants (Table 1) used PCR-generated *ARG4* and *HIS1* disruption cassettes flanked by 120 base pairs of target homology region (primer sequences are provided in supplementary material S1) as described by (Gola et al., 2003) and (Schaub et al., 2006). Independent transformants were produced and the gene replacements were verified by PCR on whole yeast cells as described previously (Gola et al., 2003; Schaub et al., 2006).

The *SMI1* (C1_07870C_A) gene was amplified using primers: SMI1 Forward and SMI1 Reverse (sequences provided in supplementary material Table S1). The resulting 1.6Kb PCR product was purified and inserted in the GTW sequences of pEntry (Gateway systemTM,

Invitrogen). Construction of the *Smi1* over-expressing strain was performed by recombination of pEntry-SMI1 plasmid and pCip10-P_{TDH3}-GTW plasmid (Chauvel et al., 2012) also using the Gateway systemTM (Invitrogen), followed by genome integration through targeted homologous recombination at the genomic *RPS10* locus in the host strain BWP17.

GFP tagging of Smi1

Smi1 was C-terminally-tagged with GFP by amplifying GFP-NAT cassette from the pGFP-NAT1 plasmid (Milne et al., 2011) using primers g*Smi1*_F^a and g*Smi1*_R^a containing 100 bp of flanking homology to the *SMI1* terminator and C-terminal of the *SMI1* ORF (C1_07870C_A), respectively (Figure X –primer table). Transformants were selected on YPD agar containing 300 µg/ml nourseothricin (Sigma). Integration of the GFP-NAT cassette was confirmed by PCR using primers s*Smi1*_F and sFP_R, which anneal to the chromosome outside the targeted region and within the cassette, respectively (primer sequences are provided in Supplementary material Table S1).

Microscopy of Smi1-GFP

Yeast were grown for 2 h at 30 °C in YNB medium containing amino acids and (NH₄)₂SO₄ (Sigma Aldrich). Morphogenesis was induced at 37 °C for 3 h in 20 % FCS, 2 % glucose. Cells were imaged on µ-slides (Ibidi, Martinsried, Germany). Images of *Smi1*-GFP localization were captured on an UltraVIEW® VoX spinning disk confocal microscope (Nikon, Surrey, UK), using a 488 nm laser. Multiple Z-stack images were acquired and Z-stack projections at maximum intensity were created using Volocity 6.3 software (Perkin Elmer).

Phenotypic Sensitivity tests:

Drop tests to evaluate the sensitivity of different strains and mutants to cell wall affecting drugs were performed as previously described (Ram et al., 1998) with minor modifications (Martin et al., 1999). Briefly, yeast cells were grown in liquid YPD to OD₆₀₀ of 1+/- 0.1, then concentrated by centrifugation and resuspended in sterile water to an OD₆₀₀ of 8. Serial dilutions of 1/1, 1/10, 1/100 and 1/1000 were then spotted on solid media containing either calcofluor white or Caspofungin at the indicated concentrations. Growth was scored and photographs taken after 48h of incubation at 30 °C, or at 37 °C for testing the sensitivity to elevated temperature.

Atomic Force Microscopy:

Sample preparation for AFM experiments.

Yeast cells were concentrated by centrifugation, washed two times in acetate buffer (18 mM CH₃COONa, 1mM CaCl₂, 1mM MnCl₂, pH5.2), resuspended in the same buffer, and immobilized on polydimethylsiloxane (PDMS) stamps prepared as described by Dague et al. (Dague et al., 2011; Formosa et al., 2014a). Briefly, freshly oxygen-activated microstructured PDMS stamps were covered with a total of 100 µl of cell suspension and allowed to stand for 15 min at room temperature. Yeast cells were then deposited into the stamps microstructures by convective (capillary) assembly.

Force curves were then recorded in acetate buffer in quantitative-imaging mode (JPK Instruments, 2011, QITM mode-quantitative imaging with Nano-Wizard 3 AFM), (Chopin et al., 2013; Formosa et al., 2014b; Smolyakov et al., 2016) with MLCT AUWH cantilevers (nominal spring constants: 0.01, 0.1, and 0.5 N/m). For imaging, cantilevers with a spring constant of 0.01 N/m were used. For force spectroscopy experiments, cantilevers with spring constants of 0.1 and 0.5 N/m were used. The applied force was kept at 1 nN for both imaging and force spectroscopy. For imaging and force spectroscopy, we used an AFM Nanowizard III (JPK Instruments, Berlin, Germany). The spring constant of each cantilever was determined by the thermal-noise method (Hutter and Bechhoefer, 1993).

For elasticity measurements, force maps of 32 by 32 or 64 by 64, hence either 1024 or 4096 force curves were recorded on a 1 µm x 1 µm square area on top of the cells. The force-distance curves recorded were transformed into force-indentation curves by subtracting the cantilever deflection on a solid surface. The indentation curves were then fitted to the Hertz model (Hertz, H., 1881).

Fluorescence Microscopy:

Yeast were grown for 2 h at 30 °C in YNB medium containing amino acids and (NH₄)₂SO₄ (Sigma Aldrich). Morphogenesis was induced at 37 °C for 3 h in 20 % FCS, 2 % glucose. Cells were imaged on µ-slides (Ibidi, Martinsried, Germany). Images of Smi1-GFP localization were captured on an UltraVIEW® VoX spinning disk confocal microscope (Nikon, Surrey, UK), using a 488 nm laser. Multiple Z-stack images were acquired and Z-stack projections at maximum intensity were created using Volocity 6.3 software (Perkin Elmer).

Results:

1. Conservation of cellular function between *Saccharomyces cerevisiae* *Knr4* and *Candida albicans* *Smi1*:

The genome of the human pathogen *Candida albicans* contains two distinct homologs of the *Saccharomyces cerevisiae* *KNR4* gene, *SMI1* (C1_07870C_A) and *SMI1B* (C3_05350C_A). Gene deletion mutants for each gene were generated and initial phenotypic analysis showed a strong phenotype (sensitivity to CFW or SDS) for the *smi1*Δ/Δ mutant but not for the *smi1B* Δ/Δ mutant (Figure S1). We therefore focused on the role of Smi1 in this study.

The *SMI1* open reading frame was amplified and cloned into a *S. cerevisiae* expression vector, under the control of the strong and constitutive *ADHI* promoter. The *SMI1* coding sequence is 1,863 bp and contains two GTC codons at positions 1717 and 1762, which have a 97 % chance of translation as a Serine in *C. albicans* through non-canonical codon usage in this fungus, compared to a Leucine in *S. cerevisiae* (White et al., 1995), although these usage frequencies appear to be sensitive to environmental stressors (Simões et al., 2016). These two codons are located within the C-terminal of Smi1, which is less conserved than the central domains among members of the fungal Knr4/Smi1 super-family of proteins. The complete sequence alignment of several fungal protein members of this super-family is shown in Figure S2. The C-terminal part of Knr4 is largely unstructured and not directly necessary for protein function in *S. cerevisiae* (Dagkessamanskaia et al., 2010; Martin-Yken et al., 2016). Hence, we tested the ability of the *C. albicans* *SMI1* gene, retaining its two ambiguous codons, to complement the cell wall-related phenotypes of *S. cerevisiae* *knr4* null mutants. Our phenotypic screen included sensitivity to Calcofluor White (CFW), SDS, caspofungin and elevated temperature in haploid and diploid *S. cerevisiae* genetic backgrounds. CFW is a compound that binds to nascent chitin fibrils and has been used since 1994 to identify fungal cell wall mutants (Ram et al., 1994). Expression of *SMI1* gene in *S. cerevisiae* was able to complement these phenotypes (Figure 1A) and also complemented the *smi1*Δ growth defect at elevated temperature (Fig.1B), a specific defect in this organism that is linked to cell cycle progression through START (Fishel et al., 1993, Martin-Yken et al., 2003). Hence, the function of the Smi1/Knr4 proteins appears to be conserved between the two species despite their phylogenetic distance and the differences between the two protein sequences, which share only 34% identity and 49% similarity (Figure S1).

2. Deletion of *SMI1* renders *C. albicans* sensitive to cell wall targeting drugs:

We further investigated the role of Smi1 in maintaining cell wall integrity using a medically relevant β -glucan synthase-targeting drug, the echinocandin Caspofungin. The influence of Smi1 on the sensitivity to CFW was also retested in parallel. Homozygous deletions of the two alleles of *SMI1* gene in the *C. albicans* BWP17 genetic background lead to a marked increase in the sensitivity towards both Caspofungin and CFW at 30 °C (Figure 2). These phenotypes are consistent with the ones observed for the *S. cerevisiae knr4* Δ mutant as well as with the role proposed for these proteins in stress signaling pathways (See discussion). Re-integration of *SMI1* open reading frame under the constitutive promoter P_{TDH3} in the *smi1* Δ/Δ deletion mutant restored the wild-type phenotype, while integration of the empty *URA3*-bearing vector did not (Fig. 2).

3. Cell Wall Biophysics

3.1 Cell Wall Strength / Elasticity

Atomic Force Microscopy under liquid conditions can be used to investigate the nanomechanical properties of live wildtype and mutant cells (Dague et al., 2010; Ene et al., 2015; Formosa et al., 2013; Liu et al., 2015). Here we first measured the cell surface resistance vs the elasticity of three *C. albicans* strains: the control strain, BWP17 AHU (which carries the functional *URA3* marker), the homozygous deletion mutant *smi1* Δ/Δ AHU and a strain over-expressing the *SMI1* gene from the strong constitutive promoter P_{TDH3} in the same BWP17 genetic background, *SMI1*-OE. Using the Atomic Force Microscope in the Force Volume mode, we collected between 1024 and 4160 force curves per cell on a minimum of 12 cells from 4 independent cultures for each strain. The elasticity of the cells was quantified from these curves by calculating the Young's Modulus (YM) as previously described (Dague et al., 2010), with $YM = 1 / \text{Elasticity}$. The higher the Young's Modulus, the stiffer the cells are. As can be seen on Figure 3, the Young's Moduli measured for the control strain BWP17 AHU showed a mean value of 782 \pm 158 kPa, significantly higher than the ones measured for the homozygous deletion mutant which had a mean value of 93 \pm 34 kPa. The Young's

Moduli measured on the cells of the over-expression strain lay between the control strain and the deletion mutant, with a mean of 298 ± 62 kPa, suggesting either a gene dosage effect for *SMII* expression or possibly an effect of uncoupling the expression of this gene from the cell cycle. These values are represented for each cell individually on Figure 3 since to allow visualization of the cell-to-cell variability. The mean values \pm SEM of the Young Moduli obtained for these three strains are also reported in Table 2. These measurements show that the deletion of both alleles of *SMII* resulted in a reduction of the cell surface stiffness by eight to ten fold, indicating a very soft cell wall for this mutant.

3.2 Adhesion:

Another cell surface feature that can be easily and very precisely measured by AFM is the ability to adhere to surfaces, using Single Molecule Force Spectroscopy (Axner et al., 2010; Benoit et al., 2000; Formosa et al., 2014a; Hinterdorfer et al., 1996; Neuman and Nagy, 2008). Here, adhesion between the cell surface and the AFM bare tips, constituted of Si_3N_4 was measured by scanning square areas on the top of yeast cells. We recorded force curves whose retract parts were then used to generate adhesion maps where the intensity of each pixel corresponds to the force required to dissociate the AFM tip from the sample, *i.e.* the adhesion force, expressed in picoNewtons pN (see Figure 4). It can be seen immediately from these heat maps that the *smi1* Δ/Δ mutant cell surface is almost not adhesive compared to the control strain, while the *SMII* overexpressing cells on the other hand display a marked increase in their surface adhesion. In addition to the visual representation by these adhesion maps, the cellular adhesion was also evaluated by three quantitative parameters: the mean adhesion force and specific energy of each adhesion event observed and the overall proportion of these adhesive events among all the recorded force curves. In order to quantify these values we defined as an adhesive event any force curve showing an adhesion force above 50pN. With this threshold level we calculated the percentage of adhesive events for each cell type and measured the area situated below the retract curves, which represent the adhesion energy of the event (Table 2). These values indicated that adhesive events were encountered more frequently at the surface of the over-expressing strain SMI1-OE (63% of the recorded 21,500 force curves) than on the control strain (46% of over 13,300 force curves), and were rare on the deletion mutant surface (19% of over 12,300 force curves). The average adhesion force measured on the control strain was of 127 ± 52 pN, calculated on 6256 adhesive force curves. This has to be compared with an average force of 70 ± 16 pN

(hence barely above the threshold) on 2300 force curves for the *smi1* Δ/Δ mutant. For the *SMII* OE-strain on the other hand, not only were the adhesive events more frequent (63 %) but they were also much stronger, with forces measured up to 2176 pN and an average value of 712 \pm 260 pN calculated on 12,257 adhesive force curves. The specific energy of these adhesive events also differed; with adhesion energies measured for the over-expressing strain an order of magnitude stronger than for the control strain, while they were about seven times lower on the surface of the deletion mutant. Hence, the homozygous deletion of *SMII* gene abrogates almost entirely the ability of the mutant cell to adhere using the chemistry described here, while *SMII* over-expression (or constitutive expression) leads to a highly adhesive phenotype.

4. Cellular Localization:

The homolog of Smi1 and Knr4 in the model filamentous fungus *Neurospora crassa*, GS-1 localizes at the growing tip of hyphae as a sphere positioned at the “Spitzenkörper” (Verdin et al., 2009). The Spitzenkörper (or apical body) is a fungal structure specific to true hyphae, located at the hyphal tip. It is comprised of the secretory vesicles that are required for continuous polarized growth (Girbardt, 1957; Harris et al., 2005). GS-1-GFP and Knr4-GFP have been imaged respectively at the tip of *N. crassa* hyphae (Riquelme et al., 2011; Sánchez-León et al., 2011) and at the tip of elongated shmoos in *S. cerevisiae* (Martin-Yken, 2012) and our own unpublished data). To test whether the *C. albicans* homolog would be similarly positioned, a GFP-tag was integrated at the C-terminus of the Smi1 protein at its chromosomal locus and its cellular localization was visualized by confocal fluorescent microscopy in yeast and hyphal cells.

In yeast cells, Smi1-GFP appeared both as punctate patches in the cytoplasm and localized transiently to nascent buds. This localization is similar to that reported for Knr4 (Dagkessamanskaia et al., 2010; Martin et al., 1999). Punctate patches and a more diffuse cytoplasmic distribution were observed in *C. albicans* hyphae, but, unlike in yeast, Smi1-GFP was consistently retained at the growing hyphal tip throughout the cell cycle. This is reminiscent of *N. crassa* GS-1 localization (see above and discussion). Smi1-GFP was occasionally observed at hyphal septa as a dim signal, but this presence did not reflect specific stages of the cell cycle. Hence, Smi1 in *C. albicans* appears to associate with intracellular

organelles and localizes to sites of new cell wall growth, a pattern which partially reflects those observed for homologs of Smi1 in *S. cerevisiae* in yeast cells and the hyphae of *N. crassa*.

DISCUSSION

Our results establish a significant conservation of Knr4/Smi1 function between *S. cerevisiae* and *C. albicans* and demonstrate the role of Smi1 in resistance to Caspofungin, regulation of cell wall integrity and cell surface adhesion properties of this major human fungal pathogen. These features suggest that Smi1 might be relevant as a new drug target for combination therapies. Previous work by Nett and colleagues identified a role for Smi1 in the production of extracellular matrix upon biofilm formation and hence the associated resistance to Fluconazole (Nett et al., 2011). Their results indicated that these effects were linked to the cell wall integrity pathway but were in fact regulated by Smi1 independently of the CWI pathway, suggesting a control pathway for Smi1 distinct from that of the PKC pathway. Lafayette and colleagues dissected the mechanisms through which PKC regulates resistance to both azoles and echinocandins in the two yeast models *C. albicans* and *S. cerevisiae* (LaFayette et al., 2010). They showed that, in *C. albicans*, Pkc1 and calcineurin signaling pathways independently regulate antifungal resistance *via* a common unknown target, which they designed as “X” (see (LaFayette et al., 2010), Fig 9B thereof). Considering the knowledge accumulated on Knr4 in budding yeast together with the data obtained for *C. albicans* and presented here, we propose that Smi1 might constitute this previously unidentified “X”, a common target of the Pkc1 and calcineurin signaling pathways. In addition, another result of this work is the very low stiffness of the *smi1Δ/Δ* mutant cell wall indicated by its Young Modulus (Fig. 3, Table 2). The severity of this phenotype compared to the milder one observed for *S. cerevisiae knr4Δ* mutant suggests a more important (central) role of Smi1 in the network of signaling pathways controlling cell wall integrity than that of Knr4 in the baker yeast, which is in agreement with these two proteins being the hypothetical “X” in the work of LaFayette and colleagues (LaFayette et al., 2010). The very soft cell wall phenotype observed for the deletion mutant *smi1Δ/Δ* is similar to, although more severe, than the one observed for the *S. cerevisiae knr4Δ* mutant whose Young’s Modulus in stationary phase was reduced three-fold compared to the control strain (Dague et al., 2010). This could

result from the reduced level of β -glucan in the cell wall of this mutant previously reported (Nett et al., 2011). The fact that the homozygous deletion mutant *smi1* Δ/Δ is so strongly affected, despite the presence in this strain of both functional alleles of *SMIB*, argues for a major role for the *SMII* gene, at least in the conditions tested (30°C, liquid rich medium, yeast form of *C. albicans* cells).

Our results show that homozygous deletion of *SMII* gene leads to a marked increase in the sensitivity of *C. albicans* to both CFW and Caspofungin at 30 °C. Since Caspofungin tolerance has been reported in *C. albicans* mutants with elevated cell-wall chitin (Perlin, 2015; Plaine et al., 2008; Walker et al., 2013; Yang et al., 2017) while elevated chitin is known to induce hypersensitivity to CFW (Elorza et al., 1983; Roncero and Durán, 1985), this finding may appear puzzling at first glance. However, this is consistent with the phenotypes observed for the *S. cerevisiae knr4* Δ mutant (Lesage et al., 2004; Martin et al., 1999). Moreover, as our cell wall stiffness measurements, these phenotypes indicate a severely modified cell wall organization of the deletion mutant *smi1* Δ/Δ compared to the control strain, coherent with a possible upstream role of Smi1 in cellular signaling pathways allowing this protein to influence simultaneously the expression level of different cell wall synthesis genes. Consistent with this, early results obtained in the bakers' yeast established a role for Knr4 in the transcriptional control of all *S. cerevisiae* chitin synthase genes (Martin et al., 1999). The complex interplay of cellular signaling pathways controlling *C. albicans* susceptibility to echinocandins has been described by Munro and colleagues (Walker et al., 2010), and it now seems possible that Smi1 is one piece of this puzzle.

A previous study investigating the effects of Caspofungin on the yeast cell wall by Atomic Force Microscopy did reveal that treatment by this echinocandin increased *C. albicans* cell wall stiffness and at the same time enhanced cell surface adhesion (Formosa et al., 2013). With this in mind, together with the remarkable sensitivity of the *smi1* Δ/Δ deletion mutant to this drug, it is tempting to postulate that *SMII* gene product is required for the establishment of the cell's response to Caspofungin. It has however not been possible to test the effect of Caspofungin treatment on the deletion mutant by AFM, precisely due to its extreme sensitivity and lethality in the conditions of the test.

The cellular localization observed for Smi1 in yeast cells and hyphae, as cytoplasmic patches and concentrated spots at the polarized growth sites, is coherent with the idea of cellular function conservation between Smi1 and *S. cerevisiae* Knr4. Indeed, similar cellular

localization has been reported for the MAP kinase Slt2 and for Calmodulin, the phosphatase which activates Calcineurin (Brockhoff and Davis, 1992; van Drogen and Peter, 2002). In the model filamentous fungus, *N. crassa*, GS1, the homolog of Sml1, localizes at the Spitzenkörper within the hyphal tip (Verdin et al., 2009; Sánchez-León et al., 2011). This localization is conserved in another ascomycete filamentous fungus, *A. nidulans* (also called *Emericella nidulans*), (Schultzhaus et al., 2015). The results presented here indicate that Sml1 also localizes in or around the Spitzenkörper of *C. albicans* hyphae in a similar manner to that observed for GS-1 in *N. crassa*. Finally, given the cell wall related phenotypes reported for GS-1 mutants of *N. crassa* (Enderlin and Selitrennikoff, 1994; Resheat-Eini et al., 2008; Seiler and Plamann, 2003), it is tempting to speculate that the function of these proteins is not only conserved between *C. albicans* and *S. cerevisiae*, but also, to some extent at least, in filamentous fungi. The role of Sml1 in the control of cell wall synthesis, cellular adhesion and drug resistance is of great interest with regard to finding new antifungal targets. The decisive advantage of Sml1 as a drug target over Hsp90, calcineurin, Pkc1 or other MAP kinases is the fact that the Knr4/Sml1 superfamily of proteins is specific of the Fungal Kingdom and absent from host cells (Martin-Yken et al., 2016). In addition, since this protein family is conserved among fungi, including other fungal pathogens of mammals (*C. glabrata* and *Aspergillus* species notably) and also plants (ex: *Magnaporthe grisea*), developing drugs that target Sml1 might lead to broader antifungal applications in domains such as agriculture.

Acknowledgements:

Pr. Christophe d'Enfert from Institut Pasteur (Paris) for strains and plasmids, Dr Jean-Luc Parrou for vector YEplac195 *PGK/CYC1*.

The FEBS society and the organizers of the Human Fungal Pathogen Schools 2015 and 2017 for a great introduction to *C. albicans* and other human fungal pathogens, as well as Formation permanente of the Centre INRA Toulouse Occitanie for financing HMY attendance to these researcher schools.

HMY is forever grateful to Frans M. Klis for his kindness throughout the years and his great advice to “*Start working on C. albicans*”.

481

482

Strain name	Genotype	References
<i>Candida albicans</i> strains :		
BWP17	<i>ura3Δ::λimm434/ura3Δ::λimm434, arg4Δ::hisG/arg4Δ::hisG, his1Δ::hisG/his1Δ::hisG</i>	(Wilson et al., 1999)
CEC161	Isogenic to BWP17 but <i>arg4Δ::hisG/ARG4, his1Δ::hisG/HIS1</i>	(Firon et al., 2007)
BWP17 AHU	Isogenic to CEC161 but <i>RPS1/rps1::Clp10-URA3</i>	(Chauvel et al., 2012)
<i>smi1Δ/Δ</i>	Isogenic to BWP17 but <i>smi1Δ::HIS1/smi1Δ::ARG4</i>	This study
<i>smi1Δ/Δ</i> - AHU	Isogenic to <i>smi1Δ/Δ</i> but <i>RPS1/rps1::Clp10-URA3</i>	This study
<i>smi1Δ/Δ</i> + P _{TDH3} <i>SMI1</i>	Isogenic to <i>smi1Δ/Δ</i> but <i>RPS1/rps1::Clp10-P_{TDH3}-SMI1</i>	This study
<i>smi1BΔ/Δ</i>	Isogenic to BWP17 but <i>smi1BΔ::HIS1/smi1BΔ::ARG4</i>	This study
SMI1-OE	Isogenic to CEC2908 but <i>RPS1/rps1::Clp10-P_{TDH3}-SMI1</i>	This study
DAY185	<i>ura3::imm434/ura3::imm434 his1::hisG::HIS1/his1::hisG arg4::hisG::ARG4-URA3/arg4::hisG</i>	Davis <i>et al.</i> , 2000
SMI1-GFP	DAY185 <i>SMI1::SMI1-GFP-NAT</i>	This study
<i>Saccharomyces cerevisiae</i> strains :		
BY4741a	MAT a; <i>his3Δ1 leu2Δ0; met15Δ0; ura3Δ0</i>	(Brachmann et al., 1998)
<i>knr4Δ</i>	BY4741a <i>YGR229c::KanMX4</i>	YKO Collection (Open Biosystems)
W303-2N	MAT a/α <i>ura3-1/ura3-1 leu2-3,112/leu2-3,112 trp1-1/trp1-1 his3-11,15/his3-11,15 ade2-1/ade2-1 can1-100/can1-100</i>	
HM1315	W303-2N <i>YGR229c::KanMX4/ YGR229c::KanMX4</i>	This study.

483

484

Table 1. Yeast strains used in this study.

485

486

487

488

Cell Type	Young Modulus (kPa)	% of Adhesive events	Mean Adhesion Force (pN)	Adhesion Energy (=Area below the force curve) (J)
-----------	---------------------	----------------------	--------------------------	---------------------------------------------------

BWP17 AHU	782 (+/- 158)	46.1	127 (+/- 52)	1.77 x10⁻¹⁷
<i>smi1</i>Δ/Δ AHU	93 (+/- 34)	19.4	70 (+/- 26)	0.26 x10⁻¹⁷
SMI1-OE	298 (+/- 62)	62.9	712 (+/- 445)	15.82 x10⁻¹⁷

Table 2: Summary of Atomic Force Microscopy measurements for BWP17 AHU, *smi1*Δ/Δ AHU, and SMI1-OE strains.

Column 1: Mean values with standard deviation of Young's Moduli calculated from force curves obtained as described above (3.1).

Column 2: Percentage of adhesive events measured by AFM, calculated from at least 12,000 force curves for each cell type, with a threshold level for the definition of an adhesive event as 50pN on the retraction curve.

Column 3: Mean values of Adhesion forces for each cell type, calculated from adhesive force curves obtained as described above.

Column 4: Mean values of the Adhesion Energy for each strain, calculated from the area below the force curves presenting an adhesion event.

LEGENDS TO FIGURES

Figure 1. *C. albicans* *SMI1* gene expression suppresses the cell wall associated phenotypes of *S. cerevisiae* *knr4* Δ mutants.

A) Transformed control strain BY4741a and mutant strain *knr4* Δ with either empty plasmid YEplac195 *PGK/CYC1* or p*SMI* bearing *C. albicans* *SMI* gene or p*KNR4* bearing *S. cerevisiae* *KNR4* gene were grown in liquid SD medium lacking uracile at 30°C to an OD₆₀₀ of 1, and concentrated to OD₆₀₀ 8 \pm 0. Serial dilutions of yeast cultures were spotted on YPD plates in the absence or presence of 150 ng caspofungin ml⁻¹. Growth was scored after 2 days at 30°C.

B) Transformed control strain W3032N and mutant strain HM1315 *knr4* Δ/Δ with either empty YEplac195 *PGK/CYC1* plasmid or p*SMI* (Yeplac195 bearing *C. albicans* *SMI* gene under P_{PGK1}) or p*KNR4* (Yeplac195 bearing *S. cerevisiae* *KNR4* gene under P_{PGK1}) were grown overnight in liquid SD medium lacking uracile at 30°C and concentrated to OD₆₀₀ 8 \pm 0. Serial dilutions of yeast cultures were spotted on YPD plates. Growth was scored after 2 days at 30°C and 37°C.

Figure 2. Calcofluor White and caspofungin sensitivity of the *C. albicans* *smi1* Δ/Δ mutant.

Control strain BWP17 and mutant strain *smi1* Δ/Δ were grown in liquid YPD medium at 30°C to an OD₆₀₀ of 1, and concentrated to OD₆₀₀ 8 \pm 0. Serial dilutions of yeast cultures were spotted on YPD plates in the absence or presence of 40mg of CFW or 150 ng caspofungin ml⁻¹. Growth was scored after 2 days at 30°C.

Figure 3. Young Moduli of *smi1* Δ/Δ mutant and *SMI1*-OE vs control strain.

Atomic Force Microscope was used to collect over 12,300 force curves for each strain on the control strain BWP17 AHU, the *smi1* Δ/Δ mutant AHU and the *SMI1*-OE strain over-expressing the *SMI1*. The Young's Moduli (YM = 1 / Elasticity) were quantified from these curves as described and are presented here as a dot on the mean YM value, with SEM for each cell. The bar represents the mean of the YM values with each SEM. Statistical analysis was done using the One-way ANOVA test, **** = p value < 0.0001.

Figure 4. Adhesion Maps recorded on the cellular surfaces of *smi* Δ/Δ mutant and SMI1-OE vs control strain.

Adhesion force measurements performed on single *C. albicans* cells by Atomic Force Microscopy. The adhesion maps presented have been recorded on three independent and representative *C. albicans* cells for each cell type. Each analyzed area covers $1 \times 1 \mu\text{m}^2$. Adhesion scales are shown and read as follows: bright yellow = maximum adhesion force at 2 nN; dark red = minimum at 0.0 nN.

Figure 5. Smi1-GFP localizes to apical growth sites in yeast and hyphae. Cells were grown on Ibidi μ -slides in YNB medium at 30°C for 2h (yeast) and 20 % FCS, 2 % glucose at 37°C for 3h (hyphae). Smi1-GFP localised transiently to emerging bud tips in yeast (arrows) and to septa in hyphae (asterisks) but was maintained consistently at hyphal/branch tips (arrows). Punctate fluorescence patches were also observed throughout yeast and hyphal cells. Images are maximum projections of individual z-stacks.

References :

- Axner, O., Björnham, O., Castelain, M., Koutris, E., Schedin, S., Fällman, E., and Andersson, M. (2010). Unraveling the secrets of bacterial adhesion organelles using single molecule force spectroscopy. In *DIVA*, (Springer Verlag), pp. 337–362.
- Benoit, M., Gabriel, D., Gerisch, G., and Gaub, H.E. (2000). Discrete interactions in cell adhesion measured by single-molecule force spectroscopy. *Nat. Cell Biol.* 2, 313–317.
- Birkaya, B., Maddi, A., Joshi, J., Free, S.J., and Cullen, P.J. (2009). Role of the Cell Wall Integrity and Filamentous Growth Mitogen-Activated Protein Kinase Pathways in Cell Wall Remodeling during Filamentous Growth. *Eukaryot. Cell* 8, 1118–1133.
- Brachmann, C.B., Davies, A., Cost, G.J., Caputo, E., Li, J., Hieter, P., and Boeke, J.D. (1998). Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* 14, 115–132.
- Brockerhoff, S.E., and Davis, T.N. (1992). Calmodulin concentrates at regions of cell growth in *Saccharomyces cerevisiae*. *J Cell Biol* 118, 619–629.
- Brown, A.J.P., Leach, M.D., and Nicholls, S. (2010). The relevance of heat shock regulation in fungal pathogens of humans. *Virulence* 1, 330–332.
- Brown, G.D., Denning, D.W., Gow, N.A.R., Levitz, S.M., Netea, M.G., and White, T.C. (2012). Hidden Killers: Human Fungal Infections. *Sci. Transl. Med.* 4, 165rv13-165rv13.
- Chauvel, M., Nesseir, A., Cabral, V., Znaidi, S., Goyard, S., Bachellier-Bassi, S., Firon, A., Legrand, M., Diogo, D., Naulleau, C., et al. (2012). A Versatile Overexpression Strategy in the Pathogenic Yeast *Candida albicans*: Identification of Regulators of Morphogenesis and Fitness. *PLoS ONE* 7, e45912.
- Chopinnet, L., Formosa, C., Rols, M.P., Duval, R.E., and Dague, E. (2013). Imaging living cells surface and quantifying its properties at high resolution using AFM in QI™ mode. *Micron* 48, 26–33.
- Dagkessamanskaia, A., El Azzouzi, K., Kikuchi, Y., Timmers, T., Ohya, Y., Francois, J.M., and Martin-Yken, H. (2010). Knr4 N-terminal domain controls its localization and function during sexual differentiation and vegetative growth. *Yeast* 27, 563–574.
- Dague, E., Bitar, R., Ranchon, H., Durand, F., Yken, H.M., and Francois, J.M. (2010). An atomic force microscopy analysis of yeast mutants defective in cell wall architecture. *Yeast* 27, 673–684.
- Dague, E., Jauvert, E., Laplatine, L., Viallet, B., Thibault, C., and Ressler, L. (2011). Assembly of live micro-organisms on microstructured PDMS stamps by convective/capillary deposition for AFM bio-experiments. *Nanotechnology* 22, 395102.
- Desai, J.V., Mitchell, A.P., and Andes, D.R. (2014). Fungal Biofilms, Drug Resistance, and Recurrent Infection. *Cold Spring Harb. Perspect. Med.* 4, a019729–a019729.

- 596 van Drogen, F., and Peter, M. (2002). Spa2p functions as a scaffold-like protein to recruit the
597 Mpk1p MAP kinase module to sites of polarized growth. *Curr Biol* 12, 1698–1703.
- 598 Enderlin, C.S., and Selitrennikoff, C.P. (1994). Cloning and characterization of a *Neurospora*
599 *crassa* gene required for (1,3) beta-glucan synthase activity and cell wall formation. *Proc Natl*
600 *Acad Sci U A* 91, 9500–9504.
- 601 Ene, I.V., Walker, L.A., Schiavone, M., Lee, K.K., Martin-Yken, H., Dague, E., Gow, N.A.R.,
602 Munro, C.A., and Brown, A.J.P. (2015). Cell Wall Remodeling Enzymes Modulate Fungal Cell
603 Wall Elasticity and Osmotic Stress Resistance. *MBio* 6, e00986-15.
- 604 Firon, A., Aubert, S., Iraqui, I., Guadagnini, S., Goyard, S., Prévost, M.-C., Janbon, G., and
605 d’Enfert, C. (2007). The SUN41 and SUN42 genes are essential for cell separation in *Candida*
606 *albicans*. *Mol. Microbiol.* 66, 1256–1275.
- 607 Fishel, B.R., Sperry, A.O., and Garrard, W.T. (1993). Yeast calmodulin and a conserved
608 nuclear protein participate in the in vivo binding of a matrix association region. *Proc Natl*
609 *Acad Sci U A* 90, 5623–5627.
- 610 Formosa, C., Schiavone, M., Martin-Yken, H., François, J.M., Duval, R.E., and Dague, E.
611 (2013). Nanoscale effects of caspofungin against two yeast species, *Saccharomyces*
612 *cerevisiae* and *Candida albicans*. *Antimicrob. Agents Chemother.* 57, 3498–3506.
- 613 Formosa, C., Pillet, F., Schiavone, M., Duval, R.E., Ressler, L., and Dague, E. (2014a).
614 Generation of living cell arrays for atomic force microscopy studies. *Nat. Protoc.* 10, 199–
615 204.
- 616 Formosa, C., Schiavone, M., Boisrame, A., Richard, M.L., Duval, R.E., and Dague, E. (2014b).
617 Multiparametric imaging of adhesive nanodomains at the surface of *Candida albicans* by
618 atomic force microscopy. *Nanomedicine Nanotechnol. Biol. Med.*
- 619 Gietz, R.D., and Sugino, A. (1988). New yeast-*Escherichia coli* shuttle vectors constructed
620 with in vitro mutagenized yeast genes lacking six-base pair restriction sites. *Gene* 74, 527–
621 534.
- 622 Gietz, R.D., and Woods, R.A. (2006). Yeast Transformation by the LiAc/SS Carrier DNA/PEG
623 Method. In *Yeast Protocol*, (Humana Press, Totowa, NJ), pp. 107–120.
- 624 Girbardt, M. (1957). Der Spitzenkörper von *Polystictus versicolor* (L.). *Planta* 50, 47–59.
- 625 Gola, S., Martin, R., Walther, A., Dünkler, A., and Wendland, J. (2003). New modules for PCR-
626 based gene targeting in *Candida albicans*: rapid and efficient gene targeting using 100 bp of
627 flanking homology region. *Yeast* 20, 1339–1347.
- 628 Harris, S.D., Read, N.D., Roberson, R.W., Shaw, B., Seiler, S., Plamann, M., and Momany, M.
629 (2005). Polarisome Meets Spitzenkörper: Microscopy, Genetics, and Genomics Converge.
630 *Eukaryot. Cell* 4, 225–229.
- 631 Hertz, H. (1881). Ueber die Berührung fester elastischer Körper. *J Für Reine Angew Math*
632 156–171.

- 633 Hinterdorfer, P., Baumgartner, W., Gruber, H.J., Schilcher, K., and Schindler, H. (1996).
 634 Detection and localization of individual antibody-antigen recognition events by atomic force
 635 microscopy. *Proc. Natl. Acad. Sci.* *93*, 3477–3481.
- 636 Hutter, J.L., and Bechhoefer, J. (1993). Calibration of atomic-force microscope tips. *Rev. Sci.*
 637 *Instrum.* *64*, 1868–1873.
- 638 Kathiravan, M.K., Salake, A.B., Chothe, A.S., Dudhe, P.B., Watode, R.P., Mukta, M.S., and
 639 Gadhwe, S. (2012). The biology and chemistry of antifungal agents: a review. *Bioorg. Med.*
 640 *Chem.* *20*, 5678–5698.
- 641 LaFayette, S.L., Collins, C., Zaas, A.K., Schell, W.A., Betancourt-Quiroz, M., Gunatilaka, A.A.L.,
 642 Perfect, J.R., and Cowen, L.E. (2010). PKC Signaling Regulates Drug Resistance of the Fungal
 643 Pathogen *Candida albicans* via Circuitry Comprised of Mkc1, Calcineurin, and Hsp90. *PLoS*
 644 *Pathog* *6*, e1001069.
- 645 Lesage, G., Sdicu, A.-M., Menard, P., Shapiro, J., Hussein, S., and Bussey, H. (2004). Analysis
 646 of beta-1,3-glucan assembly in *Saccharomyces cerevisiae* using a synthetic interaction
 647 network and altered sensitivity to caspofungin. *Genetics* *167*, 35–49.
- 648 Liu, R., Formosa, C., Dagkessamanskaia, Dague, E., Francois, J.M., and Martin-Yken (2015).
 649 Combining Atomic Force Microscopy and genetics to investigate the role of Knr4 in
 650 *Saccharomyces cerevisiae* sensitivity to K9 Killer toxin. *Lett. Appl. NanoBioscience* *4*, 306–
 651 315.
- 652 Liu, T.T., Lee, R.E.B., Barker, K.S., Lee, R.E., Wei, L., Homayouni, R., and Rogers, P.D. (2005).
 653 Genome-wide expression profiling of the response to azole, polyene, echinocandin, and
 654 pyrimidine antifungal agents in *Candida albicans*. *Antimicrob. Agents Chemother.* *49*, 2226–
 655 2236.
- 656 Markovich, S., Yekutieli, A., Shalit, I., Shadkchan, Y., and Osherov, N. (2004). Genomic
 657 Approach to Identification of Mutations Affecting Caspofungin Susceptibility in
 658 *Saccharomyces cerevisiae*. *Antimicrob. Agents Chemother.* *48*, 3871–3876.
- 659 Martin, H., Dagkessamanskaia, A., Satchanska, G., Dallies, N., and François, J. (1999). KNR4, a
 660 suppressor of *Saccharomyces cerevisiae* cwh mutants, is involved in the transcriptional
 661 control of chitin synthase genes. *Microbiol. Read. Engl.* *145 (Pt 1)*, 249–258.
- 662 Martin-Yken, H. (2012). Knr4/Smi1 family: Conserved fungal chaperones of puzzling origin.
 663 (Primosten, Croatia).
- 664 Martin-Yken, H., Dagkessamanskaia, A., Basmaji, F., Lagorce, A., and Francois, J. (2003). The
 665 interaction of Slt2 MAP kinase with Knr4 is necessary for signalling through the cell wall
 666 integrity pathway in *Saccharomyces cerevisiae*. *Mol Microbiol* *49*, 23–35.
- 667 Martin-Yken, H., François, J.M., and Zerbib, D. (2016). Knr4: a disordered hub protein at the
 668 heart of fungal cell wall signalling. *Cell. Microbiol.* *18*, 1217–1227.

- 669 Milne, S.W., Cheetham, J., Lloyd, D., Aves, S., and Bates, S. (2011). Cassettes for PCR-
 670 mediated gene tagging in *Candida albicans* utilizing nourseothricin resistance. *Yeast*
 671 *Chichester Engl.* 28, 833–841.
- 672 Nett, J.E., Sanchez, H., Cain, M.T., Ross, K.M., and Andes, D.R. (2011). Interface of *Candida*
 673 *albicans* biofilm matrix-associated drug resistance and cell wall integrity regulation.
 674 *Eukaryot. Cell* 10, 1660–1669.
- 675 Neuman, K.C., and Nagy, A. (2008). Single-molecule force spectroscopy: optical tweezers,
 676 magnetic tweezers and atomic force microscopy. *Nat. Methods* 5, 491–505.
- 677 Odds, F.C., Brown, A.J.P., and Gow, N.A.R. (2003). Antifungal agents: mechanisms of action.
 678 *Trends Microbiol.* 11, 272–279.
- 679 Perlin, D.S. (2015). Mechanisms of echinocandin antifungal drug resistance. *Ann. N. Y. Acad.*
 680 *Sci.* 1354, 1–11.
- 681 Plaine, A., Walker, L., Da Costa, G., Mora-Montes, H.M., McKinnon, A., Gow, N.A.R.,
 682 Gaillardin, C., Munro, C.A., and Richard, M.L. (2008). Functional analysis of *Candida albicans*
 683 GPI-anchored proteins: Roles in cell wall integrity and caspofungin sensitivity. *Fungal Genet.*
 684 *Biol.* 45, 1404–1414.
- 685 Ram, A.F., Kapteyn, J.C., Montijn, R.C., Caro, L.H., Douwes, J.E., Baginsky, W., Mazur, P., van
 686 den Ende, H., and Klis, F.M. (1998). Loss of the plasma membrane-bound protein Gas1p in
 687 *Saccharomyces cerevisiae* results in the release of beta1,3-glucan into the medium and
 688 induces a compensation mechanism to ensure cell wall integrity. *J Bacteriol* 180, 1418–1424.
- 689 Resheat-Eini, Z., Zelter, A., Gorovits, R., Read, N., and Yarden, O. (2008). *Neurospora crassa*
 690 colonial temperature sensitive 2, 4 and 5 (cot-2, cot-4 and cot-5) genes encode regulatory
 691 and structural proteins required for hyphal elongation and branching.
- 692 Riquelme, M., Yarden, O., Bartnicki-Garcia, S., Bowman, B., Castro-Longoria, E., Free, S.J.,
 693 Fleißner, A., Freitag, M., Lew, R.R., Mouriño-Pérez, R., et al. (2011). Architecture and
 694 development of the *Neurospora crassa* hypha – a model cell for polarized growth. *Fungal*
 695 *Biol.* 115, 446–474.
- 696 Sánchez-León, E., Verdín, J., Freitag, M., Roberson, R.W., Bartnicki-Garcia, S., and Riquelme,
 697 M. (2011). Traffic of chitin synthase 1 (CHS-1) to the Spitzenkörper and developing septa in
 698 hyphae of *Neurospora crassa*: actin dependence and evidence of distinct microvesicle
 699 populations. *Eukaryot. Cell* 10, 683–695.
- 700 Sanglard, D., and Odds, F.C. (2002). Resistance of *Candida* species to antifungal agents:
 701 molecular mechanisms and clinical consequences. *Lancet Infect. Dis.* 2, 73–85.
- 702 Sanglard, D., Kuchler, K., Ischer, F., Pagani, J.L., Monod, M., and Bille, J. (1995). Mechanisms
 703 of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients
 704 involve specific multidrug transporters. *Antimicrob. Agents Chemother.* 39, 2378–2386.

- 705 Schaub, Y., Dünkler, A., Walther, A., and Wendland, J. (2006). New pFA-cassettes for PCR-
706 based gene manipulation in *Candida albicans*. *J. Basic Microbiol.* 416–429.
- 707 Scorzoni, L., de Paula e Silva, A.C.A., Marcos, C.M., Assato, P.A., de Melo, W.C.M.A., de
708 Oliveira, H.C., Costa-Orlandi, C.B., Mendes-Giannini, M.J.S., and Fusco-Almeida, A.M. (2017).
709 Antifungal Therapy: New Advances in the Understanding and Treatment of Mycosis. *Front.*
710 *Microbiol.* 8.
- 711 Seiler, S., and Plamann, M. (2003). The genetic basis of cellular morphogenesis in the
712 filamentous fungus *Neurospora crassa*. *Mol. Biol. Cell* 14, 4352–4364.
- 713 Simões, J., Bezerra, A.R., Moura, G.R., Araújo, H., Gut, I., Bayes, M., and Santos, M.A.S.
714 (2016). The Fungus *Candida albicans* Tolerates Ambiguity at Multiple Codons. *Front.*
715 *Microbiol.* 7.
- 716 Smolyakov, G., Formosa-Dague, C., Severac, C., Duval, R.E., and Dague, E. (2016). High speed
717 indentation measures by FV, QI and QNM introduce a new understanding of
718 bionanomechanical experiments. *Micron* 85, 8–14.
- 719 Taff, H.T., Mitchell, K.F., Edward, J.A., and Andes, D.R. (2013). Mechanisms of *Candida*
720 biofilm drug resistance. *Future Microbiol.* 8.
- 721 Vanden Bossche, H., Koymans, L., and Moereels, H. (1995). P450 inhibitors of use in medical
722 treatment: focus on mechanisms of action. *Pharmacol. Ther.* 67, 79–100.
- 723 VEDIYAPPAN, G., ROSSIGNOL, T., and d'ENFERT, C. (2010). Interaction of *Candida albicans*
724 Biofilms with Antifungals: Transcriptional Response and Binding of Antifungals to Beta-
725 Glucans. *Antimicrob. Agents Chemother.* 54, 2096–2111.
- 726 Verdin, J., Bartnicki-Garcia, S., and Riquelme, M. (2009). Functional stratification of the
727 Spitzenkörper of *Neurospora crassa*. *Mol Microbiol* 74, 1044–1053.
- 728 Walker, L.A., Gow, N.A.R., and Munro, C.A. (2010). Fungal echinocandin resistance. *Fungal*
729 *Genet. Biol.* 47, 117–126.
- 730 Walker, L.A., Gow, N.A.R., and Munro, C.A. (2013). Elevated Chitin Content Reduces the
731 Susceptibility of *Candida* Species to Caspofungin. *Antimicrob. Agents Chemother.* 57, 146–
732 154.
- 733 Walther, A., and Wendland, J. (2003). An improved transformation protocol for the human
734 fungal pathogen *Candida albicans*. *Curr. Genet.* 42, 339–343.
- 735 White, T.C., Andrews, L.E., Maltby, D., and Agabian, N. (1995). The “universal” leucine codon
736 CTG in the secreted aspartyl proteinase 1 (SAP1) gene of *Candida albicans* encodes a serine
737 in vivo. *J. Bacteriol.* 177, 2953–2955.
- 738 Wilson, R.B., Davis, D., and Mitchell, A.P. (1999). Rapid Hypothesis Testing with *Candida*
739 *albicans* through Gene Disruption with Short Homology Regions. *J. Bacteriol.* 181, 1868–
740 1874.

741 Yang, F., Zhang, L., Wakabayashi, H., Myers, J., Jiang, Y., Cao, Y., Jimenez-Ortigosa, C., Perlin,
742 D.S., and Rustchenko, E. (2017). Tolerance to Caspofungin in *Candida albicans* Is Associated
743 with at Least Three Distinctive Mechanisms That Govern Expression of FKS Genes and Cell
744 Wall Remodeling. *Antimicrob. Agents Chemother.* 61.

745 Zarnowski, R., Westler, W.M., Lacmbouh, G.A., Marita, J.M., Bothe, J.R., Bernhardt, J.,
746 Lounes-Hadj Sahraoui, A., Fontaine, J., Sanchez, H., Hatfield, R.D., et al. (2014). Novel Entries
747 in a Fungal Biofilm Matrix Encyclopedia. *MBio* 5, e01333-14-e01333-14.

748
749
750