



HAL
open science

Exponential ergodicity for diffusions with jumps driven by a Hawkes process

Charlotte Dion, Sarah Lemler, Eva Löcherbach

► **To cite this version:**

Charlotte Dion, Sarah Lemler, Eva Löcherbach. Exponential ergodicity for diffusions with jumps driven by a Hawkes process. 2019. hal-02094514v2

HAL Id: hal-02094514

<https://hal.science/hal-02094514v2>

Preprint submitted on 18 Jul 2019 (v2), last revised 20 Dec 2019 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exponential ergodicity for diffusions with jumps driven by a Hawkes process

Charlotte Dion⁽¹⁾ and Sarah Lemler⁽²⁾ and Eva Löcherbach⁽³⁾

⁽¹⁾ Sorbonne Université, UMR CNRS 8001, LPSM, 75005 Paris, France

⁽²⁾ École CentraleSupélec, MICS, 91190 Gif-sur-Yvette, France

⁽³⁾ Université Paris 1, SAMM EA 4543, 75013 Paris, France

July 18, 2019

Abstract

In this paper, we introduce a new class of processes which are diffusions with jumps driven by a multivariate nonlinear Hawkes process. Our goal is to study their long-time behavior. In the case of exponential memory kernels for the underlying Hawkes process we establish conditions for the positive Harris recurrence of the couple (X, Y) , where X denotes the diffusion process and Y the piecewise deterministic Markov process (PDMP) defining the stochastic intensity of the driving Hawkes. As a direct consequence of the Harris recurrence, we obtain the ergodic theorem for X . Furthermore, we provide sufficient conditions under which the process is exponentially β -mixing.

AMS Classification: 60J35, 60F99, 60H99, 60G55

Keywords: Diffusions with jumps; nonlinear Hawkes process, Piecewise deterministic Markov processes (PDMP), Ergodicity; Foster-Lyapunov drift criteria; β -mixing.

1 Introduction

1.1 Motivation

In this work, we introduce a new class of continuous time processes $X = (X_t)_{t \geq 0}$ with jumps driven by a multivariate nonlinear Hawkes process N , satisfying the following equation

$$X_t = X_0 + \int_0^t b(X_s) ds + \int_0^t \sigma(X_s) dW_s + \int_0^t a(X_{s-}) \sum_{j=1}^M dN_s^{(j)}, \quad (1.1)$$

where W is the standard one-dimensional Brownian motion and N is a multivariate Hawkes process having M components with intensity process λ . N is supposed to be independent from W . We shall specify the dynamics of N later on, in particular, the precise form of the stochastic intensity process λ is given in (2.1) below.

The jumps of the Hawkes process impact the dynamic of the diffusion process. In this model, the structure of the jumps is different from classical jump-diffusion processes with Lévy jumps. Indeed:

1) the intensity of the jump process N does not depend on the dynamic of X ,

- 2) the Hawkes process has a special structure of time dependency: the state of the intensity at time t depends on the entire past,
- 3) the multidimensional nature of the Hawkes process can be interpreted as the influence of M subjects communicating one with each other and impacting the dynamic of X along time.

Our main motivation is to be able to describe the behavior of a set of neurons. Neurons are communicating through the emission of electrical signals, thus forming a network. When a (presynaptic) neuron sends a signal to another (postsynaptic) neuron, the membrane potential of the neuron increases sharply, and we say that the neuron spikes at this time. If we are looking at one fixed neuron at the heart of this network, its activity between two spikes of one of its presynaptic neurons can be designed by the present model. Indeed, during this interval the membrane potential can be described by a diffusion process influenced by the spikes coming from the presynaptic neurons which are modelled by the Hawkes process. Hawkes processes have been studied recently in the context of neurosciences [see 31, 17] to model the interactions between the neurons within their sequences of spikes. The new model we propose allows to use both informations: the continuous membrane potential for a fixed neuron together with the spikes of several other neurons around it.

The aim of the present article is to study the longtime behavior of the process X . Since X is not Markov on its own, we propose a study of the couple of processes (X, λ) in the case when the intensity λ of N can be described in terms of a piecewise deterministic Markov process (PDMP) Y , see (2.4) below. Such a description is possible when the memory kernels are *exponential*. We prove the positive Harris recurrence of (X, Y) together with ergodicity, and we establish the speed of β -mixing. These are important probabilistic properties which are interesting in its own right but which are also the key ingredients for the theoretical study of estimators of the model parameters. The estimation of the coefficients of the process is the issue of the companion paper [11].

1.2 State of the art

Time-homogeneous diffusion processes are well known Markov processes that have been widely studied. Let us cite for example [21] which has initiated the work on Markovian properties of diffusion processes or the by now classical references [20, 30] in which all properties of diffusions are summarized. Ergodicity and strong mixing have been established in the pioneering papers [34, 22]. Then, jump-diffusion processes have been introduced, for example in the context of risk management [see 33], either driven by a Poisson process or more generally by a Lévy process [see 1, for a review]. Their ergodicity and mixing properties have been studied from a probabilistic point of view by [26].

On the other hand, Hawkes processes [16] have been studied a lot lately. These point processes generalize the Poisson processes by introducing what is commonly called “self-excitation”: Present jumps are able to trigger (or to inhibit) future jumps, and in this way, correlations between successive jump events are introduced. [9] gives a precise synthesis of earlier results published by Hawkes. Important probabilistic results for these mutually exciting point processes, such as stability, limit theorems and large population limits, have been obtained in [6, 10, 2, 3, 8, 5]. Ergodicity for general non-linear Hawkes processes with general memory kernels has been recently established by [5]. For linear Hawkes processes, [14] proves exponential ergodicity, giving very precise estimates on the regeneration times. Finally [8] establishes exponential ergodicity in the particular case of linear Hawkes processes with exponential memory kernels.

Relying on one of these recent results on ergodicity of Hawkes processes, it is therefore reasonable to argue that the solution of (1.1) should be ergodic itself, provided we are able to ensure that the diffusion part of (1.1), that is the dynamic (1.1) with $a(\cdot) \equiv 0$, driving the motion in between successive jumps, is ergodic.

However, no such general criterion, ensuring that the ergodicity of the underlying jump process

implies the ergodicity (with exponential speed) of the associated diffusion with jumps, exists, at least not to our knowledge.

The present paper proposes therefore a short study of the ergodicity of diffusions with jumps driven by Hawkes processes with exponential memory kernels. In this case, the associated stochastic intensity processes of the jump part can be expressed in terms of particular Markov processes, PDMP's, see e.g. [12]. For these PDMP's, ergodicity has been established by [8] in the case of linear Hawkes processes, and by [12] in the case of non-linear Hawkes processes with Erlang kernels. Here, we extend these results to the general non-linear case, including also inhibitions ¹, and more importantly, we prove the joint ergodicity as well as the exponential β -mixing property for the couple (X, Y) in the case of exponential kernel functions.

1.3 Main contributions and plan of the paper

We propose a general study of the longtime behavior of a jump diffusion process, where the jumps are driven by a multivariate non-linear Hawkes process in dimension M , having exponential memory kernels. Our proof of the Harris recurrence of (X, Y) follows a well-known scheme, see e.g. [27, 28].

Firstly, we establish a local Döblin lower bound for the transition semigroup of the process (X, Y) , based on a lower bound of the joint transition density of (X, Y) . This is the content of Theorem 3.1. The ideas that we use to obtain this lower bound are very natural, and they have appeared independently of our work already in [8]. They have also been exploited in [12] and in [24]: more precisely, we use M^2 consecutive jumps which are carefully chosen to produce a part of Lebesgue density for the transition of Y . Once the jump part of the process possesses some part of Lebesgue density, we have to ensure that this density is preserved by the stochastic flow. This part of the proof relies on fine support properties of (X, Y) and provides, as a by-product, a simulation algorithm for the process. In particular we need to localize the possible positions of the diffusion part, conditionally on the jump times of the underlying Hawkes process that have already been chosen.

In a second step, a Foster-Lyapunov condition (see Proposition 3.4) ensures the control of the return times to some compact set K . We then use control arguments to deduce from this a precise control of the hitting times of the set C where the transition densities are lower-bounded according to the local Döblin lower bound. Both results together imply the positive Harris recurrence of the couple (X, Y) , as well as the ergodic theorem. This is our first main result, stated in Theorem 3.5.

Finally, following the steps of [26] where the ergodicity and exponential β -mixing bounds are established for diffusions with jumps driven by a Lévy process, we obtain the control of the β -mixing coefficient of (X, Y) in our second main theorem, Theorem 3.8.

Our paper is organised as follows. We introduce our model together with all necessary notation and assumptions in Section 2. Section 2.3 provides a simulation algorithm of the process which is interesting in its own right. In Section 3, we prove the exponential ergodicity and the exponential β -mixing of the coupled process (X, Y) .

2 Presentation of the model and first properties

2.1 The model

Throughout this paper we work on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F})$. We start by introducing the jump part of our process, that is, the multivariate Hawkes process.

This process is defined in terms of a collection of jump rate functions $f_i : \mathbb{R} \rightarrow \mathbb{R}_+, 1 \leq i \leq M$, and in terms of interaction functions (also called *memory kernels*) $h_{ij} : \mathbb{R}_+ \rightarrow \mathbb{R}, 1 \leq i, j \leq M$, which are

¹Notice that [7] has obtained a control of the τ -mixing coefficient in the inhibitory case – but this does not imply the β -mixing property of the process.

measurable functions. Let moreover $n^{(i)}, 1 \leq i \leq M$, be discrete point measures on \mathbb{R}_- satisfying that

$$\int_{\mathbb{R}_-} |h_{ij}|(t-s)n^{(j)}(ds) < \infty \text{ for all } t \geq 0.$$

We interpret them as *initial condition* of our process.

Assumption 2.1. *We suppose that each $f_i, 1 \leq i \leq M$, is Lipschitz continuous with Lipschitz constant γ_i .*

Definition 2.2. *A non-linear Hawkes process with parameters $(f_i, h_{ij})_{1 \leq i, j \leq M}$, and with initial condition $n^{(i)}, 1 \leq i \leq M$, is a multivariate counting process $N_t = (N_t^{(1)}, \dots, N_t^{(M)}), t \geq 0$, such that*

1. \mathbb{P} -almost surely, for all $i \neq j, N^{(i)}$ and $N^{(j)}$ never jump simultaneously,
2. for all $1 \leq i \leq M$, the compensator of $N_t^{(i)}$ is given by $\int_0^t \lambda_s^{(i)} ds$, where

$$\lambda_t^{(i)} = f_i \left(\sum_{j=1}^M \int_0^{t-} h_{ij}(t-u) dN_u^{(j)} + \sum_{j=1}^M \int_{-\infty}^0 h_{ij}(t-u) dn_u^{(j)} \right). \quad (2.1)$$

If the functions h_{ij} are locally integrable, the existence of the process $(N_t)_{t \geq 0}$ with the prescribed intensity (2.1) on finite time intervals follows from standard arguments, see e.g. [6] or [10].

We consider a jump diffusion process $X = (X_t)_{t \geq 0}$ taking values in \mathbb{R} , whose jumps are governed by N . More precisely, X is solution of the stochastic differential equation

$$X_t = X_0 + \int_0^t b(X_s) ds + \int_0^t \sigma(X_s) dW_s + \int_0^t a(X_{s-}) \sum_{j=1}^M dN_s^{(j)}, \quad (2.2)$$

where X_0 is a random variable independent of W and of N . Here W is the standard Brownian motion in dimension one, independent of the multivariate Hawkes process N having intensity (2.1).

Proposition 2.3. *Suppose that Assumption 2.1 holds, that the coefficients b, σ are globally Lipschitz continuous, that a is measurable and that the memory kernel functions h_{ij} are locally integrable. Then equation (2.2) admits a unique strong solution.*

Proof. Theorem 6 of [10] implies that for any $T > 0$, almost surely, N_t has only a finite number of jumps on $[0, T]$. We can therefore work conditionally with respect to the choice of these jumps and construct the solution of (2.2) by pasting together trajectories of strong solutions to the diffusion equation (that is, the solution of (2.2) with $a(\cdot) \equiv 0$) at the successive jump times (see [19] for related ideas). \square

In the sequel we shall also need estimates on the transition densities of the diffusion process underlying (2.2). We want to be able to deal with a general drift term b that is not bounded in order to include Ornstein-Uhlenbeck type processes to the class of models that we consider. Unbounded drift coefficients add however a substantial difficulty when one wishes to apply techniques from Malliavin calculus to obtain bounds on the transition densities. Therefore, we need to impose the following additional assumptions which are inspired by [13].

Assumption 2.4. *1. a is continuous.*

2. b and σ are of class \mathcal{C}^2 .

3. There exist positive constants c, q such that for all $x \in \mathbb{R}$, $|b'(x)| + |\sigma'(x)| \leq c$ and $|b''(x)| + |\sigma''(x)| \leq c(1 + |x|^q)$.
4. There exist strictly positive constants σ_0 and σ_1 such that $\sigma_0 \leq \sigma(x) \leq \sigma_1$ for all $x \in \mathbb{R}$.

Remark 2.5. We do only need the above assumption to obtain good lower bounds on the transition densities of the underlying non-jumping diffusion process, see (3.3) below.

To study the longtime behavior of X and to ensure its ergodicity, we introduce two additional conditions which are classical in the study of diffusions and which are derived from [15], see also [34].

Assumption 2.6. There exist $d \geq 0, r > 0$ such that for all x satisfying $|x| > r$, we have $xb(x) \leq -dx^2$. Moreover, one of the two following conditions holds true.

1. For all $|x| > r$, $2xa(x) + a^2(x) < 0$.
2. For all $|x| > r$, $|a(x)| \leq C|x|^\eta$ for some $\eta < 1$.

2.2 Markovian framework

Our study relies on the general theory of Markov processes. To be able to work within a Markovian framework, we impose a special structure on the interaction functions and suppose that

Assumption 2.7. For all $1 \leq i, j \leq M$,

$$h_{ij}(t) = c_{ij}e^{-\alpha_{ij}t}, c_{ij} \in \mathbb{R}, \alpha_{ij} > 0. \quad (2.3)$$

In this case we may introduce the auxiliary Markov process $Y = Y^{(ij)}$,

$$Y_t^{(ij)} = c_{ij} \int_0^t e^{-\alpha_{ij}(t-s)} dN_s^{(j)} + c_{ij} \int_{-\infty}^0 e^{-\alpha_{ij}(t-s)} dn_s^{(j)}, 1 \leq i, j \leq M. \quad (2.4)$$

The intensities can be expressed in terms of sums of these Markov processes, that is, for all $1 \leq i \leq M$,

$$\lambda_t^{(i)} = f_i \left(\sum_{j=1}^M Y_{t-}^{(ij)} \right).$$

Notice that

$$Y_0^{(ij)} = y_0^{(ij)} = c_{ij} \int_{-\infty}^0 e^{\alpha_{ij}s} dn_s^{(j)}, \quad \text{and} \quad dY_t^{(ij)} = -\alpha_{ij}Y_t^{(ij)}dt + c_{ij}dN_t^{(j)}. \quad (2.5)$$

In particular, the process

$$(X_t, Y_t^{(ij)}, 1 \leq i, j \leq M)$$

is a Markov process. Its longtime behavior is determined on the one hand by the longtime behavior of the underlying Hawkes process N_t and on the other hand on the one of the continuous diffusion process with drift b and diffusion coefficient σ . Note also that N_t is an autonomous process, that is, N_t does not depend on X .

2.3 First properties of the process (X, Y) and an associated stochastic flow.

In the sequel, we shall denote the whole process by $Z_t := (X_t, Y_t)$. It takes values in $\mathbb{R} \times \mathbb{R}^{M \times M}$. We write $(P_t)_{t \geq 0}$ for its associated transition semigroup and $z = (x, y), y = (y^{(ij)})_{1 \leq i, j \leq M}$ for generic elements of its state space.

Proposition 2.8. *Grant Assumptions 2.1 and 2.4. Then the process Z is a Feller process, that is, $P_t \Phi$ belongs to $C_b(\mathbb{R} \times \mathbb{R}^{M \times M})$ whenever $\Phi \in C_b(\mathbb{R} \times \mathbb{R}^{M \times M})$.*

The proof of this result follows from classical arguments, see e.g. the proof of Proposition 4.8 in [18], or [19]. In what follows we shall give an alternative proof relying on an explicit construction of the process Z as a *stochastic flow*. This construction is interesting in its own right and based on the fact that Z is a *piecewise continuous Markov process*, that is, a generalization of the piecewise deterministic Markov processes to those traveling in between successive jumps according to *stochastic flows* instead of deterministic ones, see [19]. We start by giving the principal elements needed to construct this flow.

The associated stochastic Brownian flow. Thanks to our assumptions on b and σ , by Theorem 4.2.5 of [23], there exists a unique stochastic flow of homeomorphisms

$$\Phi_t(x), \quad 0 \leq t < \infty, x \in \mathbb{R}$$

such that for all $t \geq 0, x \in \mathbb{R}$,

$$\Phi_t(x) = x + \int_0^t b(\Phi_u(x)) du + \int_0^t \sigma(\Phi_u(x)) dW_u.$$

In particular, we have that

$$(t, x) \mapsto \Phi_t(x) \text{ is continuous } (0 \leq t < \infty, x \in \mathbb{R}).$$

The stochastic flow Φ describes the evolution of X in between successive jumps of X (or, equivalently, of N).

The associated deterministic flow. In between successive jump events of N , the process Y evolves according to the deterministic flow

$$\varphi_t(y) = (\varphi_t^{(ij)}(y))_{1 \leq i, j \leq M}, \quad \text{where } \varphi_t^{(ij)}(y) = e^{-\alpha_{ij}t} y^{(ij)}, 1 \leq i, j \leq M. \quad (2.6)$$

We are now ready to give the

Simulation algorithm for Z . We propose a simulation algorithm for Z for any family of starting configurations $z \in K_1 \times K_2$, where $K_1 \subset \mathbb{R}, K_2 \subset \mathbb{R}^{M \times M}$ are compact sets. The first step is to construct an upper bound \bar{N}_t on the number of jumps of Z , that is, of N , during some finite time interval $[0, t]$. To do so, observe that by Lipschitz continuity of f_i with Lipschitz constant γ_i , we have

$$f_i(x) \leq f_i(0) + \gamma_i |x|.$$

Let

$$\bar{\gamma} := \max_i \gamma_i, \quad \bar{c} := \sup_{1 \leq i, j \leq M} |c_{ij}|. \quad (2.7)$$

Consider now the one-dimensional auxiliary linear Hawkes process N_t^* having intensity $\lambda_t^*, t \geq 0$, where

$$\lambda_0^* := \sup_{y \in K_2} \sup_{t \geq 0} \sum_{i=1}^M f_i \left(\sum_{j=1}^M e^{-\alpha_{ij}t} y^{(ij)} \right)$$

and

$$\lambda_t^* = \lambda_0^* + M\bar{\gamma}\bar{c} \int_0^{t-} dN_u^* = \lambda_0^* + M\bar{\gamma}\bar{c}N_{t-}^*, \quad t \geq 0.$$

Observe that N^* is a one dimensional linear Hawkes process with memory kernel $\bar{h}(t) = M\bar{\gamma}\bar{c}1_{\mathbb{R}_+}(t)$. Obviously, for all $t \geq 0$,

$$\sum_{i=1}^M N_t^{(i)} \leq N_t^*.$$

Moreover, for all $t \geq 0$, $N_t^* < \infty$ almost surely. In what follows, we shall write $T_1^* < T_2^* < \dots < T_n^* < \dots$ for the successive jump times of N^* .

We work conditionally on the realization of N^* on $[0, t]$, that is, on the event $\{N_t^* = n\}$, and on a realization of the associated jump times $0 < t_1 < t_2 \dots < t_n < t$. Our goal is to construct a version of Z , conditionally on these choices, which is continuous in the starting point z . This construction relies on the classical thinning method, also known as acceptance-rejection method.

During this construction, we choose successively random variables U_1, \dots, U_n taking values in $\{0, 1, \dots, M\}$ and define a process $z_s = z_s(z, t_1^n, U_1^n)$, depending on these choices, for $0 \leq s \leq t$. Here, $t_1^n = (t_1, \dots, t_n)$ denotes the choices of the successive jump times and $U_1^n = (U_1, \dots, U_n)$ the sequence of indices of jumping particles. This process is defined recursively as follows. Firstly, we put

$$z_s = (x_s, y_s) = (\Phi_s(x), \varphi_s(y)) \text{ for all } 0 \leq s < t_1.$$

Then, conditionally on $y_{t_1-} = y_1$, we choose a random variable $U_1 \in \{0, 1, \dots, M\}$ with law $q(y_1, t_1, \cdot)$, where

$$q(y, t, \{i\}) = \frac{f_i(\sum_j y^{(ij)})}{\lambda_t^*}, \quad 1 \leq i \leq M, \quad q(y, t, \{0\}) = 1 - \sum_{i=1}^M q(y, t, \{i\}). \quad (2.8)$$

U_1 gives the label of the particle (or component) that will jump at time t_1 , if $U_1 = 0$, no jump happens at this time. More precisely, on $\{U_1 \geq 1\}$, we put

$$x_{t_1} := x_1 + a(x_1), \quad y_{t_1}^{(ij)} := y_1^{(ij)} + c_{ij}1_{\{U_1=j\}}, \quad \text{for all } 1 \leq i, j \leq M,$$

and $z_{t_1} := z_1$ on $\{U_1 = 0\}$.

Then we put

$$x_s = \Phi_{s-t_1}(x_{t_1}), \quad y_s = \varphi_{s-t_1}(y_{t_1}) \text{ for all } t_1 \leq s < t_2,$$

and we proceed iteratively by choosing, conditionally on $y_{t_2-} = y_2$, a random variable

$$U_2 \sim q(y_2, t_2, \cdot),$$

and so on. Finally, we obtain a terminal value $x_t = \Phi_{t-t_n}(x_{t_n})$ and $y_t = \varphi_{t-t_n}(y_{t_n})$. It is easy to check that

$$\mathcal{L}(z_t(z, t_1^n, U_1^n)) = \mathcal{L}(Z_t | N_t^* = n, T_1^* = t_1, \dots, T_n^* = t_n).$$

The important point is that the above construction ensures the continuity of the application

$$z \mapsto z_t(z, t_1^n, U_1^n).$$

Alternative proof of Proposition 2.8. Noticing that the law of N^* does not depend on the starting point z but only on an upper bound of the associated intensities, the above construction implies that for any $\Phi \in C_b(\mathbb{R} \times \mathbb{R}^{M \times M})$, the mapping

$$K \ni z \mapsto P_t \Phi(z) \in \mathbb{R}$$

is continuous, implying the assertion of Proposition 2.8. Indeed, we can write

$$P_t \Phi(z) = \sum_{n \geq 0} P(N_t^* = n) \int_{[0, t]^n} \mathcal{L}(T_1^*, \dots, T_n^* | N_t^* = n)(dt_1, \dots, dt_n) \\ \sum_{i=1}^n \sum_{u_i=0}^M P(U_1 = u_1, \dots, U_n = u_n | T_1 = t_1, \dots, T_n = t_n, Z_0 = z) \mathbb{E}^W \Phi(z_t(z, u_1^n)),$$

where the last expectation \mathbb{E}^W is only taken with respect to the underlying Brownian motion W . Suppose now that $(z^k)_k \subset K_1 \times K_2$ is a sequence of initial configurations converging to z as $k \rightarrow \infty$. Then, by the structure of (2.8),

$$P(U_1 = u_1, \dots, U_n = u_n | T_1 = t_1, \dots, T_n = t_n, Z_0 = z^k) \rightarrow \\ P(U_1 = u_1, \dots, U_n = u_n | T_1 = t_1, \dots, T_n = t_n, Z_0 = z),$$

as $k \rightarrow \infty$, by continuity of $f_i, 1 \leq i \leq M$, and of $z \mapsto z_s(z, t_1^n, U_1^n)$, for all $s \leq t$. Moreover, using dominated convergence, $\mathbb{E}^W \Phi(z_t(z^k, u_1^n)) \rightarrow \mathbb{E}^W \Phi(z_t(z, u_1^n))$, as $k \rightarrow \infty$, implying that $P_t \Phi(z^k) \rightarrow P_t \Phi(z)$, as $k \rightarrow \infty$, where we have used dominated convergence once more. \square

3 Exponential ergodicity of the process (X, Y)

In this section, we start proving the positive Harris recurrence of $(X_t, Y_t)_t$ using the regeneration technique.

3.1 A Döblin type lower bound

The following theorem is the main result of this section. It states a local Döblin lower bound for the transition semigroup of the process, which is the main ingredient towards ergodicity of the process $Z = (X, Y)$.

Theorem 3.1. *Grant Assumptions 2.4 and 2.7 and suppose moreover that $f_i(x) > 0$ for all $1 \leq i \leq M$, for all $x \in \mathbb{R}$. Then there exists $T > 0$ such that for all $z^* = (x^*, y^*) \in \mathbb{R} \times \mathbb{R}^{M \times M}$ and for all $x^{**} \in \mathbb{R}$ the following holds. There exist $R > 0$, open sets $I_1 \subset \mathbb{R}$ and $I_2 \subset \mathbb{R}^{M \times M}$ with strictly positive Lebesgue measure such that*

$$x^{**} \in I_1, \tag{3.1}$$

and a constant $\beta \in (0, 1)$, depending on I_1, I_2, R and the coefficients of the system with

$$P_T(z, dz') \geq \beta 1_C(z) \nu(dz'), \tag{3.2}$$

where $C = B_R(z^*)$ is the (open) ball of radius R centred at z^* , and where ν is the uniform probability measure on $I_1 \times I_2$.

Remark 3.2. 1. Together with a Lyapunov argument implying that the process Z comes back to the set C infinitely often, the above result will imply the ν -irreducibility of the sampled chain $(Z_{kT})_{k \geq 0}$. We want to stress that the above construction implies that for any given $x^{**} \in \mathbb{R}$ and some $\varepsilon > 0$ we can construct ν such that $B_\varepsilon(x^{**})$ lies in the support of the projection on the x -variable of ν . Of course, this property is related to the support properties of the underlying diffusion process, granted by Assumption 2.4. It will imply that the invariant density of X is bounded away from 0 on each compact set, see Proposition 3.6 below.

2. A lower bound of the type (3.2) is trivial if we are only interested in the transitions of the process X . Indeed, in this case it is sufficient to consider the event where no jump has appeared up to time T and to use known lower bounds on the transition densities of the diffusion part. However, since X is not Markov on its own, we do need to work with the couple of processes (X, Y) , and therefore have to establish such lower bounds for the joint transition of X and Y .

The main idea of the proof is to use the jump noise of M^2 successive jumps of the underlying Hawkes process to create firstly a Lebesgue density for the process Y . Such ideas have already been exploited in [8] and in [12]. The important second step is then to use density estimates of the underlying diffusion to prove that a joint Lebesgue density of (X_T, Y_T) exists which is strictly lower-bounded on C .

The proof is done in several steps which are the subject of the next subsection.

3.2 Some useful properties of the underlying stochastic flow and proof of Theorem 3.1

We start by collecting useful properties of the stochastic flow governing the evolution of X in between successive jumps.

Transition densities. Due to our assumptions on b and σ , $\Phi_t(x)$ possesses a transition density $p_t(x, y)$ with explicit lower bounds. More precisely, Proposition 1.2 of [13] implies that, for some suitable constants c, C ,

$$c^{-1}t^{-1/2}e^{-C(x-y)^2/t}e^{-Cta^2} \leq p_t(x, y) \leq ct^{-1/2}e^{-\frac{1}{c}(x-y)^2/t}e^{Cta^2}. \quad (3.3)$$

Here, the constants c and C do only depend on the coefficients b and σ .

Support properties. We will use the control theorem which goes back to Stroock and Varadhan (1972) [32], see also Millet and Sanz-Sole (1994) [29], Theorem 3.5. For some time horizon $T_1 < \infty$ which is arbitrary but fixed, write \mathbf{H} for the Cameron-Martin space of measurable functions $\mathbf{h} : [0, T_1] \rightarrow \mathbb{R}$ having absolutely continuous components $\mathbf{h}(t) = \int_0^t \dot{\mathbf{h}}(s)ds$ with $\int_0^{T_1} (\dot{\mathbf{h}})^2(s)ds < \infty$. For $x \in \mathbb{R}$ and $\mathbf{h} \in \mathbf{H}$, consider the deterministic flow

$$(\varphi_t^{(\mathbf{h})}(x))_{t \geq 0} \text{ solution to } d\varphi_t^{(\mathbf{h})}(x) = \tilde{b}(\varphi_t^{(\mathbf{h})}(x))dt + \sigma(\varphi_t^{(\mathbf{h})}(x))\dot{\mathbf{h}}(t)dt, \text{ with } \varphi_0^{(\mathbf{h})}(x) = x, \quad (3.4)$$

on $[0, T_1]$. In the above formula (3.4), \tilde{b} is Stratonovich drift given by

$$\tilde{b}(x) = b(x) - \frac{1}{2}\sigma(x)\sigma'(x).$$

Denote by $Q_x^{T_1}$ the law of the solution $(\Phi_t(x))_{0 \leq t \leq T_1}$. Then for any $x \in \mathbb{R}$ and $\mathbf{h} \in \mathbf{H}$,

$$\left(\varphi_t^{(\mathbf{h})}(x)\right)_{|t \in [0, T_1]} \in \overline{\text{supp}\left(Q_x^{T_1}\right)}. \quad (3.5)$$

We are now able to give the proof of Theorem 3.1.

Proof. We suppose without loss of generality that for all $1 \leq j \leq M$,

$$c_{ij} \neq 0 \text{ for all } i, \alpha_{ij} \neq \alpha_{kj}, \text{ for all } 1 \leq i, k \leq M, i \neq k. \quad ^2$$

In what follows, we impose first M consecutive jumps of $N^{(1)}$, followed by M jumps of $N^{(2)}$ etc up to M consecutive jumps of $N^{(M)}$, and we suppose that they all happen before time T . Then we can lower-bound, for all $A \in \mathcal{B}(\mathbb{R}), B \in \mathcal{B}(\mathbb{R}^{M \times M}), z \in \mathbb{R} \times \mathbb{R}^{M \times M}$,

$$P_T(z, A \times B) \geq E_z(1_A(X_T)1_B(Y_T)), N_T^{(i)} = M, \forall 1 \leq i \leq M, \\ T_1^{(1)} < T_2^{(1)} < \dots < T_M^{(1)} < T_1^{(2)} < \dots < T_M^{(2)} < \dots < T_1^{(M)} < \dots < T_M^{(M)} < T),$$

²Otherwise, if $\alpha_{ij} = \alpha_{kj}$, we introduce the new process $Y_t^{(i+k),j} := Y_t^{ij} + Y_t^{kj}$ which is Markov again.

where $T_k^{(i)}, k \geq 1$, are the successive jump times of $N^{(i)}, 1 \leq i \leq M$. In what follows we shall write

$$A_T := \{N_T^{(i)} = M, \forall 1 \leq i \leq M, \\ T_1^{(1)} < T_2^{(1)} < \dots < T_M^{(1)} < T_1^{(2)} < \dots < T_M^{(2)} < \dots < T_1^{(M)} < \dots < T_M^{(M)} < T\}.$$

Step 1. We first deal with the process Y_t . This part of the proof follows well-known arguments that have been developed independently from our work in [8], and that we have also already used in [12] and [24]. We reproduce the main arguments here since the second step will be a control of the diffusion part conditionally on the results of this first step.

Recall that in between successive jump events of N , the process Y evolves according to the deterministic flow $\varphi_t(y)$ introduced in (2.6) above. Thus, on the event A_T , starting from $Y_0 = y \in \mathbb{R}^{M \times M}$, we first let the flow φ evolve starting from y up to some first jump time t_1 . At that jump time each particle having index $(i1)$ gains an additional value c_{i1} . We then successively choose the following inter-jump waiting times t_2, \dots, t_{M^2} under the constraint $t_1 + \dots + t_{M^2} < T$. We write

$$s_1 = T - t_1, s_2 = T - (t_1 + t_2), \dots, s_{M^2} = T - (t_1 + \dots + t_{M^2}).$$

Conditionally on $Y_0 = y$, the successive choices of $\mathbf{s} = (s_1, \dots, s_{M^2}^2)$ as above, the position of $Y_T^{(ij)}$ is given by

$$\gamma^{(ij)}(y, \mathbf{s}) = e^{-\alpha_{ij}T} y^{(ij)} + c_{ij} [e^{-\alpha_{ij}s_{jM-M+1}} + \dots + e^{-\alpha_{ij}s_{jM}}].$$

In what follows we work at fixed y and we write

$$\gamma_y : \mathbf{s} \mapsto \gamma(y, \mathbf{s}) = (\gamma^{(11)}, \dots, \gamma^{(M1)}, \gamma^{(12)}, \dots, \gamma^{(M2)}, \dots, \gamma^{(1M)}, \dots, \gamma^{(MM)})(y, \mathbf{s}).$$

We will use the jump noise which is created by the M^2 jumps, i.e., we will use a change of variables on the account of s_1, \dots, s_{M^2} . Therefore, let

$$\frac{\partial \gamma_y(\mathbf{s})}{\partial \mathbf{s}} = \left[\frac{\partial \gamma_y(\mathbf{s})}{\partial s_1}, \dots, \frac{\partial \gamma_y(\mathbf{s})}{\partial s_{M^2}} \right]$$

be the Jacobian matrix of the the map $\mathbf{s} \mapsto \gamma_y(\mathbf{s})$. This matrix does not depend on the initial position y . Indeed, one easily finds that

$$\frac{\partial \gamma_y(\mathbf{s})}{\partial \mathbf{s}} = C(\mathbf{s}) = \begin{pmatrix} C^{(1)} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & C^{(2)} & \mathbf{0} & \dots & \mathbf{0} \\ \vdots & \vdots & \vdots & \ddots & \dots \\ \mathbf{0} & \dots & \mathbf{0} & \mathbf{0} & C^{(M)} \end{pmatrix}(\mathbf{s}),$$

where for each $1 \leq j \leq M$, $C^{(j)}(\mathbf{s})$ is the $M \times M$ matrix given by

$$C^{(j)}(\mathbf{s}) = - \begin{pmatrix} c_{1j}(\alpha_{1j})^{-1} e^{-\alpha_{1j}s_{jM-M+1}} & \dots & c_{1j}(\alpha_{1j})^{-1} e^{-\alpha_{1j}s_{jM}} \\ c_{2j}(\alpha_{2j})^{-1} e^{-\alpha_{2j}s_{jM-M+1}} & \dots & c_{2j}(\alpha_{2j})^{-1} e^{-\alpha_{2j}s_{jM}} \\ \vdots & \vdots & \vdots \\ c_{Mj}(\alpha_{Mj})^{-1} e^{-\alpha_{Mj}s_{jM-M+1}} & \dots & c_{Mj}(\alpha_{Mj})^{-1} e^{-\alpha_{2j}s_{jM}} \end{pmatrix}.$$

We have to prove that each $C^{(j)}(\mathbf{s})$ is invertible. This is difficult for general choices of \mathbf{s} . But if we choose t_0 such that $M^2 t_0 < T$ and then $s_1^* = M^2 t_0, s_2^* = (M^2 - 1)t_0, \dots, s_{M^2}^* = t_0$ such that $s_k - s_{k+1} = t_0$ for all $1 \leq k < M^2$, then

$$\det C^{(j)}(\mathbf{s}^*) = \left(\prod_{i=1}^M c_{ij}(\alpha_{ij})^{-1} e^{-\alpha_{ij}s_{jM}^*} \right) \det V^{(j)}(t_0),$$

where

$$V^{(j)}(t_0) = \begin{pmatrix} e^{-(M-1)\alpha_1 j t_0} & \dots & e^{-2\alpha_1 j t_0} & e^{-\alpha_1 j t_0} & 1 \\ e^{-(M-1)\alpha_2 j t_0} & \dots & e^{-2\alpha_2 j t_0} & e^{-\alpha_2 j t_0} & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ e^{-(M-1)\alpha_M j t_0} & \dots & e^{-2\alpha_M j t_0} & e^{-\alpha_M j t_0} & 1 \end{pmatrix}.$$

Therefore, $\det V^{(j)}(t_0)$ is a Vandermonde determinant which is different from 0 since by assumption all $e^{-\alpha_{ij} t_0}$ and $e^{-\alpha_{kj} t_0}$ are different, for all $i \neq k$.

By continuity we therefore have that for any choice of $t_0 < T/M^2$ there exists $\varepsilon > 0$ such that for all $\mathbf{s} \in B_\varepsilon(\mathbf{s}^*)$, $\frac{\partial \gamma_y(\mathbf{s})}{\partial \mathbf{s}}$ is invertible.

It will be proved now that this uniform invertibility of the Jacobian matrix of the map $\mathbf{s} \mapsto \gamma_y(\mathbf{s})$ implies the first part of inequality (3.2). For that sake, we shall also need the following notation. For each couple (y, \mathbf{s}) , we write $y_0 = y$, and then define recursively for all $jM - M + 1 \leq k \leq jM$, where j varies between $1 \leq j \leq M$,

$$y_k^{(il)} = \varphi_{s_{k-1} - s_k}^{(il)}(y_{k-1}) + c_{ij} 1_{\{l=j\}}, \quad (3.6)$$

for all $1 \leq i, l \leq M$, where we put $s_0 = T$.

The sequence y_1, \dots, y_{M^2} corresponds to the positions of the process Y right after the successive jumps, starting from the initial location $y \in \mathbb{R}^{M \times M}$, induced by the inter-jump waiting times $T - s_1, s_1 - s_2, \dots, s_{M^2-1} - s_{M^2}$ which are determined by \mathbf{s} .

Introduce now for each $y \in \mathbb{R}^{M \times M}$ the total jump rate

$$\bar{f}(y) := \sum_{i=1}^M f_i \left(\sum_j y^{(ij)} \right)$$

and for each $t \geq 0$, the survival rate

$$e(y, t) = \exp \left\{ - \int_0^t \bar{f}(\varphi_s(y)) ds \right\}. \quad (3.7)$$

We define for each couple (y, \mathbf{s}) (recall that $s_0 = T$),

$$q_y(\mathbf{s}) = \left(\prod_{j=1}^M \prod_{k=jM-M}^{jM-1} f_j(\varphi_{s_k - s_{k+1}}(y_k)) e(y_k, s_k - s_{k+1}) \right) e(y_{M^2}, s_{M^2}). \quad (3.8)$$

Since $\bar{f}(y) > 0$ for all $y \in \mathbb{R}^{M \times M}$ and from the definition of $e(\cdot, \cdot)$, we deduce that for any couple (y^*, \mathbf{s}^*) there are neighborhoods $W_{\mathbf{s}^*}$ and U_{y^*} of \mathbf{s}^* and y^* respectively such that

$$\inf_{(y, \mathbf{s}) \in U_{y^*} \times W_{\mathbf{s}^*}} q_y(\mathbf{s}) > 0. \quad (3.9)$$

Let us now fix (y^*, \mathbf{s}^*) such that the matrix $\frac{\partial \gamma_{y^*}(\mathbf{s}^*)}{\partial \mathbf{s}}$ is invertible. By Lemma 6.2 of [4], there exist an open neighborhood $B_R(y^*)$ of y^* , an open set $I_2 \subset \mathbb{R}^{M \times M}$, and for any $y \in B_R(y^*)$, an open set W_y such that

$$\tilde{\gamma}_y(\mathbf{s}) : \begin{cases} W_y \rightarrow I_2 \\ \mathbf{s} \mapsto \gamma_y(\mathbf{s}) \end{cases}$$

is a diffeomorphism, with $W_y \subset W_{\mathbf{s}^*}$, and also

$$\inf_{y \in B_R(y^*)} \inf_{\mathbf{s} \in W_{\mathbf{s}^*}} \left| \det \left(\frac{\partial \gamma_y(\mathbf{s})}{\partial \mathbf{s}} \right)^{-1} \right| > 0. \quad (3.10)$$

Reducing (if necessary) R , we may assume also that $B_R(y^*) \subset U_{y^*}$. Thus we have that by (3.9) and (3.10),

$$\beta_1 := \inf_{y \in B_R(y^*)} \inf_{s \in W_y} q_y(\mathbf{s}) \left| \det \left(\frac{\partial \gamma_y(\mathbf{s})}{\partial \mathbf{s}} \right)^{-1} \right| > 0. \quad (3.11)$$

Once we have done all these steps we can conclude with the following preliminary result. For any measurable $B \in \mathcal{B}(\mathbb{R}^{M \times M})$ and for any $z = (x, y)$ such that $y \in B_R(y^*)$, using the change of variables $y = \tilde{\gamma}_y(\mathbf{s})$, we obtain the lower bound

$$\begin{aligned} E_y(1_B(Y_T), N_T^{(i)} = M, \forall 1 \leq i \leq M, T_1^{(1)} < \dots < T_M^{(1)} < T_1^{(2)} < \dots < T_M^{(M)} < T) \\ \geq \int_{W_y} q_y(\mathbf{s}) 1_B(\tilde{\gamma}_y(\mathbf{s})) ds_1 \dots ds_{M^2} \geq \beta_1 \int_{I_2 \cap B} dy_1 \dots dy_{M^2}. \end{aligned} \quad (3.12)$$

This would establish the desired result if we were only interested in the autonomous process Y .

Step 2. We now deal with the process X . Of course, we still work conditionally on the choice of s_1, \dots, s_{M^2} of the first step. Analogously to (3.6) we therefore introduce the successive jump positions of the process X which are given by $x_0 = x$ and then for all $1 \leq k \leq M^2$,

$$x_k = \Phi_{s_{k-1}-s_k}(x_{k-1}) + a(\Phi_{s_{k-1}-s_k}(x_{k-1})), \quad (3.13)$$

where $s_0 = T$ as before.

Conditionally on $X_0 = x, Y_0 = y$, the successive choices of $\mathbf{s} = (s_1, \dots, s_{M^2})$ as above, the position of X_T is then given by

$$\Gamma(z, \mathbf{s}) = \Phi_{s_{M^2}}(x_{M^2}),$$

where $z = (x, y)$. Notice that x_{M^2} depends on the choices of s_1, \dots, s_{M^2} and of course on the evolution of the stochastic flow between the successive jump times.

Step 3. Therefore, conditioning with respect to $X_{T-s_{M^2}} = x_{M^2}$, we obtain

$$\begin{aligned} P_T(z, A \times B) &\geq E_z(1_A(X_T) 1_B(Y_T), A_T) \\ &\geq \int_{W_y} q_y(\mathbf{s}) 1_B(\tilde{\gamma}_y(\mathbf{s})) \mathbb{E} \left(\int_A p_{s_{M^2}}(x_{M^2}, u) du \right) ds_1 \dots ds_{M^2}, \end{aligned} \quad (3.14)$$

where $p_{s_{M^2}}(x_{M^2}, u)$ is the transition density of (3.3).

Notice that in the last line, expectation \mathbb{E} is taken with respect to Brownian motion only, that is, with respect to the law of $X_{T-s_{M^2}}$ under the Wiener measure, conditionally on the choices s_1, \dots, s_{M^2} .

The lower bound estimate given in (3.3) implies that for fixed s_{M^2} and x_{M^2} , $u \mapsto p_{s_{M^2}}(x_{M^2}, u)$ is lower bounded in small neighborhoods of x_{M^2} . Therefore, in what follows we need to localize the possible positions of x_{M^2} , conditionally on the choices of the jumps times s_1, \dots, s_{M^2} .

Step 4. Localization of x_{M^2} .

We apply the support theorem, that is, (3.5), to our process X_t in between the successive jump times $0, T_1 := t_1, T_2 := t_1 + t_2, \dots, T_{M^2} := t_1 + \dots + t_{M^2}$, by choosing on each time interval $[T_n, T_{n+1}[$, $n = 0, \dots, M^2 - 1$, the control function $\mathbf{h} \equiv 0$. Consequently, introducing

$$\tilde{x}_1 := \varphi_{t_1}^{(0)}(x) + a(\varphi_{t_1}^{(0)}(x)), \dots, \tilde{x}_{M^2} := \varphi_{s_{M^2-1}-s_{M^2}}^{(0)}(\tilde{x}_{M^2-1}) + a(\varphi_{s_{M^2-1}-s_{M^2}}^{(0)}(\tilde{x}_{M^2-1})),$$

there exists an open neighborhood $U_{\tilde{x}_{M^2}}$ of \tilde{x}_{M^2} , such that

$$\mathbb{P}(X_{T-s_{M^2}} \in U_{\tilde{x}_{M^2}}) > 0.$$

Notice that \tilde{x}_{M^2} is a continuous function of the starting point x and of \mathbf{s} ; that is, $\tilde{x}_{M^2} = F(x, \mathbf{s})$ for some continuous function F . This implies that for any starting point x^* and for $R > 0$ sufficiently

small, reducing $W_{\mathbf{s}^*}$ if necessary, there exists a compact $K = K(x^*, \mathbf{s}^*)$ such that $F(x, \mathbf{s}) \in K$ for all $x \in B_R(x^*)$ and for all $\mathbf{s} \in W_{\mathbf{s}^*}$, whence

$$\inf_{x \in B_R(x^*)} \inf_{\mathbf{s} \in W_{\mathbf{s}^*}} \mathbb{P}(X_{T-s_{M^2}} \in J_1) > 0,$$

where

$$J_1 = \bigcup_{\tilde{x}_{M^2} \in K} U_{\tilde{x}_{M^2}}.$$

Notice that J_1 has compact closure.

Let now $x^{**} \in \mathbb{R}$ be arbitrarily chosen. The lower bound of (3.3) implies that there exists an open interval $I_1 \ni x^{**}$, such that

$$\inf_{x \in I_1, y \in I_1, \mathbf{s} \in W_{\mathbf{s}^*}} P_{s_{M^2}}(x, y) > 0.$$

Therefore,

$$\beta_2 := \inf_{x \in B_R(x^*)} \inf_{y \in B_R(y^*)} \inf_{\mathbf{s} \in W_y} \inf_{u \in J_1, v \in I_1} p_{s_{M^2}}(u, v) \mathbb{P}_x(X_{T-s_M} \in J_1) > 0. \quad (3.15)$$

Therefore, coming back to (3.14), for all $z \in B_R(x^*, y^*)$,

$$\begin{aligned} P_T(z, A \times B) &\geq E_z(1_A(X_T)1_B(Y_T), N_T^{(i)} = M, \forall 1 \leq i \leq M, T_1^{(1)} < T_2^{(1)} < \dots < T_M^{(M)} < T) \\ &\geq \int_{W_y} q_y(\mathbf{s}) 1_B(\tilde{\gamma}_y(\mathbf{s})) \mathbb{E} \left(1_{\{X_{T-s_{M^2}} \in J_1\}} \int_A p_{s_{M^2}}(X_{T-s_{M^2}}, u) du \right) ds_1 \dots ds_{M^2} \\ &\geq \int_{W_y} q_y(\mathbf{s}) 1_B(\tilde{\gamma}_y(\mathbf{s})) \mathbb{E} \left(1_{\{X_{T-s_{M^2}} \in J_1\}} \int_{A \cap I_1} \inf_{x \in J_1} p_{s_{M^2}}(x, u) du \right) ds_1 \dots ds_{M^2} \\ &\geq \beta_2 \lambda(A \cap I_1) \int_{W_y} q_y(\mathbf{s}) 1_B(\tilde{\gamma}_y(\mathbf{s})) ds_1 \dots ds_{M^2} \\ &\geq \beta_2 \lambda(A \cap I_1) \beta_1 \lambda^{M \times M}(B \cap I_2), \end{aligned}$$

where we have finally applied (3.12), and where λ and $\lambda^{M \times M}$ denote the Lebesgue measure on \mathbb{R} , $\mathbb{R}^{M \times M}$, respectively. This implies the desired result putting

$$\beta := \beta_1 \beta_2 \lambda(I_1) \lambda^{M \times M}(I_2) \quad \text{and} \quad \nu := \mathcal{U}_{I_1 \times I_2},$$

the uniform probability law on $I_1 \times I_2$. □

3.3 A Foster-Lyapunov type condition

In order to prove the positive Harris recurrence of the process Z , we need of course a stability condition which is a Lyapunov type condition. Notice that the process $Z_t = (X_t, Y_t)$ has the following extended generator, defined for sufficiently smooth test functions g by

$$\begin{aligned} \mathcal{A}^Z g(x, y) &= - \sum_{i,j=1}^M \alpha_{ij} y^{(ij)} \partial_{y^{(ij)}} g(x, y) + \partial_x g(x, y) b(x) + \frac{1}{2} \sigma^2(x) \partial_x^2 g(x, y) \\ &\quad + \sum_{j=1}^M f_j \left(\sum_{k=1}^M y^{(jk)} \right) [g(x + a(x), y + \Delta_j) - g(x, y)], \end{aligned}$$

with $(\Delta_j)^{(il)} = c_{ij} 1_{\{j=l\}}$ for all $1 \leq i, l \leq M$.

To obtain stability we first introduce the following classical stability assumption for Hawkes processes. Recall that γ_i denotes the Lipschitz constant of the rate function f_i .

Assumption 3.3. Let H be the $M \times M$ -matrix with entries $H_{ij} = \gamma_j \frac{|c_{ij}|}{\alpha_{ij}}, 1 \leq i, j \leq M$. Then its spectral radius satisfies

$$\rho := \rho(H) < 1.$$

Under the above stability condition, let $\kappa \in \mathbb{R}_+^M$ be a left eigenvector of H , associated to the eigenvalue ρ and having non-negative components, that is, for all j , $\sum_i \kappa_i H_{ij} = \rho \kappa_j \geq 0$. Such a vector κ exists thanks to the theorem of Perron-Frobenius. Following [8], we introduce a Lyapunov function by defining $m_{ij} = \frac{\kappa_i}{\alpha_{ij}}$ for all $1 \leq i, j \leq M$ and $V : \mathbb{R} \times \mathbb{R}^{M \times M} \rightarrow \mathbb{R}_+$ by

$$V(x, y) = x^2 + e^{\sum_{i,j} m_{ij} |y^{(ij)}|}. \quad (3.16)$$

Proposition 3.4. Grant Assumptions 2.4, 2.6 and 3.3. Then there exist positive constants d_1, d_2 such that the following Foster-Lyapunov type drift condition holds

$$\mathcal{A}^Z V \leq d_1 - d_2 V. \quad (3.17)$$

Proof. We write $V(x, y) = V_1(x) + V_2(y)$, where $V_1(x) = x^2$, $V_2(y) = e^{\sum_{i,j} m_{ij} |y^{(ij)}|}$, and $\mathcal{A}^Z V = \mathcal{A}_1^Z V + \mathcal{A}_2^Z V$, where

$$\mathcal{A}_1^Z V(x, y) = 2xb(x) + \sigma^2(x) + \sum_{j=1}^M f_j \left(\sum_{k=1}^M y^{(jk)} \right) [(x + a(x))^2 - x^2]$$

is the diffusion part and $\mathcal{A}_2^Z V = \mathcal{A}^Z V - \mathcal{A}_1^Z V$ is the jump part of the generator. The arguments of the proof of Proposition 4.5 of [8] imply that

$$\mathcal{A}_2^Z V(x, y) = \mathcal{A}_2^Z V_2(y) \leq -c_1 V_2(y) + c_2 1_{K_1}(y), \quad (3.18)$$

where $c_1, c_2 > 0$ and where $K_1 \subset \mathbb{R}^{M \times M}$ is some compact. Moreover, writing for short $\bar{f}(y) = \sum_i f_i(\sum_j y^{(ij)})$ for the total jump rate,

$$\mathcal{A}_1^Z V(x, y) = 2xb(x) + \sigma^2(x) + (2xa(x) + a(x)^2) \bar{f}(y). \quad (3.19)$$

Thanks to Assumption 2.6 and boundedness of σ^2 by Assumption 2.4,

$$2xb(x) + \sigma^2(x) \leq -c_3 x^2 + c_4 1_{K_2}(x), \quad (3.20)$$

where $K_2 \subset \mathbb{R}$ is a compact set. Now, suppose that Item 1. of Assumption 2.6 is satisfied. Then for all $|x| > r$, $(2xa(x) + a(x)^2) \bar{f}(y) \leq 0$ (recall that all \bar{f} is non-negative) implying that

$$\mathcal{A}_1^Z V(x, y) \leq -c_3 x^2 + c_4 1_{K_2}(x) + c_5 \bar{f}(y).$$

This implies (3.17) since

$$\bar{f}(y) \leq c_6 + c_7 \sum_{i,j} |y^{(ij)}| \leq c_6 + \tilde{c}_7 \ln(V_2(y))$$

which is thus negligible with respect to the negative term $-c_1 V_2(y)$ of (3.18).

If only Item 2. of Assumption 2.6 holds, then we upper bound the jump part of (3.19) by

$$(2xa(x) + a(x)^2) \bar{f}(y) \leq C \bar{f}(y) + C|x|^{1+\eta} \bar{f}(y).$$

Choose $1 < p < 2$ and $q > 2$ such that $(1 + \eta)p < 2$ and $\frac{1}{p} + \frac{1}{q} = 1$. Then

$$|x|^{1+\eta} \bar{f}(y) \leq \frac{1}{p} |x|^{p(1+\eta)} + \frac{1}{q} \bar{f}^q(y).$$

Since $(1 + \eta)p < 2$, the first term

$$C \frac{1}{p} |x|^{(1+\eta)p}$$

will again be negligible with respect to the negative term $-c_3 x^2$ of (3.20). Finally, the second term $C \bar{f}(y) + C \frac{1}{q} \bar{f}^q(y)$ is treated as previously. This concludes our proof. \square

3.4 Harris recurrence of Z

We do now possess all ingredients to obtain our main results.

Theorem 3.5. *Grant Assumptions 2.4, 2.6, 2.7 and 3.3 and suppose that for all $1 \leq i \leq M$, $f_i(x) > 0$ for all $x \in \mathbb{R}$. Then $(Z_t)_{t \geq 0}$ is positive Harris recurrent with unique invariant measure π . In particular, for any starting point z and any positive measurable function $g : \mathbb{R} \times \mathbb{R}^{M \times M} \rightarrow \mathbb{R}_+$, as $T \rightarrow \infty$, \mathbb{P}_z -almost surely,*

$$\frac{1}{T} \int_0^T g(Z_s) ds \rightarrow \pi(g).$$

Proof of Theorem 3.5. 1) We fix any $x^* \in \mathbb{R}$ and we wish to apply Theorem 3.1 with $y^* = 0$ and $x^* = x^{**}$. Let R be the associated radius.

By Proposition 3.4, we know that for a suitable compact set $K = K_1 \times K_2$, with $K_1 \subset \mathbb{R}, K_2 \subset \mathbb{R}^{M \times M}$, Z comes back to K infinitely often almost surely. Moreover,

$$\sup_{y \in K_2, t \geq 0} \|\varphi_t(y)\|_1 := F < \infty \quad \text{and} \quad \sup_{y \in K_2} \|\varphi_t(y)\|_1 \rightarrow 0$$

as $t \rightarrow \infty$, by the explicit form of the flow in (2.6). Therefore there exists t^* such that $\varphi_t(y) \in B_R(0)$ for all $t \geq t^*$, for all $y \in K_2$.

Applying once more the support theorem for diffusions and observing that σ is strictly positive, Equation (3.5) implies that

$$\inf_{x \in K_1} \mathbb{P}(\Phi_{t^*+s}(x) \in B_R(x^*), 0 \leq s \leq 2T) > 0$$

and thus

$$\inf_{z \in K} \mathbb{P}_z(X_{t^*+s} \in B_R(x^*), Y_{t^*+s} \in B_R(0), 0 \leq s \leq 2T) > 0.$$

Consequently, using a conditional version of the Borel-Cantelli lemma, the sampled Markov chain $(Z_{kT})_{k \in \mathbb{N}}$ visits $B_R(x^*, 0)$ infinitely often almost surely.

2) The standard regeneration technique (see e.g. [25]) allows to conclude that $(Z_{kT})_{k \in \mathbb{N}}$ and therefore $(Z_t)_t$ are Harris recurrent. This concludes the proof. \square

The following by-product of the above result will prove to be useful when dealing with statistical inference within this new model class.

Proposition 3.6. *Grant all assumptions of Theorem 3.5 and write π^X for the projection of the invariant measure π onto the X -coordinate, that is, $\pi^X(dx) = \int_{\mathbb{R}^{M \times M}} \pi(dx, dy)$. Then π^X possesses a Lebesgue density which is bounded away from zero on each compact of \mathbb{R} .*

Proof. Let $A \in \mathcal{B}(\mathbb{R})$. Then for any $t > 0$,

$$\pi^X(A) = \int_{\mathbb{R} \times \mathbb{R}^{M \times M}} \pi(dz) \mathbb{E}_z[1_A(X_t)]. \quad (3.21)$$

Let $L_t := \sup\{s \leq t : \exists j : \Delta N_s^{(j)} = 1\}$ be the last jump time of the process before time t , $L_t = 0$ if there is no such jump. Then by Fubini,

$$\mathbb{E}_z[1_A(X_t)] = \int_A \mathbb{E}_z[p_{t-L_t}(X_{L_t}, y)] dy,$$

where $p_t(x, y)$ is the transition density of (3.3). This implies the existence of the density $\pi^X(x)$ which is given by

$$\pi^X(x) = \int_{\mathbb{R} \times \mathbb{R}^{M \times M}} \pi(dz) \mathbb{E}_z[p_{t-L_t}(X_{L_t}, x)]$$

for any $t > 0$.

Notice that we do not dispose of any regularity results of $\pi^X(x)$ with respect to x . Indeed, the upper bound in (3.3) does not allow to conclude that the almost sure continuity in x of $p_{t-L_t}(X_{L_t}, x)$ survives the integration $\pi(dz) \mathbb{E}_z(\dots)$.

We are however able to prove that π^X is lower bounded on compacts $K \subset \mathbb{R}$. For that sake, fix any $x^{**} \in K$ and apply (3.2) to $(x^*, y^*) \in \text{supp}(\pi)$ such that $\pi(B_R(z^*)) = \pi(C) > 0$ and to x^{**} . Then for any measurable $A \subset I_1 = I_1(x^{**})$, applying the lower bound of (3.2) to (3.21) and taking $t = T$,

$$\int_A \pi^X(x) dx \geq \beta \pi(C) \frac{1}{\lambda(I_1)} \int_A dx$$

implying that

$$\inf_{x \in I_1} \pi^X(x) \geq \beta \pi(C) \frac{1}{\lambda(I_1)} > 0.$$

Therefore we have just shown that for all $x^{**} \in K$, there exists an open interval $I_1 = I_1(x^{**})$ containing $x^{**} \in I_1$ such that π^X is strictly lower bounded on I_1 . Since we can cover the compact K by a finite collection of such open intervals $I_1(x^{**})$, this implies the desired lower bound of π^X on compacts. \square

In the sequel, following [28], we introduce

$$\bar{V}(z) := V(z) + 1 \text{ and } \|\mu\|_{\bar{V}} := \sup_{g: |g| \leq \bar{V}} |\mu(g)|.$$

It is now straightforward to obtain our second main result.

Theorem 3.7 (Exponential ergodicity). *Grant all assumptions of Theorem 3.5. Then there exist $c_1, c_2 > 0$ such that for all $z \in \mathbb{R} \times \mathbb{R}^{M \times M}$,*

$$\|P_t(z, \cdot) - \pi\|_{\bar{V}} \leq c_1 \bar{V}(z) e^{-c_2 t}. \quad (3.22)$$

Proof. The sampled chain $(Z_{kT})_{k \geq 0}$ is Feller according to Proposition 2.8. Moreover it is ν -irreducible, where ν is the measure introduced in Theorem 3.1, associated with the point $(x^*, 0)$ and x^{**} , for any choice of $x^*, x^{**} \in \mathbb{R}$, used in the proof of Theorem 3.5. Since ν is the uniform measure on some open set of strictly positive Lebesgue measure, the support of ν has non-empty interior. Theorem 3.4 of [27] implies that all compact sets are ‘petite’ sets of the sampled chain. The Lyapunov condition established in Proposition 3.4 allows to apply Theorem 6.1 of [28] which implies the assertion. \square

3.5 Exponential β -mixing for $Z = (X, Y)$.

It is now easy to deduce from the above results the exponentially β -mixing property of the process. Recall that the β -mixing coefficient of Z is given by

$$\beta_Z(t) = \sup_{s \geq 0} \int \|P_t(z, \cdot) - \mu P_{s+t}(\cdot)\|_{TV} \mu P_s(dz),$$

where $\mu = \mathcal{L}(Z_0)$ is the law of the initial configuration and where

$$\|\mu\|_{TV} := \sup_{g: |g| \leq 1} \mu(g)$$

denotes the total variation distance. Notice that if $\mu = \pi$, then the process is in its stationary regime, and

$$\beta_Z(t) = \int \|P_t(z, \cdot) - \pi\|_{TV} \pi(dz).$$

Theorem 3.8. *Grant all assumptions of Theorem 3.5. Then Z is exponentially β -mixing, that is, there exist constants $K, \theta > 0$ such that for any initial law μ with $\mu(V) < \infty$,*

$$\beta_Z(t) \leq K e^{-\theta t}.$$

Proof. Suppose firstly that $\mu = \pi$. Then Theorem 4.3 of [28] implies that $\int V d\pi < \infty$ such that we are able to integrate (3.22) against $\pi(dx)$ to obtain

$$\beta_Z(t) \leq c_1 \pi(\bar{V}) e^{-c_2 t}.$$

Putting $K := c_1$ and $\theta = c_2$, this implies the result in this case.

In order to deal with the general non-stationary process, we apply Lemma 3.9 of [26] with $h = \bar{V}$, $\delta(t) = c_1 e^{-c_2 t}$ and

$$\kappa = \sup_{s \geq 0} \mathbb{E}(\bar{V}(Z_s)),$$

to deduce that

$$\beta_Z(t) \leq 2c_1 \kappa e^{-c_2 t}.$$

Putting $K := 2c_1 \kappa$ and $\theta = c_2$, this implies the result, if we have already shown that κ is finite. This last fact follows immediately from (3.17), following the first lines of the proof of Theorem 6.1 of [28]. Indeed, we have by Dynkin's formula that

$$e^{\alpha t} \mathbb{E}_z(V(Z_t)) \leq V(z) + \frac{\beta}{\alpha} e^{\alpha t},$$

implying that

$$\mathbb{E}_z(V(Z_t)) \leq e^{-\alpha t} V(z) + \frac{\beta}{\alpha}.$$

Integrating this last inequality with respect to $\mu(dz)$ implies the result. \square

References

- [1] D. Applebaum. *Lévy processes and Stochastic Calculus*. Cambridge University Press, 2009.
- [2] E. Bacry, S. Delattre, M. Hoffmann, and J-F Muzy. Some limit theorems for Hawkes processes and application to financial statistics. *Stochastic Processes and their Applications*, 123(7):2475–2499, 2013.
- [3] E. Bacry and J-F Muzy. Second order statistics characterization of Hawkes processes and non-parametric estimation. *arXiv preprint arXiv:1401.0903*, 2014.
- [4] M. Benaïm, S. Le Borgne, F. Malrieu, and P-A Zitt. Qualitative properties of certain piecewise deterministic Markov processes. *Ann. Inst. H. Poincaré Probab. Statist.*, 51(3):1040–1075, 08 2015.
- [5] M. Bonde Raad. Renewal time points for Hawkes processes. *arXiv*, 2019.
- [6] P. Brémaud and L. Massoulié. Stability of nonlinear Hawkes processes. *The Annals of Probability*, pages 1563–1588, 1996.

- [7] S. Chen, A. Shojaie, E. Shea-Brown, and D. Witten. The multivariate Hawkes process in high dimensions: Beyond mutual excitation. *arXiv preprint arXiv:1707.04928*, 2017.
- [8] Simon Clinet and Nakahiro Yoshida. Statistical inference for ergodic point processes and application to limit order book. *Stochastic Processes and their Applications*, 127(6):1800 – 1839, 2017.
- [9] Daryl J. Daley and D. Vere-Jones. *An introduction to the theory of point processes: volume II: general theory and structure*. Springer Science & Business Media, 2007.
- [10] S. Delattre, N. Fournier, and M Hoffmann. Hawkes processes on large networks. *The Annals of Applied Probability*, 26(1):216–261, 2016.
- [11] C. Dion and S. Lemler. Nonparametric drift estimation for diffusions with jumps driven by a Hawkes process. *HAL*, 2019.
- [12] A. Duarte, E. Löcherbach, and G. Ost. Stability, convergence to equilibrium and simulation of non-linear Hawkes processes with memory kernels given by the sum of Erlang kernels. *arXiv preprint arXiv:1610.03300*, 2016.
- [13] E. Gobet. Lan property for ergodic diffusions with discrete observations. *Ann. Inst. H. Poincaré Probab. Statist.*, 38(5):711–737, 2002.
- [14] C. Graham. Regenerative properties of the linear Hawkes process with unbounded memory. *arXiv*, 2019.
- [15] R.Z. Has'minskii. *Stochastic stability of differential equations*. Sijthoff & Noordhoff, 1980.
- [16] A.G Hawkes. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, 58(1):83–90, 1971.
- [17] P. Hodara and E. Löcherbach. Hawkes processes with variable length memory and an infinite number of components. *Advances in Applied Probability*, 49(1):84–107, 2017.
- [18] R. Höpfner and E. Löcherbach. Statistical models for Birth and Death on a Flow: Local absolute continuity and likelihood ratio processes. *Scandinavian Journal of Statistics*, 26(1):107–128, 1999.
- [19] N. Ikeda, M. Nagasawa, and S. Watanabe. A construction of Markov processes by piecing out. *Proc. Japan Acad.*, 42(4):370–375, 1966.
- [20] N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. North-Holland Mathematical Library. Elsevier Science, 2014.
- [21] K. Ito and H.P McKean. *Diffusion processes and their sample paths*, 1965.
- [22] SA. Klokov and A Yu Veretennikov. On subexponential mixing rate for Markov processes. *Theory of Probability & Its Applications*, 49(1):110–122, 2005.
- [23] H. Kunita. *Stochastic flows and stochastic differential equations*. Cambridge University Press, Cambridge, 1992.
- [24] E. Löcherbach. Absolute continuity of the invariant measure in piecewise deterministic Markov Processes having degenerate jumps. *Stoch. Proc. Appl.*, 128:1797–1829, 2018.

- [25] E. Löcherbach and D. Loukianova. On Nummelin splitting for continuous time Harris recurrent Markov processes and application to kernel estimation for multi-dimensional diffusions. *Stoch. Proc. Appl.*, 118:1301–1321, 2008.
- [26] H Masuda. Ergodicity and exponential β -mixing bounds for multidimensional diffusions with jumps. *Stochastic Processes and their Applications*, 117(1):35–56, 2007.
- [27] S.P. Meyn and R.L. Tweedie. Stability of Markovian processes I : Criteria for discrete-time chains. *Adv. Appl. Probab.*, 24:542–574, 1992.
- [28] S.P. Meyn and R.L. Tweedie. Stability of Markovian processes III : Foster-Lyapunov criteria for continuous-time processes. *Adv. Appl. Probab.*, 25:487–548, 1993.
- [29] A. Millet and M Sanz-Solé. A simple proof of the support theorem for diffusion processes. *Séminaire de probabilités de Strasbourg*, 28:36–48, 1994.
- [30] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293. Springer Science & Business Media, 2013.
- [31] P. Reynaud-Bouret, V. Rivoirard, and C. Tuleau-Malot. Inference of functional connectivity in neurosciences via Hawkes processes. In *Global Conference on Signal and Information Processing (GlobalSIP), 2013 IEEE*, pages 317–320. IEEE, 2013.
- [32] D.W. Stroock and S. R. S. Varadhan. On the support of diffusion processes with applications to the strong maximum principle. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 3: Probability Theory*, pages 333–359, Berkeley, Calif., 1972. University of California Press.
- [33] P. Tankov and E. Voltchkova. Jump-diffusion models: a practitioner’s guide. *Banque et Marchés*, 99(1):24, 2009.
- [34] A.Yu. Veretennikov. On polynomial mixing bounds for stochastic differential equations. *Stochastic Processes and their Applications*, 70(1):115 – 127, 1997.