



Observation of CP Violation in Charm Decays

Roel Aaij, Carlos Abellán Beteta, Bernardo Adeva, Marco Adinolfi, Christine Angela Aidala, Ziad Ajaltouni, Simon Akar, Pietro Albicocco, Johannes Albrecht, Federico Alessio, et al.

► To cite this version:

Roel Aaij, Carlos Abellán Beteta, Bernardo Adeva, Marco Adinolfi, Christine Angela Aidala, et al.. Observation of CP Violation in Charm Decays. Physical Review Letters, 2019, 122 (21), pp.211803. 10.1103/PhysRevLett.122.211803 . hal-02097259

HAL Id: hal-02097259

<https://hal.science/hal-02097259>

Submitted on 7 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Observation of CP violation in charm decays

LHCb collaboration[†]

Abstract

A search for charge-parity (CP) violation in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays is reported, using pp collision data corresponding to an integrated luminosity of 5.9 fb^{-1} collected at a center-of-mass energy of 13 TeV with the LHCb detector. The flavor of the charm meson is inferred from the charge of the pion in $D^{*}(2010)^+ \rightarrow D^0 \pi^+$ decays or from the charge of the muon in $\bar{B} \rightarrow D^0 \mu^- \bar{\nu}_\mu X$ decays. The difference between the CP asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays is measured to be $\Delta A_{CP} = [-18.2 \pm 3.2 \text{ (stat.)} \pm 0.9 \text{ (syst.)}] \times 10^{-4}$ for π -tagged and $\Delta A_{CP} = [-9 \pm 8 \text{ (stat.)} \pm 5 \text{ (syst.)}] \times 10^{-4}$ for μ -tagged D^0 mesons. Combining these with previous LHCb results leads to

$$\Delta A_{CP} = (-15.4 \pm 2.9) \times 10^{-4},$$

where the uncertainty includes both statistical and systematic contributions. The measured value differs from zero by more than five standard deviations. This is the first observation of CP violation in the decay of charm hadrons.

Published in Phys. Rev. Lett. **122** (2019) 211803

© 2019 CERN for the benefit of the LHCb collaboration. CC-BY-4.0 licence.

[†]Authors are listed at the end of this Letter.

The noninvariance of fundamental interactions under the combined action of charge conjugation (C) and parity (P) transformations, so-called CP violation, is a necessary condition for the dynamical generation of the baryon asymmetry of the universe [1]. The Standard Model (SM) of particle physics includes CP violation through an irreducible complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [2, 3]. The realization of CP violation in weak interactions has been established in the K - and B -meson systems by several experiments [4–12], and all results are well interpreted within the CKM formalism. However, the size of CP violation in the SM appears to be too small to account for the observed matter-antimatter asymmetry [13–15], suggesting the existence of sources of CP violation beyond the SM.

The observation of CP violation in the charm sector has not been achieved yet, despite decades of experimental searches. Charm hadrons provide a unique opportunity to measure CP violation with particles containing only up-type quarks. The size of CP violation in charm decays is expected to be tiny in the SM, with asymmetries typically of the order of 10^{-4} – 10^{-3} , but due to the presence of low-energy strong-interaction effects, theoretical predictions are difficult to compute reliably [16–34]. Motivated by the fact that contributions of beyond-the-SM virtual particles may alter the size of CP violation with respect to the SM expectation, a number of theoretical analyses have been performed [19, 27, 32, 35].

Unprecedented experimental precision can be reached at LHCb in the measurement of CP -violating asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays. The inclusion of charge-conjugate decay modes is implied throughout except in asymmetry definitions. Searches for CP violation in these decay modes have been performed by the BaBar [36], Belle [37], CDF [38, 39] and LHCb [40–44] collaborations. The corresponding CP asymmetries have been found to be consistent with zero within a precision of a few per mille.

This Letter presents a measurement of the difference of the time-integrated CP asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays, performed using pp collision data collected with the LHCb detector at a center-of-mass energy of 13 TeV, and corresponding to an integrated luminosity of 5.9 fb^{-1} .

The time-dependent CP asymmetry, $A_{CP}(f; t)$, between states produced as D^0 or $\bar{D}^0$ mesons decaying to a CP eigenstate f at time t is defined as

$$A_{CP}(f; t) \equiv \frac{\Gamma(D^0(t) \rightarrow f) - \Gamma(\bar{D}^0(t) \rightarrow f)}{\Gamma(D^0(t) \rightarrow f) + \Gamma(\bar{D}^0(t) \rightarrow f)}, \quad (1)$$

where Γ denotes the time-dependent rate of a given decay. For $f = K^- K^+$ or $f = \pi^- \pi^+$, $A_{CP}(f; t)$ can be expressed in terms of a direct component associated to CP violation in the decay amplitude and another component associated to CP violation in D^0 – $\bar{D}^0$ mixing or in the interference between mixing and decay.

A time-integrated asymmetry, $A_{CP}(f)$, can be determined, and its value will exhibit a dependence on the variation of the reconstruction efficiency as a function of the decay time. To first order in the D^0 – $\bar{D}^0$ mixing parameters, it can be written as [38, 45]

$$A_{CP}(f) \approx a_{CP}^{\text{dir}}(f) - \frac{\langle t(f) \rangle}{\tau(D^0)} A_{\Gamma}(f), \quad (2)$$

where $\langle t(f) \rangle$ denotes the mean decay time of $D^0 \rightarrow f$ decays in the reconstructed sample, incorporating the effects of the time-dependent experimental efficiency, $a_{CP}^{\text{dir}}(f)$ is the direct

CP asymmetry, $\tau(D^0)$ the D^0 lifetime and $A_\Gamma(f)$ the asymmetry between the $D^0 \rightarrow f$ and $\bar{D}^0 \rightarrow f$ effective decay widths [46, 47]. In the limit of U-spin symmetry, the direct CP asymmetry is equal in magnitude and opposite in sign for K^-K^+ and $\pi^-\pi^+$, though the size of U-spin-breaking effects at play is uncertain [19]. Taking A_Γ to be independent of final state [19, 48, 49], the difference in CP asymmetries between $D^0 \rightarrow K^-K^+$ and $D^0 \rightarrow \pi^-\pi^+$ decays is

$$\begin{aligned}\Delta A_{CP} &\equiv A_{CP}(K^-K^+) - A_{CP}(\pi^-\pi^+) \\ &\approx \Delta a_{CP}^{\text{dir}} - \frac{\Delta\langle t \rangle}{\tau(D^0)} A_\Gamma,\end{aligned}\quad (3)$$

where $\Delta a_{CP}^{\text{dir}} \equiv a_{CP}^{\text{dir}}(K^-K^+) - a_{CP}^{\text{dir}}(\pi^-\pi^+)$ and $\Delta\langle t \rangle$ is the difference of the mean decay times $\langle t(K^-K^+) \rangle$ and $\langle t(\pi^-\pi^+) \rangle$.

The D^0 mesons considered in this analysis are produced either promptly at a pp collision point (primary vertex, PV) in the strong decay of $D^{*}(2010)^+$ mesons (hereafter referred to as D^{*+}) to a $D^0\pi^+$ pair or at a vertex displaced from any PV in semileptonic $\bar{B} \rightarrow D^0\mu^-\bar{\nu}_\mu X$ decays, where $\bar{B}$ denotes a hadron containing a b quark and X stands for potential additional particles. The flavor at production of D^0 mesons from D^{*+} decays is determined from the charge of the accompanying pion (π -tagged), whereas that of D^0 mesons from semileptonic b -hadron decays is obtained from the charge of the accompanying muon (μ -tagged). The raw asymmetries measured for π -tagged and μ -tagged D^0 decays are defined as

$$\begin{aligned}A_{\text{raw}}^{\pi\text{-tagged}}(f) &\equiv \frac{N(D^{*+} \rightarrow D^0(f)\pi^+) - N(D^{*-} \rightarrow \bar{D}^0(f)\pi^-)}{N(D^{*+} \rightarrow D^0(f)\pi^+) + N(D^{*-} \rightarrow \bar{D}^0(f)\pi^-)}, \\ A_{\text{raw}}^{\mu\text{-tagged}}(f) &\equiv \frac{N(\bar{B} \rightarrow D^0(f)\mu^-\bar{\nu}_\mu X) - N(B \rightarrow \bar{D}^0(f)\mu^+\nu_\mu X)}{N(\bar{B} \rightarrow D^0(f)\mu^-\bar{\nu}_\mu X) + N(B \rightarrow \bar{D}^0(f)\mu^+\nu_\mu X)},\end{aligned}\quad (4)$$

where N is the measured signal yield for the given decay. These can be approximated as

$$\begin{aligned}A_{\text{raw}}^{\pi\text{-tagged}}(f) &\approx A_{CP}(f) + A_D(\pi) + A_P(D^*), \\ A_{\text{raw}}^{\mu\text{-tagged}}(f) &\approx A_{CP}(f) + A_D(\mu) + A_P(B),\end{aligned}\quad (5)$$

where $A_D(\pi)$ and $A_D(\mu)$ are detection asymmetries due to different reconstruction efficiencies between positive and negative tagging pions and muons, whereas $A_P(D^*)$ and $A_P(B)$ are the production asymmetries of D^* mesons and b hadrons, arising from the hadronization of charm and beauty quarks in pp collisions [50]. Owing to the smallness of the involved terms, which averaged over phase space for selected events are $\mathcal{O}(10^{-2})$ or less [50–53], the approximations in Eqs. (5) are valid up to corrections of $\mathcal{O}(10^{-6})$. The values of $A_D(\pi)$ and $A_P(D^*)$, as well as those of $A_D(\mu)$ and $A_P(B)$, are independent of the final state f , and thus cancel in the difference, resulting in

$$\Delta A_{CP} = A_{\text{raw}}(K^-K^+) - A_{\text{raw}}(\pi^-\pi^+).\quad (6)$$

This simple relation between ΔA_{CP} and the measurable raw asymmetries in K^-K^+ and $\pi^-\pi^+$ makes the determination of ΔA_{CP} largely insensitive to systematic uncertainties.

The LHCb detector is a single-arm forward spectrometer designed for the study of particles containing b or c quarks, as described in detail in Refs. [54, 55]. The LHCb

tracking system exploits a dipole magnet to measure the momentum of charged particles. Although the analysis presented in this Letter is expected to be insensitive to such effects, the magnetic-field polarity is reversed periodically during data taking to mitigate the differences of reconstruction efficiencies of particles with opposite charges. Data sets corresponding to about one half of the total integrated luminosity are recorded with each magnetic-field configuration.

The online event selection is performed by a trigger, which consists of a hardware stage based on information from the calorimeter and muon systems, followed by two software stages. In the first software stage, events used in this analysis are selected if at least one track has large transverse momentum and is incompatible with originating from any PV, or if any two-track combination forming a secondary vertex, consistent with that of a D^0 decay, is found in the event by a multivariate algorithm [56, 57]. In between the first and second software stages, detector alignment and calibration are performed and updated constants are made available to the software trigger [58]. In the second stage, D^0 candidates are fully reconstructed using kinematic, topological and particle-identification (PID) criteria. Requirements are placed on: the D^0 decay vertex, which must be well separated from all PVs in the event; the quality of reconstructed tracks; the D^0 transverse momentum; the angle between the D^0 momentum and its flight direction; PID information; and the impact-parameter significances (χ_{IP}^2) of the D^0 decay products with respect to all PVs in the event, where the χ_{IP}^2 is defined as the difference between the χ^2 of the PV reconstructed with and without the considered particle. In the analysis of the μ -tagged sample, B candidates are formed by combining a D^0 candidate with a muon under the requirement that they are consistent with originating from a common vertex. In addition, requirements on the invariant mass of the $D^0\mu$ system, $m(D^0\mu)$, and on the corrected mass (m_{corr}) are applied. The corrected mass partially recovers the missing energy of the unreconstructed particles and is defined as $m_{\text{corr}} \equiv \sqrt{m(D^0\mu)^2 + p_{\perp}(D^0\mu)^2 + p_{\perp}(D^0\mu)}$ [59], where $p_{\perp}(D^0\mu)$ is the momentum of the $D^0\mu$ system transverse to the flight direction of the b hadron, determined from the primary and $D^0\mu$ vertices.

In the offline selection, trigger signals are associated to reconstructed particles. Selection requirements are applied on the trigger decision, taking into account the information on whether the decision was taken due to the signal decay products or to other particles produced in the event. Fiducial requirements are imposed to exclude kinematic regions characterized by large detection asymmetries for the tagging pion or muon. Very large raw asymmetries, up to 100%, occur in certain kinematic regions because, for a given magnet polarity, low-momentum particles of one charge at large or small polar angles in the horizontal plane may be deflected out of the detector or into the (uninstrumented) LHC beam pipe, whereas particles with the other charge are more likely to remain within the acceptance [60]. About 35% and 10% of the selected candidates are rejected by these fiducial requirements for the π -tagged and μ -tagged samples, respectively. In the retained samples, raw asymmetries are typically at the percent level or below. For π -tagged D^0 mesons, a requirement on the D^0 χ_{IP}^2 is applied to suppress the background of D^0 mesons from B decays, and PID requirements on the D^0 decay products are further tightened. Then the D^0 and pion candidates are combined to form D^{*+} candidates by requiring a good fit quality of the D^{*+} vertex and the invariant mass of D^0 candidates to lie within a range of about ± 3 standard deviations around the known D^0 mass. The D^{*+} vertex is determined as a common vertex of D^0 and tagging π^+ candidates, and is constrained to

coincide with the nearest PV [61].

For μ -tagged mesons, the B candidates are further filtered using a dedicated boosted decision tree (BDT) to suppress the combinatorial background due to random combinations of charged kaon or pion pairs not originating from a D^0 decay. The variables used in the BDT to discriminate signal from combinatorial background are: the fit quality of the D^0 and the B decay vertices; the D^0 flight distance; the D^0 impact parameter, *i.e.*, the minimum distance of its trajectory to the nearest PV; the transverse momenta of the D^0 decay products, the significance of the distance between the D^0 and B decay vertices; the invariant mass $m(D^0\mu)$ and the corrected mass m_{corr} . To suppress background from b -hadron decays to $c\bar{c}\pi^\pm X$ ($c\bar{c}K^\pm X$), where the $c\bar{c}$ resonance decays to a pair of muons, D^0 candidates are vetoed if the invariant mass of the $\mu^\mp\pi^\pm$ ($\mu^\mp K^\pm$) pair, where the pion (kaon) is given the muon mass hypothesis, lies within a window of about $\pm 50 \text{ MeV}/c^2$ around the J/ψ or $\psi(2S)$ known masses.

The data sample includes events with multiple D^{*+} and B candidates. The majority of these events contain the same reconstructed D^0 meson combined with different tagging pions or muons. When multiple candidates are present in the event, only one is kept randomly. The fractions of events with multiple candidates are about 10% and 0.4% in the π -tagged and μ -tagged samples, respectively. A small fraction of events, of the order of per mille, belong to both the selected π -tagged and μ -tagged samples.

As the detection and production asymmetries are expected to depend on the kinematics of the reconstructed particles, the cancellation in the difference between the raw asymmetries in Eq. (6) may be incomplete if the kinematic distributions of reconstructed D^{*+} or B candidates and of the tagging pions or muons differ between the K^-K^+ and $\pi^-\pi^+$ decay modes. For this reason, a small correction to the K^-K^+ sample is applied by means of a weighting procedure [60]. For the π -tagged sample, candidate-by-candidate weights are calculated by taking the ratio between the three-dimensional background-subtracted distributions of transverse momentum, azimuthal angle and pseudorapidity of the D^{*+} meson in the K^-K^+ and $\pi^-\pi^+$ modes. An analogous procedure is followed for the μ -tagged sample, where D^0 distributions are used instead of those of the D^{*+} meson. It is then checked *a posteriori* that the distributions of the same variables for tagging pions and muons are also equalized by the weighting. The application of the weights leads to a small variation of ΔA_{CP} , below 10^{-4} for both the π -tagged and μ -tagged samples.

The raw asymmetries of signal and background components for each decay mode are free parameters determined by means of simultaneous least-square fits to the binned mass distributions of D^{*+} and D^{*-} candidates for the π -tagged sample, or D^0 and $\bar{D}^0$ candidates for the μ -tagged sample. In particular, in the analysis of the π -tagged sample the fits are performed to the $m(D^0\pi^+)$ and $m(\bar{D}^0\pi^-)$ distributions. As outlined in Ref. [38], using these distributions has the advantage that they are the same for both $D^0 \rightarrow K^-K^+$ and $D^0 \rightarrow \pi^-\pi^+$ decay modes.

The signal mass model, which is obtained from simulation, consists of the sum of three Gaussian functions and a Johnson S_U function [62], whose parameters are free to be adjusted by the fit to the data. The mean values of the Gaussian functions are distinct for positive and negative tags, whereas widths and fractions are shared. The parameters of the Johnson S_U function, which accounts for the slight asymmetric shape of the signal distribution due to the proximity of the $m(D^0) + m(\pi^+)$ threshold, are also shared. The combinatorial background is described by an empirical function of the form $[m(D^0\pi^+) - m(D^0) - m(\pi^+)]^\alpha e^{\beta m(D^0\pi^+)}$, where α and β are two free parameters which




Figure 1: Mass distributions of selected (top) $\pi^\pm$ -tagged and (bottom) $\mu^\pm$ -tagged candidates for (left) $K^- K^+$ and (right) $\pi^- \pi^+$ final states of the D^0 -meson decays, with fit projections overlaid.

are shared among positive and negative tags. In the analysis of the μ -tagged sample, the fits are performed to the $m(D^0)$ distributions. The signal is described by the sum of two Gaussian functions convolved with a truncated power-law function that accounts for final-state photon radiation effects, whereas the combinatorial background is described by an exponential function. A small contribution from $D^0 \rightarrow K^- \pi^+$ decays with a misidentified kaon or pion is also visible, which is modeled as the tail of a Gaussian function. Separate fits are performed to subsamples of data collected with different magnet polarities and in different years. All partial ΔA_{CP} values corresponding to each subsample are found to be in good agreement and then averaged to obtain the final results. If single fits are performed to the overall π -tagged and μ -tagged samples, small differences of the order of a few 10^{-5} are found. The $m(D^0 \pi^+)$ and $m(D^0)$ distributions corresponding to the entire samples are displayed in Fig. 1 (see also Ref. [60] for the corresponding asymmetries as a function of mass). The π -tagged (μ -tagged) signal yields are approximately 44 (9) million

$D^0 \rightarrow K^- K^+$ decays and 14 (3) million $D^0 \rightarrow \pi^- \pi^+$ decays. In the case of π -tagged decays, the fits to the $m(D^0 \pi^+)$ distributions do not distinguish between background that produces peaks in $m(D^0 \pi^+)$, which can arise from D^{*+} decays where the correct tagging pion is found but the D^0 meson is misreconstructed, and signal. The effect on ΔA_{CP} of residual peaking backgrounds, suppressed by selection requirements to less than 1% of the number of signal candidates, is evaluated as a systematic uncertainty.

Studies of systematic uncertainties on ΔA_{CP} are carried out independently for the π -tagged and μ -tagged samples. Several sources affecting the measurement are considered. In the case of π -tagged decays, the dominant systematic uncertainty is related to the knowledge of the signal and background mass models. It is evaluated by generating pseudoexperiments according to the baseline fit model, then fitting alternative models to those data. A value of 0.6×10^{-4} is assigned as a systematic uncertainty, corresponding to the largest variation observed using the alternative functions. Possible differences between $D^0 \pi^+$ and $\bar{D}^0 \pi^-$ invariant-mass shapes are investigated by studying a sample of 232 million $D^{*+} \rightarrow D^0(K^- \pi^+) \pi^+$ and $D^{*-} \rightarrow \bar{D}^0(K^+ \pi^-) \pi^-$ decays. The effect on ΔA_{CP} is estimated to be order of 10^{-5} at most, hence negligible. A similar study with pseudoexperiments is also performed with the μ -tagged sample and a value of 2×10^{-4} is found.

In the case of μ -tagged decays, the main systematic uncertainty is due to the possibility that the D^0 flavor is not tagged correctly by the muon charge because of misreconstruction. The probability of wrongly assigning the D^0 flavor (mistag) is studied with a large sample of μ -tagged $D^0 \rightarrow K^- \pi^+$ decays by comparing the charges of kaon and muon candidates. Mistag rates are found to be at the percent level and compatible for positively and negatively tagged decays. The corresponding systematic uncertainty is estimated to be 4×10^{-4} , also taking into account the fact that wrongly tagged decays include a fraction of doubly Cabibbo-suppressed $D^0 \rightarrow K^+ \pi^-$ and mixed $D^0 \rightarrow \bar{D}^0 \rightarrow K^+ \pi^-$ decays, calculated to be 0.39% with negligible uncertainty for both the $K^+ \pi^-$ and $K^- \pi^+$ final states using input from Ref. [63].

Systematic uncertainties of 0.2×10^{-4} and 1×10^{-4} accounting for the knowledge of the weights used in the kinematic weighting procedure are assessed for π -tagged and μ -tagged decays, respectively. Although suppressed by the requirement that the D^0 trajectory points back to the PV, a fraction of D^0 mesons from B decays is still present in the final π -tagged sample. As $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays may have different levels of contamination, the value of ΔA_{CP} may be biased because of an incomplete cancellation of the production asymmetries of b hadrons. The fractions of D^0 mesons from B decays are estimated by performing a fit to the distribution of the D^0 -candidate impact parameter in the plane transverse to the beam direction [60]. The corresponding systematic uncertainty is estimated to be 0.3×10^{-4} . A systematic uncertainty associated to the presence of background components peaking in $m(D^0 \pi)$ and not in $m(D^0)$ is determined by fits to the $m(D^0)$ distributions [60], where these components are modeled using fast simulation [64]. The main sources are the $D^0 \rightarrow K^- \pi^+ \pi^0$ decay for the $K^+ K^-$ final state, and the $D^0 \rightarrow \pi^- \mu^+ \nu_\mu$ and $D^0 \rightarrow \pi^- e^+ \nu_e$ decays for the $\pi^+ \pi^-$ final state. Yields and raw asymmetries of the peaking-background components measured from the fits are then used as inputs to pseudoexperiments designed to evaluate the corresponding effects on the determination of ΔA_{CP} . A value of 0.5×10^{-4} is assigned as a systematic uncertainty.

In the case of μ -tagged decays, the fractions of reconstructed $\bar{B}$ decays can be slightly different between the $K^- K^+$ and $\pi^- \pi^+$ decay modes, which could lead to a small bias

Table 1: Systematic uncertainties on ΔA_{CP} for π - and μ -tagged decays (in 10^{-4}). The total uncertainties are obtained as the sums in quadrature of the individual contributions.

Source	π -tagged	μ -tagged
Fit model	0.6	2
Mistag	–	4
Weighting	0.2	1
Secondary decays	0.3	–
Peaking background	0.5	–
B fractions	–	1
B reco. efficiency	–	2
Total	0.9	5

in ΔA_{CP} . Using the LHCb measurements of the b -hadron production asymmetries [50], the systematic uncertainty on ΔA_{CP} is estimated to be 1×10^{-4} . The combination of a difference in the B reconstruction efficiency as a function of the decay time between the $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ modes and the presence of neutral B -meson oscillations may also cause an imperfect cancellation of $A_P(B)$ in ΔA_{CP} . The associated systematic uncertainty is estimated to be 2×10^{-4} .

All individual contributions are summed in quadrature to give total systematic uncertainties on ΔA_{CP} of 0.9×10^{-4} and 5×10^{-4} for the π -tagged and μ -tagged samples, respectively. A summary of all systematic uncertainties is reported in Table 1. Other possible systematic uncertainties are investigated and found to be negligible.

Numerous additional robustness checks are carried out [60]. The measured value of ΔA_{CP} is studied as a function of several variables, notably including: the azimuthal angle, χ^2_{IP} , transverse momentum and pseudorapidity of π -tagged and μ -tagged D^0 mesons as well as of the tagging pions or muons; the χ^2 of the D^{*+} and B vertex fits; the track quality of the tagging pion and the charged-particle multiplicity in the event. Furthermore, the total sample is split into subsamples taken in different run periods within the years of data taking, also distinguishing different magnet polarities. No evidence for unexpected dependences of ΔA_{CP} is found in any of these tests. A check using more stringent PID requirements is performed, and all variations of ΔA_{CP} are found to be compatible within statistical uncertainties. An additional check concerns the measurement of ΔA_{bkg} , that is the difference of the background raw asymmetries in $K^- K^+$ and $\pi^- \pi^+$ final states. As the prompt background is mainly composed of genuine D^0 candidates paired with unrelated pions originating from the PV, ΔA_{bkg} is expected to be compatible with zero. A value of $\Delta A_{bkg} = (-2 \pm 4) \times 10^{-4}$ is obtained.

The difference of time-integrated CP asymmetries of $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays is measured using 13 TeV pp collision data collected with the LHCb detector and corresponding to an integrated luminosity of 5.9 fb^{-1} . The results are

$$\begin{aligned}\Delta A_{CP}^{\pi\text{-tagged}} &= [-18.2 \pm 3.2 (\text{stat.}) \pm 0.9 (\text{syst.})] \times 10^{-4}, \\ \Delta A_{CP}^{\mu\text{-tagged}} &= [-9 \pm 8 (\text{stat.}) \pm 5 (\text{syst.})] \times 10^{-4}.\end{aligned}$$

Both measurements are in good agreement with world averages [65] and previous LHCb results [42, 43].

By making a full combination with previous LHCb measurements [42, 43], the following value of ΔA_{CP} is obtained

$$\Delta A_{CP} = (-15.4 \pm 2.9) \times 10^{-4},$$

where the uncertainty includes statistical and systematic contributions. The significance of the deviation from zero corresponds to 5.3 standard deviations. This is the first observation of CP violation in the decay of charm hadrons.

The interpretation of ΔA_{CP} in terms of direct CP violation and A_Γ requires knowledge of the difference of reconstructed mean decay times for $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays normalized to the D^0 lifetime, as shown in Eq. (3). The values corresponding to the present measurements are $\Delta \langle t \rangle^{\pi^- \text{tagged}} / \tau(D^0) = 0.135 \pm 0.002$ and $\Delta \langle t \rangle^{\mu^- \text{tagged}} / \tau(D^0) = -0.003 \pm 0.001$, whereas that corresponding to the full combination is $\Delta \langle t \rangle / \tau(D^0) = 0.115 \pm 0.002$. The uncertainties include statistical and systematic contributions, and the world average of the D^0 lifetime is used [66].

By using in addition the LHCb average $A_\Gamma = (-2.8 \pm 2.8) \times 10^{-4}$ [46, 47], from Eq. (3) it is possible to derive

$$\Delta a_{CP}^{\text{dir}} = (-15.7 \pm 2.9) \times 10^{-4},$$

which shows that, as expected, ΔA_{CP} is primarily sensitive to direct CP violation. The overall improvement in precision brought by the present analysis to the knowledge of $\Delta a_{CP}^{\text{dir}}$ is apparent when comparing with the value obtained from previous measurements, $\Delta a_{CP}^{\text{dir}} = (-13.4 \pm 7.0) \times 10^{-4}$ [65].

In summary, this Letter reports the first observation of a nonzero CP asymmetry in charm decays, using large samples of $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays collected with the LHCb detector. The result is consistent with, although in magnitude at the upper end of, SM expectations, which lie in the range 10^{-4} – 10^{-3} [16–34]. In particular, the result challenges predictions based on first-principle QCD dynamics [19, 33]. It complies with predictions based on flavor-SU(3) symmetry, if one assumes a dynamical enhancement of the penguin amplitude [16, 26–30, 32]. In the next decade, further measurements with charmed particles, along with possible theoretical improvements, will help clarify the physics picture, and establish whether this result is consistent with the SM or indicates the presence of new dynamics in the up-quark sector.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities

behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom); Laboratory Directed Research and Development program of LANL (USA).

References

- [1] A. D. Sakharov, *Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe*, Pisma Zh. Eksp. Teor. Fiz. **5** (1967) 32.
- [2] N. Cabibbo, *Unitary symmetry and leptonic decays*, Phys. Rev. Lett. **10** (1963) 531.
- [3] M. Kobayashi and T. Maskawa, *CP-violation in the renormalizable theory of weak interaction*, Prog. Theor. Phys. **49** (1973) 652.
- [4] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, *Evidence for the 2π decay of the K_2^0 meson*, Phys. Rev. Lett. **13** (1964) 138.
- [5] KTeV collaboration, A. Alavi-Harati *et al.*, *Observation of direct CP violation in $K_{S,L} \rightarrow \pi\pi$ decays*, Phys. Rev. Lett. **83** (1999) 22, [arXiv:hep-ex/9905060](#).
- [6] NA48 collaboration, A. Lai *et al.*, *A precise measurement of the direct CP violation parameter $Re(\varepsilon'/\varepsilon)$* , Eur. Phys. J. **C22** (2001) 231, [arXiv:hep-ex/0110019](#).
- [7] BaBar collaboration, B. Aubert *et al.*, *Observation of CP violation in the B^0 meson system*, Phys. Rev. Lett. **87** (2001) 091801, [arXiv:hep-ex/0107013](#).
- [8] Belle collaboration, K. Abe *et al.*, *Observation of large CP violation in the neutral B meson system*, Phys. Rev. Lett. **87** (2001) 091802, [arXiv:hep-ex/0107061](#).
- [9] BaBar collaboration, B. Aubert *et al.*, *Direct CP violating asymmetry in $B^0 \rightarrow K^+\pi^-$ decays*, Phys. Rev. Lett. **93** (2004) 131801, [arXiv:hep-ex/0407057](#).
- [10] Belle collaboration, Y. Chao *et al.*, *Evidence for direct CP violation in $B^0 \rightarrow K^+\pi^-$ decays*, Phys. Rev. Lett. **93** (2004) 191802, [arXiv:hep-ex/0408100](#).
- [11] LHCb collaboration, R. Aaij *et al.*, *First observation of CP violation in the decays of B_s^0 mesons*, Phys. Rev. Lett. **110** (2013) 221601, [arXiv:1304.6173](#).
- [12] LHCb collaboration, R. Aaij *et al.*, *Observation of CP violation in $B^\pm \rightarrow DK^\pm$ decays*, Phys. Lett. **B712** (2012) 203, Erratum *ibid.* **B713** (2012) 351, [arXiv:1203.3662](#).
- [13] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, *Progress in electroweak baryogenesis*, Ann. Rev. Nucl. Part. Sci. **43** (1993) 27, [arXiv:hep-ph/9302210](#).
- [14] A. Riotto and M. Trodden, *Recent progress in baryogenesis*, Ann. Rev. Nucl. Part. Sci. **49** (1999) 35, [arXiv:hep-ph/9901362](#).

- [15] W.-S. Hou, *Source of CP violation for the baryon asymmetry of the universe*, Chin. J. Phys. **47** (2009) 134, [arXiv:0803.1234](#).
- [16] M. Golden and B. Grinstein, *Enhanced CP violations in hadronic charm decays*, Phys. Lett. **B222** (1989) 501.
- [17] F. Buccella *et al.*, *Nonleptonic weak decays of charmed mesons*, Phys. Rev. **D51** (1995) 3478, [arXiv:hep-ph/9411286](#).
- [18] S. Bianco, F. L. Fabbri, D. Benson, and I. Bigi, *A Cicerone for the physics of charm*, Riv. Nuovo Cim. **26N7** (2003) 1, [arXiv:hep-ex/0309021](#).
- [19] Y. Grossman, A. L. Kagan, and Y. Nir, *New physics and CP violation in singly Cabibbo suppressed D decays*, Phys. Rev. **D75** (2007) 036008, [arXiv:hep-ph/0609178](#).
- [20] M. Artuso, B. Meadows, and A. A. Petrov, *Charm meson decays*, Ann. Rev. Nucl. Part. Sci. **58** (2008) 249, [arXiv:0802.2934](#).
- [21] J. Brod, A. L. Kagan, and J. Zupan, *Size of direct CP violation in singly Cabibbo-suppressed D decays*, Phys. Rev. **D86** (2012) 014023, [arXiv:1111.5000](#).
- [22] H.-Y. Cheng and C.-W. Chiang, *Direct CP violation in two-body hadronic charmed meson decays*, Phys. Rev. **D85** (2012) 034036, Erratum *ibid.* **D85** (2012) 079903, [arXiv:1201.0785](#).
- [23] H.-Y. Cheng and C.-W. Chiang, *SU(3) symmetry breaking and CP violation in $D \rightarrow PP$ decays*, Phys. Rev. **D86** (2012) 014014, [arXiv:1205.0580](#).
- [24] H.-n. Li, C.-D. Lu, and F.-S. Yu, *Branching ratios and direct CP asymmetries in $D \rightarrow PP$ decays*, Phys. Rev. **D86** (2012) 036012, [arXiv:1203.3120](#).
- [25] E. Franco, S. Mishima, and L. Silvestrini, *The Standard Model confronts CP violation in $D^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^+K^-$* , JHEP **05** (2012) 140, [arXiv:1203.3131](#).
- [26] D. Pirtskhalava and P. Uttayarat, *CP Violation and flavor SU(3) breaking in D-meson decays*, Phys. Lett. **B712** (2012) 81, [arXiv:1112.5451](#).
- [27] T. Feldmann, S. Nandi, and A. Soni, *Repercussions of flavour symmetry breaking on CP violation in D-meson decays*, JHEP **06** (2012) 007, [arXiv:1202.3795](#).
- [28] J. Brod, Y. Grossman, A. L. Kagan, and J. Zupan, *A consistent picture for large penguins in $D \rightarrow \pi^+\pi^-$, K^+K^-* , JHEP **10** (2012) 161, [arXiv:1203.6659](#).
- [29] G. Hiller, M. Jung, and S. Schacht, *SU(3)-flavor anatomy of nonleptonic charm decays*, Phys. Rev. **D87** (2013) 014024, [arXiv:1211.3734](#).
- [30] Y. Grossman and D. J. Robinson, *SU(3) sum rules for charm decay*, JHEP **04** (2013) 067, [arXiv:1211.3361](#).
- [31] B. Bhattacharya, M. Gronau, and J. L. Rosner, *CP asymmetries in singly-Cabibbo-suppressed D decays to two pseudoscalar mesons*, Phys. Rev. **D85** (2012) 054014, [arXiv:1201.2351](#).

- [32] S. Müller, U. Nierste, and S. Schacht, *Sum rules of charm CP asymmetries beyond the $SU(3)_F$ limit*, Phys. Rev. Lett. **115** (2015) 251802, [arXiv:1506.04121](#).
- [33] A. Khodjamirian and A. A. Petrov, *Direct CP asymmetry in $D \rightarrow \pi^- \pi^+$ and $D \rightarrow K^- K^+$ in QCD-based approach*, Phys. Lett. **B774** (2017) 235, [arXiv:1706.07780](#).
- [34] F. Buccella, A. Paul, and P. Santorelli, *On $SU(3)_F$ breaking through final state interactions and CP asymmetries in $D \rightarrow PP$ decays*, [arXiv:1902.05564](#).
- [35] D. Atwood and A. Soni, *Searching for the origin of CP violation in Cabibbo-suppressed D-meson decays*, PTEP **2013** (2013) 093B05, [arXiv:1211.1026](#).
- [36] BaBar collaboration, B. Aubert *et al.*, *Search for CP violation in the decays $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$* , Phys. Rev. Lett. **100** (2008) 061803, [arXiv:0709.2715](#).
- [37] Belle collaboration, M. Starič *et al.*, *Search for a CP asymmetry in Cabibbo-suppressed D^0 decays*, Phys. Lett. **B670** (2008) 190, [arXiv:0807.0148](#).
- [38] CDF collaboration, T. Aaltonen *et al.*, *Measurement of CP-violating asymmetries in $D^0 \rightarrow \pi^+ \pi^-$ and $D^0 \rightarrow K^+ K^-$ decays at CDF*, Phys. Rev. **D85** (2012) 012009, [arXiv:1111.5023](#).
- [39] CDF collaboration, T. Aaltonen *et al.*, *Measurement of the difference of CP-violating asymmetries in $D^0 \rightarrow K^+ K^-$ and $D^0 \rightarrow \pi^+ \pi^-$ decays at CDF*, Phys. Rev. Lett. **109** (2012) 111801, [arXiv:1207.2158](#).
- [40] LHCb collaboration, R. Aaij *et al.*, *Evidence for CP violation in time-integrated $D^0 \rightarrow h^- h^+$ decay rates*, Phys. Rev. Lett. **108** (2012) 111602, [arXiv:1112.0938](#).
- [41] LHCb collaboration, R. Aaij *et al.*, *Search for direct CP violation in $D^0 \rightarrow h^- h^+$ modes using semileptonic B decays*, Phys. Lett. **B723** (2013) 33, [arXiv:1303.2614](#).
- [42] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP asymmetry in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays*, JHEP **07** (2014) 041, [arXiv:1405.2797](#).
- [43] LHCb collaboration, R. Aaij *et al.*, *Measurement of the difference of time-integrated CP asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays*, Phys. Rev. Lett. **116** (2016) 191601, [arXiv:1602.03160](#).
- [44] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP asymmetry in $D^0 \rightarrow K^+ K^-$ decays*, Phys. Lett. **B767** (2017) 177, [arXiv:1610.09476](#).
- [45] M. Gersabeck *et al.*, *On the interplay of direct and indirect CP violation in the charm sector*, J. Phys. **G39** (2012) 045005, [arXiv:1111.6515](#).
- [46] LHCb collaboration, R. Aaij *et al.*, *Measurement of indirect CP asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays using semileptonic B decays*, JHEP **04** (2015) 043, [arXiv:1501.06777](#).

- [47] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CP violation parameter A_Γ in $D^0 \rightarrow K^+ K^-$ and $D^0 \rightarrow \pi^+ \pi^-$ decays*, Phys. Rev. Lett. **118** (2017) 261803, [arXiv:1702.06490](#).
- [48] A. L. Kagan and M. D. Sokoloff, *Indirect CP violation and implications for D^0 - $\bar{D}^0$ and B_s - $\bar{B}_s$ mixing*, Phys. Rev. **D80** (2009) 076008, [arXiv:0907.3917](#).
- [49] D.-S. Du, *CP violation for neutral charmed meson decays to CP eigenstates*, Eur. Phys. J. **C50** (2007) 579, [arXiv:hep-ph/0608313](#).
- [50] LHCb collaboration, R. Aaij *et al.*, *Measurement of B^0 , B_s^0 , B^+ and Λ_b^0 production asymmetries in 7 and 8 TeV proton-proton collisions*, Phys. Lett. **B774** (2017) 139, [arXiv:1703.08464](#).
- [51] LHCb collaboration, R. Aaij *et al.*, *Measurement of the flavour-specific CP-violating asymmetry a_{sl}^s in B_s^0 decays*, Phys. Lett. **B728** (2014) 607, [arXiv:1308.1048](#).
- [52] LHCb collaboration, R. Aaij *et al.*, *Measurement of the $D^\pm$ production asymmetry in 7 TeV pp collisions*, Phys. Lett. **B718** (2013) 902, [arXiv:1210.4112](#).
- [53] LHCb collaboration, R. Aaij *et al.*, *Measurement of the $D_s^+ - D_s^-$ production asymmetry in 7 TeV pp collisions*, Phys. Lett. **B713** (2012) 186, [arXiv:1205.0897](#).
- [54] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [55] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [56] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [57] T. Likhomanenko *et al.*, *LHCb topological trigger reoptimization*, J. Phys. Conf. Ser. **664** (2015) 082025, [arXiv:1510.00572](#).
- [58] G. Dujany and B. Storaci, *Real-time alignment and calibration of the LHCb Detector in Run II*, J. Phys. Conf. Ser. **664** (2015) 082010.
- [59] E653 collaboration, K. Kodama *et al.*, *Measurement of the relative branching fraction $\Gamma(D^0 \rightarrow K \mu \nu)/\Gamma(D^0 \rightarrow \mu X)$* , Phys. Rev. Lett. **66** (1991) 1819.
- [60] See supplemental material for additional numerical values and plots.
- [61] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, [arXiv:physics/0503191](#).
- [62] N. L. Johnson, *Systems of frequency curves generated by methods of translation*, Biometrika **36** (1949) 149.
- [63] LHCb collaboration, R. Aaij *et al.*, *Updated determination of D^0 - $\bar{D}^0$ mixing and CP violation parameters with $D^0 \rightarrow K^+ \pi^-$ decays*, Phys. Rev. **D97** (2018) 031101, [arXiv:1712.03220](#).

- [64] G. A. Cowan, D. C. Craik, and M. D. Needham, *RapidSim: An application for the fast simulation of heavy-quark hadron decays*, Comput. Phys. Commun. **214** (2017) 239, [arXiv:1612.07489](#).
- [65] Heavy Flavor Averaging Group, Y. Amhis *et al.*, *Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2016*, Eur. Phys. J. **C77** (2017) 895, [arXiv:1612.07233](#), updated results and plots available at <https://hflav.web.cern.ch>.
- [66] Particle Data Group, M. Tanabashi *et al.*, *Review of particle physics*, Phys. Rev. **D98** (2018) 030001.

LHCb collaboration

R. Aaij²⁸, C. Abellán Beteta⁴⁶, B. Adeva⁴³, M. Adinolfi⁵⁰, C.A. Aidala⁷⁸, Z. Ajaltouni⁶,
 S. Akar⁶¹, P. Albicocco¹⁹, J. Albrecht¹¹, F. Alessio⁴⁴, M. Alexander⁵⁵, A. Alfonso Alberio⁴²,
 G. Alkhazov⁴¹, P. Alvarez Cartelle⁵⁷, A.A. Alves Jr⁴³, S. Amato², Y. Amhis⁸, L. An¹⁸,
 L. Anderlini¹⁸, G. Andreassi⁴⁵, M. Andreotti¹⁷, J.E. Andrews⁶², F. Archilli²⁸, P. d'Argent¹³,
 J. Arnau Romeu⁷, A. Artamonov⁴⁰, M. Artuso⁶³, K. Arzymatov³⁷, E. Aslanides⁷, M. Atzeni⁴⁶,
 B. Audurier²³, S. Bachmann¹³, J.J. Back⁵², S. Baker⁵⁷, V. Balagura^{8,b}, W. Baldini^{17,44},
 A. Baranov³⁷, R.J. Barlow⁵⁸, S. Barsuk⁸, W. Barter⁵⁷, M. Bartolini²⁰, F. Baryshnikov⁷⁴,
 V. Batozskaya³², B. Batsukh⁶³, A. Battig¹¹, V. Battista⁴⁵, A. Bay⁴⁵, F. Bedeschi²⁵, I. Bediaga¹,
 A. Beiter⁶³, L.J. Bel²⁸, S. Belin²³, N. Beliy⁶⁶, V. Bellec⁴⁵, N. Belloli^{21,i}, K. Belous⁴⁰,
 I. Belyaev³⁴, E. Ben-Haim⁹, G. Bencivenni¹⁹, S. Benson²⁸, S. Beranek¹⁰, A. Berezhnoff³⁵,
 R. Bernet⁴⁶, D. Berninghoff¹³, E. Bertholet⁹, A. Bertolin²⁴, C. Betancourt⁴⁶, F. Betti^{16,e},
 M.O. Bettler⁵¹, M. van Beuzekom²⁸, I. Bezshyiko⁴⁶, S. Bhasin⁵⁰, J. Bhom³⁰, M.S. Bieker¹¹,
 S. Bifani⁴⁹, P. Billoir⁹, A. Birnkraut¹¹, A. Bizzeti^{18,u}, M. Bjørn⁵⁹, M.P. Blago⁴⁴, T. Blake⁵²,
 F. Blanc⁴⁵, S. Blusk⁶³, D. Bobulska⁵⁵, V. Bocci²⁷, O. Boente Garcia⁴³, T. Boettcher⁶⁰,
 A. Bondar^{39,x}, N. Bondar⁴¹, S. Borghi^{58,44}, M. Borisyak³⁷, M. Borsato¹³, M. Boubdir¹⁰,
 T.J.V. Bowcock⁵⁶, C. Bozzi^{17,44}, S. Braun¹³, M. Brodski⁴⁴, J. Brodzicka³⁰, A. Brossa Gonzalo⁵²,
 D. Brundu^{23,44}, E. Buchanan⁵⁰, A. Buonauro⁴⁶, C. Burr⁵⁸, A. Bursche²³, J. Buytaert⁴⁴,
 W. Byczynski⁴⁴, S. Cadeddu²³, H. Cai⁶⁸, R. Calabrese^{17,g}, S. Cali¹⁷, R. Calladine⁴⁹,
 M. Calvi^{21,i}, M. Calvo Gomez^{42,m}, A. Camboni^{42,m}, P. Campana¹⁹, D.H. Campora Perez⁴⁴,
 L. Capriotti^{16,e}, A. Carbone^{16,e}, G. Carboni²⁶, R. Cardinale²⁰, A. Cardini²³, P. Carniti^{21,i},
 K. Carvalho Akiba², G. Casse⁵⁶, M. Cattaneo⁴⁴, G. Cavallero²⁰, R. Cenci^{25,p}, M.G. Chapman⁵⁰,
 M. Charles^{9,44}, Ph. Charpentier⁴⁴, G. Chatzikonstantinidis⁴⁹, M. Chefdeville⁵, V. Chekalina³⁷,
 C. Chen³, S. Chen²³, S.-G. Chitic⁴⁴, V. Chobanova⁴³, M. Chruszcz⁴⁴, A. Chubykin⁴¹,
 P. Ciambrone¹⁹, X. Cid Vidal⁴³, G. Ciezarek⁴⁴, F. Cindolo¹⁶, P.E.L. Clarke⁵⁴, M. Clemencic⁴⁴,
 H.V. Cliff⁵¹, J. Closier⁴⁴, V. Coco⁴⁴, J.A.B. Coelho⁸, J. Cogan⁷, E. Cogneras⁶, L. Cojocariu³³,
 P. Collins⁴⁴, T. Colombo⁴⁴, A. Comerma-Montells¹³, A. Contu²³, G. Coombs⁴⁴, S. Coquereau⁴²,
 G. Corti⁴⁴, C.M. Costa Sobral⁵², B. Couturier⁴⁴, G.A. Cowan⁵⁴, D.C. Craik⁶⁰, A. Crocombe⁵²,
 M. Cruz Torres¹, R. Currie⁵⁴, C. D'Ambrosio⁴⁴, C.L. Da Silva⁷⁹, E. Dall'Occo²⁸, J. Dalseno^{43,v},
 A. Danilina³⁴, A. Davis⁵⁸, O. De Aguiar Francisco⁴⁴, K. De Bruyn⁴⁴, S. De Capua⁵⁸,
 M. De Cian⁴⁵, J.M. De Miranda¹, L. De Paula², M. De Serio^{15,d}, P. De Simone¹⁹, C.T. Dean⁵⁵,
 W. Dean⁷⁸, D. Decamp⁵, L. Del Buono⁹, B. Delaney⁵¹, H.-P. Dembinski¹², M. Demmer¹¹,
 A. Dendek³¹, D. Derkach³⁸, O. Deschamps⁶, F. Dese⁸, F. Dettori²³, B. Dey⁶⁹, A. Di Canto⁴⁴,
 P. Di Nezza¹⁹, S. Didenko⁷⁴, H. Dijkstra⁴⁴, F. Dordei²³, M. Dorigo^{25,y}, A. Dosil Suárez⁴³,
 L. Douglas⁵⁵, A. Dovbnya⁴⁷, K. Dreimanis⁵⁶, L. Dufour⁴⁴, G. Dujany⁹, P. Durante⁴⁴,
 J.M. Durham⁷⁹, D. Dutta⁵⁸, R. Dzhelezhyan^{40,†}, M. Dziewiecki¹³, A. Dziurda³⁰, A. Dzyuba⁴¹,
 S. Easo⁵³, U. Egede⁵⁷, V. Egorychev³⁴, S. Eidelman^{39,x}, S. Eisenhardt⁵⁴, U. Eitschberger¹¹,
 R. Ekelhof¹¹, L. Eklund⁵⁵, S. Ely⁶³, A. Ene³³, S. Escher¹⁰, S. Esen²⁸, T. Evans⁶¹, A. Falabella¹⁶,
 N. Farley⁴⁹, S. Farry⁵⁶, D. Fazzini^{21,i}, P. Fernandez Declara⁴⁴, A. Fernandez Prieto⁴³,
 F. Ferrari^{16,e}, L. Ferreira Lopes⁴⁵, F. Ferreira Rodrigues², S. Ferreres Sole²⁸, M. Ferro-Luzzi⁴⁴,
 S. Filippov³⁶, R.A. Fini¹⁵, M. Fiorini^{17,g}, M. Firlej³¹, C. Fitzpatrick⁴⁴, T. Fiutowski³¹,
 F. Fleuret^{8,b}, M. Fontana⁴⁴, F. Fontanelli^{20,h}, R. Forty⁴⁴, V. Franco Lima⁵⁶, M. Frank⁴⁴,
 C. Frei⁴⁴, J. Fu^{22,q}, W. Funk⁴⁴, C. Färber⁴⁴, M. Féo⁴⁴, E. Gabriel⁵⁴, A. Gallas Torreira⁴³,
 D. Galli^{16,e}, S. Gallorini²⁴, S. Gambetta⁵⁴, Y. Gan³, M. Gandelman², P. Gandini²², Y. Gao³,
 L.M. Garcia Martin⁷⁶, B. Garcia Plana⁴³, J. García Pardiñas⁴⁶, J. Garra Tico⁵¹, L. Garrido⁴²,
 D. Gascon⁴², C. Gaspar⁴⁴, G. Gazzoni⁶, D. Gerick¹³, E. Gersabeck⁵⁸, M. Gersabeck⁵⁸,
 T. Gershon⁵², D. Gerstel⁷, Ph. Ghez⁵, V. Gibson⁵¹, O.G. Girard⁴⁵, P. Gironella Gironell⁴²,
 L. Giubega³³, K. Gizdov⁵⁴, V.V. Gligorov⁹, D. Golubkov³⁴, A. Golutvin^{57,74}, A. Gomes^{1,a},
 I.V. Gorelov³⁵, C. Gotti^{21,i}, E. Govorkova²⁸, J.P. Grabowski¹³, R. Graciani Diaz⁴²,

L.A. Granado Cardoso⁴⁴, E. Graugés⁴², E. Graverini⁴⁶, G. Graziani¹⁸, A. Grecu³³, R. Greim²⁸,
 P. Griffith²³, L. Grillo⁵⁸, L. Gruber⁴⁴, B.R. Gruber Cazon⁵⁹, C. Gu³, X. Guo⁶⁷, E. Gushchin³⁶,
 A. Guth¹⁰, Yu. Guz^{40,44}, T. Gys⁴⁴, C. Göbel⁶⁵, T. Hadavizadeh⁵⁹, C. Hadjivasiliou⁶,
 G. Haefeli⁴⁵, C. Haen⁴⁴, S.C. Haines⁵¹, B. Hamilton⁶², Q. Han⁶⁹, X. Han¹³, T.H. Hancock⁵⁹,
 S. Hansmann-Menzemer¹³, N. Harnew⁵⁹, T. Harrison⁵⁶, C. Hasse⁴⁴, M. Hatch⁴⁴, J. He⁶⁶,
 M. Hecker⁵⁷, K. Heinicke¹¹, A. Heister¹¹, K. Hennessy⁵⁶, L. Henry⁷⁶, E. van Herwijnen⁴⁴,
 J. Heuel¹⁰, M. Heß⁷¹, A. Hicheur⁶⁴, R. Hidalgo Charman⁵⁸, D. Hill⁵⁹, M. Hilton⁵⁸,
 P.H. Hopchev⁴⁵, J. Hu¹³, W. Hu⁶⁹, W. Huang⁶⁶, Z.C. Huard⁶¹, W. Hulsbergen²⁸, T. Humair⁵⁷,
 M. Hushchyn³⁸, D. Hutchcroft⁵⁶, D. Hynds²⁸, P. Ibis¹¹, M. Idzik³¹, P. Ilten⁴⁹, A. Inglessi⁴¹,
 A. Inyakin⁴⁰, K. Ivshin⁴¹, R. Jacobsson⁴⁴, S. Jakobsen⁴⁴, J. Jalocha⁵⁹, E. Jans²⁸, B.K. Jashal⁷⁶,
 A. Jawahery⁶², F. Jiang³, M. John⁵⁹, D. Johnson⁴⁴, C.R. Jones⁵¹, C. Joram⁴⁴, B. Jost⁴⁴,
 N. Jurik⁵⁹, S. Kandybei⁴⁷, M. Karacson⁴⁴, J.M. Kariuki⁵⁰, S. Karodia⁵⁵, N. Kazeev³⁸,
 M. Kecke¹³, F. Keizer⁵¹, M. Kelsey⁶³, M. Kenzie⁵¹, T. Ketel²⁹, B. Khanji⁴⁴, A. Kharisova⁷⁵,
 C. Khurewathanakul⁴⁵, K.E. Kim⁶³, T. Kirn¹⁰, V.S. Kirsebom⁴⁵, S. Klaver¹⁹,
 K. Klimaszewski³², S. Koliiev⁴⁸, M. Kolpin¹³, R. Kopečna¹³, P. Koppenburg²⁸, I. Kostiuik^{28,48},
 S. Kotriakhova⁴¹, M. Kozeiha⁶, L. Kravchuk³⁶, M. Kreps⁵², F. Kress⁵⁷, S. Kretzschmar¹⁰,
 P. Krokovny^{39,x}, W. Krupa³¹, W. Krzemien³², W. Kucewicz^{30,l}, M. Kucharczyk³⁰,
 V. Kudryavtsev^{39,x}, G.J. Kunde⁷⁹, A.K. Kuonen⁴⁵, T. Kvaratskheliya³⁴, D. Lacarrere⁴⁴,
 G. Lafferty⁵⁸, A. Lai²³, D. Lancierini⁴⁶, G. Lanfranchi¹⁹, C. Langenbruch¹⁰, T. Latham⁵²,
 C. Lazzeroni⁴⁹, R. Le Gac⁷, A. Leflat³⁵, R. Lefèvre⁶, F. Lemaitre⁴⁴, O. Leroy⁷, T. Lesiak³⁰,
 B. Leverington¹³, H. Li⁶⁷, P.-R. Li^{66,ab}, X. Li⁷⁹, Y. Li⁴, Z. Li⁶³, X. Liang⁶³, T. Likhomanenko⁷³,
 R. Lindner⁴⁴, P. Ling⁶⁷, F. Lionetto⁴⁶, V. Lisovsky⁸, G. Liu⁶⁷, X. Liu³, D. Loh⁵², A. Loi²³,
 I. Longstaff⁵⁵, J.H. Lopes², G. Loustau⁴⁶, G.H. Lovell⁵¹, D. Lucchesi^{24,o}, M. Lucio Martinez⁴³,
 Y. Luo³, A. Lupato²⁴, E. Luppi^{17,g}, O. Lupton⁵², A. Lusiani²⁵, X. Lyu⁶⁶, R. Ma⁶⁷,
 F. Machefert⁸, F. Maciuc³³, V. Macko⁴⁵, P. Mackowiak¹¹, S. Maddrell-Mander⁵⁰, O. Maev^{41,44},
 K. Maguire⁵⁸, D. Maisuzenko⁴¹, M.W. Majewski³¹, S. Malde⁵⁹, B. Malecki⁴⁴, A. Malinin⁷³,
 T. Maltsev^{39,x}, H. Malygina¹³, G. Manca^{23,f}, G. Mancinelli⁷, D. Marangotto^{22,q}, J. Maratas^{6,w},
 J.F. Marchand⁵, U. Marconi¹⁶, C. Marin Benito⁸, M. Marinangeli⁴⁵, P. Marino⁴⁵, J. Marks¹³,
 P.J. Marshall⁵⁶, G. Martellotti²⁷, M. Martinelli^{44,21}, D. Martinez Santos⁴³, F. Martinez Vidal⁷⁶,
 A. Massafferri¹, M. Materok¹⁰, R. Matev⁴⁴, A. Mathad⁴⁶, Z. Mathe⁴⁴, V. Matiunin³⁴,
 C. Matteuzzi²¹, K.R. Mattioli⁷⁸, A. Mauri⁴⁶, E. Maurice^{8,b}, B. Maurin⁴⁵, M. McCann^{57,44},
 A. McNab⁵⁸, R. McNulty¹⁴, J.V. Mead⁵⁶, B. Meadows⁶¹, C. Meaux⁷, N. Meinert⁷¹,
 D. Melnychuk³², M. Merk²⁸, A. Merli^{22,q}, E. Michielin²⁴, D.A. Milanese⁷⁰, E. Millard⁵²,
 M.-N. Minard⁵, L. Minzoni^{17,g}, D.S. Mitzel¹³, A. Mogini⁹, R.D. Moise⁵⁷, T. Mombächer¹¹,
 I.A. Monroy⁷⁰, S. Monteil⁶, M. Morandin²⁴, G. Morello¹⁹, M.J. Morello^{25,t}, J. Moron³¹,
 A.B. Morris⁷, R. Mountain⁶³, F. Muheim⁵⁴, M. Mukherjee⁶⁹, M. Mulder²⁸, C.H. Murphy⁵⁹,
 D. Murray⁵⁸, A. Mödden¹¹, D. Müller⁴⁴, J. Müller¹¹, K. Müller⁴⁶, V. Müller¹¹, P. Naik⁵⁰,
 T. Nakada⁴⁵, R. Nandakumar⁵³, A. Nandi⁵⁹, T. Nanut⁴⁵, I. Nasteva², M. Needham⁵⁴,
 N. Neri^{22,q}, S. Neubert¹³, N. Neufeld⁴⁴, R. Newcombe⁵⁷, T.D. Nguyen⁴⁵, C. Nguyen-Mau^{45,n},
 S. Nieswand¹⁰, R. Niet¹¹, N. Nikitin³⁵, N.S. Nolte⁴⁴, D.P. O'Hanlon¹⁶, A. Oblakowska-Mucha³¹,
 V. Obraztsov⁴⁰, S. Ogilvy⁵⁵, R. Oldeman^{23,f}, C.J.G. Onderwater⁷², J. D. Osborn⁷⁸,
 A. Ossowska³⁰, J.M. Otalora Goicochea², T. Ovsiannikova³⁴, P. Owen⁴⁶, A. Oyanguren⁷⁶,
 P.R. Pais⁴⁵, T. Pajero^{25,t}, A. Palano¹⁵, M. Palutan¹⁹, G. Panshin⁷⁵, A. Papanestis⁵³,
 M. Pappagallo⁵⁴, L.L. Pappalardo^{17,g}, W. Parker⁶², C. Parkes^{58,44}, G. Passaleva^{18,44},
 A. Pastore¹⁵, M. Patel⁵⁷, C. Patrignani^{16,e}, A. Pearce⁴⁴, A. Pellegrino²⁸, G. Penso²⁷,
 M. Pepe Altarelli⁴⁴, S. Perazzini¹⁶, D. Pereima³⁴, P. Perret⁶, L. Pescatore⁴⁵, K. Petridis⁵⁰,
 A. Petrolini^{20,h}, A. Petrov⁷³, S. Petrucci⁵⁴, M. Petruzzo^{22,q}, B. Pietrzyk⁵, G. Pietrzyk⁴⁵,
 M. Pikić³⁰, M. Pili⁵⁹, D. Pinci²⁷, J. Pinzino⁴⁴, F. Pisani⁴⁴, A. Piućci¹³, V. Placinta³³,
 S. Playfer⁵⁴, J. Plews⁴⁹, M. Plo Casasus⁴³, F. Polci⁹, M. Poli Lener¹⁹, M. Poliakov⁶³,
 A. Poluektov⁷, N. Polukhina^{74,c}, I. Polyakov⁶³, E. Polycarpo², G.J. Pomery⁵⁰, S. Ponce⁴⁴,

A. Popov⁴⁰, D. Popov^{49,12}, S. Poslavskii⁴⁰, E. Price⁵⁰, C. Prouve⁴³, V. Pugatch⁴⁸,
A. Puig Navarro⁴⁶, H. Pullen⁵⁹, G. Punzi^{25,p}, W. Qian⁶⁶, J. Qin⁶⁶, R. Quagliani⁹, B. Quintana⁶,
N.V. Raab¹⁴, B. Rachwal³¹, J.H. Rademacker⁵⁰, M. Rama²⁵, M. Ramos Pernas⁴³, M.S. Rangel²,
F. Ratnikov^{37,38}, G. Raven²⁹, M. Ravonel Salzgeber⁴⁴, M. Reboud⁵, F. Redi⁴⁵, S. Reichert¹¹,
A.C. dos Reis¹, F. Reiss⁹, C. Remon Alepuz⁷⁶, Z. Ren³, V. Renaudin⁵⁹, S. Ricciardi⁵³,
S. Richards⁵⁰, K. Rinnert⁵⁶, P. Robbe⁸, A. Robert⁹, A.B. Rodrigues⁴⁵, E. Rodrigues⁶¹,
J.A. Rodriguez Lopez⁷⁰, M. Roehrken⁴⁴, S. Roiser⁴⁴, A. Rollings⁵⁹, V. Romanovski⁴⁰,
A. Romero Vidal⁴³, J.D. Roth⁷⁸, M. Rotondo¹⁹, M.S. Rudolph⁶³, T. Ruf⁴⁴, J. Ruiz Vidal⁷⁶,
J.J. Saborido Silva⁴³, N. Sagidova⁴¹, B. Saitta^{23,f}, V. Salustino Guimaraes⁶⁵, C. Sanchez Gras²⁸,
C. Sanchez Mayordomo⁷⁶, B. Sanmartin Sedes⁴³, R. Santacesaria²⁷, C. Santamarina Rios⁴³,
M. Santimaria^{19,44}, E. Santovetti^{26,j}, G. Sarpis⁵⁸, A. Sarti^{19,k}, C. Satriano^{27,s}, A. Satta²⁶,
M. Saur⁶⁶, D. Savrina^{34,35}, S. Schael¹⁰, M. Schellenberg¹¹, M. Schiller⁵⁵, H. Schindler⁴⁴,
M. Schmelling¹², T. Schmelzer¹¹, B. Schmidt⁴⁴, O. Schneider⁴⁵, A. Schopper⁴⁴, H.F. Schreiner⁶¹,
M. Schubiger⁴⁵, S. Schulte⁴⁵, M.H. Schune⁸, R. Schwemmer⁴⁴, B. Sciascia¹⁹, A. Sciubba^{27,k},
A. Semennikov³⁴, E.S. Sepulveda⁹, A. Sergi^{49,44}, N. Serra⁴⁶, J. Serrano⁷, L. Sestini²⁴,
A. Seuthe¹¹, P. Seyfert⁴⁴, M. Shapkin⁴⁰, T. Shears⁵⁶, L. Shekhtman^{39,x}, V. Shevchenko⁷³,
E. Shmanin⁷⁴, B.G. Siddi¹⁷, R. Silva Coutinho⁴⁶, L. Silva de Oliveira², G. Simi^{24,o},
S. Simone^{15,d}, I. Skiba¹⁷, N. Skidmore¹³, T. Skwarnicki⁶³, M.W. Slater⁴⁹, J.G. Smeaton⁵¹,
E. Smith¹⁰, I.T. Smith⁵⁴, M. Smith⁵⁷, M. Soares¹⁶, I. Soares Lavra¹, M.D. Sokoloff⁶¹,
F.J.P. Soler⁵⁵, B. Souza De Paula², B. Spaan¹¹, E. Spadaro Norella^{22,q}, P. Spradlin⁵⁵,
F. Stagni⁴⁴, M. Stahl¹³, S. Stahl⁴⁴, P. Stefko⁴⁵, S. Stefkova⁵⁷, O. Steinkamp⁴⁶, S. Stemmler¹³,
O. Stenyakin⁴⁰, M. Stepanova⁴¹, H. Stevens¹¹, A. Stocchi⁸, S. Stone⁶³, S. Stracka²⁵,
M.E. Stramaglia⁴⁵, M. Straticiu³³, U. Straumann⁴⁶, S. Strokov⁷⁵, J. Sun³, L. Sun⁶⁸, Y. Sun⁶²,
K. Swientek³¹, A. Szabelski³², T. Szumlak³¹, M. Szymanski⁶⁶, S. T'Jampens⁵, Z. Tang³,
T. Tekampe¹¹, G. Tellarini¹⁷, F. Teubert⁴⁴, E. Thomas⁴⁴, J. van Tilburg²⁸, M.J. Tilley⁵⁷,
V. Tisserand⁶, M. Tobin⁴, S. Tolk⁴⁴, L. Tomassetti^{17,g}, D. Tonelli²⁵, D.Y. Tou⁹,
R. Tourinho Jadallah Aoude¹, E. Tournefier⁵, M. Traill⁵⁵, M.T. Tran⁴⁵, A. Trisovic⁵¹,
A. Tsaregorodtsev⁷, G. Tuci^{25,44,p}, A. Tully⁵¹, N. Tuning²⁸, A. Ukleja³², A. Usachov⁸,
A. Ustyuzhanin^{37,38}, U. Uwer¹³, A. Vagner⁷⁵, V. Vagnoni¹⁶, A. Valassi⁴⁴, S. Valat⁴⁴,
G. Valenti¹⁶, H. Van Hecke⁷⁹, C.B. Van Hulse¹⁴, R. Vazquez Gomez⁴⁴, P. Vazquez Regueiro⁴³,
S. Vecchi¹⁷, M. van Veghel²⁸, J.J. Velthuis⁵⁰, M. Veltri^{18,r}, A. Venkateswaran⁶³, M. Vernet⁶,
M. Veronesi²⁸, M. Vesterinen⁵², J.V. Viana Barbosa⁴⁴, D. Vieira⁶⁶, M. Vieites Diaz⁴³,
H. Viemann⁷¹, X. Vilasis-Cardona^{42,m}, A. Vitkovskiy²⁸, M. Vitti⁵¹, V. Volkov³⁵, A. Vollhardt⁴⁶,
D. Vom Bruch⁹, B. Voneki⁴⁴, A. Vorobyev⁴¹, V. Vorobyev^{39,x}, N. Voropaev⁴¹, J.A. de Vries²⁸,
C. Vázquez Sierra²⁸, R. Waldi⁷¹, J. Walsh²⁵, J. Wang⁴, M. Wang³, Y. Wang⁶⁹, Z. Wang⁴⁶,
D.R. Ward⁵¹, H.M. Wark⁵⁶, N.K. Watson⁴⁹, D. Websdale⁵⁷, A. Weiden⁴⁶, C. Weisser⁶⁰,
M. Whitehead¹⁰, G. Wilkinson⁵⁹, M. Wilkinson⁶³, I. Williams⁵¹, M.R.J. Williams⁵⁸,
M. Williams⁶⁰, T. Williams⁴⁹, F.F. Wilson⁵³, M. Winn⁸, W. Wislicki³², M. Witek³⁰,
G. Wormser⁸, S.A. Wotton⁵¹, K. Wyllie⁴⁴, D. Xiao⁶⁹, Y. Xie⁶⁹, H. Xing⁶⁷, A. Xu³, M. Xu⁶⁹,
Q. Xu⁶⁶, Z. Xu³, Z. Xu⁵, Z. Yang³, Z. Yang⁶², Y. Yao⁶³, L.E. Yeomans⁵⁶, H. Yin⁶⁹, J. Yu^{69,aa},
X. Yuan⁶³, O. Yushchenko⁴⁰, K.A. Zarebski⁴⁹, M. Zavertyaev^{12,c}, M. Zeng³, D. Zhang⁶⁹,
L. Zhang³, W.C. Zhang^{3,z}, Y. Zhang⁴⁴, A. Zhelezov¹³, Y. Zheng⁶⁶, X. Zhu³, V. Zhukov^{10,35},
J.B. Zonneveld⁵⁴, S. Zucchelli^{16,e}.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴Institute Of High Energy Physics (ihep), Beijing, China

⁵Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁶Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁷Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

- ⁸LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France
- ⁹LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France
- ¹⁰I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany
- ¹¹Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany
- ¹²Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany
- ¹³Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ¹⁴School of Physics, University College Dublin, Dublin, Ireland
- ¹⁵INFN Sezione di Bari, Bari, Italy
- ¹⁶INFN Sezione di Bologna, Bologna, Italy
- ¹⁷INFN Sezione di Ferrara, Ferrara, Italy
- ¹⁸INFN Sezione di Firenze, Firenze, Italy
- ¹⁹INFN Laboratori Nazionali di Frascati, Frascati, Italy
- ²⁰INFN Sezione di Genova, Genova, Italy
- ²¹INFN Sezione di Milano-Bicocca, Milano, Italy
- ²²INFN Sezione di Milano, Milano, Italy
- ²³INFN Sezione di Cagliari, Monserrato, Italy
- ²⁴INFN Sezione di Padova, Padova, Italy
- ²⁵INFN Sezione di Pisa, Pisa, Italy
- ²⁶INFN Sezione di Roma Tor Vergata, Roma, Italy
- ²⁷INFN Sezione di Roma La Sapienza, Roma, Italy
- ²⁸Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands
- ²⁹Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands
- ³⁰Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland
- ³¹AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland
- ³²National Center for Nuclear Research (NCBJ), Warsaw, Poland
- ³³Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania
- ³⁴Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia, Moscow, Russia
- ³⁵Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia
- ³⁶Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia
- ³⁷Yandex School of Data Analysis, Moscow, Russia
- ³⁸National Research University Higher School of Economics, Moscow, Russia
- ³⁹Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia
- ⁴⁰Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia
- ⁴¹Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia , St.Petersburg, Russia
- ⁴²ICCUB, Universitat de Barcelona, Barcelona, Spain
- ⁴³Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain
- ⁴⁴European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ⁴⁵Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
- ⁴⁶Physik-Institut, Universität Zürich, Zürich, Switzerland
- ⁴⁷NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
- ⁴⁸Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
- ⁴⁹University of Birmingham, Birmingham, United Kingdom
- ⁵⁰H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
- ⁵¹Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- ⁵²Department of Physics, University of Warwick, Coventry, United Kingdom
- ⁵³STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
- ⁵⁴School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- ⁵⁵School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- ⁵⁶Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
- ⁵⁷Imperial College London, London, United Kingdom

- ⁵⁸ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁹ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶⁰ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶¹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶² *University of Maryland, College Park, MD, United States*
- ⁶³ *Syracuse University, Syracuse, NY, United States*
- ⁶⁴ *Laboratory of Mathematical and Subatomic Physics , Constantine, Algeria, associated to ²*
- ⁶⁵ *Pontificia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶⁶ *University of Chinese Academy of Sciences, Beijing, China, associated to ³*
- ⁶⁷ *South China Normal University, Guangzhou, China, associated to ³*
- ⁶⁸ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³*
- ⁶⁹ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³*
- ⁷⁰ *Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁹*
- ⁷¹ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹³*
- ⁷² *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ²⁸*
- ⁷³ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³⁴*
- ⁷⁴ *National University of Science and Technology “MISIS”, Moscow, Russia, associated to ³⁴*
- ⁷⁵ *National Research Tomsk Polytechnic University, Tomsk, Russia, associated to ³⁴*
- ⁷⁶ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ⁴²*
- ⁷⁷ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom, Bristol, United Kingdom*
- ⁷⁸ *University of Michigan, Ann Arbor, United States, associated to ⁶³*
- ⁷⁹ *Los Alamos National Laboratory (LANL), Los Alamos, United States, associated to ⁶³*
- ^a *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*
- ^b *Laboratoire Leprince-Ringuet, Palaiseau, France*
- ^c *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^d *Università di Bari, Bari, Italy*
- ^e *Università di Bologna, Bologna, Italy*
- ^f *Università di Cagliari, Cagliari, Italy*
- ^g *Università di Ferrara, Ferrara, Italy*
- ^h *Università di Genova, Genova, Italy*
- ⁱ *Università di Milano Bicocca, Milano, Italy*
- ^j *Università di Roma Tor Vergata, Roma, Italy*
- ^k *Università di Roma La Sapienza, Roma, Italy*
- ^l *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- ^m *LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁿ *Hanoi University of Science, Hanoi, Vietnam*
- ^o *Università di Padova, Padova, Italy*
- ^p *Università di Pisa, Pisa, Italy*
- ^q *Università degli Studi di Milano, Milano, Italy*
- ^r *Università di Urbino, Urbino, Italy*
- ^s *Università della Basilicata, Potenza, Italy*
- ^t *Scuola Normale Superiore, Pisa, Italy*
- ^u *Università di Modena e Reggio Emilia, Modena, Italy*
- ^v *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ^w *MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*
- ^x *Novosibirsk State University, Novosibirsk, Russia*
- ^y *Sezione INFN di Trieste, Trieste, Italy*
- ^z *School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China*
- ^{aa} *Physics and Micro Electronic College, Hunan University, Changsha City, China*
- ^{ab} *Lanzhou University, Lanzhou, China*

[†] *Deceased*

Supplemental material

Arithmetic average of mean decay times

The arithmetic average of the reconstructed mean decay times for $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays, $\langle t \rangle$, can be useful when interpreting the measurement of ΔA_{CP} . The values corresponding to the present measurements are $\langle t \rangle^{\pi\text{-tagged}}/\tau(D^0) = 1.74 \pm 0.10$ and $\langle t \rangle^{\mu\text{-tagged}}/\tau(D^0) = 1.21 \pm 0.01$, whereas that corresponding to the combination with previous LHCb measurements is $\langle t \rangle/\tau(D^0) = 1.71 \pm 0.10$. The uncertainties include statistical and systematic contributions, and the world average of the D^0 lifetime is used.

Additional plots




Figure 2: Raw asymmetries of the tagging pion for the π -tagged $D^0 \rightarrow K^- K^+$ sample, with the polarity of the magnet (top) up and (bottom) down. The plots on the left include only candidates with $|p_y/p_z| < 0.02$, *i.e.*, close to the horizontal plane, and the fiducial requirements used to exclude the kinematic region surrounding the beam pipe, characterized by large values of the raw asymmetry, are indicated as black lines (in addition to the forementioned requirement $|p_y/p_z| < 0.02$). The plots on the right include all candidates except those excluded by the beam-pipe fiducial requirements, and the black lines indicate the fiducial requirements used to exclude regions at the boundary of the detector acceptance, which are also characterized by large values of the raw asymmetry. Distributions for the $D^0 \rightarrow \pi^- \pi^+$ sample are very similar.




Figure 3: Background-subtracted distributions of momentum (p), transverse momentum (p_T), azimuthal angle (φ) and pseudorapidity (η) of D^{*+} mesons for the prompt sample: (left column) before and (right column) after the weighting procedure for $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays, as indicated in the legends. The distributions are normalized to unit area.




Figure 4: Background-subtracted distributions of momentum (p), transverse momentum (p_T), azimuthal angle (φ) and pseudorapidity (η) of D^0 mesons for the semileptonic sample: (left column) before and (right column) after the weighting procedure for $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays, as indicated in the legends. The distributions are normalized to unit area.




Figure 5: Raw asymmetries in the π -tagged sample for (left) $D^0 \rightarrow K^- K^+$ and (right) $D^0 \rightarrow \pi^- \pi^+$ candidates as a function of $m(D^0 \pi^+)$.




Figure 6: Raw asymmetries in the μ -tagged sample for (left) $D^0 \rightarrow K^- K^+$ and (right) $D^0 \rightarrow \pi^- \pi^+$ candidates as a function of $m(D^0)$.




Figure 7: Distributions of the signed D^0 impact parameter in the plane transverse to the beam direction, TIP, in bins of reconstructed D^0 decay time, for the π -tagged $D^0 \rightarrow K^- K^+$ sample: (top left) $\tilde{t} < 1.5$, (top right) $1.5 < \tilde{t} < 3.0$, (bottom left) $3.0 < \tilde{t} < 4.5$ and (bottom right) $\tilde{t} > 4.5$, where $\tilde{t} \equiv t/\tau(D^0)$. The fit results are overlaid and the contributions from prompt and secondary decays are shown, as indicated in the legends. Distributions for the $D^0 \rightarrow \pi^- \pi^+$ sample are very similar.




Figure 8: Invariant-mass distributions of (top) $D^0 \rightarrow K^- K^+$ and (bottom) $D^0 \rightarrow \pi^- \pi^+$ candidates in the prompt sample with fit results overlaid. These fits are used to determine the yields and raw asymmetries of (top) $D^0 \rightarrow K^- \pi^+ \pi^0$ and (bottom) $D^0 \rightarrow \pi^- l^+ \nu_l$ backgrounds, whose mass shapes extend to the D^0 -signal mass region. The various components included in the fit model are indicated in the legends.




Figure 9: Measurements of ΔA_{CP} in time-ordered data-taking subsamples (referred to as run blocks) for (left) prompt and (right) semileptonic samples. The uncertainties are statistical only. The horizontal red-dashed lines show the central values of the nominal results.




Figure 10: Measurements of ΔA_{CP} in bins of D^0 decay time for (left) prompt and (right) semileptonic samples. In each plot, the last bin on the right also includes a few overflow candidates. The uncertainties are statistical only. The horizontal red-dashed lines show the central values of the nominal results.




Figure 11: Measurements of ΔA_{CP} in bins of (top left) pseudorapidity, (top right) azimuthal angle, (bottom left) transverse momentum and (bottom right) χ^2_{IP} of tagging pions for the prompt sample. In each plot but that of the azimuthal angle, the last bin on the right also includes a few overflow candidates. The uncertainties are statistical only. The horizontal red-dashed lines show the central value of the nominal result.




Figure 12: Measurements of ΔA_{CP} in bins of (top left) pseudorapidity, (top right) azimuthal angle, (bottom left) transverse momentum and (bottom right) χ^2_{IP} of tagging muons for the semileptonic sample. In each plot but that of the azimuthal angle, the last bin on the right also includes a few overflow candidates. The uncertainties are statistical only. The horizontal red-dashed lines show the central value of the nominal result.




Figure 13: Measurements of ΔA_{CP} in bins of $D^0 \chi^2_{IP}$ for (left) prompt and (right) semileptonic samples. In each plot, the last bin on the right also includes a few overflow candidates. The uncertainties are statistical only. The horizontal red-dashed lines show the central values of the nominal results.