



**HAL**  
open science

## Gut Microbiota Modifies Olfactory-Guided Microbial Preferences and Foraging Decisions in *Drosophila*

Adam Chun-Nin Wong, Qiao-Ping Wang, Juliano Morimoto, Alistair M. Senior, Mathieu Lihoreau, G. Gregory Neely, Stephen J. Simpson, Fleur Ponton

► **To cite this version:**

Adam Chun-Nin Wong, Qiao-Ping Wang, Juliano Morimoto, Alistair M. Senior, Mathieu Lihoreau, et al.. Gut Microbiota Modifies Olfactory-Guided Microbial Preferences and Foraging Decisions in *Drosophila*. *Current Biology - CB*, 2017, 27 (15), pp.2397-2404.e4. 10.1016/j.cub.2017.07.022 . hal-02105100

**HAL Id: hal-02105100**

**<https://hal.science/hal-02105100>**

Submitted on 22 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Gut Microbiota Modifies Olfactory-Guided Microbial Preferences and Foraging Decisions in *Drosophila*

Adam Chun-Nin Wong,<sup>1,6,7,\*</sup> Qiao-Ping Wang,<sup>1</sup> Juliano Morimoto,<sup>1,2</sup> Alistair M. Senior,<sup>1,3</sup> Mathieu Lihoreau,<sup>1,4</sup> G. Gregory Neely,<sup>1</sup> Stephen J. Simpson,<sup>1</sup> and Fleur Ponton<sup>1,5</sup>

<sup>1</sup>Charles Perkins Centre and School of Life and Environmental Sciences, Johns Hopkins Drive, Camperdown, NSW 2006, Australia  
<sup>2</sup>Programa de Pós-Graduação em Ecologia e Conservação, Federal University of Paraná, Caixa Postal 19031, CEP 81531-990 Curitiba, Brazil  
<sup>3</sup>The University of Sydney, School of Mathematics and Statistics, Eastern Ave, Camperdown, NSW 2006, Australia

<sup>4</sup>Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI), University Paul Sabatier, CNRS, UPS, 118 Route de Narbonne, 31062 Toulouse, France

<sup>5</sup>Department of Biological Sciences, 205A Culloden Road, Macquarie University, Sydney, NSW 2109, Australia

<sup>6</sup>Present address: University of Florida Entomology and Nematology Department, 1881 Natural Area Drive, Gainesville, FL 32608, USA

<sup>7</sup>Lead Contact

\*Correspondence: [cw442@cornell.edu](mailto:cw442@cornell.edu) <http://dx.doi.org/10.1016/j.cub.2017.07.022>

## SUMMARY

The gut microbiota affects a wide spectrum of host physiological traits, including development [1–5], germline [6], immunity [7–9], nutrition [4, 10, 11], and longevity [12, 13]. Association with microbes also influences fitness-related behaviors such as mating [14] and social interactions [15, 16]. Although the gut microbiota is evidently important for host wellbeing, how hosts become associated with particular assemblages of microbes from the environment remains unclear. Here, we present evidence that the gut microbiota can modify microbial and nutritional preferences of *Drosophila melanogaster*. By experimentally manipulating the gut microbiota of flies subjected to behavioral and chemosensory assays, we found that fly-microbe attractions are shaped by the identity of the host microbiota. Conventional flies exhibit preference for their associated *Lactobacillus*, a behavior also present in axenic flies as adults and marginally as larvae. By contrast, fly preference for *Acetobacter* is primed by early-life exposure and can override the innate preference. These microbial preferences are largely olfactory guided and have profound impact on host foraging, as flies continuously trade off between acquiring beneficial microbes and balancing nutrients from food. Our study shows a role of animal microbiota in shaping host fitness-related behavior through their chemosensory responses, opening a research theme on the interrelationships between the microbiota, host sensory perception, and behavior.

## RESULTS

### Flies' Microbial Preference Depends on Their Microbiota

Based on previous studies [17, 18] and our 16S pyrosequencing survey, the fly egg surface inherited a high abundance of *Acetobacter* and *Lactobacillus*, especially *Acetobacter pomorum* (*AP*) and *Lactobacillus plantarum* (*LP*), which made up 75%–95% reads in our samples (Figure 1A; Table S1). These bacteria have been known to persist throughout developmental stages [2, 3, 19]. At the population level, *Lactobacillus* abundance is generally higher in eggs and at early development but declines, whereas *Acetobacter* abundance increases, as flies age [18, 19]. At the individual level, the relative abundances of *Lactobacillus* and *Acetobacter* vary greatly among flies [20] and are easily perturbed by environmental conditions such as diet and population density [18, 21]. Here, we report a similar trend (Figure 1A; Figures S1A and S1B). Flies deprived of food overnight showed a 9-fold reduction in *Acetobacter* and *Lactobacillus* colony forming unit (CFU) (Figure S1C), pointing to a possibility that *Drosophila* need to replenish their gut microbiota [22]. Based on this assumption, we investigated the chemosensory and behavioral responses of *Drosophila* toward cues associated with beneficial bacteria present in food and tested whether the gut microbiota plays a role in shaping these responses.

We monitored flies' behavioral responses to microbes using a foraging assay (Figure S2A; see STAR Methods). Food-deprived flies were allowed to forage in an arena containing seven patches of yeast-sucrose diet that was either unseeded or seeded with one of six bacteria species isolated from fly guts or bodies: *AP*, *Acetobacter tropicalis* (*AT*), *Lactobacillus casei* (*LC*), *Lactobacillus pantheris* (*LPa*), *LP*, and *Staphylococcus saprophyticus* (*S*). Foraging assays were conducted 1 hr after the bacteria were added to the diets to avoid substantial food modifications by microbial activities. We tested flies raised from conventional, axenic, or monoassociated eggs with the dominant *LP* or *AP*. Conventional flies showed a strong preference for foods seeded with both *Acetobacter* strains tested, *AP* and *AT*, as well as *LP* relative to unseeded food (Figure 1B1; Data S1A). A slight but significant preference was also observed for *LC*, a strain detected at low abundances in fly eggs and adults (0.15%–1.2% and 0.25%–3.9% of pyrosequencing reads, respectively; Figure 1A; Table S1). Flies showed no preference for food seeded with *LPa* or *S*, a strain isolated from fly whole bodies. Both *LPa* and *S* were undetected in our pyrosequencing dataset.

Microbial preferences were dramatically altered in axenic flies. The preference for *AT* was diminished and the preference for *AP* abolished. Nonetheless, axenic flies retained preference for the two *Lactobacillus* strains (*LC* and *LP*) but displayed a higher chance of foraging on the unseeded food (Figure 1B2; Data S1A). Flies raised in monoassociation with *AP* and *LP* showed increased preferences for foods seeded with the corresponding bacteria (Figures 1B3 and 1B4; Data S1A). Interestingly, *AP* monoassociation overrode fly preference for *LP*, although *AP* flies showed preferences toward *AT*, *LC*, and *S*. Conversely, *LP* monoassociation did not show the same antagonistic effects on flies' preference for *Acetobacter* strains. Together, our results demonstrate that flies' microbial preferences are influenced by their microbiota identity.

#### Flies' Microbial Preferences Depend on Early-Life Exposure

The preferences of adult *Drosophila* toward their associated bacteria may be innate or exposure dependent, for instance the result of associative learning between food stimuli and microbial cues. To disentangle between these possible mechanisms, we developed a larval food choice assay (Figure S2B; see STAR Methods). Newly emerged larvae from conventional, axenic, and *AP*- or *LP*-inoculated eggs were given the choices of yeast-sucrose medium either unseeded or seeded with *AP* or *LP*. Larvae from conventional and inoculated eggs promptly sought out media seeded with *AP* or *LP* from day 1 (Figure 2A; Data S1B). In contrast, larvae from axenic eggs evenly spread among all media on day 1. On day 2, larvae from axenic eggs began to show preferences for *AP* and *LP*, although the magnitude of preference was smaller compared to larvae associated with *AP* and *LP*, and a notable proportion of the larvae remained in unseeded medium (Figure 2B; Data S1B). Larvae from conventional and *AP*-inoculated eggs showed the strongest preference for *AP* medium on day 2. Larvae from *LP*-inoculated eggs preferred *LP* and *AP* medium comparably, on both days (Figures 2A and 2B; Data S1B). Together, our results suggest that early-life microbial exposure influences host microbial preference.

#### Fly Attraction toward Beneficial Microbes Is Guided by Olfaction

In the previous assays, fly adults and larvae were allowed to migrate freely in and out of the food patches. Hence, it remained unclear whether the host microbial preferences were mediated via close contact with the microbes (e.g., taste), via volatile cues over a distance (i.e., olfaction), or both. To test for the contribution of close contact, we used a proboscis extension response (PER) assay in which conventional flies were presented single bacterial suspensions (*AP* or *LP*) across a series of doses onto the tarsi (Figure S2C; see STAR Methods). Responses were measured at different recovery times (i.e., 0, 20, 40, and 60 min). We used bacteria re-suspended in water along with a water-only negative control and a sugar solution as positive control. Flies showed a non-linear, positive dose-dependent PER to *LP* (Figure 3A; Data S1C) with a maximum response to CFU of  $1.17 \times 10^9$  per ml. However, PER to *AP* did not differ from the water-only control (Figure 3A, lower panel), suggesting that direct sampling or other close contact mechanisms are unlikely to contribute to fly attraction toward *Acetobacter*.

We then tested the contribution of olfaction by setting up a two-choice trap assay, in which flies were given a choice between an unseeded medium and a *AP*- or *LP*-seeded medium (Figure S2D; see STAR Methods). Conventional flies were strongly attracted to the *LP*-seeded medium and, to a lesser but significant extent, the *AP*-seeded medium (Figure 3B; Data S1D), suggesting that attraction to *LP* and *AP* cues are mediated by olfaction. We further tested olfactory-guided microbial preference in axenic flies. Parallel to our adult fly foraging data, axenic flies preferred the *LP*-seeded medium, but not the *AP*-seeded medium, over the unseeded medium (Figure 3B; Data S1D). Together, our results suggest that olfaction plays an important role in *Drosophila* microbial preferences, but that other chemosensory mechanisms (such as taste) can also be at play for different microbes.

Foraging Decisions Involve Balancing Cues from Both Microbes and Nutrients Animals sense and respond to a changing nutritional environment by adjusting their food choices and consumption patterns [23]. Studies using semi-defined diets have shown that insects can behaviorally balance their food intake to specific protein-to-carbohydrate ratios (P:C) that maximize correlates of evolutionary fitness (e.g., lifetime egg production [24, 25] and resistance to infection [26–28]). Given the importance of the gut microbiota to host nutrition and our observed differing fly responses toward bacterial cues on food, we questioned how these microbes affect fly foraging decisions.

We first tested whether the gut microbiota influences flies' nutritional preference. We offered adult *Drosophila* five unseeded yeast-sucrose diets with varying nutritional contents (P:C 2:1, 1:1, 1:2, 1:4, and 1:8) in the foraging assay (see STAR Methods). As expected (see [29]), conventional flies preferred foraging on the balanced diet (P:C 1:2) the most. Imbalanced diets, either high in protein (P:C 2:1) or carbohydrate (P:C 1:8) (see [24, 29]), were least preferred (Figure 4A1; Data S1E). By manipulating the fly microbiota, we observed subtle yet notable changes on flies' nutritional preferences. Axenic flies retained the greatest preference for P:C 1:2 diet, although the relative preference strength diminished because a higher proportion foraged on the high protein diet (P:C 2:1) (Figure 4A2). In contrast, *AP* flies had a reduced tendency to choose the high-protein diet (P:C 2:1) (Figure 4A3), whereas *LP* flies shifted their greatest preference toward the diet higher in carbohydrate (P:C 1:4) (Figure 4A4). These results indicate that the microbiota can modify flies' nutritional preference (see also [30]).

An important ecological question arising from our findings is how flies prioritize and respond to nutritional and microbial signals in food. To address this, we examined the foraging pattern of conventional flies subjected to binary choices between the highly preferred P:C 1:2 diet and the least preferred P:C 2:1 diet that was unseeded or seeded with *LP* or *AP* (see STAR Methods). Without microbial supplementation, the majority of flies promptly foraged on the P:C 1:2 diet (Figure 4B), and the time spent on the P:C 1:2 diet was three times higher than on the P:C 2:1 diet (Figure 4C; Data S1F). The presence of *AP* or *LP* on the food dramatically altered flies' foraging pattern, promoting both the proportion of flies and their time spent on the high protein diet (P:C 2:1) when seeded, such that the average time spent between the diets was no longer different (Figures 4B and 4C; Data S1F). Our results suggest flies exhibit behavioral tradeoffs between acquiring beneficial microbes and balancing nutrients in foraging.

#### DISCUSSION

Our study provides new evidence that the gut microbiota can modify host chemosensory responses and behavior. Flies show preferences for beneficial bacteria, but these preferences vary depending on host-microbial history and identity. Members of the gut microbiota also affect flies' nutritional preference and can drive behavioral tradeoffs in foraging, as flies have to accommodate both microbial acquisition and nutritional balance. Our results corroborate recent proposals that the gut microbiota controls animal feeding preferences [30–33].

Olfaction is the central mechanism through which flies sense environmental cues and adjust their behaviors. Our study suggests that olfaction participates in host-microbe recognition, potentially facilitating processes such as replenishment of the gut microbiota [22] and symbiont dispersal and transmission [34, 35]. We further show that the effects of microbial exposure on host microbial preference begins at early life stages, signifying maternal microbiota deposition on eggs and offspring, observed in many animals [36], may promote symbiotic association. Together, our findings open new questions about the evolutionary processes that shape animal microbial recognition and foraging behaviors.

In flies, different olfactory receptors and pathways have been implicated in sensing and responding to microbes including pathogens [37], beneficial *Acetobacter*, and yeast [35]. Candidate metabolites that can attract flies through olfaction have recently been suggested [38], raising the question as to whether the gut microbiota can modify host responses to these microbial fermentation products. One possibility is that the gut bacteria produce metabolites that shape the gut-brain axis in flies, like in mammals [39]. The conservation of these microbial-recognition mechanisms and the significance of gut microbiota in other animals remain to be explored.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes two figures, two tables, and one data file and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2017.07.022>.

#### AUTHOR CONTRIBUTIONS

A.C.-N.W., Q.-P.W., S.J.S., and F.P. designed the experiments. A.C.-N.W., Q.-P.W., J.M., M.L., and F.P. ran the experiments. A.C.-N.W., Q.-P.W., J.M., A.M.S., M.L., and F.P. analyzed the results. A.C.-N.W., J.M., A.M.S., M.L., G.G.N., S.J.S., and F.P. wrote the manuscript.

#### ACKNOWLEDGMENTS

We thank Tamara Pulpitel, David Eno Corleto, Ama Sharada Ranasinghe, Ying Shen, Eva Shteinman, Mervyn Wang, and Doris Yau for technical support. The study was funded by the Allen Foundation for nutrition research (2014.483 to A.C.-N.W. and S.J.S.), the University of Sydney J&D Coffey Fellowship (to A.M.S.), the French National Agency for Research (ANR 16CE0200201 to M.L.), the Australian Research Council (DP130103222 to S.J.S. and F.P.), and a CNPq scholarship to J.M. (211668/2013-3).

#### REFERENCES

- Ridley, E.V., Wong, A.C., Westmiller, S., and Douglas, A.E. (2012). Impact of the resident microbiota on the nutritional phenotype of *Drosophila melanogaster*. *PLoS ONE* 7, e36765.
- Shin, S.C., Kim, S.H., You, H., Kim, B., Kim, A.C., Lee, K.A., Yoon, J.H., Ryu, J.H., and Lee, W.J. (2011). *Drosophila* microbiome modulates host developmental and metabolic homeostasis via insulin signaling. *Science* 334, 670–674.
- Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J., and Leulier, F. (2011). *Lactobacillus plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals through TOR-dependent nutrient sensing. *Cell Metab.* 14, 403–414.
- Wong, A.C., Dobson, A.J., and Douglas, A.E. (2014). Gut microbiota dictates the metabolic response of *Drosophila* to diet. *J. Exp. Biol.* 217, 1894–1901.
- Schwarzer, M., Makki, K., Storelli, G., Machuca-Gayet, I., Srutkova, D., Hermanova, P., Martino, M.E., Balmand, S., Hudcovic, T., Heddi, A., et al. (2016). *Lactobacillus plantarum* strain maintains growth of infant mice during chronic undernutrition. *Science* 351, 854–857.
- Elgart, M., Stern, S., Salton, O., Gnainsky, Y., Heifetz, Y., and Soen, Y. (2016). Impact of gut microbiota on the fly's germ line. *Nat. Commun.* 7, 11280.
- Buchon, N., Broderick, N.A., Poidevin, M., Pradervand, S., and Lemaitre, B. (2009). *Drosophila* intestinal response to bacterial infection: activation of host defense and stem cell proliferation. *Cell Host Microbe* 5, 200–211.
- Sansone, C.L., Cohen, J., Yasunaga, A., Xu, J., Osborn, G., Subramanian, H., Gold, B., Buchon, N., and Cherry, S. (2015). Microbiota-dependent priming of antiviral intestinal immunity in *Drosophila*. *Cell Host Microbe* 18, 571–581.
- Lee, K.A., Kim, S.H., Kim, E.K., Ha, E.M., You, H., Kim, B., Kim, M.J., Kwon, Y., Ryu, J.H., and Lee, W.J. (2013). Bacterial-derived uracil as a modulator of mucosal immunity and gut-microbe homeostasis in *Drosophila*. *Cell* 153, 797–811.
- Newell, P.D., and Douglas, A.E. (2014). Interspecies interactions determine the impact of the gut microbiota on nutrient allocation in *Drosophila melanogaster*. *Appl. Environ. Microbiol.* 80, 788–796.
- Teffit, M.A., and Leulier, F. (2017). *Lactobacillus plantarum* favors the early emergence of fit and fertile adult *Drosophila* upon chronic undernutrition. *J. Exp. Biol.* 220, 900–907.
- Clark, R.I., Salazar, A., Yamada, R., Fitz-Gibbon, S., Morselli, M., Alcaraz, J., Rana, A., Rera, M., Pellegrini, M., Ja, W.W., and Walker, D.W. (2015). Distinct shifts in microbiota composition during *Drosophila* aging impair intestinal function and drive mortality. *Cell Rep.* 12, 1656–1667.
- Guo, L., Karpac, J., Tran, S.L., and Jasper, H. (2014). PGRP-SC2 promotes gut immune homeostasis to limit commensal dysbiosis and extend lifespan. *Cell* 156, 109–122.
- Sharon, G., Segal, D., Zilber-Rosenberg, I., and Rosenberg, E. (2011). Symbiotic bacteria are responsible for diet-induced mating preference in *Drosophila melanogaster*, providing support for the hologenome concept of evolution. *Gut Microbes* 2, 190–192.
- Lize, A., McKay, R., and Lewis, Z. (2014). Kin recognition in *Drosophila*: the importance of ecology and gut microbiota. *ISME J.* 8, 469–477.
- Venu, I., Durisko, Z., Xu, J., and Dukas, R. (2014). Social attraction mediated by fruit flies' microbiome. *J. Exp. Biol.* 217, 1346–1352.
- Broderick, N.A., and Lemaitre, B. (2012). Gut-associated microbes of *Drosophila melanogaster*. *Gut Microbes* 3, 307–321.
- Wong, A.C., Luo, Y., Jing, X., Franzenburg, S., Bost, A., and Douglas, A.E. (2015). The host as the driver of the microbiota in the gut and external environment of *Drosophila melanogaster*. *Appl. Environ. Microbiol.* 81, 6232–6240.
- R Core Team (2014). R: a language and environment for statistical computing (R Foundation for Statistical Computing). <http://www.R-project.org/>.
- Ridley, E.V., Wong, A.C., and Douglas, A.E. (2013). Microbe-dependent and nonspecific effects of procedures to eliminate the resident microbiota from *Drosophila melanogaster*. *Appl. Environ. Microbiol.* 79, 3209–3214.

22. Lihoreau, M., Poissonnier, L.A., Isabel, G., and Dussutour, A. (2016). *Drosophila* females trade off good nutrition with high-quality oviposition sites when choosing foods. *J. Exp. Biol.* *219*, 2514–2524.
23. Venables, W.N., and Ripley, B.D. (2002). *Modern Applied Statistics with S*, Fourth Edition (Springer).
24. Bates, D., Machler, M., Bolker, B.M., and Walker, S.C. (2015). Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* *67*, 1–48.
25. Hadfield, J.D. (2010). MCMC methods for multi-response general- ized linear mixed models: the MCMCglmm R package. *J. Stat. Softw.* *33*, 1–22.
26. Langsrud, O. (2003). ANOVA for unbalanced data: used type II instead of type III sums of squares. *Stat. Comput.* *13*, 163–167.
27. Wong, C.N., Ng, P., and Douglas, A.E. (2011). Low-diversity bacterial community in the gut of the fruitfly *Drosophila melanogaster*. *Environ. Microbiol.* *13*, 1889–1900.
28. Wong, A.C., Chaston, J.M., and Douglas, A.E. (2013). The inconstant gut microbiota of *Drosophila* species revealed by 16S rRNA gene analysis. *ISME J.* *7*, 1922–1932.
29. Chandler, J.A., Lang, J.M., Bhatnagar, S., Eisen, J.A., and Kopp, A. (2011). Bacterial communities of diverse *Drosophila* species: ecological context of a host-microbe model system. *PLoS Genet.* *7*, e1002272.
30. Blum, J.E., Fischer, C.N., Miles, J., and Handelsman, J. (2013). Frequent replenishment sustains the beneficial microbiome of *Drosophila melanogaster*. *MBio* *4*, e00860-13.
31. Simpson, S.J., and Raubenheimer, D. (2012). *The Nature of Nutrition: A Unifying Framework from Animal Adaptation to Human Obesity* (Princeton University Press).
32. Lee, K.P., Simpson, S.J., Clissold, F.J., Brooks, R., Ballard, J.W., Taylor, P.W., Soran, N., and Raubenheimer, D. (2008). Lifespan and reproduction in *Drosophila*: new insights from nutritional geometry. *Proc. Natl. Acad. Sci. USA* *105*, 2498–2503.
33. Lihoreau, M., Clarke, I.M., Buhl, J., Sumpter, D.J., and Simpson, S.J. (2016). Collective selection of food patches in *Drosophila*. *J. Exp. Biol.* *219*, 668–675.
34. Graham, R.I., Deacutus, J.M., Pulpitel, T., Ponton, F., Simpson, S.J., and Wilson, K. (2014). Locusts increase carbohydrate consumption to protect against a fungal biopesticide. *J. Insect Physiol.* *69*, 27–34.
35. Ponton, F., Lalubin, F., Fromont, C., Wilson, K., Behm, C., and Simpson, S.J. (2011). Hosts use altered macronutrient intake to circumvent parasite-induced reduction in fecundity. *Int. J. Parasitol.* *41*, 43–50.
36. Povey, S., Cotter, S.C., Simpson, S.J., and Wilson, K. (2014). Dynamics of macronutrient self-medication and illness-induced anorexia in virally infected insects. *J. Anim. Ecol.* *83*, 245–255.
37. Lee, K.P. (2015). Dietary protein:carbohydrate balance is a critical modulator of lifespan and reproduction in *Drosophila melanogaster*: a test using a chemically defined diet. *J. Insect Physiol.* *75*, 12–19.
38. Leitaˆo-Goncˆalves, R., Carvalho-Santos, Z., Francisco, A.P., Fioreze, G.T., Anjos, M., Baltazar, C., Elias, A.P., Itskov, P.M., Piper, M.D.W., and Ribeiro, C. (2017). Commensal bacteria and essential amino acids control food choice behavior and reproduction. *PLoS Biol.* *15*, e2000862.
39. Alcock, J., Maley, C.C., and Aktipis, C.A. (2014). Is eating behavior manipulated by the gastrointestinal microbiota? Evolutionary pressures and potential mechanisms. *BioEssays* *36*, 940–949.
40. Fetissov, S.O. (2017). Role of the gut microbiota in host appetite control: bacterial growth to animal feeding behaviour. *Nat. Rev. Endocrinol.* *13*, 11–25.
41. Wong, A.C., Holmes, A., Ponton, F., Lihoreau, M., Wilson, K., Raubenheimer, D., and Simpson, S.J. (2015). Behavioral microbiomics: a multi-dimensional approach to microbial influence on behavior. *Front. Microbiol.* *6*, 1359.
42. Bright, M., and Bulgheresi, S. (2010). A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol.* *8*, 218–230.
43. Christiaens, J.F., Franco, L.M., Cools, T.L., De Meester, L., Michiels, J., Wenseleers, T., Hassan, B.A., Yaksi, E., and Verstrepen, K.J. (2014). The fungal aroma gene ATF1 promotes dispersal of yeast cells through insect vectors. *Cell Rep.* *9*, 425–432.
44. Funkhouser, L.J., and Bordenstein, S.R. (2013). Mom knows best: the universality of maternal microbial transmission. *PLoS Biol.* *11*, e1001631.
45. Stensmyr, M.C., Dweck, H.K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe, V., Steck, K., Lavista-Llanos, S., et al. (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in *Drosophila*. *Cell* *151*, 1345–1357.
46. Fischer, C.N., Trautman, E.P., Crawford, J.M., Stabb, E.V., Handelsman, J., and Broderick, N.A. (2017). Metabolite exchange between microbiome members produces compounds that influence *Drosophila* behavior. *eLife* *6*, e18855.
47. Mayer, E.A., Tillisch, K., and Gupta, A. (2015). Gut/brain axis and the microbiota. *J. Clin. Invest.* *125*, 926–938.
48. Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.L., et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* *7*, 335–336.





Figure 1. Microbiota Affects Adult *Drosophila* Microbial Preference

(A) 16S rRNA gene pyrosequencing of *Drosophila* eggs. Duplicate samples of eggs are shown. See also [Figure S1](#) and [Table S1](#). (B) Foraging of adult *Drosophila* on food patches unseeded or seeded with six bacteria isolated from flies: *Acetobacter pomorum* (AP), *Acetobacter tropicalis* (AT), *Lactobacillus casei* (LC), *Lactobacillus pantheris* (LPA), *Lactobacillus plantarum* (LP), and *Staphylococcus saprophyticus* (S). Multiple fly microbiota lines (conventional [Conv], axenic [Ax], and AP or LP monocolonized) were tested; n indicates the number of flies tested. Circle sizes are proportional to the total observations of flies. Bars indicate multinomial generalized linear model estimates of the log odds of a fly selecting a given inoculated food over the unseeded food (see [Data S1A](#) for model coefficients). Error bars indicate the SE. Statistical significance inferred from confidence intervals (CIs) is indicated by asterisks (\*, 95% CI or p % 0.05; \*\*, 99% CI or p % 0.01; \*\*\*, 99.9% CI or p % 0.001). See also [Figure S2A](#) and [Data S1A](#).



**Figure 2. Early Microbial Exposure Influences Drosophila Larvae Microbial Preference**

Proportion of newly emerged larvae observed on food patches unseeded or seeded with *AP* or *LP* on day 1 (A) and day 2 (B); n indicates the total number of observations of larvae on food. Data were analyzed using a multinomial generalized linear mixed model (GLMM) with a random effect accounting for repeated measures of the same cohorts of larvae. Asterisks indicate a significantly greater number of observations on a given food relative to the unseeded food based on credible intervals (\*, 95% CI; \*\*, 99% CI; \*\*\*, 99.9% CI). See also [Figure S2B](#) and [Data S1B](#).




Figure 3. *Drosophila* Chemosensory Responses toward Beneficial Bacteria

(A) Proboscis extension response (PER) of adult *Drosophila* toward substrates as a function of dosage of *LP* (top) and *AP* (bottom) and test time (bar colors); *n* indicates the number of flies tested. Data were analyzed using binomial generalized linear mixed models (GLMMs), with a random effect accounting for repeated-measurements of individual flies. The red curve indicates the GLMM estimated probability of PER as a function of concentration of bacterial cells in the suspension. See also [Figure S2C](#) and [Data S1C](#). (B) Attraction index (AI) indicating olfactory preference of conventional and axenic flies toward medium seeded with *LP* and *AP*. AI was calculated as follows: number of flies in *LP*- or *AP*-seeded trap minus number of flies in unseeded trap divided by the total number of flies placed in the foraging arena. Statistical significance was assigned by a two-way type II ANOVA. Different letters indicate significant differences (Student-Newman-Keuls [SNK] post hoc test). See also [Figure S2D](#) and [Data S1D](#).




Figure 4. *Drosophila* Foraging Decisions Are Compounded by the Host and Food Microbiota

(A) Foraging of adult *Drosophila* on yeast-sucrose diets at five different protein-to-carbohydrate ratios (P:C). Multiple fly microbiota lines (conventional [Conv], axenic [Ax], and AP and LP monocolonized) were tested. n indicates the number of flies tested. Circle sizes are proportional to the total observations of flies on each food type. Bars indicate multinomial generalized linear model estimates of the log odds of a fly selecting a given P:C ratio relative to the 1:2 diet. Error bars indicate the SE. Statistical significance was inferred from CIs and is indicated by asterisks (\*, 95% CI or p % 0.05; \*\*, 99% CI or p % 0.01; \*\*\*, 99.9% CI or p % 0.001). See also [Data S1E](#).

(B) Real-time foraging dynamics of flies toward P:C 1:2 and P:C 2:1 diets unseeded or seeded with AP or LP, indicated as the proportion of flies on each diet every 3 min for one hour. n indicates the number of flies tested.

(C) Average time spent per fly upon making a foraging decision on a given food during the 1 hr assay. Error bars indicate the SE. A paired Wilcoxon test (repeated measures) was used to test whether time spent per fly differed between the two foods in each of the three assay groups (1:2 versus 2:1, 1:2 versus 2:1+AP, 1:2 versus 2:1+LP). The asterisk indicates statistical significance, whereby p values < 0.016 are significant after Bonferroni correction for multiple comparisons. See also [Data S1F](#)