



HAL
open science

Control of axial chirality in absence of transition metals based on arynes

Frédéric R. Leroux, Armen Panossian, David Augros

► **To cite this version:**

Frédéric R. Leroux, Armen Panossian, David Augros. Control of axial chirality in absence of transition metals based on arynes. *Comptes Rendus. Chimie*, 2017, 20 (6), pp.682-692. 10.1016/j.crci.2016.12.001 . hal-02105315

HAL Id: hal-02105315

<https://hal.science/hal-02105315>

Submitted on 20 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Control of axial chirality in absence of transition metals based on arynes

Contrôle de la chiralité axiale à l'aide d'arynes et en absence de métaux de transition

Frédéric R. Leroux,* Armen Panossian, David Augros,

Université de Strasbourg, CNRS, LCM UMR 7509, 67000 Strasbourg, France

Phone: +33 3 68 85 26 40, email: frederic.leroux@unistra.fr

Abstract

The modular construction of enantioenriched biaryl derivatives is presented. This approach is based on (a) an almost quantitative access to polybrominated precursors *via* a transition metal-free aryl-aryl coupling, the ARYNE-coupling, (b) the regioselective introduction of a traceless chiral auxiliary (an enantiopure *para*-tolylsulfinyl group), (c) the chemoselective functionalization of this auxiliary and (d) subsequent regioselective functionalization of the remaining bromine atoms without any racemization during these steps.

Next, the atropo-selective coupling of *in situ* generated arynes and aryllithiums bearing various chiral auxiliaries (*tert*-butylsulfoxide, *para*-tolylsulfoxide, tartrate-derived chiral diethers and oxazolines) is described and applied to the formal synthesis of (-)-steganacin.

La construction modulaire de dérivés biaryliques énantioenrichis est présentée. Cette approche est basée sur (a) un accès quasi quantitatif aux précurseurs polybromés *via* un

couplage aryle-aryle sans métaux de transition, le couplage ARYNE, (b) l'introduction régiosélective d'un auxiliaire chiral (un groupe *para*-tolylsulfinyl énantiopur), (c) la fonctionnalisation chimiosélective de cet auxiliaire et (d) la fonctionnalisation régiosélective subséquente des atomes de brome restants sans racémisation au cours de ces étapes.

Ensuite, le couplage atropo-sélectif à l'aide d'arynes générés *in situ* et d'aryllithiums portant divers auxiliaires chiraux (*tert*-butylsulfoxyde, *para*-tolylsulfoxyde, diéthers dérivés du tartrate et des oxazolines chirales) est décrit et appliqué à la synthèse formelle de la (-)-stéganacine.

Keywords

Aryne; Biaryl; Organolithium; Selectivity; Chirality; Synthetic methods; steganacin

1. Introduction

Biaryls are key structural feature of numerous natural products, biologically active molecules, tools for asymmetric synthesis, pharmaceuticals, agrochemicals and other materials (Figure 1). Various approaches have been developed towards efficient methods for their atropo-selective synthesis.^[1] Three important issues have to be addressed for their synthesis, namely the regio- and stereoselective control in biaryl synthesis, as well as the minimization of metal amounts in these processes.




Figure 1. Biaryls as important structural motifs.

In recent years, classical methods for creating aryl-aryl bonds^[2] have been complemented by direct arylation with the carbon-hydrogen bond used as a functional group (Figure 2).^[3] The major drawback in these approaches is the use of heavy metals, which may cause contamination of the products, requiring purification for biological applications. Due to potentially toxic contamination of pharmaceutical products, effective removal of the metal (Pt, Pd, Ir, Rh, Ru, Os) in active pharmaceutical ingredients (API) causes acute problems for pharmaceutical companies.^[4]





Figure 2. Access routes towards biaryls.

In the present account, we report on our contributions in this field. The transition metal-free ARYNE-coupling will be presented towards first racemic biaryl scaffolds and then our developments for the synthesis of enantiopure- or enriched biaryls.

2. Results and Discussion

2.1. The aryne route to access biaryls

The use of arynes to access biaryls or polyaryls has been nicely reviewed recently.^[5] The first example of such a process was described by G. Wittig in 1940 by means of phenyllithium and fluorobenzene.^[6] Later, H. Gilman reported in the 1950s on the reaction of *ortho*-dihalobenzenes with butyllithium leading to a biphenyl backbone, presumably via a 1,2-didehydroarene, i.e. an *ortho*-aryne (Scheme 1).^[7] This halogen/lithium interconversion-based approach remained dormant for several decades except for a procedure improvement.^[8]



Scheme 1. Gilman's observation on the reaction of *ortho*-dihalobenzenes with butyllithium.

Upper line with stoichiometric amounts, lower line with half a molar equivalent.

We started reinvestigating this reaction in 2001 and then extended its scope to access various biaryls, and we clarified the mechanism of the reaction. The reaction of aryllithium compounds with *ortho*-dihaloarenes, proceeding *via* the *in situ* formation of arynes allows the preparation of *ortho*-halobiaryls. This 'ARYNE coupling' has become a robust method for aryl-aryl coupling, and combines several advantages, as the use of cheap and/or easily accessible halogeno aromatic compounds, the access to biaryls bearing two distinct aromatic units and to poly-halogenated biaryls —which can be functionalized further—, the use of lithium reagents (i.e. in the absence of transition metals), and multi-gram reaction scales (Figure 3).^[9]




Figure 3. The ARYNE coupling towards *ortho,ortho'*-di-, tri- and tetra-substituted biaryls.

The mechanism is based on a subtle interplay of several organometallic species and their relative basicities (Figure 4). The chain reaction proceeds as follows: **(1)** A thermodynamically stable organolithium intermediate is formed, of which **(2)** a small amount performs a halogen/metal exchange on the coupling partner *via* an ate complex, generating a thermodynamically unstable *ortho*-halophenyllithium intermediate (*initiation*). The latter **(3)** eliminates spontaneously lithium halide affording an aryne; then, **(4)** the transient aryne species undergoes nucleophilic addition of the organolithium precursor and **(5)** the resulting 2-biaryl lithium intermediate is finally stabilized by in situ transfer of bromine or iodine from the starting material, affording again some *ortho*-halophenyllithium to pursue the chain reaction (*propagation steps*). The *termination* of the chain reaction proceeds by hydrolysis of aryllithium species (by abstraction of a proton of the solvent or during aqueous work-up) and by dimerization or trimerization of residual aryne traces (the corresponding side-products being sometimes detected by GC-MS analysis).





Figure 4. Mechanism of the ARYNE coupling.

The next logical step was then to apply this ARYNE coupling to the preparation of axially chiral biaryls. The synthesis of axially stereo-enriched biaryls remains a challenging and extensively studied research field due to the above-cited importance of the biaryl motif. The most important approaches are the resolution of pre-constructed biaryls and the atropo-selective coupling of two aromatic partners.^[1a-d,10a,10b,1e,1f,10c,10d]

We first focused on the resolution strategy by taking advantage of a specificity of the ARYNE coupling, namely the regeneration of a carbon-exchangeable halogen bond in position 2 of the biaryl product which can be further functionalized.

2.2. Regioselective Halogen/Metal Exchange

Systematic investigations by our laboratory have shown that differences exist in the stabilities of aryllithium intermediates generated by butyllithium-promoted halogen/metal interconversion on bromobenzene and heterosubstituted close congeners. For example, in an intramolecular competition experiment we could show that 1,3-dibromo-5-[(3-bromophenyl)methoxy)methyl]benzene underwent with butyllithium exclusively a halogen/metal permutation at the multiply substituted ring (Scheme 2).




Scheme 2. Intramolecular competition experiments revealing a regioselective bromine/lithium interconversion.

Similarly, 2,2',6-tribromobiphenyl, obtained on multi-gram scale via the ARYNE coupling in almost quantitative yield^[11,9c] undergoes successive and completely regioselective bromine/lithium interconversion starting on the higher substituted phenyl ring.^[12] Such an effective discrimination between two bromine atoms as a function of their chemical environment has so far been observed only sporadically in such kind of processes.^[13] Thus, based on the ARYNE coupling, giving rise to a large number of diversely substituted coupling products, and the regioselective bromine/lithium interconversion and trapping with various electrophiles, we were able to access a multitude of functionalized biaryls in a modular way.^[14,9d]

2.3. Post-ARYNE coupling desymmetrization or deracemization of achiral biaryls

The regioselectivity issue being addressed, the next step was then to control the axial chirality. Therefore, we decided to introduce after the first bromine/lithium exchange a traceless chiral auxiliary. In fact, the sulfinyl group presents numerous synthetic advantages:^[15] (i) its high optical stability; (ii) the existence of a large number of sulfinylating agents in both enantiomeric forms, (iii) its well-known *ortho* directing ability and the efficiency as a carrier of the chiral information, (iv) its electronic properties, which enhance the contrast between the physical properties of the two stereoisomers, mainly because of dipole orientation,^[15a] (iv) it is a very useful as traceless resolving agents, as it may undergo sulfoxide/lithium exchange affording new organometallic intermediates, which can subsequently be trapped by various electrophiles.^[16a,12a,16b]

When we submitted 2,2',6-tribromobiphenyl or other analogues to the regioselective bromine/lithium interconversion^[12b,12c] followed by trapping with enantiomerically pure (1*R*,2*S*,5*R*)-(-)-menthyl-(*S*)-*p*-toluenesulfinate,^[17] the atropo-diastereomeric biarylsulfoxides were obtained in excellent yields. Simple crystallization allowed for the separation of both atropo-diastereoisomers (*S*,*aR*) and (*S*,*aS*) in 54% and 70% yield based on the theoretically possible yield of each diastereoisomer (Scheme 3).




Scheme 3. Desymmetrization of achiral or deracemization of racemic *ortho*-dibromobiphenyls by means of enantiomerically pure *p*-tolylsulfoxide as chiral auxiliary.

The most challenging question was, whether a chemoselective functionalization of the atropo-diastereomerically pure biaryls is possible without any racemization. Both substituents, the sulfoxide group as well as the bromine atoms may undergo exchange reactions. We observed with standard organolithium reagents (butyllithium or *tert*-butyllithium) no chemoselectivity between both types of substituents. In contrast, when employing P. Knochel's *i*PrMgCl•LiCl reagent, a clean sulfoxide/magnesium interconversion occurred at -50 °C. Unfortunately, when the intermediate biaryl Grignard reagent was


treated with an electrophile, the temperature had to be raised to 0 °C, which caused partial racemization of the biaryl axis. Thus, we had to find an exchange reagent which (a) allows the discrimination between the sulfoxide group and the bromine atom(s) and (b) which leads to an intermediate which is reactive enough for trapping at lower temperature. Phenyllithium appeared as the reagent of choice to fulfill all these requirements.

When atropo-diastereomeric biaryl sulfoxides were treated with PhLi at -78 °C, a clean sulfoxide/Li interconversion occurred selectively within 10 min. Subsequent trapping with various electrophiles afforded enantioenriched biaryls with excellent er's ($\geq 96:4$, Scheme 4). Furthermore, since 2 atropodiastereomers were initially generated upon sulfinylation, this method allowed to access both enantiomers of the same biphenyl with identical yield and enantiopurity after the sulfoxide/lithium exchange-electrophilic trapping sequence (see the enantiomers of 2,2'-dibromo-6-methylbiphenyl in Scheme 4).



Scheme 4. Chemoselective functionalization of atropo-isomerically pure biaryl sulfoxides.

In order to show the scope and the modularity of this approach, we decided to perform a complete functionalization of the biaryl *via* subsequent sulfoxide– and bromine/lithium interconversions and to introduce three types of different substituents. We succeeded in functionalizing 2,2',6-tribromobiphenyl and to introduce a methoxy-, a methyl group and a carboxylic function after prior desymmetrization according to the sulfoxide protocol (Scheme 5). A slight erosion of enantiopurity was noticed in the last bromine/lithium exchange, and was expected due to the small size of all *ortho*-substituents on the corresponding biphenyllithium intermediate.^[18] Yet the strategy was validated, and enantiopure biphenyls could be obtained from 2,2',6-tribromobiphenyl after sequential element/lithium exchanges provided that the new substituents introduced are bulky enough.



Scheme 5. Modular atropo-enantiopure biaryl construction starting initially from achiral 2,2',6-tribromobiphenyl.

2.4. Atropo-diastereoselective ARYNE coupling

We next wondered if an atropo-diastereoselective ARYNE coupling, where a coupling partner bears a chiral auxiliary prior to the coupling, might be possible allowing to yield atropo-diastereomers of the desired biaryl.^[9i,9j,9l] The choice of the chiral auxiliary and its location onto the coupling partners revealed to be particularly critical. First, the auxiliary should be able to bind to lithium, in order to produce more rigid transition states and, consequently, lead to higher stereoselectivity. Therefore, an oxygen- and/or nitrogen-bearing auxiliary was needed. Second, to avoid regioselectivity issues during the addition of the aryllithium nucleophile onto the aryne, the auxiliary should be located on the former, not on the latter.

2.4.1. Aryl *tert*-butyl sulfoxides

Our first attempt towards the atropo-diastereoselective ARYNE coupling employed aryl *tert*-butyl sulfoxides.^[9i] After a preliminary study with model substrates prepared from 3-chloro or 3-methoxyphenyllithium with (*S*)-di-*tert*-butyl thiosulfinate, we soon realized that the usual ARYNE coupling conditions (1 equiv. of BuLi to generate the aryllithium nucleophile, followed by addition of 1 equiv. of a 1,2-dibromobenzene as aryne precursor) were ineffective to produce the expected biaryls. We assumed a low reactivity of the 2-lithioaryl sulfoxide and/or of the 2-lithio-2¹-(*tert*-butylsulfinyl)-1,1'-biphenyl in the halogen/lithium exchange reaction with the brominated aryne precursor, thus blocking respectively the initiation and/or the propagation of the chain reaction (Figure 5).





Figure 5. Attempts of atropo-diastereoselective ARYNE coupling based on *tert*-butyl sulfoxides.


This problem could be solved by replacing the dibrominated arylene precursors by the corresponding 1-bromo-2-iodobenzenes and adding a small amount of additional BuLi after introduction of the aryne precursor so as to trigger the generation of the aryne. However, it appeared difficult to achieve simultaneously high yield and reasonable diastereoselectivity.



Scheme 6. Improved coupling conditions for the atropo-diastereoselective ARYNE coupling with *tert*-butyl sulfoxides.

2.4.2. Aryl *para*-tolyl sulfoxides

We considered then aryl *para*-tolyl sulfoxides as appealing alternatives in the atropo-diastereoselective ARYNE coupling. Unlike the corresponding *tert*-butyl sulfoxides, aryl *p*-tolyl sulfoxides can be easily converted into diversely functionalized arenes by sulfoxide/metal exchange using lithium or magnesium bases (for leading reviews, see refs.^[19,16a,12a,16b]) —a valuable asset for the derivatization of biaryls, as others and ourselves already demonstrated.^[20,9g] Unfortunately, application of the procedure developed for *tert*-butyl sulfoxides (*vide supra*)^[9i] proved inefficient. Consequently, we forced the conditions by following the introduction of 1-bromo-2-iodobenzene onto the solution of aryllithium by the addition of a second equivalent of BuLi. We expected a higher amount of aryne electrophile to be produced, in order to increase the trapping of the *ortho*-lithioaryl sulfoxide. The reaction mixture was allowed to stir for 12 h before addition of iodomethane in order to trap the expected biaryllithium intermediate. However, the desired biaryl was not observed but instead an unexpected *ortho*-terphenyl, apart from the methylated arylsulfoxide (Scheme 7). A possible explanation for its formation is depicted in Figure 6. Internal attack of the biaryllithium onto the sulfinyl group would lead, *via* an oxysulfurane intermediate, to a ligand-coupling reaction,^[19] possibly by 1,2-migration of the tolyl group from sulfur to the aryl carbon in an S_NAr fashion, providing a sulfenate anion, which would finally be trapped by iodomethane.



Scheme 7. Diverging course of the atropo-diastereoselective ARYNE coupling with the *p*-tolyl sulfinyl group as chiral auxiliary.





Figure 6. Tentative explanation for the formation of tolyl migration product.

2.4.3. Aryl diethers

Besides the use of sulfur-based chiral auxiliaries, we also studied oxygen-based ones like chiral diethers derived from hydrobenzoin or tartaric acid, inspired by the work of B. Lipshutz on atroposelective intramolecular oxidative aryl-aryl couplings.^[21]

When we opposed a tartrate-derived alkoxyaryl iodide to *ortho*-bromiodobenzenes (Scheme 8), we were pleased to obtain the desired coupling products in good yields. The compounds were obtained with modest diastereomeric ratios of 60:40 and 62:38. The moderate diastereomeric ratios in comparison with *tert*-butylsulfoxide could be caused by the dynamic structure of the aryllithium and biaryllithium intermediates, which could exist

as different species in equilibrium, where the tartrate-derived polyether auxiliary could coordinate lithium through different oxygen atoms.




Scheme 8. Atropo-diastereoselective coupling of chiral diether-bearing substrates.

2.4.4. Aryloxazolines

After these studies on sulfur- and oxygen-based chiral auxiliaries for the atropo-diastereoselective ARYNE coupling, we studied chiral oxazolines. The oxazoline group is a well-known chiral inductor, and a well-known directing group in polar organometallic chemistry.^[22] Moreover, it was used successfully as chiral auxiliary in aryl-aryl couplings *via* an S_NAr process involving 2-(2-magnesioaryl)oxazolines, the Meyers reaction.^[22b-d,23] We decided to introduce the oxazoline group onto the aryl nucleophilic part, i.e. the future aryllithium intermediate instead of the aryne, so as to overcome any regioselectivity issues and to better induce atropo-selectivity by creating a stereoisomerically well-defined metallacycle by coordination of the oxazoline to lithium. The coordination was expected to take place *via* the nitrogen atom of the oxazoline in ethereal solvents, as shown by J. J. Reich.^[24]

We prepared and assessed various aryloxazolines in the ARYNE coupling. The expected biaryls were formed along with a side product, the corresponding fluorenone resulting from

the cyclisation of the biaryllithium intermediate followed by hydrolysis during workup. Considering that the fluorenone byproduct and the desired biaryl are both formed from the same biaryllithium, the total coupling yield has to be taken into account, which ranges from 48 to 79%. Unfortunately, axial chirality is lost in the planar fluorenone, which thus constitutes an issue. Nevertheless, the formed biaryls showed axial stereo-enrichment, with a narrow 66:34–71:29 range of d.r. The diastereoselectivity seems to be slightly influenced by the size of the R² group (Scheme 9).



Scheme 9. Atropo-diastereoselective ARYNE coupling of chiral aryloxazolines with aryne.

The results indicate that with the uncongested aryne precursor a 2.3:1 ratio of biaryl with regard to fluorenone is obtained, while the sterically hindered TMS analogue afforded a much better 5.9:1 ratio in favor of the desired biaryl. This difference might indicate that cyclisation is disturbed by the proximity of the TMS group and the resulting oxazolidine (Figure 7).






Figure 7. The cyclization issue and the role of bulky substituents in the vicinity of the aryne triple bond.

2.5. Application of the atropo-diastereoselective ARYNE coupling to the formal synthesis of (-)-steganacin

Based on these findings, we imagined introducing two trimethylsilyl groups on each side of the triple bond of the aryne. The second TMS group shall avoid competitive regioselectivity of the nucleophilic addition onto the aryne. Additionally, the versatile silicon-carbon bond can be transformed towards various derivatives. Bis-TMS-substituted benzodioxole-derived aryne precursors were thus prepared, and evaluated in the coupling with aryloxazolines. Gratifyingly, the iodinated aryne precursor led to the expected biaryls without any cyclized product with 45–57% yields, which is acceptable for such congested biphenyls.



Aryl oxazoline



Scheme 10. ARYNE coupling of chiral aryloxazolines with a bis-silylated aryne precursor.

Moderate diastereoselectivities were obtained due to presumed opposing substituent effects in competing transition states. Fortunately, atropo-diastereomers could be easily separated in several cases. One of them was then successfully converted into Harayama and Abe's key intermediate towards (-)-steganacin after desilylation, hydrolysis of the oxazoline auxiliary into a protected benzalcohol and formylation of the iodine-bearing carbon *via* iodine/lithium exchange. All the steps, performed on diastereomerically pure precursors (d.r. > 98:2) led, without any racemization, to both enantiomers of the target molecule in e.r. > 99:1.^[25,9],9]

3. Conclusion

The "ARYNE"-adventure which started in 2001 led to an efficient major method for accessing *ortho,ortho'*-di-, tri- and tetrasubstituted biaryls without any transition metals on multi gram quantities. We showed that regioselective bromine/lithium interconversions allow for their functionalization at will. In order to access enantiopure biphenyls, we developed a desymmetrization/deracemization strategy based on an enantiopure sulfinyl group as chiral auxiliary.

Most challenging, we developed then an atropo-selective coupling of *in situ* generated arynes and aryllithiums bearing various chiral auxiliaries *ortho* to lithium. *tert*-Butylsulfoxides gave low to high yields of the coupling product with variable diastereoselectivity. *para*-Tolylsulfoxides, on the other hand, were unable to produce the desired biaryls, mainly because of a lack of reactivity of the 2-lithiophenyl *p*-tolyl sulfoxide intermediates towards halogen/lithium exchange and/or trapping by the aryne. Tartrate-derived chiral diethers yielded the coupling products with moderate diastereoselectivity. Oxazolines gave the most abundant results in terms of coupling, after an evaluation of different reaction parameters.

Finally, we carried out the synthesis of Harayama and Abe's intermediate of (-)-steganacin in 9 steps, 22% overall yield, excellent control of chirality (>98% e.e., with retention of axial stereo-enrichment at each stage of the process), and in the absence of transition metals. Interestingly, the enantiomer of the target compound could be obtained with similar efficiency. The current work enforces the interest in heavy metal-free biaryl synthesis applied to the preparation of potential pharmaceutically relevant ingredients.

4. Acknowledgement

This work was supported by the French Centre National de la Recherche Scientifique (CNRS) and the University of Strasbourg. The French Agence Nationale pour la Recherche (ANR) (grant number ANR-14-CE06-0003-01, ChirNoCat) is gratefully acknowledged for financial support.

5. References

- [1] (a) G. Bringmann, A. J. Price Mortimer, P. A. Keller, M. J. Gresser, J. Garner, M. Breuning, *Angew. Chem. Int. Ed.* 44 (2005) 5384; (b) T. W. Wallace, *Org. Biomol. Chem.* 4 (2006) 3197; (c) M. C. Kozlowski, B. J. Morgan, E. C. Linton, *Chem. Soc. Rev.* 38 (2009) 3193; (d) G. Bringmann, T. Gulder, T. A. M. Gulder, M. Breuning, *Chem. Rev.* 111 (2011) 563; (e) G. Ma, M. P. Sibi, *Chem. Eur. J.* 21 (2015) 11644; (f) J. Wencel-Delord, A. Panossian, F. R. Leroux, F. Colobert, *Chem. Soc. Rev.* 44 (2015) 3418.
- [2] J. Hassan, M. Sévignon, C. Gozzi, E. Schulz, M. Lemaire, *Chem. Rev.* 102 (2002) 1359.
- [3] (a) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* 107 (2007) 174; (b) I. V. Seregin, V. Gevorgyan, *Chem. Soc. Rev.* 36 (2007) 1173; (c) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* 42 (2009) 1074; (d) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem. Int. Ed.* 48 (2009) 5094; (e) L. Ackermann, R. Vicente, A. R. Kapdi, *Angew. Chem. Int. Ed.* 48 (2009) 9792; (f) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* 110 (2010) 624; (g) T. W. Lyons, M. S. Sanford, *Chem. Rev.* 110 (2010) 1147; (h) J. Wencel-Delord, C. Nimphius, F. W. Patureau, F. Glorius, *Angew. Chem. Int. Ed.* 51 (2012) 2247.
- [4] (a) K. Königsberger, G.-P. Chen, R. R. Wu, M. J. Girgis, K. Prasad, O. Repic, T. J. Blacklock, *Org. Proc. Res. Dev.* 7 (2003) 733; (b) C. J. Pink, H. T. Wong, F. C. Ferreira, A. G. Livingston, *Org. Proc. Res. Dev.* 12 (2008) 589; (c) J.-P. Huang, X.-X. Chen, S.-X. Gu, L. Zhao, W.-X. Chen, F.-E. Chen, *Org. Process Res. Dev.* 14 (2010) 939; (d) J. F. Toczko, in: H. Yamamoto and E. M. Carreira (Eds), *9.9 Catalyst Recovery and Recycle: Metal Removal Techniques*, Elsevier Ltd., 2012, p. 209; (e) J. Magano, in: J. Magano and J. R. Dunetz (Eds), *Large-Scale Applications of Transition Metal Removal Techniques in the Manufacture of Pharmaceuticals*, Wiley-VCH Verlag GmbH & Co. KGaA, 2013, p. 313; (f) H. Miyamoto, C. Sakumoto, E. Takekoshi, Y. Maeda, N. Hiramoto, T. Itoh, Y. Kato, *Org. Process Res. Dev.* 19 (2015) 1054.
- [5] J.-A. Garcia-Lopez, M. F. Greaney, *Chem. Soc. Rev.* (2016) DOI: 10.1039/c6cs00220j.
- [6] G. Wittig, G. Pieper, G. Fuhrmann, *Ber. Dtsch. Chem. Ges.* 73 (1940) 1193.
- [7] (a) H. Gilman, R. D. Gorsich, *J. Am. Chem. Soc.* 78 (1956) 2217; (b) H. Gilman, B. J. Gaj, *J. Org. Chem.* 22 (1957) 447.
- [8] T. K. Dougherty, K. S. Y. Lau, F. L. Hedberg, *J. Org. Chem.* 48 (1983) 5273.

- [9] (a) F. Leroux, M. Schlosser, *Angew. Chem. Int. Ed.* 41 (2002) 4272; (b) F. R. Leroux, L. Bonnafoux, C. Heiss, F. Colobert, D. A. Lanfranchi, *Adv. Synth. Catal.* 349 (2007) 2705; (c) L. Bonnafoux, F. Colobert, F. R. Leroux, *Synlett* (2010) 2953; (d) V. Diemer, M. Begaud, F. R. Leroux, F. Colobert, *Eur. J. Org. Chem.* (2011) 341; (e) L. Bonnafoux, R. Gramage-Doria, F. Colobert, F. R. Leroux, *Chem. Eur. J.* 17 (2011) 11008; (f) V. Diemer, A. Berthelot, J. Bayardon, S. Jugé, F. R. Leroux, F. Colobert, *J. Org. Chem.* 77 (2012) 6117; (g) F. R. Leroux, A. Berthelot, L. Bonnafoux, A. Panossian, F. Colobert, *Chem. Eur. J.* 18 (2012) 14232; (h) A. Oukhrib, L. Bonnafoux, A. Panossian, S. Waifang, D. H. Nguyen, M. Urrutigoity, F. Colobert, M. Gouygou, F. R. Leroux, *Tetrahedron* 70 (2014) 1431; (i) A. Berthelot-Bréhier, A. Panossian, F. Colobert, F. R. Leroux, *Org. Chem. Front.* 2 (2015) 634; (j) B. Yalcouye, A. Berthelot-Bréhier, D. Augros, A. Panossian, S. Choppin, M. Chessé, F. Colobert, F. R. Leroux, *Eur. J. Org. Chem.* (2016) 725; (k) M. J. Fer, J. Cinqualbre, J. Bortoluzzi, M. Chessé, F. R. Leroux, A. Panossian, *Eur. J. Org. Chem.* (2016) 4545; (l) D. Augros, B. Yalcouye, A. Berthelot-Bréhier, M. Chessé, S. Choppin, A. Panossian, F. R. Leroux, *Tetrahedron* 72 (2016) 5208.
- [10] (a) G. Bringmann, T. Gulder, T. A. Gulder, M. Breuning, *Chem. Rev.* 111 (2011) 563; (b) D. Zhang, Q. Wang, *Coord. Chem. Rev.* 286 (2015) 1; (c) P. Loxq, E. Manoury, R. Poli, E. Deydier, A. Labande, *Coord. Chem. Rev.* 308 (2016) 131; (d) O. M. Demchuk, K. Kapłan, A. Kačka, K. M. Pietrusiewicz, *Phosphorus Sulfur Silicon Relat. Elem.* 191 (2016) 180.
- [11] F. R. Leroux, L. Bonnafoux, F. Colobert, WO 2008037440, LONZA AG.
- [12] (a) F. Leroux, M. Schlosser, E. Zohar, I. Marek, in: Z. Rappoport (Eds), *The preparation of organolithium reagents and intermediates*, Wiley, Chichester, 2004, p. 435; (b) F. Leroux, N. Nicod, L. Bonnafoux, B. Quissac, F. Colobert, *Lett. Org. Chem.* 3 (2006) 165; (c) L. Bonnafoux, F. R. Leroux, F. Colobert, *Beilstein J. Org. Chem.* 7 (2011) 1278.
- [13] (a) M. Schlosser, *Angew. Chem. Int. Ed.* 44 (2005) 376; (b) M. Schlosser, *Organometallics in synthesis third manual*, Wiley, Hoboken, 2013.
- [14] V. Diemer, F. R. Leroux, F. Colobert, *Eur. J. Org. Chem.* (2011) 327.
- [15] (a) J. Clayden, D. Mitjans, L. H. Youssef, *J. Am. Chem. Soc.* 124 (2002) 5266; (b) J. Clayden, P. M. Kubinski, F. Sammiceli, M. Helliwell, L. Diorazio, *Tetrahedron* 60 (2004) 4387; (c) J. Clayden, S. P. Fletcher, J. J. W. McDouall, S. J. M. Rowbottom, *J. Am. Chem. Soc.* 131 (2009) 5331.
- [16] (a) J. Clayden, *Organolithiums: Selectivity for Synthesis*, Pergamon, 2002; (b) A. Panossian, F. R. Leroux, in: J. Mortier (Eds), *The Halogen-Metal Interconversion and Related Processes (M = Li, Mg)*, Wiley, Hoboken, 2016, p. 813.
- [17] (a) K. K. Andersen, *Tetrahedron Lett.* 3 (1962) 93; (b) G. Solladié, J. Hutt, A. Girardin, *Synlett* (1987) 173.
- [18] (a) A. Mazzanti, L. Lunazzi, M. Minzoni, J. E. Anderson, *J. Org. Chem.* 71 (2006) 5474; (b) R. Ruzziconi, S. Spizzichino, L. Lunazzi, A. Mazzanti, M. Schlosser, *Chem. Eur. J.* 15 (2009) 2645; (c) A. Mazzanti, L. Lunazzi, R. Ruzziconi, S. Spizzichino, M. Schlosser, *Chem. Eur. J.* 16 (2010) 9186; (d) R. Ruzziconi, S. Spizzichino, A. Mazzanti, L. Lunazzi, M. Schlosser, *Org. Biomol. Chem.* 8 (2010) 4463; (e) L. Lunazzi, M. Mancinelli, A. Mazzanti, S. Lepri, R. Ruzziconi, M. Schlosser, *Org. Biomol. Chem.* 10 (2012) 1847.
- [19] N. Furukawa, S. Sato, *Top. Curr. Chem.* 205 (1999) 89.
- [20] (a) J. Clayden, P. M. Kubinski, F. Sammiceli, M. Helliwell, L. Diorazio, *Tetrahedron* 60 (2004) 4387; (b) P. Knochel, T. Thaler, F. Geittner, *Synlett* (2007) 2655; (c) J. Clayden, S. P. Fletcher, J. J. W. McDouall, S. J. M. Rowbottom, *J. Am. Chem. Soc.* 131 (2009) 5331.
- [21] (a) B. H. Lipshutz, F. Kayser, Z.-P. Liu, *Angew. Chem. Int. Ed.* 33 (1994) 1842; (b) G.-Q. Lin, M. Zhong, *Tetrahedron Lett.* 38 (1997) 1087; (c) L. Guo-Qiang, Z. Min, *Tetrahedron: Asymmetry* 8 (1997) 1369; (d) R. V. Kyasnoor, M. V. Sargent, *Chem.*

- Commun. (1998) 2713; (e) B. H. Lipshutz, P. Müller, D. Leinweber, *Tetrahedron Lett.* 40 (1999) 3677.
- [22] (a) A. I. Meyers, *Acc. Chem. Res.* 11 (1978) 375; (b) M. Reuman, A. I. Meyers, *Tetrahedron* 41 (1985) 837; (c) T. G. Gant, A. I. Meyers, *Tetrahedron* 50 (1994) 2297; (d) A. I. Meyers, *J. Org. Chem.* 70 (2005) 6137.
- [23] J. Mortier, *Curr. Org. Chem.* 15 (2011) 2413.
- [24] K. L. Jantzi, I. A. Guzei, H. J. Reich, *Organometallics* 25 (2006) 5390.
- [25] D. Augros, B. Yalcouye, S. Choppin, M. Chessé, A. Panossian, F. R. Leroux, accepted; DOI: 10.1002/ejoc.201601239.