



HAL
open science

Comment on "Inverse Doppler shift and control field as coherence generators for the stability in superluminal light"

Bruno Macke, Bernard Ségard

► **To cite this version:**

Bruno Macke, Bernard Ségard. Comment on "Inverse Doppler shift and control field as coherence generators for the stability in superluminal light". *Physical Review A : Atomic, molecular, and optical physics* [1990-2015], 2019, 99 (4), 10.1103/PhysRevA.99.047801 . hal-02111063

HAL Id: hal-02111063

<https://hal.science/hal-02111063>

Submitted on 25 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comment on “Inverse Doppler shift and control field as coherence generators for the stability in superluminal light”

Bruno Macke and Bernard Ségard*

Université de Lille, CNRS, UMR 8523, Physique des Lasers, Atomes et Molécules, F-59000 Lille, France

(Dated: April 25, 2019)

In their study of inverse Doppler shift and superluminal light [Phys. Rev. A **91**, 053807 (2015)], Ghafoor *et al.* consider a three-level atomic arrangement with transitions in the optical domain. In fact, the values they give to the parameters lead to a probe wavelength lying in the decimeter band. We point out that the Doppler shifts are then negligible and remark that the simulations performed by Ghafoor *et al.* do not evidence any superluminal effect.

In their study of inverse Doppler shift and superluminal light [1], Ghafoor *et al.* consider a three-level atomic arrangement with transitions in the optical domain, referring in particular to the sodium D_2 line at 586.9 nm. On the other hand, they specify in the caption of their Fig.2 that all the (angular) frequencies are given in units of $\Gamma = 2\pi \times 1$ MHz and that the frequency of the probe transition $\omega_{ac} = 1000\Gamma$. The corresponding wavelength is thus $\lambda = 30$ cm (in the decimeter band).

The first consequence of the large value of the probe wavelength is that the Doppler broadening V_D is very small, in the order of $2\pi \times 1$ kHz. The consideration of V_D going from 2 to 12 MHz as made in [1] is meaningless and the so-called inverse Doppler shift, claimed as the novelty of the article, is in fact negligible.

A second point is that the calculations developed in [1] lead to fully unrealistic values of the atomic number density N . As correctly given in the article, the electric susceptibility for the probe reads in SI units:

$$\chi = \frac{2N |\wp_{ac}|^2 \rho_{ac}}{\varepsilon_0 \hbar \Omega_p} \quad (1)$$

where a (c) is the upper (lower) level of the probe transition, \wp_{ac} (ρ_{ac}) is the corresponding matrix element of the dipole moment (of the density operator) and Ω_p is the Rabi (angular) frequency of the probe. From the involved discussion following this equation, it results that

$$\Gamma = \frac{|\wp_{ac}|^2 \omega_{ac}^3}{\varepsilon_0 \hbar c^3} = O\left(\frac{N |\wp_{ac}|^2}{\varepsilon_0 \hbar}\right) \quad (2)$$

and that

$$N = O\left(\frac{8\pi^3}{\lambda^3}\right) \quad (3)$$

For $\lambda = 30$ cm, we get an atomic number density in the order of 10^{-2}cm^{-3} , which is 12 orders of magnitude lower than those attainable in the best vacuum devices.

As a third point, we remark that, contrary to the claim made in the article title, the simulations made in [1] do not evidence any superluminal effect, namely an advance of the intensity profile of the transmitted pulse on that of the incident one (see Fig.5). The calculation itself raises

some questions. The transmitted field is actually the inverse Fourier transform of $S_{in}(\omega)H(\omega)$, where $S_{in}(\omega)$ and $H(\omega)$ are, respectively, the Fourier transform of the incident field and the transfer function of the medium. Insofar as $S_{in}(\omega)$ is Gaussian and $H(\omega)$ is the exponential of a polynomial of degree 3, the result cannot be that given by Eq.(15) in [1] but necessarily involves an Airy function. We also note that the transfer function $H(\omega)$ considered by Ghafoor *et al.* neglects the frequency dependence of the medium transmission that can considerably affect the profile of the transmitted pulse [2].

For completeness, we mention that some equations in [1] seem to be dimensionally inhomogeneous, that the Einstein’s coefficient given below Eq.(4) is erroneous (see [3] for its exact value in SI units) and that Eq.(1) and Eq.(8) mix results that hold, respectively, in SI and in electrostatic units (the corresponding susceptibilities differ by a factor of 4π).

We finally point out that the atomic number density given by Eq.(3), anomalously weak in the conditions considered in [1], raises on the contrary to values $N = O(10^{15} \text{cm}^{-3})$ which are too large when the probe wavelength λ is that of the sodium D_2 line. On another hand, the fixed ratio $\omega_{ac}/\Gamma = 1000$ leads then to lifetimes of the excited atomic states which are fully unrealistic (in the subpicosecond range).

We thank Shubhrangshu Dasgupta from the Indian Institute of Technology Ropar for helpful information. This work has been partially supported by the Ministère de l’Enseignement Supérieur, de la Recherche et de l’Innovation, the Conseil Régional des Hauts de France and the European Regional Development Fund (ERDF) through the Contrat de Projets État-Région (CPER) 2015–2020, as well as by the Agence Nationale de la Recherche through the LABEX CEMPI project (ANR-11-LABX-0007).

* Electronic address: bernard.segard@univ-lille.fr

[1] F. Ghafoor, B.A. Bacha, and S. Khan, Inverse Doppler shift and control field as coherence generators for the stability in superluminal light, Phys. Rev. A **91**, 053807

- (2015).
- [2] B. Macke and B. Ségard, On-resonance material fast light, Phys. Rev. A **97**, 063830 (2018).
- [3] R.C. Hilborn, Einstein coefficients, cross sections, values, dipole moments, and all that, arXiv: physics/0202029. See Eq.(40).