



HAL
open science

Rab35 GTPase and cancer: Linking membrane trafficking to tumorigenesis

Ronan Shaughnessy, Arnaud Echard

► **To cite this version:**

Ronan Shaughnessy, Arnaud Echard. Rab35 GTPase and cancer: Linking membrane trafficking to tumorigenesis. *Traffic*, 2018, 19 (4), pp.247-252. 10.1111/tra.12546 . hal-02114022

HAL Id: hal-02114022

<https://hal.science/hal-02114022v1>

Submitted on 29 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rab35 GTPase and Cancer: Linking membrane trafficking to tumorigenesis

Ronan Shaughnessy¹ and Arnaud Echard^{1,2,*}

1. Membrane Traffic and Cell Division Lab, Cell Biology and Infection Department, Institut Pasteur, 25-28 rue du Dr Roux, 75724 Paris, France.
2. Centre National de la Recherche Scientifique, UMR3691, 75015 Paris, France.

*Corresponding author:

Arnaud Echard, PhD
Membrane Traffic and Cell Division Lab,
Institut Pasteur,
CNRS UMR3691,
28 Rue de Dr Roux,
75015 Paris, France.
Phone: +33 1 44 38 94 09
E-mail: arnaud.echard@pasteur.fr

Running title: Role for Rab35 in membrane trafficking and cancer

Keywords: Rab35, Rab protein, Membrane trafficking, Cancer, Cytokinesis, Cell signaling, Cell migration, Tumorigenesis

Brief synopsis: Rab35 is a small GTPase that was recently reported to possess oncogenic activating mutations in human tumors. Conversely, Rab35 depletion inverts apico-basal cell polarity and promotes cell migration. In the present review we describe how Rab35's known functions in membrane trafficking, signaling, cell division and cell migration could explain its role in tumorigenesis.

Abstract

Rab35 is a small GTPase that is involved in many cellular processes, including membrane trafficking, cell polarity, lipid homeostasis, immunity, phagocytosis and cytokinesis. Recent studies showed that activating mutations confer Rab35 with oncogenic properties. Conversely downregulation of Rab35 inverts apico-basal cell polarity and promotes cell migration. Here we review Rab35's known functions in membrane trafficking and signaling, cell division, and cell migration in cancer cells and discuss the importance of Rab35-dependent membrane trafficking in cancer progression.

Introduction

Intracellular membrane trafficking regulates uptake of nutrients and signaling from extracellular molecules by controlling the plasma membrane localization of transporters and receptors, and thus plays a key role in cell differentiation and proliferation. In addition, cells rely on membrane trafficking to maintain cell architecture, cell shape and cell polarity, as well as organ structure and function^{1,2}. Cancer cells often have altered membrane trafficking that can lead to the upregulation of growth factor receptors at the cell surface (e.g. epidermal growth factor receptor; EGFR) and abnormal levels of adhesion molecules (e.g. β 1-integrin, cadherins), which can promote epithelial-mesenchymal transition (EMT) and an invasive phenotype during cancer progression³⁻⁵.

Rab proteins are evolutionarily-conserved GTP-binding proteins that regulate intracellular membrane trafficking in all eukaryotic cells⁶⁻⁸. They act as molecular switches between an inactive GDP and active GTP form, where the latter form interacts with specific intracellular membranes and controls local recruitment of downstream effectors. Guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs) regulate Rab activation and deactivation, respectively⁹. Importantly, Rab proteins control the endocytosis, recycling and lysosomal degradation of growth factor receptors including those overexpressed in many cancers⁴, such as the EGFR¹⁰, platelet derived growth factor receptor (PDGFR)¹¹, vascular endothelial growth factor receptor (VEGFR)¹², c-Met¹³ and integrins^{14,15}. Manipulating the endocytic trafficking and recycling of these receptors has emerged as an attractive method to counteract cancer cell proliferation and survival. For example, pharmacologically-induced accumulation of ligand-activated EGFR in endosomes activates apoptosis selectively in cancer cells^{16,17}. Taken together, this suggests the potential for targeting endocytic trafficking to counteract cancer cell proliferation.

There is increasing evidence to implicate a select few Rab proteins in cancer progression (reviewed in ref.⁷). Here we will focus on Rab35, a GTPase¹⁹ first described as being important for endocytic recycling and cytokinesis¹⁸. Rab35 is unique in that it contains an evolutionarily-conserved polybasic C-terminal tail and localizes both at the plasma membrane and on endosomes^{18,20}. Its polybasic tail facilitates its interaction with PtdIns(4,5)P₂ and PtdIns(3,4,5)P₃ at the plasma membrane²⁰. Rab35 has attracted a lot of attention in the past ten years in the endocytic community and beyond, as reviewed recently in ref.²¹. Rab35 has many functions in various cellular processes including membrane trafficking (see below), autophagy²², cell polarity^{23,24}, cytokinesis^{25,26}, membrane lipid homeostasis^{25,27}, neurite outgrowth²⁸⁻³¹, axonal elongation³², immunity³³⁻³⁶, phagocytosis^{37,38}, pathogen infection^{39,40}, myoblast fusion⁴¹, exosome secretion⁴² and Weibel-Palade bodies secretion⁴³. Moreover, the mechanisms controlling Rab35 activation in endocytosis is well understood with the identification of Rab35 GEFs (DENND1A-C/Connecdenn1-3^{27,44,47}) and GAPs (TBC1D10A-C/EPI64A-C^{33,36,48}, TBC1D13⁴⁹, TBC1D24⁵⁰), and reviewed in detail in²¹ and⁵¹.

A recent study described Rab35 oncogenic somatic mutations in human tumors, which activate Rab35 and have transforming capabilities in mouse fibroblasts *in vitro*¹¹ (see Rab35 tumor somatic mutations in Table 1). In this context, Rab35 regulates PDGFR trafficking leading to cell transformation through a PI3K/AKT-dependent mechanism¹¹. Rab35 was initially linked to cancer by its interaction with the oncogenic nucleophosmin-anaplastic lymphoma fusion kinase (NPM-ALK)⁵² and the p53-related protein kinase (PRPK), where it negatively regulates PRPK-mediated p53 transcriptional activity^{32,53}. However, the relevance of Rab35 function in lymphoma or p53-dependent transformation remains to be established.

Here, we review the recent data suggesting a direct role for Rab35-dependent membrane trafficking in multiple and intertwined aspects of cancer progression by controlling cell division, polarity and migration, as well as survival and proliferative intracellular signaling.

Rab35 in membrane trafficking and cell division

Cytokinesis (the last step of cell division leading to the physical separation of the daughter cells) and cell polarity are two key processes deregulated in cancer and Rab35 plays a role in both. An RNAi screen looking for Rabs involved in cytokinesis identified Rab35 and showed that depletion of this Rab generated binucleated cells and perturbed a fast recycling pathway¹⁸. The first isolated effector of Rab35 was the PtdIns(4,5)P₂ phosphatase Oculo-Cerebro-Renal syndrome of Lowe (OCRL), which is necessary for successful cytokinetic abscission²⁵. Of note, OCRL traffics in and out of the cytokinetic bridge on Rab35-positive vesicles²⁵. Another Rab35 effector, MICAL1, which belongs to an emerging class of oxidation-reduction enzymes that depolymerize F-actin through selective oxidation^{54,55}, facilitates ESCRT-III recruitment at cytokinetic bridges and thus abscission²⁶. Both OCRL and MICAL1 localize to cytokinetic bridges in a Rab35-dependent manner, and their depletion causes the accumulation of F-actin at this location^{25,26}. This suggests a parallel, and perhaps synergistic, mechanism of F-actin removal from cytokinetic bridges, which appears to be essential for successful cytokinesis. Importantly, studies have established that cytokinetic failure leading to tetraploidy promotes tumorigenesis *in vivo*⁵⁶, as shown experimentally in primary mouse mammary epithelial cells⁵⁷.

In addition to a mechanistic role for Rab35 and its effectors in cytokinesis by clearing F-actin, Rab35 plays a role in connecting cytokinesis with apical polarity initiation in MDCK cells grown in Matrigel to form cysts²³. In this study Rab35 was found to tether vesicles containing important apical proteins, such as Cdc42, Crumbs3, aPKC, and the lumen promoting factor GP135/Podocalyxin (PODXL) at the cytokinetic abscission site to deliver them to the future apical membrane initiation site (AMIS). Consequently, Rab35 depletion or deactivation prevents AMIS and lumen formation, and leads to a complete inversion of apico-basal polarity^{23,24}. As changes in apico-basal polarity are believed to favor cancer development, this suggests that loss of function mutations of Rab35 might promote tumorigenesis. Interestingly, Rab35 is downregulated in several tumors including grade IV gliomas, breast and renal carcinomas^{58,59}.

In summary, the function of Rab35 in the final step of cell division and in apico-basal polarity establishment might be important to explain a role for this GTPase in cancer initiation.

Rab35 in membrane trafficking and cell signaling

Rab35 plays a central role in intracellular trafficking by promoting endocytic recycling of various cargoes, including the transferrin receptor (TfR)^{18,40}, yolk receptor⁶⁰, Shiga toxin⁴⁸,

major histocompatibility complex class-I (MHC-I) and MHC-II^{33,34}, T-cell receptor³⁶, Ca²⁺-activated K⁺ channel KCa2.3⁶¹, Megalin⁶², GLUT4⁴⁹, M- and N-Cadherin^{41,58} and β 1-integrin^{58,63}. Moreover, Rab35 knockdown inhibits trafficking of CI-mannose-6-phosphate receptors from endosomes to the trans Golgi network (TGN)²⁷.

Recent studies showed that the tumor suppressor folliculin (FLCN) interacts with GDP-bound Rab35 *in vitro* and negatively regulates EGFR signaling after EGF stimulation^{64,65}, possibly acting as a Rab35 GEF⁶⁶. Moreover, FLCN-mediated EGFR degradation in HeLa cells requires Rab35 activation⁶⁴. In these cells, depletion of Rab35 increases AKT activation after EGF stimulation and cell proliferation⁶⁴, as previously observed in COS-7 cells⁵⁸. Rab35 is downregulated in some cancers^{58,59}, which could lead to increased EGFR receptor recycling and signaling, a common hallmark of many malignancies³. Surprisingly, a recent study reported that depletion of Rab35 in HeLa cells rather diminished AKT signaling, at least after 10% serum stimulation¹¹. The authors attributed cell context and experimental conditions as a reason for this discrepancy (perhaps serum vs EGF stimulation⁶⁴) but this needs to be clarified. Nevertheless, two Rab35 activating mutations (A151T and F161L) found in human cancers (lung, uterus and lymphoid tissues; see Table 1) lead to upregulation of the PI3K/AKT signaling pathway via mTORC2, as well as mouse fibroblast transformation and apoptosis evasion¹¹. This associates with increased signaling and trafficking of the PDGFR- α into Lamp2-positive vesicles¹¹. This study thus suggests a pivotal role for Rab35 activating mutations in endocytic deregulation and PDGFR-mediated oncogenic signaling in cancer progression.

It is not known, however, how exactly Rab35 controls EGFR and PDGFR trafficking in FLCN-depleted cells or in cells expressing activating mutations of Rab35. Given the number of cargoes whose trafficking is regulated by Rab35 (see above), it is likely that Rab35 plays a fundamental and general role in the endosomal pathway. As during cytokinesis, GTP-bound Rab35 binds to the OCRL lipid phosphatase²⁷, which promotes PtdIns(4,5)P₂ hydrolysis on newborn endosomes/clathrin coated vesicles recently pinched from the plasma membrane⁶⁷. This promotes the generation and maintenance of PtdIns4P on endosomes, an important step for normal cargo sorting and recycling towards the plasma membrane or targeting to the trans-Golgi network⁶⁷. In addition to PtdIns4P vs. PtdIns(4,5)P₂ homeostasis, active Rab35 might regulate PtdIns(3,4,5)P₃ levels and thus AKT activation, since Rab35 co-immunoprecipitates with PI3K¹¹. Therefore, abnormal levels of GTP-bound Rab35 could affect the lipid composition of endosomes in cancer cells and dictate growth factor receptor fate.

In parallel to phosphoinositide regulation, Rab35 controls recycling of cargoes from ARF6-positive endosomes back to the plasma membrane through the recruitment of two additional effectors: the ARF6-GAP ACAP2/CentaurinB2^{28,29} and MICAL-L1⁶⁸, a scaffolding protein that, contrary to MICAL1 described above, has no redox activity^{54,55}. For detailed reviews on ARF proteins and cancer, please refer to refs.^{69,70}. Upon activation, Rab35 inactivates ARF6, which promotes endosomal recycling^{28,29,68,71}. In addition, Rab35 recruits MICAL-L1, which facilitates tubular endosomal fission by directly interacting with the dynamin-like protein EHD1^{28,29,72}. Finally, MICAL-L1 recruits other small GTPases (Rab8, Rab13 and Rab36) that also promote endosomal recycling⁷³. Of note, the TP53 status of cancer cells could also influence recycling in this pathway since p53 regulates MICAL-L1 transcription and tubular endosome biogenesis⁷⁴.

Altogether, abnormal levels of Rab35 activation impair endosomal recycling and phosphoinositide homeostasis by

several mechanisms, which could explain growth factor receptor and survival signaling observed in cancer situations with perturbed Rab35.

Rab35 in cell migration

Cell migration and dissemination play a critical role in tumor metastasis. Several lines of evidence indicate that Rab35 regulates cell migration, however this depends on the cell type and the influence of external cues. In COS-7 cells, depletion of Rab35 induces a phenotype with hallmarks of EMT *in vitro*⁵⁸, with increased cell migration and decreased cell-cell adhesion. This can be explained by the fact that Rab35-dependent recycling is required for maintaining cadherins at the cell surface in COS-7 cells⁵⁸, as in myoblasts⁴¹. In addition, Rab35 depletion increases β 1-integrin levels at the cell surface and thus promotes cell migration, since Rab35 inhibits ARF6-mediated β 1-integrin recycling through the ARF6-GAP ACAP2⁵⁸ (see above). Rab35 depletion in HeLa and MDA-MB-231 cells increases β 1-integrin levels, however authors attributed this to impaired internalization rather than increased recycling⁶³. In any case, overexpression of miRNA-720 (a marker for colorectal cancer⁷⁵ and multiple myeloma⁷⁶, and which targets Rab35) also resulted in increased cell migration in HeLa cells⁷⁷.

Conversely, Rab35 activation after prolonged EGF stimulation reduces the interaction between the G protein-coupled receptor kinase interacting ARFGAP 2 (GIT2) and its binding partner RUSC2 and consequently reduces directional cell migration in non-small cell lung cancer (NSCLC)⁷⁸. However, contrary to the studies mentioned above, Rab35 activation downstream of Wnt5 signaling promotes cell migration in breast cancer MCF-7 cells via a Dvl2/Rab35/Rac1 signaling pathway⁷⁹, perhaps due to additional dominant effects downstream of Wnt5.

Finally, EGF-mediated Rab35 activation leads to the activation of the enzyme MICAL1, and this associates with reactive oxygen species (ROS) production, AKT phosphorylation, and cell invasion in a Matrigel Transwell assay in breast cancer cells⁸⁰. Since GTP-Rab35 binding to MICAL1 releases an inhibitory intramolecular folding, activates the redox activity of MICAL1 and selectively depolymerizes F-actin²⁶, it would be interesting to characterize whether Rab35 remodels cortical actin during invasion.

These seemingly contradictory results suggest that depending on the migration assay (with or without extracellular matrix), the cell type and on the means by which Rab35 is activated or inhibited, Rab35 plays different roles in migration vs. invasion.

Concluding remarks

Abnormal activation of Rab35 modifies actin dynamics, cell division and polarity. In addition, Rab35 regulates the endocytic recycling of key cargoes involved in cell migration, such as β 1-integrin as well as growth factor receptor recycling and signaling through PDGFR- α , EGFR and AKT (Figure 1). Based on the *in vitro* studies reviewed above, modification of Rab35 activation observed in several cancers might influence EMT and cancer cell migration/metastasis. However, additional work is needed to address experimentally the consequences of Rab35 upregulation or downregulation on tumorigenesis *in vivo*.

Rab proteins have gained interest as targets for the development of treatments for diseases^{81,82}. As Rab35 is known to play an important role in endocytic recycling, an interesting approach would be to target the Rab35-specific effectors responsible for this function instead of the Rab35 molecule itself, which is involved beyond, and thus specifically

affecting the endocytic trafficking of cancer-involved growth factor receptors. Perhaps pharmacologically targeting the Rab35 effector OCRL could be an attractive possibility. Interestingly, a bistable switch between Rab35 and ARF6 has been postulated²¹, since GTP-Rab35 interacts with ACAP2/CentaurinB2 (an ARF6 GAP)^{28,29}, while GTP-ARF6 interacts with TBC1D10/EPI64 (a Rab35 GAP)³³. Thus, activating ARF6 in the endocytic pathway may also prove interesting for inactivating Rab35-dependent trafficking of growth factor receptors.

While both Rab35 activating mutations¹¹ and Rab35 downregulation^{58,59} are described in cancer, these phenomena could play complementary roles at different stages of cancer development through a process known as clonal evolution⁶³. One can indeed speculate that during initial tumorigenic events, activating mutations described to upregulate PDGFR trafficking and AKT activation would lead to cell survival and transformation. In contrast, Rab35 downregulation leads to a migratory or invasive EMT phenotype and could be the consequence of gene silencing at a later stage, after cell transformation.

Taken together, Rab35 emerges as a key player in cancer and future studies regarding Rab35 should shed light on how this GTPase contributes to oncogenesis, linking the importance of membrane trafficking to tumorigenesis.

Acknowledgements

Work from A. Echard's laboratory received financial support from Institut Pasteur, the Centre National de la Recherche Scientifique (CNRS), Institut National du Cancer (INCa grant 2014-1-PLB IO-04) and ANR (grant AbsyStem 15-CE13-0001-02 and grant CYTOSIGN 16-CE13-004-02).

References

- Bryant DM, Mostov KE. From cells to organs: building polarized tissue. *Nat Rev Mol Cell Biol.* 2008;9(11):887-901.
- Rodriguez-Boulan E, Macara IG. Organization and execution of the epithelial polarity programme. *Nat Rev Mol Cell Biol.* 2014;15(4):225-242.
- Mosesson Y, Mills GB, Yarden Y. Derailed endocytosis: an emerging feature of cancer. *Nat Rev Cancer.* 2008;8(11):835-850.
- Sigismund S, Confalonieri S, Ciliberto A, Polo S, Scita G, Di Fiore PP. Endocytosis and signaling: cell logistics shape the eukaryotic cell plan. *Physiol Rev.* 2012;92(1):273-366.
- De Franceschi N, Hamidi H, Alanko J, Sahgal P, Ivaska J. Integrin traffic - the update. *J Cell Sci.* 2015;128(5):839-852.
- Ioannou MS, McPherson PS. Rab-mediated membrane trafficking and the control of epithelial cell polarity. *J Cell Biol.* 2016;213(3):301-303.
- Zhen Y, Stenmark H. Cellular functions of Rab GTPases at a glance. *J Cell Sci.* 2015;128(17):3171-3176.
- Wandinger-Ness A, Zerial M. Rab proteins and the compartmentalization of the endosomal system. *Cold Spring Harb Perspect Biol.* 2014;6(11):a022616.
- Pylypenko O, Hammich H, Yu IM, Houdusse A. Rab GTPases and their interacting protein partners: Structural insights into Rab functional diversity. *Small GTPases.* 2017;1-27.
- Tomas A, Futter CE, Eden ER. EGF receptor trafficking: consequences for signaling and cancer. *Trends Cell Biol.* 2014;24(1):26-34.
- Wheeler DB, Zoncu R, Root DE, Sabatini DM, Sawyers CL. Identification of an oncogenic RAB protein. *Science.* 2015;350(6257):211-217.
- Jopling HM, Odell AF, Hooper NM, Zachary IC, Walker JH, Ponnambalam S. Rab GTPase regulation of VEGFR2 trafficking and signaling in endothelial cells. *Arterioscler Thromb Vasc Biol.* 2009;29(7):1119-1124.
- Abella JV, Parachoniak CA, Sangwan V, Park M. Dorsal ruffle microdomains potentiate Met receptor tyrosine kinase signaling and down-regulation. *J Biol Chem.* 2010;285(32):24956-24967.
- Muller PA, Caswell PT, Doyle B, et al. Mutant p53 drives invasion by promoting integrin recycling. *Cell.* 2009;139(7):1327-1341.
- Christoforides C, Rainero E, Brown KK, Norman JC, Toker A. PKD controls alphavbeta3 integrin recycling and tumor cell invasive migration through its substrate Rabaptin-5. *Dev Cell.* 2012;23(3):560-572.
- Rush JS, Quinalty LM, Engelman L, Sherry DM, Ceresa BP. Endosomal accumulation of the activated epidermal growth factor receptor (EGFR) induces apoptosis. *J Biol Chem.* 2012;287(1):712-722.
- Shaughnessy R, Retamal C, Oyanadel C, et al. Epidermal growth factor receptor endocytic traffic perturbation by phosphatidate phosphohydrolase inhibition: new strategy against cancer. *FEBS J.* 2014;281(9):2172-2189.
- Kouranti I, Sachse M, Arouche N, Goud B, Echard A. Rab35 regulates an endocytic recycling pathway essential for the terminal steps of cytokinesis. *Curr Biol.* 2006;16(17):1719-1725.
- Langemeyer L, Nunes Bastos R, Cai Y, Itzen A, Reinisch KM, Barr FA. Diversity and plasticity in Rab GTPase nucleotide release mechanism has consequences for Rab activation and inactivation. *Elife.* 2014;3:e01623.
- Heo WD, Inoue T, Park WS, et al. PI(3,4,5)P3 and PI(4,5)P2 lipids target proteins with polybasic clusters to the plasma membrane. *Science.* 2006;314(5804):1458-1461.
- Klinkert K, Echard A. Rab35 GTPase: A Central Regulator of Phosphoinositides and F-actin in Endocytic Recycling and Beyond. *Traffic.* 2016;17(10):1063-1077.

22. Minowa-Nozawa A, Nozawa T, Okamoto-Furuta K, Kohda H, Nakagawa I. Rab35 GTPase recruits NPD52 to autophagy targets. *EMBO J*. 2017;36(18):2790-2807.
23. Klinkert K, Rocancourt M, Houdusse A, Echard A. Rab35 GTPase couples cell division with initiation of epithelial apico-basal polarity and lumen opening. *Nat Commun*. 2016;7:11166.
24. Mrozowska PS, Fukuda M. Regulation of podocalyxin trafficking by Rab small GTPases in 2D and 3D epithelial cell cultures. *J Cell Biol*. 2016;213(3):355-369.
25. Dambournet D, Machicoane M, Chesneau L, et al. Rab35 GTPase and OCRL phosphatase remodel lipids and F-actin for successful cytokinesis. *Nat Cell Biol*. 2011;13(8):981-988.
26. Fremont S, Hammich H, Bai J, et al. Oxidation of F-actin controls the terminal steps of cytokinesis. *Nat Commun*. 2017;8:14528.
27. Cauvin C, Rosendale M, Gupta-Rossi N, et al. Rab35 GTPase Triggers Switch-like Recruitment of the Lowe Syndrome Lipid Phosphatase OCRL on Newborn Endosomes. *Curr Biol*. 2016;26(1):120-128.
28. Kobayashi H, Fukuda M. Rab35 regulates Arf6 activity through centaurin-beta2 (ACAP2) during neurite outgrowth. *J Cell Sci*. 2012;125(Pt 9):2235-2243.
29. Kobayashi H, Fukuda M. Rab35 establishes the EHD1-association site by coordinating two distinct effectors during neurite outgrowth. *J Cell Sci*. 2013;126(Pt 11):2424-2435.
30. Kobayashi H, Etoh K, Fukuda M. Rab35 is translocated from Arf6-positive perinuclear recycling endosomes to neurite tips during neurite outgrowth. *Small GTPases*. 2014;5:e29290.
31. Chevallier J, Koop C, Srivastava A, Petrie RJ, Lamarche-Vane N, Presley JF. Rab35 regulates neurite outgrowth and cell shape. *FEBS Lett*. 2009;583(7):1096-1101.
32. Villarroel-Campos D, Henriquez DR, Bodaleo FJ, et al. Rab35 Functions in Axon Elongation Are Regulated by P53-Related Protein Kinase in a Mechanism That Involves Rab35 Protein Degradation and the Microtubule-Associated Protein 1B. *J Neurosci*. 2016;36(27):7298-7313.
33. Chesneau L, Dambournet D, Machicoane M, et al. An ARF6/Rab35 GTPase cascade for endocytic recycling and successful cytokinesis. *Curr Biol*. 2012;22(2):147-153.
34. Walseng E, Bakke O, Roche PA. Major histocompatibility complex class II-peptide complexes internalize using a clathrin- and dynamin-independent endocytosis pathway. *J Biol Chem*. 2008;283(21):14717-14727.
35. Yang CW, Hojer CD, Zhou M, et al. Regulation of T Cell Receptor Signaling by DENND1B in TH2 Cells and Allergic Disease. *Cell*. 2016;164(1-2):141-155.
36. Patino-Lopez G, Dong X, Ben-Aissa K, et al. Rab35 and its GAP EPI64C in T cells regulate receptor recycling and immunological synapse formation. *J Biol Chem*. 2008;283(26):18323-18330.
37. Egami Y, Fukuda M, Araki N. Rab35 regulates phagosome formation through recruitment of ACAP2 in macrophages during FcγR-mediated phagocytosis. *J Cell Sci*. 2011;124(Pt 21):3557-3567.
38. Deschamps C, Echard A, Niedergang F. Phagocytosis and cytokinesis: do cells use common tools to cut and to eat? Highlights on common themes and differences. *Traffic*. 2013;14(4):355-364.
39. Mukherjee S, Liu X, Arasaki K, McDonough J, Galan JE, Roy CR. Modulation of Rab GTPase function by a protein phosphocholine transferase. *Nature*. 2011;477(7362):103-106.
40. Dikshit N, Bist P, Fenlon SN, et al. Intracellular Uropathogenic E. coli Exploits Host Rab35 for Iron Acquisition and Survival within Urinary Bladder Cells. *PLoS Pathog*. 2015;11(8):e1005083.
41. Charrasse S, Comunale F, De Rossi S, Echard A, Gauthier-Rouviere C. Rab35 regulates cadherin-mediated adherens junction formation and myoblast fusion. *Mol Biol Cell*. 2013;24(3):234-245.
42. Hsu C, Morohashi Y, Yoshimura S, et al. Regulation of exosome secretion by Rab35 and its GTPase-activating proteins TBC1D10A-C. *J Cell Biol*. 2010;189(2):223-232.
43. Biesemann A, Gorontzi A, Barr F, Gerke V. Rab35 protein regulates evoked exocytosis of endothelial Weibel-Palade bodies. *J Biol Chem*. 2017;292(28):11631-11640.
44. Marat AL, McPherson PS. The connectenn family, Rab35 guanine nucleotide exchange factors interfacing with the clathrin machinery. *J Biol Chem*. 2010;285(14):10627-10637.
45. Allaire PD, Marat AL, Dall'Armi C, Di Paolo G, McPherson PS, Ritter B. The Connectenn DENN domain: a GEF for Rab35 mediating cargo-specific exit from early endosomes. *Mol Cell*. 2010;37(3):370-382.
46. Wu X, Bradley MJ, Cai Y, et al. Insights regarding guanine nucleotide exchange from the structure of a DENN-domain protein complexed with its Rab GTPase substrate. *Proc Natl Acad Sci U S A*. 2011;108(46):18672-18677.
47. Yoshimura S, Gerondopoulos A, Linford A, Rigden DJ, Barr FA. Family-wide characterization of the DENN domain Rab GDP-GTP exchange factors. *J Cell Biol*. 2010;191(2):367-381.
48. Fuchs E, Haas AK, Spooner RA, Yoshimura S, Lord JM, Barr FA. Specific Rab GTPase-activating proteins define the Shiga toxin and epidermal growth factor uptake pathways. *J Cell Biol*. 2007;177(6):1133-1143.
49. Davey JR, Humphrey SJ, Junutula JR, et al. TBC1D13 is a RAB35 specific GAP that plays an important role in GLUT4 trafficking in adipocytes. *Traffic*. 2012;13(10):1429-1441.

50. Uytterhoeven V, Kuenen S, Kasproicz J, Miskiewicz K, Verstreken P. Loss of skywalker reveals synaptic endosomes as sorting stations for synaptic vesicle proteins. *Cell*. 2011;145(1):117-132.
51. Chaîneau M, Ioannou MS, McPherson PS. Rab35: GEFs, GAPs and effectors. *Traffic*. 2013;14(11):1109-1117.
52. Crockett DK, Lin Z, Elenitoba-Johnson KS, Lim MS. Identification of NPM-ALK interacting proteins by tandem mass spectrometry. *Oncogene*. 2004;23(15):2617-2629.
53. Abe Y, Takeuchi T, Imai Y, et al. A Small Ras-like protein Ray/Rab1c modulates the p53-regulating activity of PRPK. *Biochem Biophys Res Commun*. 2006;344(1):377-385.
54. Giridharan SS, Caplan S. MICAL-family proteins: Complex regulators of the actin cytoskeleton. *Antioxid Redox Signal*. 2014;20(13):2059-2073.
55. Fremont S, Romet-Lemonne G, Houdusse A, Echard A. Emerging roles of MICAL family proteins - from actin oxidation to membrane trafficking during cytokinesis. *J Cell Sci*. 2017;130(9):1509-1517.
56. Zack TI, Schumacher SE, Carter SL, et al. Pan-cancer patterns of somatic copy number alteration. *Nat Genet*. 2013;45(10):1134-1140.
57. Fujiwara T, Bandi M, Nitta M, Ivanova EV, Bronson RT, Pellman D. Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null cells. *Nature*. 2005;437(7061):1043-1047.
58. Allaire PD, Seyed Sadr M, Chaîneau M, et al. Interplay between Rab35 and Arf6 controls cargo recycling to coordinate cell adhesion and migration. *J Cell Sci*. 2013;126(Pt 3):722-731.
59. Jones J, Otu H, Spentzos D, et al. Gene signatures of progression and metastasis in renal cell cancer. *Clin Cancer Res*. 2005;11(16):5730-5739.
60. Sato M, Sato K, Liou W, Pant S, Harada A, Grant BD. Regulation of endocytic recycling by *C. elegans* Rab35 and its regulator RME-4, a coated-pit protein. *EMBO J*. 2008;27(8):1183-1196.
61. Gao Y, Balut CM, Bailey MA, Patino-Lopez G, Shaw S, Devor DC. Recycling of the Ca²⁺-activated K⁺ channel, KCa_{2.3}, is dependent upon RME-1, Rab35/EPI64C, and an N-terminal domain. *J Biol Chem*. 2010;285(23):17938-17953.
62. Shah M, Baterina OY, Jr., Taupin V, Farquhar MG. ARH directs megalin to the endocytic recycling compartment to regulate its proteolysis and gene expression. *J Cell Biol*. 2013;202(1):113-127.
63. Argenzio E, Margadant C, Leyton-Puig D, et al. CLIC4 regulates cell adhesion and beta1 integrin trafficking. *J Cell Sci*. 2014;127(Pt 24):5189-5203.
64. Zheng J, Duan B, Sun S, Cui J, Du J, Zhang Y. Folliculin Interacts with Rab35 to Regulate EGF-Induced EGFR Degradation. *Front Pharmacol*. 2017;8:688.
65. Laviolette LA, Mermoud J, Calvo IA, et al. Negative regulation of EGFR signalling by the human folliculin tumour suppressor protein. *Nat Commun*. 2017;8:15866.
66. Nookala RK, Langemeyer L, Pacitto A, et al. Crystal structure of folliculin reveals a hidDenn function in genetically inherited renal cancer. *Open Biol*. 2012;2(8):120071.
67. Nandez R, Balkin DM, Messa M, et al. A role of OCRL in clathrin-coated pit dynamics and uncoating revealed by studies of Lowe syndrome cells. *Elife*. 2014;3:e02975.
68. Rahajeng J, Giridharan SS, Cai B, Naslavsky N, Caplan S. MICAL-L1 is a tubular endosomal membrane hub that connects Rab35 and Arf6 with Rab8a. *Traffic*. 2012;13(1):82-93.
69. Donaldson JG, Johnson DL, Dutta D. Rab and Arf G proteins in endosomal trafficking and cell surface homeostasis. *Small GTPases*. 2016;7(4):247-251.
70. D'Souza-Schorey C, Chavrier P. ARF proteins: roles in membrane traffic and beyond. *Nat Rev Mol Cell Biol*. 2006;7(5):347-358.
71. Dutta D, Donaldson JG. Sorting of Clathrin-Independent Cargo Proteins Depends on Rab35 Delivered by Clathrin-Mediated Endocytosis. *Traffic*. 2015;16(9):994-1009.
72. Giridharan SS, Cai B, Vitale N, Naslavsky N, Caplan S. Cooperation of MICAL-L1, syndapin2, and phosphatidic acid in tubular recycling endosome biogenesis. *Mol Biol Cell*. 2013;24(11):1776-1790, S1771-1715.
73. Kobayashi H, Etoh K, Ohbayashi N, Fukuda M. Rab35 promotes the recruitment of Rab8, Rab13 and Rab36 to recycling endosomes through MICAL-L1 during neurite outgrowth. *Biol Open*. 2014;3(9):803-814.
74. Takahashi Y, Tanikawa C, Miyamoto T, et al. Regulation of tubular recycling endosome biogenesis by the p53-MICALL1 pathway. *Int J Oncol*. 2017;51(2):724-736.
75. Nonaka R, Miyake Y, Hata T, et al. Circulating miR-103 and miR-720 as novel serum biomarkers for patients with colorectal cancer. *Int J Oncol*. 2015;47(3):1097-1102.
76. Jones CI, Zabolotskaya MV, King AJ, et al. Identification of circulating microRNAs as diagnostic biomarkers for use in multiple myeloma. *Br J Cancer*. 2012;107(12):1987-1996.
77. Tang Y, Lin Y, Li C, et al. MicroRNA-720 promotes in vitro cell migration by targeting Rab35 expression in cervical cancer cells. *Cell Biosci*. 2015;5:56.
78. Duan B, Cui J, Sun S, et al. EGF-stimulated activation of Rab35 regulates RUSC2-GIT2 complex formation to stabilize GIT2 during directional lung cancer cell migration. *Cancer Lett*. 2016;379(1):70-83.
79. Zhu Y, Shen T, Liu J, et al. Rab35 is required for Wnt5a/Dvl2-induced Rac1 activation and cell migration in MCF-7 breast cancer cells. *Cell Signal*. 2013;25(5):1075-1085.

80. Deng W, Wang Y, Gu L, et al. MICAL1 controls cell invasive phenotype via regulating oxidative stress in breast cancer cells. *BMC Cancer*. 2016;16:489.
81. Qin X, Wang J, Wang X, Liu F, Jiang B, Zhang Y. Targeting Rabs as a novel therapeutic strategy for cancer therapy. *Drug Discov Today*. 2017;22(8):1139-1147.
82. Stein MP, Dong J, Wandinger-Ness A. Rab proteins and endocytic trafficking: potential targets for therapeutic intervention. *Adv Drug Deliv Rev*. 2003;55(11):1421-1437.
83. Greaves M, Maley CC. Clonal evolution in cancer. *Nature*. 2012;481(7381):306-313.




Figure 1. Rab35 functions in intracellular signaling, apico-basal polarity, cytokinesis and cell migration. Rab35's functions in these processes are described. Moreover, known and possible effectors involved in each process are indicated.

Table 1. Pathogenic Rab35 somatic mutations found in human tumor samples according to the Catalogue of Somatic Mutations in Cancer (COSMIC) database. We indicated synonymous mutations as they can be sometimes pathogenic.

Tissue	Amino acid change	Mutation
Bone	V187V	Synonymous
Brain	V90I	Missense
Breast	A29V F45L G18V	Missense Missense Missense
Cervix	F45F	Synonymous
Colon	A139V E94K F33F G80E G83R G140R R27C R71C R79Q R127Q R196Q S150S V188M -	Missense Missense Synonymous Missense Missense Missense Missense Missense Missense Missense Missense Synonymous Missense Intronic substitution
Endometrium	A151T S22N	Missense Missense
Esophagus	I14I V97V	Synonymous Synonymous
Hematopoietic and lymphoid	A151T	Missense
Liver	A65V	Missense
Lung	A29V F70F F161L R101Q	Missense Synonymous Missense Missense
Pancreas	G80R	Missense
Prostate	N156S	Missense
Stomach	R27H R196*	Missense Nonsense
Thyroid	G18S	Missense
Upper aerodigestive tract	E159Q T191M	Missense Missense