

The Volume-limited A-Star (VAST) survey - II. Orbital motion monitoring of A-type star multiples

R. J. de Rosa, J. Patience, Arthur Vigan, P. A. Wilson, A. Schneider, N. J. Mcconnell, S. J. Wiktorowicz, C. Marois, I. Song, B. Macintosh, et al.

► To cite this version:

R. J. de Rosa, J. Patience, Arthur Vigan, P. A. Wilson, A. Schneider, et al.. The Volume-limited A-Star (VAST) survey - II. Orbital motion monitoring of A-type star multiples. Monthly Notices of the Royal Astronomical Society, 2012, 422 (4), pp.2765-2785. 10.1111/j.1365-2966.2011.20397.x . hal-02118392

HAL Id: hal-02118392 https://hal.science/hal-02118392

Submitted on 2 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Mon. Not. R. Astron. Soc. 422, 2765-2785 (2012)

doi:10.1111/j.1365-2966.2011.20397.x

The Volume-limited A-Star (VAST) survey – II. Orbital motion monitoring of A-type star multiples

R. J. De Rosa,^{1*†} J. Patience,¹ A. Vigan,¹ P. A. Wilson,¹ A. Schneider,² N. J. McConnell,³ S. J. Wiktorowicz,^{3,4} C. Marois,⁵ I. Song,² B. Macintosh,⁶ J. R. Graham,^{3,7} M. S. Bessell,⁸ R. Dovon⁹ and O. Lai¹⁰

¹College of Engineering, Mathematics and Physical Sciences, University of Exeter, Physics Building, Stocker Road, Exeter EX4 40L

²Physics and Astronomy, University of Georgia, 240 Physics, Athens, GA 30602, USA

³Department of Astronomy, University of California at Berkeley, Berkeley, CA 94720, USA

⁴Department of Astronomy, University of California at Santa Cruz, Santa Cruz, CA 95064, USA

⁵NRC Herzberg Institute of Astrophysics, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

⁶Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550, USA

⁷ Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St George Street, Toronto, ON M55 3H8, Canada

⁸Mount Stromlo and Siding Spring Observatories, Institute of Advanced Studies, The Australian National University, Weston Creek P.O., ACT 2611, Australia

⁹Dépt de Physique, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, QC H3C 3J7, Canada

¹⁰Canda–France–Hawaii Telescope, 65-1238 Mamalahoa Highway, Kamuela, HI 96745, USA

Accepted 2011 December 14. Received 2011 December 13; in original form 2011 November 29

ABSTRACT

As a part of our ongoing Volume-limited A-Star (VAST) adaptive optics survey, we have obtained observations of 26 binary systems with projected separations <100 au, 13 of which have sufficient historical measurements to allow for refinement of their orbital elements. For each system with an estimated orbit, the dynamical system mass obtained was compared with the system mass estimated from mass-magnitude relations. Discrepancies between the dynamical and theoretical system mass can be explained by the presence of a previously unresolved spectroscopic component or by a non-solar metallicity of the system. Using this approach to infer the presence of additional companions, a lower limit to the fraction of binaries, triples and quadruples can be estimated as 39, 46 and 15 per cent for systems with at least one companion within 100 au. The fraction of multiple systems with three or more components shows a relative increase compared to the fraction for solar-type primaries resolved in previous volume-limited surveys. The observations have also revealed a pair of potentially young (<100 Myr) M dwarf companions, which would make an ideal benchmark for the theoretical models during the pre-main-sequence contraction phase for M dwarfs. In addition to those systems with orbit fits, we report 13 systems for which further orbital monitoring observations are required, 11 of which are newly resolved as a part of the VAST survey.

Key words: techniques: high angular resolution - binaries: close - binaries: general.

*E-mail: derosa@astro.ex.ac.uk

†Based on observations obtained at the Canada-France-Hawaii Telescope (CFHT) under programme IDs 2001AF11, 2008AC22, 2009BC06, 2010AC14 and 2011AC11. Based on observations made with ESO telescopes at the La Silla Paranal Observatory under programme IDs 272.D-5068(A), 074.D-0180(A), 076.C-0270(A), 076.D-0108(A), 077.D-0147(A) and 080.D-0348(A). Based on observations obtained at the Gemini Observatory under programme IDs GN2008A-Q-74, GN-2008B-Q-119 and GN-2010A-Q-75.

1 INTRODUCTION

The orbits of binary stars offer one of the few techniques to determine stellar masses, or masses and radii in cases with a favourable geometry (e.g. Andersen et al. 1991). Orbits of binaries with young ages are particularly important to test theoretical evolutionary models, and examples include a double-lined eclipsing binary in Orion (e.g. Stassun et al. 2008). Low-mass stars and brown dwarfs also represent a regime requiring empirical calibration, and the visual orbits of nearby M, L and T dwarfs have been used to measure system masses and compare with theoretical mass-luminosity relations (Dupuy, Liu & Bowler 2009).



Visual orbits also provide a method to search for evidence of additional components and determine higher order multiplicity, by identifying systems with dynamical masses significantly in excess of the theoretical predictions. These visual binaries with an indication of unresolved companions can be monitored with spectroscopy or interferometry to determine the properties such as period and mass ratio of the closer pair and augment the statistics compiled from catalogues (Tokovinin 2008). The properties of higher order multiple systems represent tests of formation scenarios including fragmentation of cores (e.g. Pringle 1989; Bonnell 2001) and discs (e.g. Stamatellos, Hubber & Whitworth 2007; Kratter, Murray-Clay & Youdin 2010) and may be influenced by processes such as accretion (e.g. Bate 2000) and dynamical interactions (e.g. McDonald & Clarke 1995; Lodato et al. 2007)

In this paper, a subset of the systems resolved by our ongoing Volume-limited A-Star (VAST) survey are used to determine dynamical system masses from orbit fits and to compare the results with theoretical models and search for additional unresolved stellar companions. The sample of adaptive optics (AO)-imaged binaries considered in this study is detailed in Section 2, and a short summary of the new observations is given in Section 3. The data analysis, including the AO image processing to determine the relative positions and the subsequent orbit determination from the compilation of position data, is explained in Section 4. Section 5 reports the astrometric results from the new measurements and the orbital elements and masses based on the orbit fits. The discussion in Section 6 covers a comparison with theoretical mass-magnitude relations, an assessment of the higher order multiplicity and the identification of a set of targets for continued monitoring. Finally, Section 7 provides a summary and future directions for the project.

2 SAMPLE

The sample of binaries is drawn from the ongoing VAST survey (De Rosa et al. 2011), an AO imaging survey of A stars within 75 pc, and includes the 26 systems with projected separations less than 100 au (Fig. 1). The angular separations of the binaries range from 0.094 to 4.66 arcsec, and 11 are newly resolved. Fig. 2 plots the measured magnitude difference as a function of separation for the complete sample. Of the 26 systems, 13 have a substantial number of previous measurements, and these systems comprise the orbit subsample. For the remaining 13 systems, there is insufficient coverage to fit an orbit, and these binaries comprise the monitoring subsample. Table 1 lists each observed binary in the two subsamples, along with basic parameters for each such as distance (van Leeuwen 2007), Tycho2 $V_{\rm T}$ -band and Two Micron All Sky Survey (2MASS) $K_{\rm s}$ -band photometry (ESA 1997; Skrutskie et al. 2006), and spectral type listed within the SIMBAD data base.

For the brightest stars within the sample, the shortest 2MASS exposures saturate, requiring a different method for measuring the photometry, resulting in significantly larger uncertainties on the estimated magnitudes (Skrutskie et al. 2006). Therefore for three of these brighter targets, near-infrared photometry was obtained from alternative sources (Morel & Magnenat 1978; Ducati 2002) and converted into the 2MASS photometric bands using empirical colour transformations (Carpenter 2001). The distribution of the sample on the colour–magnitude diagram (CMD) is plotted in Fig. 1. The contamination from the secondary to the apparent magnitude differences measured within this study. Given the rapid evolution of massive stars off the main sequence, the position of an A star on the CMD provides a method to estimate the age of the system based



Figure 1. A CMD of the 27 stars discussed within this work, plotted alongside three different sets of theoretical isochrones at 100, 500 and 800 Myr. Those targets with new or refined orbits are plotted in blue with a diamond symbol (see Table 4), and those targets for which further measurements are required are plotted in red with a cross symbol (see Table 5 and Section 6.3). Three different sets of theoretical isochrones are plotted at ages of 100, 500 and 750 Myr; (top) Lejeune & Schaerer (2001), (middle) Marigo et al. (2008) and (bottom) Siess, Dufour & Forestini (2000).

on a comparison with theoretical isochrones. The inferred age of the system from the CMD is combined with the dynamical system mass from the orbit and system photometry from the literature to test mass–magnitude relations at the corresponding age.

3 OBSERVATIONS

With AO systems operating on telescopes ranging in diameter from the 3-m Shane to the 8-m Gemini and Very Large Telescope (VLT), near-infrared images were obtained on all targets. In most cases, the filter used for the observations was a narrow- or broad-band filter within the *K* bandpass, though some images were taken within the *J* and *H* bandpasses. Both the primary and secondary of the pairs were unsaturated in the AO images, simplifying the astrometry measurements. Table 2 details the key characteristics of the instruments used to acquire the new AO observations, with the measured pixel scale and orientation for each camera. A subset of the observations were obtained from the Canada–France–Hawaii Telescope (CFHT) and ESO Science Archive Facilities. One measurement obtained at the Southern Observatory for Astrophysical Research (SOAR) as a part



Figure 2. The magnitude difference between primary and secondary for each binary system within this study as a function of angular separation. The targets are divided into two distinct subsamples, those with new or refined orbits estimated within this study (filled circles) and those for which further measurements are required before an orbital determination can be attempted (filled stars). The majority of the systems within the second subsample were newly resolved as a part of the VAST survey, demonstrating the higher dynamic range possible with AO imaging. The filled circles connected with a solid line represent companions within the same hierarchical triple (HIP 44127 and HIP 82321).

of the VAST survey, and used within this study, has been recently published in Hartkopf, Tokovinin & Mason (2012).

4 DATA ANALYSIS

4.1 AO image processing

The AO science images obtained were processed with standard image reduction steps including dark subtraction, flat-fielding, interpolation over bad pixels and sky subtraction. To align all the images, the centroid of the bright primary was obtained in each exposure by fitting a Gaussian to the core of the central point spread function (PSF). For each system resolved within the observations, an empirical PSF was determined from the radial profile of the primary after masking any close companion. The empirical PSF was then fitted to the position and intensity of both components of the system, providing a measure of the separation, position angle and magnitude difference. Uncertainties within the photometry and astrometry were estimated from the standard deviations of the photometric and astrometric measurements from each individual exposure before combination.

To ensure accurate determination of the separation and position angle, the pixel scale and orientation of the detectors were calibrated based on observations of the Trapezium cluster, with the exception of data obtained with IRCAL and Palomar High Angular Resolution Observer (PHARO). Depending on the total field of view of the detector, the positions of 20–40 Trapezium members were compared

HIP	Name	HR	HD	ADS	Spectral type	Distance (pc)	V _T (mag)	K _s (mag)	
Orbit subsample									
5300	v Phe	331	6767		A3IV	57.0 ± 2.0	5.230 ± 0.001	4.78 ± 0.02	
9480	48 Cas	575	12111	1598	A3IV	35.3 ± 0.6	4.533 ± 0.002	4.08 ± 0.13^{a}	
11569	ι Cas	707	15089	1860	A5p	40.7 ± 1.3	4.496 ± 0.003	4.25 ± 0.03	
17954		1188	23985	2799	A2V+	56.5 ± 1.8	5.259 ± 0.004	4.81 ± 0.02	
28614	μ Ori	2124	40932	4617		47.5 ± 1.5	4.150 ± 0.002	3.64 ± 0.26	
36850	Castor	2891/2890	60178J	6175	A2Vm	15.6 ± 0.9	1.590 ± 0.020	1.47 ± 0.03^a	
44127	ι UMa	3569	76644		A7V	14.51 ± 0.03	3.159 ± 0.002	2.67 ± 0.03^b	
47479		3863	84121		A3IV	72.6 ± 2.2	5.334 ± 0.003	4.80 ± 0.02	
76952	γ CrB	5849	140436	9757	B9IV+	44.8 ± 1.0	3.819 ± 0.002	3.67 ± 0.23	
77660	b Ser	5895	141851		A3Vn	49.8 ± 0.8	5.112 ± 0.002	4.70 ± 0.02	
80628	v Oph	6129	148367		A3m	41.0 ± 1.5	4.657 ± 0.003	4.17 ± 0.04	
82321	52 Her	6254	152107	10227	A2Vspe	55.3 ± 1.0	4.833 ± 0.003	4.57 ± 0.02	
93506	ζ Sgr	7194	176687	11950	A2.5Va	27.0 ± 0.6	2.617 ± 0.003	2.29 ± 0.23	
				Monito	oring subsampl	e			
128			224890		Am	70.8 ± 1.7	6.521 ± 0.004	6.02 ± 0.02	
2381		118	2696		A3V	53.1 ± 0.8	5.188 ± 0.003	4.83 ± 0.02	
2852	BG Cet	151	3326		A5m	48.9 ± 0.8	6.093 ± 0.004	5.42 ± 0.01	
5310	79 Psc	328	6695		A3V	47.3 ± 1.7	5.581 ± 0.003	5.22 ± 0.02	
18217		1192	24141		A5m	50.5 ± 1.1	5.806 ± 0.004	5.37 ± 0.02	
29852		2265	43940		A2V	61.9 ± 1.0	5.895 ± 0.003	5.44 ± 0.02	
51384		4062	89571		F0IV	40.6 ± 0.6	5.549 ± 0.003	4.85 ± 0.02	
65241	64 Vir	5040	116235		A2m	65.9 ± 1.4	5.897 ± 0.003	5.62 ± 0.02	
66223		5108	118156	8956	F0IV	69.8 ± 2.0	6.394 ± 0.004	5.88 ± 0.02	
103298	16 Del	8012	199254	14429	A4V	60.5 ± 1.1	5.555 ± 0.003	5.19 ± 0.02	
109667		8464	210739		A3V	63.5 ± 2.1	6.209 ± 0.005	5.74 ± 0.02	
110787	ρ^1 Cep	8578	213403		A2m	63.2 ± 0.9	5.857 ± 0.004	5.54 ± 0.03	
116611	75 Peg	8963	222133		A1Vn	71.4 ± 1.4	5.483 ± 0.003	5.42 ± 0.02	

Table 1. Basic properties of sample members.

 ^{a}K -band photometry from Ducati (2002).

^bK-band photometry from Morel & Magnenat (1978).

Telescope (instrument)	Date	Filter	Plate scale (mas pixel ⁻¹)	True north (°)	Proposal ID
CFHT	2001 May 5-7	H continuum	34.87 ± 0.14	1.47 ± 0.19	2001AF11
(KIR)	2001 May 6	J continuum	34.82 ± 0.15	1.51 ± 0.17	2001AF11
	2008 June 13	H ₂ ($v = 1 - 0$)	34.81 ± 0.10	-2.42 ± 0.13	2008AC22
	2008 June 14	Feп	34.76 ± 0.08	-2.40 ± 0.11	2008AC22
	2009 August 30 to September 1	H ₂ ($v = 1 - 0$)	34.78 ± 0.10	-2.40 ± 0.20	2009BC06
	2010 February 4-5	H ₂ ($v = 1 - 0$)	34.75 ± 0.08	-2.37 ± 0.11	2010AC14
	2011	H ₂ ($v = 1 - 0$)	34.77 ± 0.06	-2.35 ± 0.09	2011AC11
Gemini N	2008 September 8 to November 14	Brγ	21.28 ± 0.12	0.52 ± 0.29	GN-2008A-Q-74
(NIRI)	2009 December 19	Brγ	21.28 ± 0.12	0.52 ± 0.29	GN-2008B-Q-119
	2010 June 25 to August 26	Brγ	21.28 ± 0.12	0.52 ± 0.29	GN-2010A-Q-75
Lick	2008 July 24 to 2009 July 25	Brγ	74.55 ± 1.00	1.68 ± 0.50	-
(IRCAL)	2008 October 16-17	Brγ	74.55 ± 1.00	1.68 ± 0.50	-
VLT-UT4	2004 June 30	IB218	27.02 ± 0.14	0.24 ± 0.30	272.D-5068(A)
(NACO)	2005 January 10	IB218	27.02 ± 0.15	0.26 ± 0.30	074.D-0180(A)
	2005 February 8	Ks	27.02 ± 0.15	0.26 ± 0.30	074.D-0180(A)
	2005 November 8	Ks	26.98 ± 0.08	0.21 ± 0.20	076.C-0270(A)
	2005 November 10	Ks	26.98 ± 0.08	0.21 ± 0.20	076.D-0108(A)
	2005 December 6	IB218	26.98 ± 0.08	0.21 ± 0.20	076.D-0108(A)
	2006 January 4	K _s	27.01 ± 0.13	0.31 ± 0.29	076.D-0108(A)
	2006 January 7	IB218	27.01 ± 0.13	0.31 ± 0.29	076.D-0108(A)
	2006 April 27	IB218	27.00 ± 0.12	0.09 ± 0.26	077.D-0147(A)
	2007 September 20	IB218	26.99 ± 0.12	-0.23 ± 0.21	080.D-0348(A)
	2007 September 20	Ks	26.99 ± 0.12	-0.23 ± 0.21	080.D-0348(A)
	2007 September 26	IB218	26.99 ± 0.12	-0.23 ± 0.21	080.D-0348(A)
	2007 November 4	Ks	27.00 ± 0.10	-0.18 ± 0.20	080.D-0348(A)
	2008 February 15	IB218	27.01 ± 0.14	-0.12 ± 0.29	080.D-0348(A)
	2008 February 24	Ks	27.01 ± 0.14	-0.12 ± 0.29	080.D-0348(A)
Palomar	2008 April 11	CH_4	25.00 ± 1.00	0.70 ± 1.00	-
(PHARO)	2008 April 12	Brγ	25.00 ± 1.00	0.70 ± 1.00	-
	2008 July 12	CH_4	25.00 ± 1.00	0.70 ± 1.00	-
	2008 July 13	Brγ	25.00 ± 1.00	0.70 ± 1.00	-

Table 2. Observing run details with calculated astrometric calibration values.

Notes. KIR – (Doyon et al. 1998); NIRI (Near-InfraRed Imager and Spectrometer) – (Hodapp et al. 2003); IRCAL – (Lloyd et al. 2000); NACO (Nasmyth Adaptive Optics System/Near-Infrared Imager and Spectrograph) – (Lenzen et al. 2003; Rousset et al. 2003); PHARO (Palomar High Angular Resolution Observer) – (Hayward et al. 2001).

with the coordinates reported in McCaughrean & Stauffer (1994). The average derived pixel scale and orientation was computed, and the standard deviation of these values was used as the associated errors; the results are given in Table 2. For the data obtained with IRCAL and PHARO, the pixel scale and orientation were calibrated from binary systems also observed with instruments calibrated with Trapezium measurements.

4.2 Orbital determination

For the 13 binaries with sufficient coverage of the orbit, a fit was performed for the orbital elements, and an estimate of the dynamical mass was determined. The measurements presented within this study were combined with previous measurements contained within the Washington Double Star (WDS; Mason et al. 2001) catalogue. These archive measurements were obtained using a variety of observational techniques and date back to the 18th century. As in some cases the statistical uncertainties were not provided in the WDS catalogue, we searched the literature for the formal errors for each individual measurement, and only separation and position angle values for which uncertainties could be assigned were included within the fitting procedure. A detailed listing of the individual measurements used for the orbital determination will be made available at the Strasbourg astronomical Data Center (CDS; http://cds.u-strasbg.fr).

Our orbit fitting approach utilizes the method presented by Hilditch (2001), and demonstrated by an application to measurements of the T Tau S system (Köhler et al. 2008). This method is similar to the grid-based search technique developed by Hartkopf, McAlister & Franz (1989). At each epoch of observation t_i , the x_i, y_i position of the companion with respect to the primary is measured in the observed tangent plane. These values are related to the true position of the secondary in the orbital plane (x'_i, y'_i) through the following equations:

$$x_i = Ax'_i + Fy'_i,$$

$$y_i = Bx'_i + Gy'_i,$$
(1)

where A, B, F and G are the orbital Thiele–Innes elements, with

 $A = a(\cos \omega \cos \Omega - \sin \omega \sin \Omega \cos i),$ $B = a(\cos \omega \sin \Omega + \sin \omega \cos \Omega \cos i),$ $F = a(-\sin \omega \cos \Omega - \cos \omega \sin \Omega \cos i),$ $G = a(-\sin \omega \sin \Omega + \cos \omega \cos \Omega \cos i),$ (2)

where *a* is the semimajor axis of the orbit, ω the longitude of periastron, Ω the longitude of the ascending node and *i* the inclination – four of the seven orbital elements. The position of the companion in the orbital plane (x'_i, y'_i) can also be expressed through the remaining orbital elements (e, P, T_0) as

$$\begin{aligned} x'_i &= \cos E - e, \\ y'_i &= \sqrt{1 - e^2} \sin E, \end{aligned} \tag{3}$$

where e is the eccentricity of the system. The eccentric anomaly (E) can be determined from a numerical solution to Kepler's equation,

$$M = E - e \sin E$$

= $(2\pi/P)(t_i - T_0),$ (4)

where *M* is the mean anomaly, *P* the period of the system and T_0 the epoch of periastron passage. At each epoch of observation, the position of the component in the orbital plane can be defined using just three of the orbital elements (*e*, *P* and T_0). The orbital position at each epoch can then be converted into the observed position using the equations in equation (1) through a least-squares determination of the four Thiele–Innes elements.

An initial estimation of the orbital parameters of each system can be determined through an iterative three-dimensional grid search of e, P and T_0 . A wider range of parameter values were searched, with 100 linear steps searched over a range of $0 \le e < 1,500$ linear steps in the range $1 \le \log (P/yr) \le 3$ and T_0 initially distributed between the years 2000.0 - (P/2) and 2000.0 + (P/2). At each position within this three-dimensional grid, the fit orbital positions (x'_i, y'_i) were directly calculated (equation 3), with the four remaining orbital parameters (a, i, ω, Ω) estimated from a least-squares fit to the observed positions using equations (1) and (2). After the χ^2 statistic was calculated at each position within the grid, the range of T_0 values searched was reduced by a factor of 10 centred on the optimum value of T_0 found within the previous iteration. This process was repeated until the step size in T_0 was reduced to less than 1 d. The values for a, i, ω and Ω can be determined from an inversion of equation (2) (Green 1985):

$$\Omega = \frac{1}{2} \left[\arctan\left(\frac{B-F}{G+A}\right) - \arctan\left(\frac{B+F}{G-A}\right) \right],$$

$$\omega = \left[\arctan\left(\frac{B-F}{G+A}\right) - \Omega \right],$$

$$i = 2 \arctan\left[\sqrt{\frac{-(B+F)\sin(\omega+\Omega)}{(B-F)\sin(\omega-\Omega)}}\right],$$

$$a = \frac{B-F}{\sin(\omega+\Omega)[1+\cos(i)]}.$$
(5)

The orbital parameters calculated at each position within the (*P*, *e*) grid are then used as a starting point for a Levenberg–Marquardt minimization to ensure the minimum of the χ^2 distribution is found. The set of orbital parameters with the minimum χ^2 statistic was then used as the orbit solution for the system.

The shape of the χ^2 distribution in the vicinity of the global minimum can be used to determine the 1σ uncertainties of each parameter (Press et al. 1992). By perturbing an individual parameter away from the global minimum, and optimizing the remaining parameters, a region of the χ^2 distribution can be calculated where the χ^2

Table 3. Summary of theoretical model grids.

Grid reference	Mass range (M_{\odot})	Metallicity (Z)
Lejeune & Schaerer (2001)	0.80 < M < 5.0	0.02
Marigo et al. (2008)	0.15 < M < 5.0	0.02
Siess et al. (2000)	0.10 < M < 5.0	0.02
Baraffe et al. (1998)	0.08 < M < 1.5	0.02

statistic is less than χ^2_{min} + 1. This region encloses 68 per cent of the probability distribution and is not necessarily symmetric about the minimum χ^2 value. Our implementation of the orbit fitting method was tested against four well-studied systems (Liu, Dupuy & Ireland 2008; Bonnefoy et al. 2009; Dupuy, Liu & Bowler 2009), with the resulting parameters being within the published 1 σ uncertainties.

4.3 Theoretical mass-magnitude relations

We obtained four different grids of evolutionary models with which the dynamical system masses estimated from the fitted orbital



Figure 3. Mass–magnitude relations were constructed from each of the four model grids. The evolution of the mass–magnitude relation is shown for each grid, with the 10-Myr (black line), 100-Myr (blue line) and 1-Gyr (red line) relations plotted. For reference the dashed line indicates the faintest A-type star within the sample, and the dotted line indicates the zero-age main-sequence magnitude of an M0 star. A-type stars typically have an absolute *K*-band magnitude ranging between $M_K = 0$ and 2, where the mass is significantly dependent on the age of the star. Within the Lejeune & Schaerer (2001) and Marigo et al. (2008) models, the mass–magnitude relation of stars fainter than $M_K \sim 2$ is not dependent on the age of the star. The Siess et al. (2000) and Baraffe et al. (1998) models include a description of the contraction phase of lower mass stars on to the main sequence during the early portion of its life.

2770 *R. J. De Rosa et al.*

Table 4.	Measured	binary j	position	angle and	separation	for previously	resolved syste	ms.
----------	----------	----------	----------	-----------	------------	----------------	----------------	-----

HIP	Comp.	WDS designation	Discoverer designation	Previous orbit	Instrument	Epoch	θ (°)	ρ (arcsec)	ΔK (mag)
5300	ΔR	01078_/129	R\$T3352	Söderhielm (1000)	NACO	2005.02	15.73 ± 0.76	0.089 ± 0.001	0.507 ± 0.020
5500	AD	01070-4127	K515552	Sodernjenn (1999)	NACO	2005.02	13.73 ± 0.70 126.69 ± 0.38	0.009 ± 0.001 0.114 ± 0.001	$a_{0.507} \pm 0.020$
9480	A-B	02020 ± 7054	BU 513	Mason Douglass & Hartkonf (1999)	IRCAL	2008.79	293.97 ± 0.30	0.679 ± 0.001	1.110 ± 0.056
2100	n b	02020 703	DC 515	Muson, Dougluss & Markopi (1999)	KIR	2009.67	297.36 ± 0.22	0.666 ± 0.002	1.172 ± 0.013
					NIRI	2010.65	302.11 ± 0.29	0.644 ± 0.004	$a_{1.137} \pm 0.007$
11569	Aa. Ab	02291+6724	CHR 6	Drummond et al. (2003)	IRCAL	2008.79	43.50 ± 0.80	0.575 ± 0.006	$a_{1.906 \pm 0.026}$
					KIR	2010.10	41.52 ± 2.61	0.602 ± 0.023	1.771 ± 0.301
17954	AB	03503+2535	STT 65	Docobo & Ling (2007)	NIRI	2008.87	194.76 ± 0.31	0.200 ± 0.001	$a0.232 \pm 0.005$
28614	AB	06024+0939	A 2715	Muterspaugh et al. (2008)	NIRI	2009.96	21.95 ± 0.29	0.403 ± 0.002	$a^{a}1.008 \pm 0.006$
					KIR	2011.31	21.31 ± 1.06	0.397 ± 0.006	0.896 ± 0.120
36850	AB	07346+3153	STF1110	Docobo & Costa (1985)	KIR	2010.09	57.46 ± 0.14	4.663 ± 0.013	$a0.698 \pm 0.050$
44127	BC	08592+4803	HU 628	Eggen (1967)	PHARO	2008.28	242.01 ± 1.04	0.433 ± 0.017	$a0.005 \pm 0.031$
					KIR	2010.10	228.98 ± 0.27	0.568 ± 0.003	0.027 ± 0.023
					KIR	2011.30	223.37 ± 0.31	0.656 ± 0.003	-0.011 ± 0.039
47479	AB	09407-5759	B 780	Söderhjelm (1999)	NACO	2008.12	300.22 ± 0.58	0.099 ± 0.001	$^{a}0.082 \pm 0.017$
76952	AB	15427+2618	STF1967	Muterspaugh et al. (2010)	PHARO	2008.28	112.65 ± 1.07	0.683 ± 0.028	_
					KIR	2011.31	111.37 ± 0.33	0.620 ± 0.005	$^{a}1.056 \pm 0.030$
77660	AB	15513-0305	CHR 51	Docobo et al. (2010)	KIR	2001.34	82.72 ± 2.42	0.188 ± 0.018	_
					KIR	2001.34	81.97 ± 1.59	0.197 ± 0.008	-
					KIR	2001.35	83.38 ± 3.67	0.185 ± 0.016	-
					NACO	2004.50	71.87 ± 0.31	0.247 ± 0.001	$a^{a}1.487 \pm 0.006$
					NACO	2006.32	69.74 ± 0.42	0.274 ± 0.003	1.472 ± 0.097
					KIR	2010.10	66.69 ± 0.47	0.330 ± 0.004	1.483 ± 0.058
					KIR	2011.30	64.21 ± 2.57	0.340 ± 0.014	1.455 ± 0.194
80628	Ab,Ab	16278-0822	RST3949	-	PHARO	2008.28	22.56 ± 1.05	0.674 ± 0.027	$a^{a}2.072 \pm 0.066$
					IRCAL	2008.56	23.53 ± 0.71	0.685 ± 0.009	1.978 ± 0.057
					KIR	2011.31	34.37 ± 0.29	0.771 ± 0.003	2.237 ± 0.038
82321	BC	16492+4559	A 1866	Popovic (1969)	PHARO	2008.53	247.33 ± 1.03	0.286 ± 0.012	-
					KIR	2011.30	260.51 ± 0.53	0.301 ± 0.005	" 0.299 ± 0.044
93506	AB	19026-2953	HDO 150	Mason et al. (1999)	KIR	2008.45	31.05 ± 2.31	0.194 ± 0.010	-
					KIR	2011.31	285.59 ± 0.71	0.304 ± 0.004	$a0.320 \pm 0.043$

^aPhotometry measurement used to determine the magnitude of each component in Table 6.

Table 5. Measured binary position angle and separation for potential orbital monitoring targets.

HIP	Instrument	Epoch	θ (°)	ρ (arcsec)	ΔK (mag)	Projected separation (au)
128	NIRI	2008.72	80.64 ± 0.29	0.983 ± 0.006	$a_{3.521 \pm 0.021}$	69.59 ± 1.72
	IRCAL	2008.79	80.73 ± 0.52	0.975 ± 0.013	3.355 ± 0.037	68.98 ± 1.90
2381	NACO	2005.93	279.51 ± 0.26	1.765 ± 0.006	$a6.206 \pm 0.088$	93.75 ± 1.48
	NACO	2007.73	279.13 ± 0.22	1.767 ± 0.008	6.229 ± 0.055	93.86 ± 1.51
	NIRI	2008.79	278.54 ± 0.29	1.759 ± 0.013	7.401 ± 0.711	93.39 ± 1.59
2852	NIRI	2008.79	260.55 ± 0.30	0.931 ± 0.006	$a^{a}5.001 \pm 0.056$	45.51 ± 0.82
5310	NIRI	2008.79	174.99 ± 0.30	0.357 ± 0.002	$a3.719 \pm 0.023$	16.88 ± 0.62
18217	IRCAL	2008.79	64.78 ± 0.65	1.037 ± 0.012	2.301 ± 0.029	52.37 ± 1.24
	NIRI	2008.87	64.96 ± 0.29	1.033 ± 0.006	$a2.41 \pm 0.010$	52.13 ± 1.12
29852	NACO	2005.85	210.76 ± 0.22	0.218 ± 0.001	$a^{a}2.003 \pm 0.013$	13.49 ± 0.21
	NACO	2005.86	210.80 ± 0.21	0.217 ± 0.001	1.983 ± 0.012	13.44 ± 0.21
51384	PHARO	2008.18	212.43 ± 1.05	2.078 ± 0.084	$a4.543 \pm 0.169$	84.44 ± 3.66
65241	NACO	2005.10	196.95 ± 0.62	0.328 ± 0.003	3.061 ± 0.231	21.65 ± 0.50
	NACO	2008.15	212.75 ± 0.57	0.258 ± 0.003	$a^{a}3.025 \pm 0.097$	17.01 ± 0.42
66223	PHARO	2008.53	187.71 ± 1.02	1.381 ± 0.056	$a^{a}5.660 \pm 0.105$	96.42 ± 4.73
103298	NIRI	2008.69	115.72 ± 0.30	0.220 ± 0.001	$a^{a}2.962 \pm 0.007$	13.29 ± 0.25
109667	NIRI	2008.69	285.17 ± 0.30	1.117 ± 0.007	$a^{a}3.997 \pm 0.011$	70.88 ± 2.38
	KIR	2009.66	284.66 ± 0.25	1.108 ± 0.006	4.051 ± 0.073	70.28 ± 2.35
	NIRI	2010.48	284.38 ± 0.29	1.101 ± 0.007	4.044 ± 0.103	69.84 ± 2.34
110787	NIRI	2008.71	211.11 ± 0.35	0.291 ± 0.002	$a^{a}3.891 \pm 0.043$	18.38 ± 0.29
116611	NIRI	2008.75	173.11 ± 0.32	0.950 ± 0.006	$a^{a}5.930 \pm 0.089$	67.78 ± 1.38
	NIRI	2010.48	172.11 ± 0.34	0.943 ± 0.006	5.794 ± 0.076	67.31 ± 1.34

^aPhotometry measurement used to determine the magnitude of each component in Table 6.

Table 6. Derived parameters of individual components.

HIP	Comp.	V _T (mag)	K _S (mag)	V - K	M_V (mag)	M_K (mag)	Additional components	SB type	Reference
					Orbit subsample	e			
5300	А	5.51 ± 0.02	528 ± 0.02	0.23 ± 0.03	1.73 ± 0.08	1.50 ± 0.08			
2200	B	6.85 ± 0.02	5.20 ± 0.02 5.85 ± 0.02	0.29 ± 0.09 0.99 ± 0.08	3.07 ± 0.11	2.08 ± 0.08			
9480 ^a	Ā	4.69 ± 0.01	4.40 ± 0.13	0.29 ± 0.13	1.95 ± 0.03	1.67 ± 0.13			
	В	6.72 ± 0.01	5.54 ± 0.13	1.18 ± 0.13	3.98 ± 0.04	2.80 ± 0.13			
11569 ^b	Aa	4.67 ± 0.01	4.79 ± 0.03	-0.12 ± 0.03	1.62 ± 0.07	1.74 ± 0.08			
	Ab	8.65 ± 0.16	6.71 ± 0.03	1.94 ± 0.16	5.60 ± 0.17	3.66 ± 0.08			
17954	А	5.81 ± 0.04	5.46 ± 0.02	0.35 ± 0.04	2.05 ± 0.08	1.70 ± 0.07			
	В	6.26 ± 0.06	5.69 ± 0.02	0.57 ± 0.06	2.50 ± 0.09	1.93 ± 0.07			
28614	А	4.32 ± 0.01	4.00 ± 0.26	0.33 ± 0.26	0.94 ± 0.07	0.61 ± 0.27	Aa, Ab	SB1	Fekel et al. (2002)
	В	6.22 ± 0.02	5.01 ± 0.26	1.22 ± 0.26	2.84 ± 0.07	1.62 ± 0.27	Ba, Bb	SB2	Fekel et al. (2002)
36850	А	1.98 ± 0.02	1.93 ± 0.03	0.05 ± 0.04	1.02 ± 0.13	0.97 ± 0.13	Aa, Ab	SB1	Vinter Hansen et al. (1940)
	В	2.88 ± 0.03	2.63 ± 0.04	0.25 ± 0.05	1.92 ± 0.13	1.66 ± 0.13	Ba, Bb	SB1	Vinter Hansen et al. (1940)
44127	A	3.16 ± 0.02	2.71 ± 0.03	0.45 ± 0.04	2.35 ± 0.02	1.90 ± 0.03			
	В	10.88 ± 0.11	6.90 ± 0.04	3.98 ± 0.12	10.07 ± 0.11	6.09 ± 0.04			
17.170	C	11.08 ± 0.12	6.90 ± 0.04	4.18 ± 0.12	10.27 ± 0.12	6.09 ± 0.04			
4/4/9	A	5.82 ± 0.01	5.51 ± 0.02	0.31 ± 0.02	1.51 ± 0.07	1.21 ± 0.07			
76052	В	6.45 ± 0.01	5.59 ± 0.02	0.85 ± 0.03	2.14 ± 0.07	1.29 ± 0.07			
70932	A D	4.03 ± 0.01 5.60 ± 0.02	4.02 ± 0.23 5.07 ± 0.23	0.03 ± 0.23 0.52 \pm 0.23	0.80 ± 0.03	0.70 ± 0.23 1.82 ± 0.23			
77660	D A	5.00 ± 0.02 5.21 ± 0.01	3.07 ± 0.23	0.35 ± 0.23	2.33 ± 0.03 1.72 ± 0.04	1.82 ± 0.23 1.46 ± 0.04			
//000	R	5.21 ± 0.01 7.82 ± 0.04	4.94 ± 0.02 6.43 ± 0.02	0.20 ± 0.02 1 38 ± 0.04	1.72 ± 0.04 4.33 ± 0.05	1.40 ± 0.04 2.95 ± 0.04			
80628	Δ 2	7.82 ± 0.04 4.68 ± 0.01	0.43 ± 0.02	1.33 ± 0.04 0.37 ± 0.04	4.53 ± 0.03 1.62 ± 0.08	2.95 ± 0.04 1 25 ± 0.09	Aa1 Aa2	SB2	Gutmann (1965)
00020	Ab	4.00 ± 0.01 8.80 ± 0.10	4.32 ± 0.04 6.39 ± 0.07	2.41 ± 0.12	5.74 ± 0.03	1.23 ± 0.09 3.33 ± 0.11	Adl, Ad2	502	Outmain (1903)
82321	A	4.87 ± 0.01	4.73 ± 0.07	0.14 ± 0.02	1.16 ± 0.04	1.02 ± 0.04			
02021	B	9.19 ± 0.11	7.33 ± 0.03	1.86 ± 0.11	5.47 ± 0.12	3.62 ± 0.05			
	C	9.29 ± 0.11	7.63 ± 0.04	1.66 ± 0.12	5.57 ± 0.12	3.92 ± 0.05			
93506	A	3.27 ± 0.01	2.90 ± 0.23	0.37 ± 0.23	1.11 ± 0.05	0.74 ± 0.24			
	В	3.48 ± 0.01	3.22 ± 0.24	0.26 ± 0.24	1.32 ± 0.05	1.06 ± 0.24			
				М	onitoring subsam	nple			
128	AB	-	6.06 ± 0.02	-	-	1.81 ± 0.06	Α, Β	SB1	Carquillat et al. (2003)
	С	-	9.58 ± 0.03	-	-	5.33 ± 0.06			
2381	A	-	4.83 ± 0.02	-	-	1.21 ± 0.04			
2052	В	-	11.04 ± 0.09	-	-	7.41 ± 0.10			
2852	A	_	5.43 ± 0.02	-	_	1.98 ± 0.04			
5210	В	-	10.43 ± 0.06	-	-	0.98 ± 0.07			
5510	A B	_	3.23 ± 0.02 8.07 ± 0.03	_	_	1.88 ± 0.08 5.60 ± 0.08			
18217	Δ	_	5.97 ± 0.03 5.48 ± 0.02	_	_	3.00 ± 0.08 1.97 ± 0.05			
10217	B	_	7.90 ± 0.02	_	_	4.38 ± 0.05			
29852	A	_	5.60 ± 0.02	_	_	1.64 ± 0.04			
27032	B	_	7.60 ± 0.02	_	_	3.64 ± 0.04			
51384	Ā	_	4.87 ± 0.02	_	_	1.82 ± 0.04			
	В	_	9.41 ± 0.17	_	_	6.37 ± 0.17			
65241	А	_	5.69 ± 0.02	_	_	1.59 ± 0.05			
	В	_	8.71 ± 0.09	_	_	4.62 ± 0.11			
66223	Aa	_	5.89 ± 0.02	_	_	1.67 ± 0.06			
	Ab	_	11.55 ± 0.11	-	_	7.33 ± 0.12			
103298	Aa	_	5.26 ± 0.02	-	_	1.35 ± 0.04			
	Ab	-	8.22 ± 0.02	-	-	4.31 ± 0.04			
109667	А	_	5.76 ± 0.02	-	_	1.75 ± 0.07			
	В	-	9.76 ± 0.02	-	-	5.75 ± 0.07			
110787	A	_	5.57 ± 0.03	-	_	1.57 ± 0.04			
11///	В	-	9.46 ± 0.05	-	-	5.46 ± 0.06			D 111 1 (2005)
116611	Aa	-	5.42 ± 0.02	-	-	1.16 ± 0.05	Aa1, Aa2	SBI	Rucinski et al. (2005)
	AD	_	11.33 ± 0.09	_	_	1.09 ± 0.10			

^aThe spectroscopic binary reported by Abt (1965) is the system resolved within the AO data.

^bThe individual component magnitudes may be significantly biased due to the presence of additional companions within the resolution limit of the Tycho2 and 2MASS observations.



Figure 4. The age of each system can be estimated based on the position of each component on the CMD [top – Lejeune & Schaerer (2001), middle – Marigo et al. (2008) and bottom – Siess et al. (2000)]. This procedure, while demonstrated for only two systems in this figure, was repeated for all the remaining members of the orbit subsample. For each system, the age estimated from each of the three model grids is reported in Table 10.

parameters were compared. Table 3 lists the mass range, metallicity and literature reference for each of the four grids. The Lejeune & Schaerer (2001), Marigo et al. (2008) and Siess et al. (2000) grids cover a significant portion of the lifespan of a typical A-type star, and as such we applied a maximum age cut-off at 1 Gyr. In addition to these grids, models from Baraffe et al. (1998) were obtained in order to study a pair of lower mass companions presented in Section 6.1.2. Each grid was converted into the photometric systems used within this study – Tycho V and 2MASS K_S (Carpenter 2001), before producing a high-resolution ($dM/M_{\odot} = 0.001$) mass–magnitude relation, created through cubic interpolation of the grid data, as shown in Fig. 3.

The absolute V- and K-band magnitudes were calculated from the V-band magnitude differences obtained from the literature and the K-band magnitude differences presented within this study (Tables 4 and 5). The individual component magnitudes and V - Kcolour for each system are presented in Table 6. Within the A-type star mass range, the mass–magnitude relations significantly change



Figure 5. The mass-magnitude relation for a hypothetical binary system with $M_K(A) = 1.25 \pm 0.05$ and $M_K(B) = 5.50 \pm 0.10$, as a function of system age. The evolution of the mass-magnitude relation is shown for both components (A – blue region, B – red region) and the system as a whole (grey region). The extent of the region in each case represents the uncertainty in the mass estimate due to the uncertainties of the magnitudes of each component.

Table 7. Visual magnitude differences for a subset of the sample.

HIP	ΔV	Reference
5300	1.34 ± 0.10^{a}	Horch, Ninkov & Franz (2001)
9480	2.03 ± 0.01	Fabricius & Makarov (2000)
11569	3.98 ± 0.16	Christou & Drummond (2006)
17954	0.45 ± 0.10^{a}	Horch, Meyer & van Altena (2004)
28614	1.90 ± 0.02	Fabricius & Makarov (2000)
36850	0.90 ± 0.03	Worley (1969)
44127 A,BC	7.06 ± 0.10^a	Baize & Petit (1989)
44127 BC	0.20 ± 0.10^a	Mason et al. (2001)
47479	0.63 ± 0.02	Mason et al. (2001)
76952	1.55 ± 0.02	Fabricius & Makarov (2000)
77660	2.61 ± 0.04	Docobo et al. (2010)
80628	4.12 ± 0.10^a	Mason et al. (2001)
82321 A,BC	3.61 ± 0.10^a	Mason et al. (2001)
83231 BC	0.10 ± 0.10^a	Mason et al. (2001)
93506	0.21 ± 0.02	Fabricius & Makarov (2000)

^{*a*}For ΔV measurements without uncertainties, 0.10 is used.

as a function of the age of the system due to the rapid evolution of A-type stars across the CMD. The age of each system is therefore estimated, based on the position of the primary on the CMD (Fig. 4), before a mass for each component is estimated from the mass-magnitude relations. The estimated masses of each component are summed to produce an estimation of the system mass, hereafter called the theoretical system mass.

To demonstrate the analysis procedure, a hypothetical binary system of magnitudes $M_K(A) = 1.25$ and $M_K(B) = 5.50$ was used to construct the mass–age relation for each component and their corresponding sum (Fig. 5). The evolution of the mass–magnitude relation as a function of age can be then visualized as a continuous function for both components within the system. For this example a set of models were used which include the pre-main-sequence contraction phase of lower mass stars, as demonstrated by the increase in the derived mass as a function of age for the companion. The two mass–age curves can then be summed to produce a mass–age relation for the system in question. Using the age estimated for the

Table 8. Estimated orbital parameters and corresponding 1σ uncertainties.

HIP	N. meas.	Period	Semimajor axis	Inclination	Longitude of node	Epoch of periastron	Eccentricity	Longitude of periastron	System mass
		<i>P</i> (yr)	a (arcsec)	<i>i</i> (°)	Ω (°)	T_0 (yr)	е	ω (°)	(M_{\bigodot})
5300	19	$28.36\substack{+0.04 \\ -0.04}$	$0.2396^{+0.0005}_{-0.0005}$	$65.3_{-0.3}^{+0.3}$	$142.7^{+0.3}_{-0.3}$	$2007.94^{+0.02}_{-0.02}$	$0.424^{+0.002}_{-0.002}$	$333.9^{+0.4}_{-0.3}$	3.16 ± 0.34
9480	66	$61.14_{-0.05}^{+0.05}$	$0.614_{-0.002}^{+0.002}$	$16.7^{+0.9}_{-0.9}$	$48.2^{+3.3}_{-3.1}$	$1964.35\substack{+0.09\\-0.09}$	$0.355\substack{+0.001\\-0.001}$	$19.5^{+3.5}_{-3.7}$	2.72 ± 0.13
11569	11	$50.2^{+1.0}_{-1.0}$	$0.429^{+0.007}_{-0.007}$	$149.0^{+1.7}_{-1.6}$	$180.0^{+2.7}_{-2.8}$	$1993.24_{-0.08}^{+0.08}$	$0.642^{+0.009}_{-0.009}$	$331.3^{+2.6}_{-2.7}$	2.12 ± 0.25
17954	44	$61.2_{-0.1}^{+0.1}$	$0.442^{+0.002}_{-0.002}$	$84.7_{-0.1}^{+0.1}$	$26.4^{+0.2}_{-0.2}$	$1998.30\substack{+0.08\\-0.08}$	$0.628\substack{+0.002\\-0.002}$	$340.3_{-0.7}^{+0.7}$	4.15 ± 0.39
28614	69	$18.641\substack{+0.008\\-0.008}$	$0.2741\substack{+0.0001\\-0.0001}$	$96.59^{+0.02}_{-0.02}$	$24.76_{-0.02}^{+0.02}$	$2003.742^{+0.004}_{-0.004}$	$0.744^{+0.001}_{-0.001}$	$217.08^{+0.08}_{-0.08}$	6.36 ± 0.62
36850	207	$466.8^{+6.3}_{-6.1}$	$6.78\substack{+0.05\\-0.05}$	$113.56\substack{+0.09\\-0.09}$	$41.2_{-0.1}^{+0.1}$	$1957.3_{-0.3}^{+0.3}$	$0.333\substack{+0.007\\-0.006}$	$249.3_{-0.5}^{+0.6}$	5.42 ± 0.97
44127	9	$39.4_{-0.1}^{+0.1}$	$0.70\substack{+0.01\\-0.01}$	$111.6^{+1.1}_{-1.1}$	$24.5^{+1.2}_{-1.3}$	$1999.1_{-0.7}^{+0.6}$	$0.35\substack{+0.02\\-0.02}$	$354.2^{+4.4}_{-4.7}$	0.68 ± 0.04
47479	5	$10.74_{-0.04}^{+0.04}$	$0.1207\substack{+0.0006\\-0.0006}$	$128.4^{+1.5}_{-1.4}$	$87.4_{-0.5}^{+0.5}$	$2005.89\substack{+0.03\\-0.04}$	$0.365\substack{+0.009\\-0.009}$	$18.6^{+1.1}_{-1.2}$	5.83 ± 0.53
76952	104	$91.2_{-0.4}^{+0.4}$	$0.729^{+0.006}_{-0.006}$	$94.45_{-0.07}^{+0.07}$	$111.75_{-0.09}^{+0.09}$	$1931.6_{-0.3}^{+0.3}$	$0.48\substack{+0.01\\-0.01}$	$103.8_{-0.5}^{+0.6}$	4.19 ± 0.30
80628	8	$82.8^{+1.5}_{-1.3}$	$0.79_{-0.02}^{+0.03}$	$31.2^{+5.4}_{-6.0}$	$86.8^{+8.1}_{-5.6}$	$1994.1_{-1.1}^{+0.9}$	$0.45\substack{+0.03\\-0.03}$	$177.9^{+6.8}_{-7.9}$	4.99 ± 0.75
82321	12	$56.4_{-0.3}^{+0.3}$	$0.279^{+0.005}_{-0.005}$	$37.4_{-2.8}^{+2.6}$	$57.5^{+5.9}_{-6.1}$	$1991.2^{+2.3}_{-2.1}$	$0.13^{+0.02}_{-0.02}$	$69.4^{+8.0}_{-7.5}$	1.16 ± 0.09
93506	42	$21.00\substack{+0.01 \\ -0.01}$	$0.489\substack{+0.001\\-0.001}$	$111.1\substack{+0.1 \\ -0.1}$	$74.0\substack{+0.1 \\ -0.1}$	$2005.99\substack{+0.03\\-0.03}$	$0.211\substack{+0.001\\-0.001}$	$7.2^{+0.6}_{-0.6}$	5.26 ± 0.37

system based upon its position on the CMD, a theoretical system mass can be estimated and compared with the dynamical system mass determined from the orbital elements.

5 RESULTS

5.1 Astrometric results

The astrometry and photometry measurements of the two subsamples are given in Tables 4 and 5. Both tables contain the *Hipparcos* designation of the primary, the components of the system under investigation, the instrument and epoch of observation, and the measured astrometric values with the corresponding uncertainties. For the orbit subsample, the WDS designation and discoverer code are also listed for reference.

Combining the K-band magnitudes of the sample with the ΔK values reported in Tables 4 and 5, and the Hipparcos parallax, allowed for an estimation of the K-band apparent and absolute magnitudes of the resolved components (Table 6). For systems with ΔV measurements reported within the literature (Table 7), the corresponding estimated V-band apparent and absolute magnitudes and V - K colours for each resolved component are reported. Seven members of the overall sample are hierarchical systems with at least one of the components resolved within our data set consisting of multiple subcomponents, indicated in Table 6. An example of this is HIP 128; our AO data are able to resolve a previously unknown binary companion within this system (HIP 128 C at ~1.0 arcsec), but are of insufficient angular resolution to resolve the previously known spectroscopic component HIP 128 B. Without an estimate of the ΔV or ΔK between HIP 128 A and HIP 128 B, the individual magnitudes cannot be estimated, and therefore only the blended magnitudes of the two components are listed.

5.2 Orbital elements and dynamical masses

The orbital parameters for each system with sufficient orbital coverage are listed in Table 8, alongside a system mass estimated from Kepler's third law (hereafter the dynamical system mass) and the number of measurements used to fit the orbit. In Fig. 6, the refined orbital fits incorporating the new data are plotted along with the previously reported orbit (references listed within Table 4). The resulting orbits span a range of periods from 10.78 to 467.4 yr and a range of semimajor axes between 0.12 and 6.78 arcsec. The relatively short period of each system allows for orbital motion to be resolved over very short baselines, typically of the order of months. The changes to the estimated dynamical system mass between the previously published orbit fit and the refined fit presented within this study are shown in Table 9.

6 DISCUSSION

We have presented high-resolution observations obtained for 26 nearby multiple systems with A-type primaries with projected separations within 100 au. The subset of 12 targets with orbit fits have been further divided into four distinct categories primarily based on a comparison between the dynamical mass and the theoretical mass estimated from the mass-magnitude relations, as shown in Fig. 7. The theoretical mass estimates for each system using the Lejeune & Schaerer (2001), Marigo et al. (2008) and Siess et al. (2000) grids are shown in Table 10. Those systems with only two known components are discussed in Section 6.1.1, where the importance of metallicity is described. Two hierarchical systems resolved within our data are discussed in Sections 6.1.2 and 6.1.3, which allow for a comparison to the models within the K- to M-type spectral range. For systems with a dynamical mass excess, significantly higher than the mass predicted from the theoretical mass-magnitude relations, the subset with known spectroscopic components is discussed in Section 6.2.1, and we present three systems with evidence suggestive of an additional unresolved component in Section 6.2.2. The remaining targets are discussed in the context of continued monitoring of the orbital motion in Section 6.3; of these, 11 are newly resolved as a part of the VAST survey, two were resolved in recent multiplicity surveys of nearby southern A-type stars (Ivanov et al. 2006; Ehrenreich et al. 2010) and one resolved within a large speckle interferometry survey (McAlister et al. 1987).

6.1 Comparison to theoretical models

6.1.1 A-type binaries

Four targets within the orbit subsample are systems where the two known components have been resolved within our high-resolution data: HIP 5300, HIP 9480, HIP 11569 and HIP 76952. For each of



Figure 6. Combining our high-resolution observations with historical measurements, refined orbits for 12 binary systems are plotted. The previous orbital fit, obtained from the Sixth Orbit Catalogue, is plotted for reference with a dashed line. Each plot uses a similar symbol scheme to the Sixth Orbit Catalogue, with open blue circles representing eyepiece interferometry, filled blue circles speckle interferometry, green crosses micrometrical observations and violet asterisks photographic measurements. Our high-resolution observations presented within this study are plotted as filled red stars. For each measurement, the corresponding O - C line is plotted, showing the difference between expected and actual position within the orbital path. Symbols in grey represent those measurements presented without formal errors, and are not used while estimating the orbital parameters. Within each plot, the 57-mas-radius black disc represents the resolution limit for *K*-band observations at an 8-m telescope.



Figure 6 – continued

these four targets, the mass-magnitude relations were used to determine how the system mass changes as a function of our estimate of the system age, shown in Fig. 8. For one system, HIP 76952, the dynamical system mass is consistent with the theoretical system

mass (Fig. 8, top panel). The ages of the systems estimated from the position of the primary on the CMD are consistent with their position within the local interstellar bubble (LIB). The minimum age of a star within the LIB, excluding those with relatively high

A6 (orcseconds)

Að (arcseconds)

<u>A</u>6 (orcseconds)

Table 9. Previous and revised dynamical system masses.

HIP	Old estimate (M⊙)	New estimate (M _☉)
5300	3.45	3.16 ± 0.34
9480	2.98 ± 0.39	2.72 ± 0.13
11569	2.12 ± 0.33	2.12 ± 0.25
17954	3.37 ± 0.35	4.15 ± 0.39
28614	6.33 ± 0.62	6.36 ± 0.62
36850	5.51	5.42 ± 0.97
44127	0.61 ± 0.03	0.68 ± 0.04
47479	7.50	5.83 ± 0.53
76952	4.14 ± 0.28	4.19 ± 0.30
80628	_	4.99 ± 0.75
82321	1.00	1.16 ± 0.09
93506	5.20 ± 0.36	5.26 ± 0.37



Figure 7. Using the method introduced in Section 4.4, a comparison can be made between the dynamical mass determined from the orbit and the mass estimated from theoretical mass–magnitude relations for the stars within the orbit subsample [the Marigo et al. (2008) models are used for this example]. The systems with an A-type star primary which are known to consist only of two components are denoted as black points (Section 6.1.1). Two hierarchical systems were fully resolved with our data, and the lower mass pair of each system are in green (Section 6.1.2). The systems which have a significantly discrepant dynamical mass can be explained by the presence of an unresolved spectroscopic companion within our data. The targets with known spectroscopic components are plotted in red (Section 6.2.1), while those systems with evidence suggesting a previously unknown spectroscopic component are plotted in blue (Section 6.2.2). The dashed line denotes the equivalence between the dynamical and theoretical mass.

space motions (e.g. the β Pic moving group – Ortega et al. 2002), has been shown to be 160 Myr (Abt 2011).

The dynamical system masses of the three remaining systems are consistently lower than their theoretical system masses. One possible explanation for the apparently low dynamical masses is a non-solar metallicity. Varying the metallicity has a significant effect on the system age estimate and the mass–magnitude relations. As an example, a $2 M_{\odot}$ star with supersolar metallicity will be more luminous and have a redder V - K colour index than a similar-mass star of solar metallicity. A supersolar metallicity star will appear to be significantly older based on its position on a CMD if solar metallicity models are used. To explore the effect of our

assumption of solar metallicity for the entire sample, the HIP 5300 system (Fig. 8, second panel) was studied at varying metallicity values. Using the solar metallicity models, the dynamical system mass of this system is significantly lower than the theoretical system mass (Fig. 9, top panel). Increasing the metallicity causes the star to appear both younger, and less massive (Fig. 9, bottom panel). Only eight of the targets included within this study have metallicity measurements, either from spectroscopic analysis (e.g. Erspamer & North 2003) or derived from Strömgren photometry (e.g. Song et al. 2001), demonstrating the need for further study in this area.

6.1.2 K- and M-type binaries – HIP 44127

The detection of two hierarchical systems, with pairs of lower mass companions in a wide orbit around an A-type primary, allows for a comparison of the theoretical models in the low-mass regime where the models differ in the treatment of the contraction phase of these objects (e.g. Fig. 3). Our observation of the HIP 44127 system, shown in Fig. 10, resolves three components to this hierarchical system, with an A star primary (A) separated by ~4 arcsec from two fainter, gravitationally bound, companions (BC).

The orbit presented in the Sixth Orbit Catalogue of the BC pair around the A-type primary is in disagreement with our recent observations of this system. Although the phase coverage is insufficient for a robust orbital determination, the high proper motion of the primary ($\Delta \alpha = -441.1 \text{ mas yr}^{-1}$, $\Delta \delta = -215.2 \text{ mas yr}^{-1}$) suggests that the BC pair is comoving. In addition, radial velocity variations detected within the spectra of the primary suggest the presence of a spectroscopic component to this system with a period of $\sim 11 \text{ vr}$ (Abt 1965). The separation of this component was anticipated to be between 0.2 and 0.6 arcsec, reaching maximum separation in the middle of 2007 (Docobo & Andrade 2006). We do not resolve any companion in any epoch of our observations which are consistent with these predictions, placing an upper limit to the separation of 0.08, 0.10 and 0.10 arcsec in 2008, 2010 and 2011, respectively. The magnitude difference between the primary and the suggested companion, estimated to be $\Delta m = 1.2$, is well within our detection limits at the expected separation (Docobo & Andrade 2006).

A refined orbit of HIP 44127 BC is presented in Table 8 and Fig. 6, with an estimated dynamical system mass of $0.68 \pm 0.04 \, M_{\odot}$. The small magnitude difference between the two components ($\Delta K \sim 0$) suggests a mass ratio close to unity. Assuming individual masses of $\sim 0.34 \,\mathrm{M_{\odot}}$, the components would be of early M to mid-M spectral type (Baraffe & Chabrier 1996), a region of particular disagreement between theoretical models. The Baraffe et al. (1998) and Siess et al. (2000) models both predict that for a binary consisting of two stars of magnitudes equal to the measured magnitudes of the two components ($M_K = 5.87$), the system mass will increase from 0.3 to $0.9 \,\mathrm{M_{\odot}}$ between 10 and 100 Myr (Fig. 10), as these models take into account the phase of contraction on to the main sequence for lower mass stars. No such change is predicted by the Marigo et al. (2008) models, with the system mass remaining unchanged between 10 Myr and 1 Gyr (Fig. 10). The theoretical system masses are only consistent with the dynamical mass at \sim 30 Myr in the Baraffe et al. (1998) and Siess et al. (2000) models, and are not consistent with any age in the Marigo et al. (2008) models. The metallicity of this system has been measured to be almost solar (Wu et al. 2010), removing the degeneracy which exists between metallicity and the mass and age of the system (e.g. Section 6.2.6).

The position of the primary on the CMD suggests a relatively young age of the system between 50 and 250 Myr. The V - K

	Dynamical	Lejeun	Lejeune & Schaerer		rigo et al.	Si	Siess et al.	
HIP	system mass (M _☉)	Age (Myr)	System mass (M_{\odot})	Age (Myr)	System mass (M _☉)	Age (Myr)	System mass (M_{\odot})	
		Systems	identified as dou	bles (Sectior	n 6.1.1)			
5300	3.16 ± 0.34	280	3.56 ± 0.13	400	3.47 ± 0.13	355	3.69 ± 0.16	
9480	2.72 ± 0.13	250	3.15 ± 0.20	280	3.12 ± 0.20	160	3.25 ± 0.31	
11569 AaAb ^a	2.12 ± 0.25	100	2.45 ± 0.07	100	2.89 ± 0.15	100	2.97 ± 0.25	
76952	4.19 ± 0.30	315	4.38 ± 0.45	400	4.10 ± 0.40	315	4.38 ± 0.45	
	Low-mass b	inary pair w	vithin hierarchica	l system (Se	ctions 6.1.2 and 6	.1.3)		
44127 BC	0.68 ± 0.04	_	_	250	0.88 ± 0.01	50	0.83 ± 0.01	
82321 BC	1.16 ± 0.09	400	1.91 ± 0.02	445	1.87 ± 0.02	500	1.90 ± 0.02	
	Systems v	vith one unr	esolved spectroso	copic compo	nent (Section 6.2	.1)		
80628	4.99 ± 0.75	630	3.03 ± 0.08	630	3.06 ± 0.18	445	3.15 ± 0.13	
	Systems v	ith two unr	esolved spectroso	copic compo	nent (Section 6.2	.1)		
28614	6.36 ± 0.62	630	3.98 ± 0.32	630	3.96 ± 0.28	500	4.18 ± 0.34	
36850	5.42 ± 0.97	280	4.26 ± 0.28	355	4.15 ± 0.24	315	4.34 ± 0.31	
	Systems with e	evidence of	unresolved spect	roscopic cor	nponents (Sectior	n 6.2.2)		
17954	4.15 ± 0.39	400	3.49 ± 0.11	400	3.44 ± 0.12	225	3.67 ± 0.16	
47479	5.83 ± 0.53	560	3.99 ± 0.10	560	3.96 ± 0.10	445	4.17 ± 0.14	
93506	5.26 ± 0.37	710	4.11 ± 0.25	710	4.07 ± 0.23	500	4.45 ± 0.36	

Table 10. Comparison of dynamical and theoretical system masses for each system within the orbit subsample.

^aThe ages and theoretical masses may be significantly biased due to the presence of additional companions within the resolution limit of the Tycho2 and 2MASS observations.

colours of the BC pair also suggest a young age, between 40 and 100 Myr using the Siess et al. (2000) and Baraffe et al. (1998) models. The inferred young age of the system is not inconsistent with the minimum age of stars found within the LIB (Abt 2011), given the relatively high *UVW* space velocity of the HIP 44127 system (Palous 1983). The detection of X-ray emission from this system is also of interest. Previous studies have shown that A7 stars such as the primary should not emit X-rays and that any detection of X-rays from the position of the star can be indicative of a lower mass companion (e.g. De Rosa et al. 2011). The *ROSAT* source J085913.0+480227 is coincident with the optical position of HIP 44127 (Voges et al. 1999).

This hierarchical triple system warrants further study, primarily in order to refine the orbital fit as the BC pair approaches apastron passage in 2018. This system also makes an ideal candidate for future spectroscopic observations to search for the narrow spectral lines of the two faint companions. With a double-lined spectroscopic orbit fit, model-independent masses can be calculated for the individual components. If the young age suggested by both the position on the CMD and the system mass estimated from the mass–magnitude relations is correct, the BC pair would be an ideal calibrator for the theoretical models in the young, low-mass regime.

6.1.3 K- and M-type binaries - HIP 82321

The hierarchical triple system HIP 82321 was resolved within a single epoch of our AO observations. The A-type primary (A) is $\sim 2 \operatorname{arcsec}$ from a binary pair of two lower mass companions (BC) in a wide orbit. The significant proper motion of the primary ($\Delta \alpha = 22.8 \operatorname{mas} \operatorname{yr}^{-1}$, $\Delta \delta = -51.4 \operatorname{mas} \operatorname{yr}^{-1}$) and near constant separation of the three stars suggest that the lower mass pair is comoving with the primary. Spectroscopic measurements of the lower mass

components of this system are possible given the small magnitude difference of the pair with respect to the primary ($\Delta K \sim 2.5$); such spectra would allow for a determination of the individual masses independent of the distance to the system. The *Hipparcos* parallax and 2MASS *K*-band magnitude measurements of this system allow for a tight constraint of the system age to between 300 and 400 Myr, based on the position of the primary on the CMD. The primary is also a possible member of the Ursa Major moving group, an association of solar metallicity stars with an age between 300 Myr (Soderblom & Mayor 1993) and 500 Myr (King et al. 2003).

A refined orbit of the BC pair is presented in Table 8, with an estimated dynamical system mass of $1.32 \pm 0.05 \,M_{\odot}$. The theoretical system mass, assuming the distance to each component is the same as the primary, is significantly higher at $\sim 2.0 \,M_{\odot}$ (Fig. 11). The sparse coverage of the orbit (Fig. 6) suggests that the orbit fit could be poor, resulting in an incorrect dynamical system mass. The orbit fit would be significantly improved with subsequent observations. An alternative scenario is that the pair are a background binary with a proper motion similar to the primary, which would bring the dynamical and photometrically derived masses into agreement.

6.2 Higher order multiplicity

6.2.1 Known spectroscopic binaries

The presence of an unresolved spectroscopic binary can have a significant impact on the magnitude assigned to each component of the multiple system, increasing the estimated component magnitude by as much as 0.75 mag for an unresolved equal-mass spectroscopic binary. For each target within this study, the literature was searched for references to additional components resolved through spectroscopic or interferometric observation which would influence any



Figure 8. The system mass as a function of age based on the mass–magnitude relations derived from three of the model grids. Each panel is similar in nature to Fig. 5, with the spread in the mass–age relation introduced by uncertainties in both the measured *K*-band magnitude, and the distance determination. The horizontal dot–dashed line indicates the dynamical mass determined from the system orbit, with the uncertainty denoted by the line shaded region. The vertical dotted line indicates the age of the system derived from the isochrones, with the line-shaded region denoting the range of ages consistent with the uncertainties in the position of the primary on the CMD. The primary of the HIP 11569 system is significantly bluer than expected for a main-sequence star, and as such its age has been assigned to 100 Myr, with no corresponding uncertainty. The presence of additional components to the HIP 11569 system, within the resolution limit of both the Tycho2 and 2MASS observations, is the likely cause of this bias.

comparison made between the dynamical system mass and the theoretical system mass. Of the targets within the orbit subsample, three are known to have additional spectroscopic components (HIP 28614, HIP 36850 and HIP 80628). The spectroscopic component to HIP 9480 resolved by Abt (1965) is resolved within our AO observations. Similarly, for the monitoring subsample, two are known spectroscopic binaries (HIP 128 and HIP 116611). The greater frequency of spectroscopic binaries within the orbit sample, relative to the monitoring sample, can be explained by the narrow spectral lines of the former sample due to their relatively low radial velocities,



Figure 9. Top panel: the dynamical mass of the HIP 5300 system (horizontal line-shaded region) is significantly below that of the system mass estimated from mass–magnitude relations (solid shaded region) calculated from the solar metallicity models from Lejeune & Schaerer (2001). Middle panel: assuming a higher metallicity (Z = 0.04), the position of the components on the CMD suggests a younger age (denoted by the vertical line-shaded region), and the mass–magnitude relation derived from the metal-enhanced models suggests a systematically lower system mass, although the dynamical mass is still discrepant. Bottom panel: with a metallicity of Z = 0.10, the age estimate is younger still (<10 Myr), and the discrepancy between the dynamical mass and the mass estimated from the mass–magnitude relation is removed.

 $\langle v \sin i \rangle = 71 \,\mathrm{km \, s^{-1}}$ compared with $\langle v \sin i \rangle = 115 \,\mathrm{km \, s^{-1}}$ for the latter sample (Abt & Morrell 1995). The magnitudes listed for the resolved components of these systems (Table 6) are the blended magnitudes of the listed spectroscopic components. In addition to these known spectroscopic binaries, three members of the orbit subsample are resolved as hierarchical triples within our high-resolution observations (HIP 11569, HIP 44128 and HIP 82321). A schematic representation of the higher order multiplicity systems is given in Fig. 12.

For the three systems with known spectroscopic components within the orbit sample, the dynamical system mass includes the mass of each component, regardless of whether it is resolved within our data. This causes a significant discrepancy when the dynamical system mass is compared with the theoretical system mass (e.g. Fig. 7), with the dynamical system mass being systematically higher. The discrepancy between the two values cannot be directly converted into a mass for the unresolved components however, as a blended magnitude would have been used when determining the mass from the mass–magnitude relations.

6.2.2 Evidence of spectroscopic components

Systems with significantly higher dynamical masses than theoretical masses obtained from mass-magnitude relations are strong candidates for multiple systems with unresolved components which have not been detected in spectroscopic observations. We find two such systems with the signature of a possible unresolved component: HIP 17954 and HIP 93506 (Fig. 13), with masses of the order of 0.5 and $1.5 \,\mathrm{M_{\odot}}$. Sensitive spectroscopic observations of both systems may lead to the detection of the spectral lines from an unresolved lower mass component. A similar phenomenon is observed for the HIP 47479 system, although the number of measurements used to determine the orbit is particularly low, making the dynamical mass less certain. The narrow spectral lines of HIP 47479, implied by the low measured stellar rotational velocity (Royer, Zorec & Gómez 2007), make it an ideal candidate for spectroscopic follow-up. Previous spectroscopic observations of this system reveal vsin i variations with a magnitude of 40 km s^{-1} (Moore 1932).

From the total sample of 26 systems, and only considering stellar companions within 100 au, a lower limit on the higher order multiplicity of A-type stars can be estimated. Assuming the suspected unresolved companions described earlier in this section are true, there are five double, six triple and two quadruple systems within the orbit subsample, corresponding to frequencies of 38, 46 and 16 per cent, respectively. This lower limit shows an enhancement on the higher order multiplicity of A-type stars when compared with solar-type primaries (74 per cent double, 20 per cent triple and 6 per cent quadruple or higher order – Raghavan et al. 2010), and is more consistent with the fraction reported for more massive O-type primaries (46 per cent double and 54 per cent triple or higher order – Mason et al. 2009).

6.3 Continued monitoring targets

6.3.1 Newly resolved binaries

We have identified 13 binary systems with projected separations ranging between 13 and 96 au which would make ideal candidates for future orbital monitoring projects (Table 5). The 100-au projected separation cut-off was applied to select only systems for which orbital motion could be detected with several years of observations. The binaries resolved within the monitoring subsample typically have lower mass ratios than for the orbit subsample, a demonstration of the effectiveness of AO observations at detecting high-contrast binaries. Based on their position on the CMD, two members of the monitoring subsample (HIP 5310 and HIP 18217) appear to lie on the zero-age main sequence, and the measured magnitude difference between primary and secondary would correspond to a late K or early M-type companion in each case. These companions are of particular interest as they will allow for further tests of the theoretical models within the young, low-mass regime. A gallery of the observations obtained of each target within the monitoring subsample is shown in Fig. 14.



Figure 10. Top row (left): an AO observation of HIP 44127 triple system. The orbit of the lower mass pair (BC) taken from Table 8 is plotted as a solid line for reference. The bright point source to the lower-right of the primary PSF is a known filter ghost. The axis labels are given in arcseconds, with each arcsecond equal to approximately 14.5 au. (Centre): a CMD highlighting the position of the primary with respect to the theoretical models (Marigo et al. 2008 – solid line; Siess et al. 2000 – dotted line), while the Baraffe et al. (1998) models do not extend to high enough masses. The age of the system was estimated using the isochrone which best fits the position of the primary. (Right): a CMD highlighting the position of the two lower mass components (B – blue, C – red). The line styles are the same as the centre panel. Bottom row: the theoretical system mass plotted as a function of system age for each of the three models. The vertical dotted line within each panel indicates the age of the system estimated from using each of the models, while the horizontal dot–dashed line indicates the dynamical system mass estimated from the orbital fit. As the Baraffe et al. (1998) models do not extend to the A-type star mass range, an age estimate is not possible.



Figure 11. The theoretical system mass plotted as a function of system age for each of the three models for the HIP 82321 BC system. The vertical dotted line within each panel indicates the age of the system estimated from using each of the models, while the horizontal dot–dashed line indicates the dynamical system mass estimated from the orbital fit. The clear discrepancy between the dynamical and theoretical system masses can be explained by the lack of complete coverage of the orbit presented in Fig. 6. Further high angular resolution observations of this system will allow for an improved determination of the orbital parameters.



Hierarchical Systems

Figure 12. A schematic of the seven hierarchical systems within both subsamples, including only components within a projected separation of 100 au to the A-type primary. The schematic diagram does not include the suspected spectroscopic binaries described in Section 6.2.2. Those systems which were newly resolved as a part of the VAST survey are highlighted in blue. Each filled circle represents an individual component, with the period of the binary pair denoted by its vertical position. Our high-resolution observations are only sensitive to systems with orbital periods greater than \sim 10 yr, denoted by the dashed horizontal line. For wide separation systems with indeterminate orbital periods, such as HIP 128 AB–C, the orbital period has been arbitrarily set at 1000 yr, denoted by the dotted horizontal line.

6.3.2 HIP 77660

An advantage of monitoring binary systems for orbital motion with AO instrumentation over interferometric techniques is the elimination of the quadrant ambiguity. In some cases, the output of the image processing of speckle data results in a 180° ambiguity on the position angle measurement (Bagnuolo et al. 1992). This may lead to a scenario where orbital motion is thought to exist for a binary pair, when in fact this motion is the product of such a quadrant uncertainty combined with the true linear motion of the companion. The ambiguity can be resolved by observing the system using AO imaging, where no reconstruction is required to obtain the final science image.

Based on our AO observations of the binary system HIP 77660, it appears that such a quadrant ambiguity has occurred, making linear motion appear as orbital motion, as shown in Fig. 15. The significant proper motion of the system ($\Delta \alpha = -91$ mas yr⁻¹, $\Delta \delta =$ -28 mas yr⁻¹), as measured by *Hipparcos*, is inconsistent with a stationary background object. While future measurements of this system will be able to resolve the presence of orbital motion after a sufficient time baseline, spectroscopy or multicolour photometry will allow for a rapid characterization of the properties of both components.

7 SUMMARY

We have presented high-resolution observations of 26 nearby multiple systems with A-type primaries with projected separations



Figure 13. The six systems for which spectroscopic components are either known to be present through previous spectroscopic observations (HIP 28614, HIP 36850 and HIP 80628) or thought to exist due to a significant discrepancy between the theoretical system mass and the dynamical system mass. For clarity only one model has been used to estimate the theoretical system mass (Lejeune & Schaerer 2001). The horizontal displacement between the intersection of the dynamical system mass (dot–dashed line) and the age estimate of the system (dotted line), and the theoretical system mass curve (solid shaded region) gives an order of magnitude estimate of the mass of the unresolved spectroscopic components.



Figure 14. Observations of the 13 systems with projected separations <100 au suggested as future orbital monitoring targets. The companion in each image is highlighted, with a logarithmic image scale between 1 (white) and 10^{-6} (black) relative to the peak intensity of the primary. An observation of one of the 13 systems with projected separations <100 au suggested as future orbital monitoring targets. The companion in each image is highlighted, with a logarithmic image scale between 1 (white) and 10^{-6} (black) relative to the peak intensity of the primary is highlighted, with a logarithmic image scale between 1 (white) and 10^{-6} (black) relative to the peak intensity of the primary.



Figure 14 – continued

<100 au, 11 of which are binaries newly resolved as a part of the VAST survey. For those systems with sufficient orbital motion coverage, refined orbital parameters were calculated and the estimated dynamical system mass was compared to masses derived from theoretical models. Due to their rapid evolution across the CMD, removing the significant age degeneracy for lower mass solar-type stars, binaries with A-type components are ideal targets with which to test theoretical models. Four such systems were investigated, with one system having consistent dynamical and theoretical system mass estimates. Of the remaining three systems, each had a dynamical mass significantly lower than that predicted from the models. While this discrepancy may be indicative of a true divergence between the models and observations, the lack of metallicity measurements for these systems provides another explanation. Future orbital monitoring observations of A star binary systems will provide further refinement to the orbital parameters and, combined with refinement of the magnitude, metallicity and parallax measurements, will improve the analysis performed within this study.

Observations of two hierarchical systems, consisting of an A-type primary and a tight low-mass binary pair in a wide orbit, allowed us to extend this analysis to the lower mass regime. The primary of the triple HIP 44127 suggests a young system age (<100 Myr), making the M dwarf pair interesting for comparison with evolutionary models. We have also shown that a dynamical system mass significantly higher than the theoretical system mass is suggestive of an unresolved spectroscopic component within the system. Demonstrated on several known spectroscopic binaries, systems which exhibit this discrepancy are ideal candidates for future spectroscopic or interferometric observations in an attempt to detect these hypothetical companions. Interferometric observations may be required to resolve these purported companions, as the rotationally broadened spectral lines of the rapidly rotating members of the monitoring sample (Abt & Morrell 1995) may preclude the detection of additional components using spectroscopy. Including the three systems with evidence of an unresolved close companion, a lower limit on the higher order multiplicity can be estimated from the 13 systems within the orbit subsample as 39 per cent double, 46 per cent triple and 15 per cent quadruple. The frequency of single A-type stars will be explored in an upcoming publication within the VAST survey paper series. The remaining systems for which an orbit could not be determined are candidates for orbital monitoring projects with ground-based high-resolution observations. A number of these systems are of particular interest, based on age estimates derived from the position of the primary on the CMD and the magnitude difference between the two components.



Figure 15. Top panel: the historical measurement of the system HIP 77660, plotted with the orbital fit estimated by Docobo et al. (2010). This system demonstrates the quadrant ambiguity limitation of the speckle interferometry technique. The binary companion was measured by Tokovinin, Mason & Hartkopf (2010) using speckle interferometry to be at a position angle of 248°.4 to 246°.8 between mid-2007 and mid-2009 (blue filled circles, top right). Our AO images, which do not suffer from the same quadrant ambiguity, show the binary at a position angle of 69°.7-67°.3 between mid-2006 and mid-2011 (red filled stars, left). Bottom panel: the relative position of the secondary, with 180° added to the four speckle interferometry measurements which were inconsistent with the observations presented within this study. The solid line connects the first and the last observations, showing a deviation away from linear motion. Further observations are required to determine if this deviation is indicative of the true orbital path of the secondary about the primary.

ACKNOWLEDGMENTS

The authors wish to express their gratitude for the constructive comments received from the referee, H. A. Abt. The authors also wish to express their gratitude to A. Tokovinin, B. Mason and W. Hartkopf for comments which helped improve the paper. We gratefully acknowledge several sources of funding. RJDR is funded through a studentship from the Science and Technology Facilities Council (STFCT) (ST/F 007124/1). JP is funded through support from the Leverhulme Trust (F/00144/BJ) and the STFC (ST/F003277/1, ST/H002707/1). RJDR gratefully acknowledges financial support received from the Royal Astronomical Society to fund collaborative visits. Portions of this work were performed under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory in part under contract W-7405-Eng-48 and in part under contract DE-AC52-07NA27344, and also supported in part by the NSF Science and Technology CfAO, managed by the UC Santa Cruz under cooperative agreement AST 98-76783. This work was supported, through JRG, in part by University of California Lab Research Programme 09-LR-118057-GRAJ and NSF grant AST-0909188. This study is based on observations obtained at the Canada-France-Hawaii Telescope (CFHT), which is operated by the National Research Council of Canada, the Institut National des Sciences de l'Univers of the Centre National de la Recherche Scientifique of France and the University of Hawaii. This study is also based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Science and Technology Facilities Council (United Kingdom), the National Research Council (Canada), CON-ICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência e Tecnologia (Brazil) and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina). The authors also wish to extend their gratitude to the staff at the Palomar Observatory and the UCO/Lick Observatory for their support and assistance provided during the course of the observations. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has made use of the Washington Double Star Catalogue maintained at the US Naval Observatory.

REFERENCES

- Abt H. A., 1965, ApJS, 11, 429
- Abt H. A., 2011, AJ, 141, 165
- Abt H. A., Morrell N. I., 1995, ApJS, 99, 135
- Andersen J., Clausen J. V., Nordstrom B., Tomkin J., Mayor M., 1991, A&A, 246, 99
- Bagnuolo W. G. J., Mason B. D., Barry D. J., Hartkopf W. I., McAlister H. A., 1992, AJ, 103, 1399
- Baize P., Petit M., 1989, A&AS, 77, 497
- Baraffe I., Chabrier G., 1996, ApJ, 461, L51
- Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1998, A&A, 337, 403
- Bate M. R., 2000, MNRAS, 314, 33
- Bonnefoy M. et al., 2009, A&A, 506, 799
- Bonnell I. A., 2001, in Zinnecker E., Mathieu R. D., eds, Proc. IAU Symp. 200, The Formation of Binary Stars. Astron. Soc. Pac., San Francisco, p. 23
- Carpenter J. M., 2001, AJ, 121, 2851
- Carquillat J.-M., Ginestet N., Prieur J. L., Debernardi Y., 2003, MNRAS, 346, 555
- Christou J. C., Drummond J. D., 2006, AJ, 131, 3100
- De Rosa R. J. et al., 2011, MNRAS, 415, 854
- Docobo J. A., Andrade M., 2006, ApJ, 652, 681
- Docobo J. A., Costa J. M., 1985, Circ. Inf., 96
- Docobo J. A., Ling J. F., 2007, AJ, 133, 1209
- Docobo J. A., Tamazian V. S., Balega Y. Y., Melikian N. D., 2010, AJ, 140, 1078
- Doyon R., Nadeau D., Vallee P., Starr B. M., Cuillandre J. C., Beuzit J.-L., Beigbeder F., Brau-Nogue S., 1998, Proc. SPIE, 3354, 760
- Drummond J., Milster S., Ryan P., RobertsL C., Jr, 2003, ApJ, 585, 1007
- Ducati J. R., 2002, Stellar Photometry in Johnson's 11-Color System. CDS, Strasbourg

- Dupuy T. J., Liu M. C., Bowler B. P., 2009, ApJ, 706, 328
- Eggen O. J., 1967, ARA&A, 5, 105
- Ehrenreich D., Lagrange A.-M., Montagnier G., Chauvin G., Galland F., Beuzit J.-L., Rameau J., 2010, A&A, 523, 73
- Erspamer D., North P., 2003, A&A, 398, 1121
- ESA 1997, ESA SP-1200, The Hipparcos and Tycho Catalogues. ESA Publications, Noordwjik
- Fabricius C., Makarov V. V., 2000, A&A, 356, 141
- Fekel F. C., Scarfe C. D., Barlow D. J., Hartkopf W. I., Mason B. D., McAlister H. A., 2002, AJ, 123, 1723
- Green R. M., 1985, Spherical Astronomy. Cambridge Univ. Press, Cambridge, p. 533
- Gutmann F., 1965, Publ. Dom. Astrophys. Obser. Victoria, 12, 391
- Hartkopf W. I., McAlister H. A., Franz O. G., 1989, ApJ, 98, 1014
- Hartkopf W. I., Tokovinin A., Mason B. D., 2012, AJ, 143, 2
- Hayward T. L., Brandl B., Pirger B., Blacken C., Gull G. E., Schoenwald J., Houck J. R., 2001, PASP, 113, 105
- Hilditch R. W., 2001, An Introduction to Close Binary Stars. Cambridge Univ. Press, Cambridge
- Hodapp K. W. et al., 2003, PASP, 115, 1388
- Horch E., Ninkov Z., Franz O. G., 2001, AJ, 121, 1583
- Horch E. P., Meyer R. D., van Altena W. F., 2004, AJ, 127, 1727
- Ivanov V. D., Chauvin G., Foellmi C., Hartung M., Huélamo N., Melo C., Nürnberger D., Sterzik M., 2006, Ap&SS, 304, 247
- King J. R., Villarreal A. R., Soderblom D. R., Gulliver A. F., Adelman S. J., 2003, AJ, 125, 1980
- Köhler R., Ratzka T., Herbst T. M., Kasper M., 2008, A&A, 482, 929
- Kratter K. M., Murray-Clay R. A., Youdin A. N., 2010, ApJ, 710, 1375
- Lejeune T., Schaerer D., 2001, A&A, 366, 538
- Lenzen R. et al., 2003, Proc. SPIE, 4841, 944
- Liu M. C., Dupuy T. J., Ireland M. J., 2008, ApJ, 689, 436
- Lloyd J. P., Liu M. C., Macintosh B. A., Severson S. A., Deich W. T., Graham J. R., 2000, Proc. SPIE, 4008, 814
- Lodato G., Meru F., Clarke C. J., Rice W. K. M., 2007, MNRAS, 374, 590
- McAlister H. A., Hartkopf W. I., Hutter D. J., Franz O. G., 1987, AJ, 93, 688
- McCaughrean M. J., Stauffer J. R., 1994, AJ, 108, 1382
- McDonald J. M., Clarke C. J., 1995, MNRAS, 275, 671
- Marigo P., Girardi L., Bressan A., Groenewegen M. A. T., Silva L., Granato G. L., 2008, A&A, 482, 883
- Mason B. D., Douglass G. G., Hartkopf W. I., 1999, AJ, 117, 1023
- Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2001, AJ, 122, 3466
- Mason B. D., Hartkopf W. I., Gies D. R., Henry T. J., Helsel J. W., 2009, AJ, 137, 3358
- Moore J. H., 1932, Publ. Lick Obser., 18, 1
- Morel M., Magnenat P., 1978, A&AS, 34, 477
- Muterspaugh M. W. et al., 2008, AJ, 135, 766
- Muterspaugh M. W. et al., 2010, AJ, 140, 1623
- Ortega V. G., de la Reza R., Jilinski E., Bazzanella B., 2002, ApJ, 575, L75
- Palous J., 1983, Bull. Astron. Inst. Czechoslovakia, 34, 286
- Popovic G. M., 1969, Bull. Obser. Astron. Belgrade, 27, 33
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1992, Numerical Recipes in FORTRAN. The Art of Scientific Computing, 2nd ed. Cambridge Univ. Press, Cambridge
- Pringle J. E., 1989, MNRAS, 239, 361
- Raghavan D. et al., 2010, ApJS, 190, 1
- Rousset G. et al., 2003, in SPIE, 4839, 140
- Royer F., Zorec J., Gómez A. E., 2007, A&A, 463, 671
- Rucinski S. M. et al., 2005, AJ, 130, 767
- Siess L., Dufour E., Forestini M., 2000, A&A, 358, 593
- Skrutskie M. F. et al., 2006, AJ, 131, 1163 Soderblom D. R., Mayor M., 1993, AJ, 105, 226
- Soderblom D. K., Mayor M., 1995, AJ, J
- Söderhjelm S., 1999, A&A, 341, 121
- Song I., Caillault J.-P., Navascues D. B. y., Stauffer J. R., 2001, ApJ, 546, 352
- Stamatellos D., Hubber D. A., Whitworth A. P., 2007, MNRAS, 382, L30

Stassun K. G., Mathieu R. D., Cargile P. A., Aarnio A. N., Stempels E., Geller A., 2008, Nat, 453, 1079

- Tokovinin A., 2008, MNRAS, 389, 925
- Tokovinin A., Mason B. D., Hartkopf W. I., 2010, AJ, 139, 743
- van Leeuwen F., 2007, A&A, 474, 653
- Vinter Hansen J. M., Neubauer F. J., Roosen-Raad D., 1940, Lick Obser. Bull., 19, 89

Voges W. et al., 1999, A&A, 349, 389 Worley C. E., 1969, AJ, 74, 764 Wu Y., Singh H. P., Prugniel P., Gupta R., Koleva M., 2010, A&A, 525, A71

This paper has been typeset from a $T_{\ensuremath{E}} X/I \!\! \ensuremath{\Delta} T_{\ensuremath{E}} X$ file prepared by the author.