



**HAL**  
open science

## Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016

M. Zemp, M. Huss, Emmanuel Thibert, Nicolas Eckert, R. McNabb, J. Huber, M. Barandun, H. Machguth, S. Nussbaumer, I. Gärtner-Roer, et al.

► **To cite this version:**

M. Zemp, M. Huss, Emmanuel Thibert, Nicolas Eckert, R. McNabb, et al.. Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016. *Nature*, 2019, 568 (7752), pp.382-386. 10.1038/s41586-019-1071-0 . hal-02121653

**HAL Id: hal-02121653**

**<https://hal.science/hal-02121653v1>**

Submitted on 24 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016

Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., & Cogley, J. G. (2019). Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016. *Nature*, 568(7752), 382-386.  
<https://doi.org/10.1038/s41586-019-1071-0>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Nature

**Publication Status:**  
Published (in print/issue): 18/04/2019

**DOI:**  
[10.1038/s41586-019-1071-0](https://doi.org/10.1038/s41586-019-1071-0)

**Document Version**  
Author Accepted version

### **General rights**

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

### **Take down policy**

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk)

## Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016

M. Zemp<sup>1\*</sup>, M. Huss<sup>2,3</sup>, E. Thibert<sup>4</sup>, N. Eckert<sup>4</sup>, R. McNabb<sup>5</sup>, J. Huber<sup>1</sup>, M. Barandun<sup>3</sup>, H. Machguth<sup>1,3</sup>, S. U. Nussbaumer<sup>1,3</sup>, I. Gärtner-Roer<sup>1</sup>, L. Thomson<sup>6</sup>, F. Paul<sup>1</sup>, F. Maussion<sup>7</sup>, S. Kutuzov<sup>8</sup> & J. G. Cogley<sup>9</sup>

<sup>1</sup>Department of Geography, University of Zurich, Switzerland.

<sup>2</sup>Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Switzerland.

<sup>3</sup>Department of Geosciences, University of Fribourg, Switzerland.

<sup>4</sup>Université Grenoble Alpes, Irstea, UR ETGR, Grenoble, France.

<sup>5</sup>Department of Geosciences, University of Oslo, Norway.

<sup>6</sup>Department of Geography and Planning, Queen's University, Kingston, Ontario, Canada. [Authors: OK?]

<sup>7</sup>Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria.

<sup>8</sup>Department of Glaciology, Institute of Geography, Russian Academy of Sciences, Moscow, Russia.

<sup>9</sup>Department of Geography, Trent University, Peterborough, Ontario, Canada. [Authors: OK?]

\*e-mail: michael.zemp@geo.uzh.ch

**Glaciers distinct from the Greenland and Antarctic ice sheets cover an area of approximately 706,000 square kilometres globally<sup>1</sup>, with an estimated total volume of 170,000 cubic kilometres, or 0.4 metres of potential sea-level-rise equivalent<sup>2</sup>. Retreating and thinning glaciers are icons of climate change<sup>3</sup> and affect regional runoff<sup>4</sup> as well as global sea level<sup>5,6</sup>. In reports thus far from the Intergovernmental Panel on Climate Change, estimates of changes in ice-sheet mass [Authors: OK?] were based on the multiplication of averaged or interpolated results from available observations of a few hundred glaciers with regional glacier areas<sup>7-10</sup>. [Authors: could you say briefly what is meant by 'regional glacier areas' here?] For data-scarce regions, these results had to be complemented with estimates based on satellite altimetry and gravimetry<sup>11</sup>. These past**

**approaches were challenged by the small number and heterogeneous spatiotemporal distribution of in situ measurement series and their often unknown ability to represent respective mountain ranges, as well as by the spatial limitations of present satellite altimetry (for which point data only are available) and gravimetry (with its coarse resolution). [Authors: OK?] Here we use an extrapolation of glaciological and geodetic observations to show that glaciers contributed  $27 \pm 22$  millimetres to global mean sea-level rise from 1961 to 2016. Region-specific mass-change rates for 2006–2016 range from  $-0.1$  metres to  $-1.2$  metres of water equivalent per year, resulting in a global sea-level contribution of  $335 \pm 144$  gigatonnes, or  $0.92 \pm 0.39$  millimetres, per year. Although statistical uncertainty ranges overlap, our conclusions suggest that glacier mass loss may be larger than previously reported<sup>11</sup>: it is equivalent to the sea-level contribution of the Greenland Ice Sheet<sup>12</sup>, clearly exceeds the loss from the Antarctic Ice Sheet<sup>13</sup>, and accounts for 25–30 per cent of the total observed sea-level rise<sup>14</sup>. [Authors: OK?] Present mass-loss rates indicate that glaciers could almost disappear in some mountain ranges in this century, while heavily glacierized regions will continue to contribute to sea-level rise beyond 2100. [Authors: please check first paragraph carefully]**

Changes in glacier volume and mass are observed by geodetic and glaciological methods<sup>15</sup>. The glaciological method provides glacier-wide changes in surface mass by using point measurements from seasonal or annual in situ campaigns, extrapolated to unmeasured regions of the glacier. The geodetic method determines glacier-wide volume changes by repeated mapping and differencing of glacier surface elevations from in situ, airborne and spaceborne surveys, usually over multiyear to decadal periods.

In this study, we used glaciological and geodetic data from the World Glacier Monitoring Service (WGMS)<sup>16</sup>, complemented by new and as-yet-unpublished geodetic assessments for glaciers in Africa, Alaska, the Caucasus, Central Asia, the Greenland

periphery, Iceland, New Zealand, Scandinavia, Svalbard and the Russian Arctic. At present, this data set includes observations from 450 and 19,130 glaciers for the glaciological and the geodetic samples, respectively, which correspond to sample sizes of, respectively, less than 1% and 9% of the total number of glaciers<sup>1</sup>. **[Authors: OK?]** We estimated regional mass changes for the 19 first-order regions of the Randolph Glacier Inventory (RGI)<sup>1</sup> (Fig. 1). The observational coverage ranges from less than 1% to 54% of the total glacier area per region for the glaciological sample, and from less than 1% to 79% for the geodetic sample (Extended Data Fig. 1). In each region, we combined the temporal variability from the glaciological sample—obtained using a spatiotemporal variance decomposition—with the glacier-specific values of the geodetic sample (see Methods). We then extrapolated the calibrated annual time series from the observational to the full glacier sample to assess regional mass changes, taking into account regional rates of area change (see Methods). **[Authors: OK?]** Uncertainties originate from four independent error sources. These relate to the temporal changes assessed from the glaciological sample, to the long-term geodetic values, to the extrapolation to unmeasured glaciers, and to estimates of regional glacier area. To estimate regional mass changes, we spatially interpolated the specific mass changes from the observational sample to all glaciers in the region. We estimated the related error from the deviations of this approach to regional (specific) mass changes, calculated as arithmetic averages or as area-weighted averages of the observational sample (see Methods). **[Authors: OK?]**

Over the full observation period from 1961 to 2016, global glacier mass changes cumulated to  $-9,625 \pm 7,975$  Gt ( $1 \text{ Gt} = 10^{12} \text{ kg}$ ). This corresponds to a contribution of  $27 \pm 22$  mm to global sea level, or a contribution of  $0.5 \pm 0.4$  mm per year when a linear rate is assumed. The total mass change excluding peripheral glaciers in Greenland and Antarctica sums to  $-8,305 \pm 5,115$  Gt, corresponding to a contribution to sea level of  $0.4 \pm 0.3$  mm per

year. Cumulative mass changes and corresponding contributions to global sea level were largest from the heavily glacierized regions, with approximately one third originating from Alaska (Fig. 1). Additionally, large contributions originate from regions with less glacierization but strongly negative specific mass changes, such as Western Canada and the USA (Extended Data Fig. 2). South Asia West was the only region that exhibited mass gain over the full observation period. Cumulative specific mass changes over the full observation period from 1961 to 2016 were most negative in the Southern Andes, followed by Alaska, the Low Latitudes, Western Canada and the USA, New Zealand, the Russian Arctic and Central Europe (Extended Data Fig. 2a). When annual rates are averaged over pentads (that is, periods of five years; Fig. 2), sea-level contributions ranged between  $0.2 \pm 0.5$  mm and  $0.3 \pm 0.4$  mm per year until the 1980s, and then increased continuously to reach  $1.0 \pm 0.4$  mm in the latest pentad (2011–2016). Over corresponding periods, our estimates show that global glacier mass loss is approximately equivalent to various mass-loss estimates from the Greenland Ice Sheet (with periods **[Authors: could you explain what is meant by ‘with periods’ in this context?]** between 2003 and 2012)<sup>12</sup>, and it exceeds present contributions to sea-level rise from the Antarctic Ice Sheet (2012–2017:  $219 \pm 43$  Gt per year, including the Antarctic Peninsula)<sup>13</sup> by 62%. Hence, glaciers contributed between 25% and 30% of the observed global mean sea-level rise, which ranged between 2.6 mm and  $2.9 \pm 0.4$  mm per year over the satellite altimetry era (1993 to mid-2014)<sup>14</sup>.

Glacier mass changes were negative in all regions over the latest observational decade, from 2006 to 2016 (Table 1)—that is, covering the hydrological years<sup>15</sup> from 2006/07 to 2015/16. Glaciers in South America had the most negative specific mass changes, with rates exceeding  $-1.0$  m water equivalent (w.e.) per year, followed by glaciers in the Caucasus, Central Europe, Alaska, and Western Canada and the USA, with rates of less than  $-0.8$  m w.e. per year (Fig. 3a; 1 m w.e. =  $1,000 \text{ kg m}^{-2}$ ). The least negative specific mass

changes were found for glaciers in the Antarctic periphery ( $-0.1$  m w.e. per year) and in South Asia West, with glaciers close to balanced-budget conditions<sup>17,18</sup>. Again, regions with large ice cover and negative specific mass changes showed the largest total losses (Fig. 3b). Record mass losses are thus found in Alaska, with rates of  $-73$  Gt per year, followed by other heavily glacierized regions (that is, with glacier areas of more than  $29,000$  km<sup>2</sup>) such as Arctic Canada North ( $-60$  Gt per year), the Greenland periphery ( $-51$  Gt per year), and the Southern Andes ( $-34$  Gt per year; Table 1). Exceptions are Central Asia and South Asia West, with limited mass losses ( $-7$  Gt and  $-1$  Gt per year) despite their large glacier areas. Of the regions with smaller glacierization, Western Canada and the USA and Iceland lost the most mass, at rates of  $-12$  Gt and  $-8$  Gt per year, respectively.

We calculated the relative annual ice loss (Extended Data Fig. 3) by comparing present mass-change rates (2006–2016) with total estimated ice volumes for each region<sup>2</sup>. Nine out of nineteen regions lost between 0.5% and 3% of their total ice volume per year. The other regions featured smaller loss rates. Under present ice-loss rates, most of today's glacier volume would thus vanish in the Caucasus, Central Europe, the Low Latitudes, Western Canada and the USA, and New Zealand in the second half of this century. However, the heavily glacierized regions would continue contributing to sea-level rise beyond this century, as glaciers in these regions would persist but continue to lose mass. It is worthwhile noting that a substantial part of the future ice loss is already committed owing to the imbalance of most glaciers with the present climate<sup>19,20</sup>, and that numerical models are required to fully assess future glacier changes in view of climate-change scenarios<sup>20,21</sup>.

Overall, error bars related to regional mass changes (Fig. 3b) reflect different sources. **[Authors: OK?]** In most regions, the geodetic error accounts for the largest contribution, followed by the error related to temporal changes assessed from the glaciological sample (Extended Data Fig. 4). The extrapolation to unmeasured glaciers contributes substantially to

the overall error only in regions with large differences between interpolation methods. The reasons for these differences are region specific and depend on various factors, such as the observational sample, the glacier size distribution, or a bias towards large tidewater or surge-type glaciers. Uncertainties related to glacier areas and their changes contribute only minimally to the overall error. However, considering area changes is important despite their small contribution to random errors, as a constant glacier area over time would result in a systematic error that increases with the length of the time series and the rate of the area change<sup>22,23</sup>. **[Authors: OK?]**

Our new approach, in combination with major advances in observational evidence, allows for a sound assessment of global glacier mass changes independently of satellite altimetry and gravimetry. **[Authors: OK?]** This is a basic requirement for the comparison of regional results and the detection of potential biases over the satellite era. By comparison with the Intergovernmental Panel on Climate Change (IPCC)'s Fifth Assessment Report (AR5)<sup>11,24</sup>, the greatest improvement herein is in the geodetic sample: it has been boosted from a few hundred glaciers<sup>7</sup> to more 19,000 globally, with an observational coverage of more than 45% of the glacier area in 11 out of 19 regions (Extended Data Fig. 1). **[Authors: OK?]** Our approach, combining the temporal variability from the glaciological sample with large-scale observations from the geodetic sample, facilitates the inference of mass changes at annual resolution for all regions, back to the hydrological year 1961/62. This represents a major development compared with IPCC AR5<sup>11,24</sup>, which had to focus on the satellite altimetry and gravimetry era (2003–2009) and relied on estimates modelled using climate data or on interpolated values from scarce and mostly uncalibrated observational samples for earlier time periods (see Methods). **[Authors: OK?]**

Our central estimate for the global rate of glacier mass loss is 47 Gt (or 18%) per year larger than that reported in IPCC AR5 (Section 4.3.3.3, Table 4.4 of AR5)<sup>11,24</sup> for the period

2003 to 2009 (Extended Data Fig. 5). **[Authors: OK?]** A direct comparison of our results is possible for the seven regions (all with less than 15,000 km<sup>2</sup> of ice cover) with estimates based on glaciological and geodetic samples in IPCC AR5<sup>11,24</sup>. In these regions, our mass-change estimates are systematically less negative (Extended Data Fig. 5a). This suggests that our new approach of calibrating regional glaciological mass-change time series with geodetic observations has overcome an earlier reported negative bias in the glaciological sample<sup>11</sup>. Regions with estimates based on satellite altimetry and gravimetry in IPCC AR5<sup>11,24</sup> featured absolute differences of the same order of magnitude but with varying signs. The more negative global mass changes result mainly from heavily glacierized regions where we estimate larger mass losses (for example, Alaska, peripheral Greenland and Antarctic, Russian Arctic, and Arctic Canada North), and are partly offset by smaller mass-loss estimates for a few other regions with abundant ice cover (for example, Arctic Canada South, Iceland, South Asia West, and Central Asia; Extended Data Fig. 5b) and by the above-mentioned bias in regions with less glacierization. Our error bars are considerably larger than and overlap with those reported in IPCC AR5 (Section 4.3.3.3, Table 4.4)<sup>11,24</sup>. However, a direct comparison is challenging, because the error bars of the earlier study<sup>11</sup> were based on a combination of regionally different methods and data sources. A detailed comparison will require a regional assessment of glacier changes and related uncertainties, including scaling issues from glacier-wide observations (this study) and results from satellite altimetry (regional averages of repeat-path measurements) and gravimetry (coarse resolution of sensor and hydrological models). However, our error estimates are methodologically consistent and consider all known relevant sources of potential errors. We consider the relative differences of our error bars between the regions to be plausible and their absolute values to be upper bounds.

Improvements in global glacier mass-change assessments are still possible and necessary. First, the observational database needs to be extended in both space and time. We currently see the most urgent need for closing observational gaps being in regions where glaciers dominate runoff during warm/dry seasons, such as in the tropical Andes and in Central Asia<sup>4</sup>, and in regions that dominate the glacier contribution to future sea-level rise, that is, Alaska, Arctic Canada, the Russian Arctic, and peripheral glaciers in Greenland and Antarctica. Second, a systematic assessment of regional area-change rates<sup>25</sup> will improve the estimate of corresponding impacts on regional mass changes. Finally, more research is required to better constrain the observational uncertainties at individual glaciers<sup>26</sup> and for regional mass-change assessments. Despite these remaining challenges, our assessment of global glacier mass changes provides a new observational baseline for a sound comparison with estimates based on other methods<sup>27</sup>, as well as for future modelling studies of glacier contributions to regional runoff and global sea-level rise.

**Online content** Any Methods, including any statements of data availability and Nature Research reporting summaries, along with any additional references and Source Data files, are available in the online version of the paper at

Received 28 September 2018; accepted 24 January 2019.

trueunknown<unknown1true0false0>true

\_id="b1"trueunknown<unknown1true0false0>true1. Arendt, A. et al. *Randolph Glacier Inventory (v.2.0): A Dataset of Global Glacier Outlines. Global Land Ice Measurements from Space, Boulder, Colorado USA* (GLIMS Technical Report, 2017).trueunknown</unknown>true trueunknown</unknown>true **[Authors: OK?]**

truejrn<jrn1true0false0>true \_doi="10.1029/2012JF002523" \_id="b2" \_issn="0148-0227"truejrn<jrn1true0false0>true2. Huss, M. & Farinotti, D. Distributed ice

thickness and volume of all glaciers around the globe. *J. Geophys. Res.* 117, F04010 (2012).

[\\_doi="10.1175/BAMS-D-13-00047.1"](https://doi.org/10.1175/BAMS-D-13-00047.1) [\\_id="b3"](#) [\\_issn="0003-0007"](#) 3. Bojinski, S. et al. The concept of essential climate variables in support of climate research, applications, and policy. *Bull. Am. Meteorol. Soc.* 95, 1431–1443 (2014). [CrossRef](#)

[\\_doi="10.1038/s41558-017-0049-x"](https://doi.org/10.1038/s41558-017-0049-x) [\\_id="b4"](#) [\\_issn="1758-678X"](#) 4. Huss, M. & Hock, R. Global-scale hydrological response to future glacier mass loss. *Nat. Clim. Chang.* 8, 135–140 (2018).

[\\_doi="10.1126/science.1254702"](https://doi.org/10.1126/science.1254702) [\\_id="b5"](#) [\\_issn="0036-8075"](#) 5. Marzeion, B., Cogley, J. G., Richter, K. & Parkes, D. Attribution of global glacier mass loss to anthropogenic and natural causes. *Science* 345, 919–921 (2014).

[\\_doi="10.1007/s00382-013-1719-7"](https://doi.org/10.1007/s00382-013-1719-7) [\\_id="b6"](#) [\\_issn="0930-7575"](#) 6. Radić, V. et al. Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models. *Clim. Dyn.* 42, 37–58 (2014).

[\\_doi="10.3189/172756409787769744"](https://doi.org/10.3189/172756409787769744) [\\_id="b7"](#) [\\_issn="0260-3055"](#) 7. Cogley, J. G. Geodetic and direct mass-balance measurements: comparison and joint analysis. *Ann. Glaciol.* 50, 96–100 (2009).

truejrn<jrn1true0false0>true \_doi="10.1029/2006GL027511" \_id="b8" \_issn="0094-8276"truejrn<jrn1true0false0>true8. Kaser, G., Cogley, J. G., Dyurgerov, M. B., Meier, M. F. & Ohmura, A. Mass balance of glaciers and ice caps: consensus estimates for 1961–2004. *Geophys. Res. Lett.* 33, L19501 (2006).truejrn</jrn>true

truebok<bok1true0false0>true \_id="b9"truebok<bok1true0false0>true9. Dyurgerov, M. B. & Meier, M. F. *Glaciers and the Changing Earth System: A 2004 Snapshot*. (Instaar, 2005).truebok</bok>true **[Authors: OK?]**

trueedb<edb1true0false0>true \_id="b10"trueedb<edb1true0false0>true10. Ohmura, A. in *The State of the Planet: Frontiers and Challenges in Geophysics* vol. 150 (eds Sparks, R. S. J. & Hawkesworth, C. J.) 239–257 (American Geophysical Union, 2004).trueedb</edb>true **[Authors: OK?]**

truejrn<jrn1true0false0>true \_doi="10.1126/science.1234532" \_id="b11" \_issn="0036-8075"truejrn<jrn1true0false0>true11. Gardner, A. S. et al. A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009. *Science* 340, 852–857 (2013).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1088/0034-4885/78/4/046801" \_id="b12" \_issn="0034-4885"truejrn<jrn1true0false0>true12. Khan, S. A. et al. Greenland ice sheet mass balance: a review. *Rep. Prog. Phys.* 78, 046801 (2015).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1038/s41586-018-0179-y" \_id="b13" \_issn="0028-0836"truejrn<jrn1true0false0>true13. IMBIE. Mass balance of the Antarctic Ice Sheet from 1992 to 2017. *Nature* 558, 219–222 (2018).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1038/nclimate2635" \_id="b14" \_issn="1758-678X"truejrn<jrn1true0false0>true14. Watson, C. S. et al. Unabated global mean sea-level rise over the satellite altimeter era. *Nat. Clim. Chang.* 5, 565–568 (2015).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_id="b15" \_issn="1350-5084"truejrn<unknown1true0false0>true15. Working Group on Mass-Balance Terminology and Methods of the International Association of Cryosphere. Glossary of Glacier Mass Balance and Related Terms. *UNESCO Digital Library* <https://unesdoc.unesco.org/ark:/48223/pf0000192525> (2011).truejrn</jrn>truejrn</jrn>true **[Authors: reference details OK?]**

trueunknown<unknown1true0false0>true \_id="b16"trueunknown<unknown1true0false0>true16. Working Group on Mass-Balance Terminology and Methods of the International Association of Cryosphere. Global Glacier Change Bulletin No. 2 (2014–2015). *UNESCO Digital Library* [https://wgms.ch/downloads/WGMS\\_GGCB\\_02.pdf](https://wgms.ch/downloads/WGMS_GGCB_02.pdf) (2017).trueunknown</unknown>truejrn</jrn>true **[Authors: reference details OK?]**

truejrn<jrn1true0false0>true \_doi="10.1038/ngeo2999" \_id="b17" \_issn="1752-0894"truejrn<jrn1true0false0>true17. Brun, F., Berthier, E., Wagnon, P., Kääb, A. & Treichler, D. A spatially resolved estimate of High Mountain Asia glacier mass balances from 2000 to 2016. *Nat. Geosci.* 10, 668–673 (2017); correction 11, 543 (2018).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-9-557-2015" \_id="b18"truejrn<jrn1true0false0>true18. Kääb, A., Treichler, D., Nuth, C. & Berthier, E. Contending estimates of 2003–2008 glacier mass balance over the

Pamir–Karakoram–Himalaya. *Cryosphere* 9, 557–564

(2015).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-7-1565-2013"

\_id="b19"truejrn<jrn1true0false0>true19. Mernild, S. H., Lipscomb, W. H., Bahr,

D. B., Radić, V. & Zemp, M. Global glacier changes: a revised assessment of  
committed mass losses and sampling uncertainties. *Cryosphere* 7, 1565–1577

(2013).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1038/s41558-018-0093-1" \_id="b20" \_issn="1758-

678X"truejrn<jrn1true0false0>true20.

Marzeion, B., Kaser, G., Maussion, F. &  
Champollion, N. Limited influence of climate change mitigation on short-term  
glacier mass loss. *Nat. Clim. Chang.* 8, 305–308

(2018).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_id="b21" \_issn="2095-0195"truejrn<jrn1true0false0>true21.

Huss, M. & Hock, R. A new model for global glacier change and sea-level  
rise. *Front. Earth Sci.* 3, <https://doi.org/10.3389/feart.2015.00054>

(2015).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3189/2012JoG11J216" \_id="b22" \_issn="0022-

1430"truejrn<jrn1true0false0>true22.

Huss, M., Hock, R., Bauder, A. & Funk,  
M. Conventional versus reference-surface mass balance. *J. Glaciol.* 58, 278–286

(2012).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-4-569-2010"

\_id="b23"truejrn<jrn1true0false0>true23.

Paul, F. The influence of changes in  
glacier extent and surface elevation on modeled mass balance. *Cryosphere* 4, 569–

581 (2010).truejrn</jrn>truejrn</jrn>true

24. Vaughan, D. G. et al. in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)* (eds Stocker, T. F. et al.) 317–382 (Cambridge Univ. Press, 2013).

25. Cogley, J. G. Glacier shrinkage across High Mountain Asia. *Ann. Glaciol.* 57, 41–49 (2016).

26. Zemp, M. et al. Reanalysing glacier mass balance measurement series. *Cryosph.* 7, 1227–1245 (2013).

27. Marzeion, B., Leclercq, P. W., Cogley, J. G. & Jarosch, A. H. Global reconstructions of glacier mass change during the 20th century are consistent. *Cryosphere* 9, 2399–2404 (2015).

**Acknowledgements** This paper is published in recognition of J. Graham Cogley, a pioneer in glacier mass-balance studies. We thank the national correspondents and principal investigators of the WGMS network as well as the Global Land Ice Measurements from Space (GLIMS) and RGI communities for free and open access to their data sets. We thank B. Armstrong for polishing the English. Arctic digital elevation model (DEM) strips were provided by the Polar Geospatial Center under National Science Foundation (NSF) Office of Polar Programs (OPP) awards 1043681, 1559691 and 1542736. This study was enabled by support from the Swiss Global Climate Observing System (GCOS) Office at the Federal Office of Meteorology and Climatology MeteoSwiss, the Cryospheric Commission of the Swiss Academy of Science, the Copernicus Climate Change

Service (C3S) implemented by the European Centre for Medium-range Weather Forecasts (ECMWF) on behalf of the European Commission, and the European Space Agency (ESA) project Glaciers\_cci (4000109873/14/I-NB). **[Authors: acknowledgements section OK?]**

**Reviewer information** *Nature* thanks A. Rowan and the other anonymous reviewer(s) for their contribution to the peer review of this work.

**Author contributions** M.Z. initiated and coordinated the study and wrote the manuscript; the basic concept was jointly developed during a workshop in the Swiss pre-Alps. M.Z., S.U.N., H.M., I.G.-R., J.H., F.P., L.T., S.K. and J.G.C. **[OK?]** compiled data from the research community and the literature; R.M., J.H. and M.B. computed additional geodetic results. M.H., M.B. and F.M. defined clusters and regions used in the analysis. E.T. and N.E. ran the variance decomposition model. M.H. performed the calibration of the glaciological signal to the geodetic series and the extrapolation to regional changes. M.Z., I.G.-R., S.U.N., E.T. and J.H. produced the figures. All authors commented on the manuscript. **[Authors: Author contributions section OK?]**

**Competing interests:** The authors declare no competing interests.

Additional information

**Extended data** is available for this paper at

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Correspondence and requests for materials** should be addressed to M.Z.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**[Author: Please ensure that the following information is included in ALL figure legends where relevant. Sample size (exact  $n$  number); a statement of replicability (how many times was experiment replicated in the lab); description of sample collection; state the statistical test used and give  $P$  values; define centre values (median or average) and error bars. For figures/images that are reproduced or adapted from a third party, it is important that you confirm that permission has been obtained and that appropriate acknowledgement of the copyright holder is given.]**

**Fig. 1 | Regional glacier contributions to sea-level rise from 1961 to 2016.** The cumulative regional and global mass changes (in Gt, represented by the volume of the bubbles) are shown for the 19 first-order regions<sup>1</sup> (outlined with bold black lines). Specific mass-change rates (m w.e.  $\text{yr}^{-1}$ ) are indicated by the colours of the bubbles. In the background, the

locations of glaciological and geodetic data samples are plotted over the glacier polygons from RGI 6.0. **[Authors: what do the plus signs represent?]** As an example, glaciers in Alaska (ALA) show the largest contribution to sea-level rise, with a total mass change of approximately  $-3,000$  Gt or 8 mm sea-level equivalent (s.l.e.) from 1961 to 2016, because of a strongly negative specific mass-change rate ( $-0.6$  m w.e.  $\text{yr}^{-1}$ ) combined with a large regional glacier area. Note that South Asia West (ASW, blue bubble) is the only region in which glaciers slightly gained mass. ACN, Arctic Canada North; ACS, Arctic Canada South; ANT, Antarctic and SubAntarctic; ASC, Central Asia; ASE, South Asia East; ASN, North Asia; CAU, Caucasus and Middle East; CEU, Central Europe; GRL, Greenland; ISL, Iceland; NZL, New Zealand; RUA, Russian Arctic; SAN, Southern Andes; SCA, Scandinavia; SJM, Svalbard and Jan Mayen; TRP, Low Latitudes; WNA, Western Canada and USA (see **Table 1**). **[Authors: it seems helpful to have the abbreviations here as well, as the figure and table may end up on different page spreads.]**

**Fig. 2 | Global glacier contributions to sea-level rise from 1961 to 2016.** Annual and pentadal mass-change rates (in  $\text{Gt yr}^{-1}$ ; left y-axis) and equivalents of mean global sea-level rise (in  $\text{mm yr}^{-1}$ ; right y-axis) are shown with related error bars **[Authors: are these the grey and pale blue envelopes?]** corresponding to 95% confidence intervals. Annual errors originate from independent sources: glaciological sample, geodetic sample, spatial interpolation and glacier area. Over the five-year periods, the individual error terms are cumulated separately and then the multi-year terms are combined according to the law of random error propagations, and divided by the number of years (see Methods). **[Authors: OK?]**

**Fig. 3 | Regional estimates of glacier mass change for the period 2006–2016.** **a, b**, Annual mass-change rates in  $\text{m w.e. yr}^{-1}$  (**a**) and in  $\text{Gt yr}^{-1}$  (**b**) as estimated from spatial interpolation (blue circles), area-weighting (black triangles) and arithmetic averaging (orange squares).

The spatial-interpolation approach is the reference, with results provided in [Table 1](#). Error bars correspond to 95% confidence intervals, and consider uncertainties related to the temporal variability of the glaciological sample, the geodetic value, the regional interpolation, the regional glacier area, and a second-order crossed term. We estimate the error related to the regional interpolation from the differences between the three interpolation approaches (see Methods section ‘Uncertainty estimates’).

**Table 1 | Annual rates of glacier change per region from 2006 to 2016**

Region (code)	Total area (km <sup>2</sup> )	Total volume (km <sup>3</sup> )	Specific mass change (m w.e. yr <sup>-1</sup> )	Mass change (Gt yr <sup>-1</sup> )
01 Alaska (ALA)	86,725	18,429	-0.85±0.19	-73±17
02 Western Canada & USA (WNA)	14,524	1,048	-0.83±0.40	-12±6
03 Arctic Canada North (ACN)	105,111	29,721	-0.57±0.80	-60±84
04 Arctic Canada South (ACS)	40,888	8,948	-0.57±0.70	-23±28
05 Greenland (GRL)	89,717	15,780	-0.63±0.21	-51±17
06 Iceland (ISL)	11,060	3,520	-0.71±0.43	-8±5
07 Svalbard and Jan Mayen (SJM)	33,959	8,076	-0.47±0.23	-16±8
08 Scandinavia (SCA)	2,949	306	-0.49±0.27	-1±1
09 Russian Arctic (RUA)	51,592	15,449	-0.47±0.37	-24±19
10 North Asia (ASN)	2,410	146	-0.37±0.31	-1±1
11 Central Europe (CEU)	2,092	116	-0.87±0.07	-2±0
12 Caucasus and Middle East (CAU)	1,307	63	-0.90±0.57	-1±1
13 Central Asia (ASC)	49,303	3,483	-0.15±0.12	-7±6
14 South Asia West (ASW)	33,568	3,092	-0.03±0.12	-1±4
15 South Asia East (ASE)	14,734	906	-0.35±0.12	-5±2
16 Low Latitudes (TRP)	2,341	80	-1.03±0.83	-2±2
17 Southern Andes (SAN)	29,429	5,518	-1.18±0.38	-34±11
18 New Zealand (NZL)	1,162	61	-0.68±1.15	-1±1
19 Antarctic and Subantarctic (ANT)	132,867	46,801	-0.11±0.87	-14±108
Total, excl. GRL and ANT	483,155	98,962	-0.56±0.04	-270±19
Global total	705,73	161,543	-0.48±0.20	-335±144

The table shows present-day regional and global glacier areas and volumes, with specific mass changes (in m w.e. yr<sup>-1</sup>) and mass-change rates from spatial interpolation (in Gt yr<sup>-1</sup>) for the period from 2006 to 2016. Regional glacier areas are from the RGI 6.0 and refer to the first decade of the twenty-first century<sup>1</sup>. Regional estimates for glacier volumes are based on ref. <sup>2</sup>, updated to the glacier outlines of RGI 6.0. Global totals are calculated as sums of regions for area, volume and mass change. Global specific mass changes are calculated by dividing the global mass-change rate by the global glacier area. Error bars correspond to 95% confidence intervals and originate from independent sources: glaciological sample, geodetic sample, spatial interpolation and glacier area (see Methods section on ‘Uncertainty estimates’). **[Authors: OK?]**

[Author: Please ensure that the following information is provided in the Methods section where relevant. We recommend that detailed protocols are deposited in Protocol Exchange, or a similar repository. If custom computer code has been used and is central to the conclusions of this paper, please insert a section into the Methods titled ‘Code availability’ and indicate within this section whether and how the code can be accessed, including any restrictions to access. If unpublished data are used, please obtain permission.]

## METHODS

### Glaciological and geodetic mass changes

The glaciological method usually provides glacier-wide surface mass balance ( $B_{\text{sfc}}$ ) over an annual period related to the hydrological year. In line with ref. <sup>15</sup>, we use the unit m w.e. for the specific mass change (1 m w.e. = 1,000 kg m<sup>-2</sup>) and the unit Gt for the mass change (1 Gt = 10<sup>12</sup> kg), with mass balance and mass change as synonymous terms. Results are reported as cumulative values over a period of record or as annual change rates (yr<sup>-1</sup>). The geodetic balance is the result of surface (sfc), internal (int) and basal (bas) mass changes and—in the case of marine-terminating or lacustrine glaciers—of calving ( $D$ ) in the unit m w.e.:

$$B_{\text{geod}} = \Delta M = B_{\text{sfc}} + B_{\text{int}} + B_{\text{bas}} + D$$

In practice, the geodetic (specific) mass change is calculated as the volume change,  $\Delta V$ , over a survey period between  $t_0$  and  $t_1$ , from differencing of digital elevation models (DEMs), over the glacier area multiplied by a volume-to-mass conversion factor:

$$B_{\text{geod}} = \frac{\Delta V}{S} \times \frac{1}{(t_1 - t_0)} \times \frac{\bar{\rho}}{\rho_{\text{water}}}$$

where  $\bar{S}$  is the average glacier area of the two survey times ( $t_0, t_1$ ) assuming a linear change through time<sup>26</sup>, and  $\bar{\rho}$  is the average density of  $\Delta V$  with a commonly applied value<sup>28</sup> of  $850 \pm 60 \text{ kg m}^{-3}$ . The glaciological method is able to satisfactorily capture the temporal variability of the glacier mass change even with only a small observational sample<sup>29,30</sup>. However, its cumulative amount over a given time span is sensitive to systematic errors, which accumulate with the number of annual measurements<sup>31,32</sup>. The geodetic method provides mass changes covering the entire glacier area and large glacier samples. However, the method requires a density conversion and surveys are typically carried out at multi-annual to decadal intervals only. For both measurements, we use the latest version (wgms-fog-2018-06) of the Fluctuations of Glaciers (FoG) database from the WGMS<sup>16</sup>. The glaciological sample was recently updated with observations from latest years, consolidated by adding results from approximately 100 additional glaciers<sup>7</sup>, and by replacing entire mass-balance series after reanalysis<sup>33–37</sup>. The geodetic sample was increased recently by the inclusion of large-scale assessments from several mountain regions<sup>17,38–42</sup>.

For the present study, we complemented the data set from the WGMS with an additional 70,873 geodetic volume change observations computed for 6,551 glaciers in Africa, Alaska, the Caucasus, Central Asia, Greenland's periphery, Iceland, New Zealand, the Russian Arctic, Scandinavia and Svalbard (**Extended Data Table 1**). This was achieved by calculating geodetic mass changes from ASTER DEMs processed using MMASTER<sup>43</sup> which were co-registered using off-glacier elevations from the Ice, Cloud, and land Elevation Satellite (ICESat) as common frame following ref. <sup>44</sup>. Where available, we used ArcticDEM 2-m strips, SPOT5-based DEMs from IPY-SPIRIT or the High Mountain Asia 8-m DEMs<sup>45</sup> to increase spatial and temporal coverage, after resampling to 30-m resolution to match the resolution of the ASTER DEMs. Pairs of DEMs (for example, ASTER/ASTER or ASTER/ArcticDEM) were automatically chosen on the basis of at least 40% overlap and a

time separation of at least eight years. This time separation, together with the selection of DEMs towards the end of the ablation period (**Extended Data Table 1**), aims to reduce the effect of seasonal variations in the surface elevation and minimizes differences to glaciological survey dates. On the basis of the selected DEM pairs, glacier elevation changes were computed for various time periods between 2000 and 2018. We used the local hypsometric method<sup>46</sup> to fill voids in the DEMs. For each glacier outline with an area of at least 0.6 km<sup>2</sup> from the RGI 6.0, we calculated the glacier hypsometry using 100-m elevation bins. For each DEM pair, we calculated the mean elevation difference per elevation bin, and multiplied this by the glacier hypsometry to obtain a volume change. **[Authors: OK?]** The longest available differences with at least 70% data coverage were then used for each glacier to obtain geodetic mass change. For the peripheral glaciers in Western Greenland, geodetic mass changes were calculated using the Aero DEM<sup>47</sup> from 1985 and a prerelease of the 2010–14 TanDEM-X Global DEM. Before differencing, all DEMs were co-registered to each other<sup>44</sup>. We estimated the glacier volume change with the local hypsometric method<sup>46</sup>, using elevations derived from TanDEM-X and the RGI 6.0 outlines. Again, only glaciers with at least 70% data coverage were used. We estimated uncertainties in geodetic mass changes on the basis of differences between the two DEMs after co-registration, following the approach of ref. <sup>17</sup>.

### **Glacier inventory**

We derived the global distribution of glaciers from the RGI<sup>1,48</sup>, which is a snapshot glacier inventory derived from the Global Land Ice Measurements from Space (GLIMS) database<sup>49</sup> and a large compilation of national and regional sources compiled by the RGI consortium<sup>1</sup>. **[Authors: OK?]** We used glacier area and its distribution with elevation (that is, glacier hypsometry) for the 215,547 glaciers in the RGI 6.0, covering a total area of 705,739 km<sup>2</sup>, mainly for survey years between 2000 and 2010. **[Authors: OK?]** Improvements with

respect to earlier RGI versions as used in IPCC AR5<sup>11,24</sup> (168,331 glaciers, 726,258 km<sup>2</sup>) include the separation of glacier complexes (for example, ice fields or ice caps) into individual glaciers, replacement of nominal glaciers (that is, size-equivalent circles) by real glacier outlines, assignment of glacier-specific survey dates, and the introduction of glacier-specific hypsometries (Extended Data Fig. 1). The latter come as a list of elevation-band areas (at a resolution of 50 m in height) in the form of integer thousands of the glacier's total area<sup>1</sup>. Note that at present the RGI includes peripheral glaciers surrounding the Greenland Ice Sheet<sup>50</sup> but not the peripheral glaciers on the Antarctic Peninsula<sup>51</sup> and in the McMurdo Dry Valleys<sup>52</sup>. For future versions of the RGI, the inclusion of these peripheral glaciers in Antarctica should be considered to reach global completeness and consistency with the classification of peripheral glaciers in Greenland<sup>50</sup>.

### **Changes in glacier area**

For hydrological and sea-level applications, it is the conventional mass balance that is relevant—that is, the mass change calculated over a constantly changing area and hypsometry of a glacier<sup>15</sup>. While the changes in hypsometry are implicitly captured, the changes in glacier area need to be explicitly accounted for by both the glaciological and the geodetic methods<sup>26</sup>. In contrast with earlier approaches, we considered the impact of changes in glacier area over time on regional mass-change estimates. Therefore, we used a collection of relative area changes from IPCC AR5 (Chapter 4, Fig. 4.10, Table 4.SM.1 of ref. <sup>24</sup>), extended with additional literature<sup>53–55</sup> to obtain area change rates for all first-order glacier regions.

**[Authors: OK?]**

### **Glacier volume estimates**

Regional estimates for glacier volumes are based on ref. <sup>2</sup>, updated to the glacier outlines of RGI 6.0.

### **Spatial regionalization**

For regional analysis, it is convenient to group glaciers by proximity. We achieved this by using the latest version of glacier regions as available from the Global Terrestrial Network for Glaciers<sup>56</sup>. These 19 first-order and more than 90 second-order regions derive ultimately from glacier regions proposed by the GLIMS project around the year 2000 and from studies dealing with global glacier distribution<sup>57,58</sup>, and are implemented in both the RGI and the FoG databases. **[Authors: OK?]** For mass-balance studies, the 19 first-order regions seem to be appropriate because of their manageable number and their geographical extent, which is close to the spatial correlation distance of glacier mass-balance variability (that is, several hundred kilometres)<sup>59,60</sup>. We further divided these regions for areas that are known to feature large diversities in mass-balance gradients and where sufficient data coverage allowed (**Extended Data Table 2**).

### **Extraction of temporal variability from the glaciological sample**

In a first step, we subdivided the sample of glaciological series into spatial clusters. We started from the smallest possible units (second-order glacier regions) and then extended them until the number and completeness of the time series was acceptable to ensure a proper variance decomposition based on visual and quantitative criteria, such as a common mass-balance temporal variability (that is, a high correlation between annual mass-balance series) and spatial consistency (that is, a cluster cannot be geographically too wide). The resulting 20 regional clusters correspond to first- and second-order glacier regions or a combination thereof (**Extended Data Table 2**). For half of these clusters, the available mass-balance series cover the full survey period with only minor data gaps of a few years. For the other half of the clusters, we complemented the glaciological sample with a few long-term series from neighbouring regions that feature a similar mass-balance variability (**Extended Data Table 2**). For the few clusters without glaciological data before the mid-1970s, we use the mean value

of the geodetic sample (that is, neglecting interannual variability) for these years and set the related uncertainty to twice the average value of the first decade with glaciological observations.

In the second step, we extracted the temporal mass-balance variability for each cluster using a variance decomposition model<sup>61</sup>, which is a further development of the approach of ref. <sup>30</sup>, based on Bayesian techniques<sup>62,63</sup> and applied to a regional sample of glacier-wide mass balances instead of to a series of point measurements. **[Authors: OK?]** For this model, we defined the specific mass change for a given glacier  $i$  and year  $t$  as:

$$B_{\text{glac},i,t} = \alpha_0 + \alpha_i + g(t) + z(t) + \varepsilon_{i,t} \quad (1)$$

where  $\alpha_0$  is the cluster's annual average and  $\alpha_i$  is the glacier-specific site deviation of the (specific) mass change from the cluster's average. The variables  $g(t)$  and  $z(t)$  are the long-term trend and annual fluctuations, respectively, of the time deviation from the average, and  $\varepsilon_{i,t}$  are residuals. The variable  $g(t)$  was taken as a smooth nonparametric trend and  $z(t)$  as a white-noise term. Their sum is the annual deviation of the glaciological sample from the average  $\alpha_0$ , which is further used in the analysis:

$$B_{\text{glac,cluster}} = g(t) + z(t) \quad (2)$$

Model inference was performed using Bayesian simulation techniques, giving access, for any parameter or combination of parameters, to a point estimate and to a credibility interval quantifying the related uncertainty. This especially applies to cumulated temporal

deviations  $\sum_{t=1}^{t_2} (g(t) + z(t))$  over any time interval  $[t_1, t_2]$ , such as the full period from 1961 to 2016 (Extended Data Fig. 6)

### Calibration to mass-change values from the geodetic sample

For each cluster (Extended Data Table 2), we calibrated the temporal mass-balance variability as derived from the glaciological sample  $B_{\text{glac,cluster}}$  to the values from the geodetic methods<sup>26</sup>. Owing to the differences in length of the geodetic survey periods, we carried out the calibration individually for all glaciers with available geodetic balances. If more than one geodetic survey was available per glacier, we combined those with the longest survey periods by arithmetic averaging of annual change rates. For each glacier  $i$ , we calculated the mean annual deviation  $\overline{\beta}_t$  between the glaciological balance of the cluster  $B_{\text{glac,cluster}}$  and the glacier-specific geodetic balance  $B_{\text{geod},i}$  over a common time period of  $N$  years between  $t_0$  and  $t_1$ :

$$\overline{\beta}_t = \frac{B_{\text{geod},i} - \sum_{t_0}^{t_1} B_{\text{glac,cluster}}}{N} \quad (3)$$

The annual calibrated specific mass change for every glacier  $i$  and year  $t$  was then calculated as:

$$\Delta M_{\text{cal},i,t} = B_{\text{glac,cluster},t} + \overline{\beta}_t \quad (4)$$

As a result, for each glacier with available geodetic data we obtained a calibrated specific mass-change series that features the temporal variability of the glaciological cluster but is adjusted to the glacier-specific geodetic value (Extended Data Fig. 7).

### Regional mass changes and contributions to sea level [Authors: Methods subheadings need to fit one line of typeset text; is this shortened suggestion OK?]

To estimate the total mass change, we need to scale the results from the sample with available (geodetic) data to all glaciers of a region (from RGI 6.0). We followed three different approaches to calculate the regional specific mass change  $\Delta M_{\text{region}}$  (in units of m w.e. yr<sup>-1</sup>): arithmetic averaging  $\Delta M_{\text{region,AVG}}$ , area-weighting  $\Delta M_{\text{region,AW}}$ , and spatial interpolation  $\Delta M_{\text{region,INT}}$ . For the approaches  $\Delta M_{\text{region,AVG}}$  and  $\Delta M_{\text{region,AW}}$ , we assigned the arithmetic and

glacier area-weighted average, respectively, of the annual specific mass change of the observational sample to all unobserved glaciers in the region. For our reference approach  $\Delta M_{\text{region,INT}}$ , we spatially interpolated the individual specific mass-changes to all glacier locations in the region using an inverse distance weighting function. For all approaches, we calculated the regional mass change  $\Delta M_{\text{region}}$  (in units of  $\text{Gt yr}^{-1}$ ) as the product of the specific mass change multiplied by the regional glacier area from RGI 6.0 and applying the relative area change rates of the corresponding region. Global mass changes,  $\Delta M_{\text{global}}$ , were calculated as the sum of all regional mass changes. For conversion to sea-level equivalent, we assumed a total area of the ocean of  $362.5 \times 10^6 \text{ km}^2$  (ref. [64](#)).

### Uncertainty estimates

The random error of the regional mass change,  $\sigma_{\text{regional}}$ , is composed of the errors related to: first, the temporal changes in the regional glaciological sample  $\sigma_{\text{glac}}$ ; second, the geodetic values of the individual glaciers  $\sigma_{\text{geod}}$ ; third, the extrapolation from the observational to the full sample  $\sigma_{\text{extrapolation}}$ ; fourth, the glacierized area  $\sigma_{\text{area}}$  of the region; and fifth, a second-order crossed term related to the calculation of the regional mass change (as the product of specific mass change multiplied by the glacierized area):

$$\sigma_{\text{regional}} = \sqrt{\sigma_{\text{glac}}^2 + \sigma_{\text{geod}}^2 + \sigma_{\text{extrapolation}}^2 + \sigma_{\text{area}}^2 + \sigma_{\text{crossed}}^2} \quad (5)$$

The variable  $\sigma_{\text{glac}}$  can be rigorously estimated from the variance decomposition ([Extended Data Fig. 6](#)). However, we used a less computationally intensive approach to estimate it for any subperiod (pentad, decade) from the full study period. Specifically, the annual standard deviations of the temporal deviation ( $g(t) + z(t)$ ) as obtained from the variance decomposition model were summed up according to the law of random error propagation. Hence, the standard deviation of any subperiod was evaluated as if annual deviations would be independent. The variable  $\sigma_{\text{glac}}$  implicitly accounts for errors related to

differences in the glaciological survey period, because the sample contains results from various time systems (for example, fixed-date, floating-date and stratigraphic)<sup>15</sup>.

The variable  $\sigma_{\text{geod}}$  is the uncertainty from the geodetic method. We calculated the annual values as rates—that is, dividing the reported (multiyear) uncertainties by the number of years between the two surveys. It includes the observation uncertainty  $\sigma_{\text{geod.observation}}$  as reported with the geodetic results. In addition, we considered the uncertainty introduced by calibrating annual mass-balance variability with geodetic values,  $\sigma_{\text{calibration}}$  (Equation (4)), which was inferred for each glacier individually on the basis of randomly superimposing  $\sigma_{\text{glac}}$  and extracting the standard deviation of average balances over the reference period. The uncertainties related to density conversion factor  $\sigma_{\text{density}}$  were set to  $\pm 60 \text{ kg m}^{-3}$  according to ref. <sup>28</sup>. The overall geodetic uncertainty was calculated from these terms, assuming them to be uncorrelated, and was divided by the square root of the number of independent items  $n$  of information in the sample:

$$\sigma_{\text{geod}} = \sqrt{\frac{\sigma_{\text{geod.observation}}^2 + \sigma_{\text{calibration}}^2 + \sigma_{\text{density}}^2}{n}}$$

In the ideal case,  $n$  would be equal to the number of geodetic series in the regional sample. For spaceborne surveys, however, the geodetic uncertainty is usually derived from the stable terrain in between a group of glaciers. We thus assumed geodetic uncertainties uncorrelated for samples larger than 50 glaciers, and estimated  $n$  by dividing the regional geodetic sample size by 50. Note that, for the geodetic sample, we do not explicitly formulate uncertainties related to differences in the survey date. For individual glaciers, a corresponding rigorous estimate would be possible using seasonal mass-balance information, meteorological data, and numerical modelling<sup>7,26,33</sup>. These studies show that the corresponding uncertainties can be relevant for individual years but tend towards zero for longer period of records and larger samples.

To estimate  $\sigma_{\text{extrapolation}}$ , we used the regional mass change from spatial interpolation ( $\Delta M_{\text{region,INT}}$ ) as a best guess and calculated the extrapolation uncertainty as 1.96 standard deviations of the results from the three approaches ( $\Delta M_{\text{region,INT}}$ ,  $\Delta M_{\text{region,AVG}}$  and  $\Delta M_{\text{region,AW}}$ ). As for  $\sigma_{\text{glac}}$ , we evaluated  $\sigma_{\text{extrapolation}}$  over any subperiod by the square root of the number of survey years, assuming that annual values are uncorrelated.

For the regional glacier area, we assumed a general uncertainty of  $\pm 5\%$  for the total area derived from the RGI 6.0, given earlier single-glacier and basin-scale uncertainty estimates<sup>48</sup> and in line with the latest GCOS product requirements (Table 25 of ref. <sup>65</sup>; terrestrial essential climate variable (ECV) product requirements). **[Authors: OK?]** This uncertainty was combined with an error related to the regional area changes  $\sigma_{\text{area,change}}$ , which was estimated as 1.96 standard deviations of different ways to calculate regional change rates. **[Authors: OK?]** For a given region, the first approach, used as reference, weights multiple published change rates by the total ice cover of the corresponding glacier samples. The second approach weights multiple results by the length of the survey periods. The uncertainties related to the total area and to area changes were assumed to be uncorrelated and, hence, cumulated according the law of random error propagation.

Over multiyear periods, in contrast with  $\sigma_{\text{glac}}$ ,  $\sigma_{\text{extrapolation}}$  and  $\sigma_{\text{crossed}}$ , the error related to the geodetic values and glacier areas ( $\sigma_{\text{geod}}$  and  $\sigma_{\text{area}}$ ) cumulates linearly. **[Authors: OK?]** Consequently, the individual terms need to be cumulated separately, followed by a combination of the multiyear terms according to the law of random error propagation (see Equation (5)). **[Authors: OK?]** For global sums, the overall error was calculated by cumulating the regional errors according to the law of random error propagation.

### **Code availability**

The analytical scripts are available from the authors on request.

### Data availability

The temporal variabilities for the glaciological clusters as well as the regional and global mass-change results have been deposited in the zenodo repository (<http://doi.org/10.5281/zenodo.1492141>). The full sample of glaciological and geodetic observations for individual glaciers is publicly available from the WGMS (<http://doi.org/10.5904/wgms-fog-2018-11>).

truejrn<jrn1true0false0>true \_doi="10.5194/tc-7-877-2013"  
\_id="b28"truejrn<jrn1true0false0>true28. Huss, M. Density assumptions for  
converting geodetic glacier volume change to mass change. *Cryosphere* 7, 877–887  
(2013).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1111/j.0435-3676.1999.00084.x" \_id="b29"  
\_issn="0435-3676"truejrn<jrn1true0false0>true29. Fountain, A. G. & Vecchia, A.  
How many stakes are required to measure the mass balance of a glacier? *Geogr.  
Ann., Ser. A* 81, 563–573 (1999).truejrn</jrn>true **[Authors:  
corrected volume number OK?]**

truejrn<jrn1true0false0>true \_doi="10.1017/S0022143000023169" \_id="b30" \_issn="0022-  
1430"truejrn<jrn1true0false0>true30. Lliboutry, L. Multivariate statistical  
analysis of glacier annual balances. *J. Glaciol.* 13, 371–392  
(1974).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3189/172756504781829855" \_id="b31" \_issn="0022-  
1430"truejrn<jrn1true0false0>true31. Cox, L. H. & March, R. S. Comparison  
of geodetic and glaciological mass-balance techniques, Gulkana Glacier, Alaska,  
U.S.A. *J. Glaciol.* 50, 363–370 (2004).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3189/002214308785837093" \_id="b32" \_issn="0022-1430"truejrn<jrn1true0false0>true32. Thibert, E., Blanc, R., Vincent, C. & Eckert, N. Glaciological and volumetric mass-balance measurements: error analysis over 51 years for Glacier de Sarennes, French Alps. *J. Glaciol.* 54, 522–532 (2008).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3189/172756409787769627" \_id="b33" \_issn="0260-3055"truejrn<jrn1true0false0>true33. Huss, M., Bauder, A. & Funk, M. Homogenization of long-term mass-balance time series. *Ann. Glaciol.* 50, 198–206 (2009).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-10-535-2016" \_id="b34"truejrn<jrn1true0false0>true34. Andreassen, L. M., Elvehøy, H., Kjøllmoen, B. & Engeset, R. V. Reanalysis of long-term series of glaciological and geodetic mass balance for 10 Norwegian glaciers. *Cryosphere* 10, 535–552 (2016).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_id="b35" \_issn="0022-1430"truejrn<jrn1true0false0>true35. Thomson, L. I., Zemp, M., Copland, L., Cogley, J. G. & Ecclestone, M. A. Comparison of geodetic and glaciological mass budgets for White Glacier, Axel Heiberg Island, Canada. *J. Glaciol.* 63, 55–66 (2016).truejrn</jrn>truejrn</jrn>true **[Authors: reference details OK?]**

truejrn<jrn1true0false0>true \_doi="10.1016/j.gloplacha.2014.01.001" \_id="b36" \_issn="0921-8181"truejrn<jrn1true0false0>true36. Wang, P., Li, Z., Li, H., Wang, W. & Yao, H. Comparison of glaciological and geodetic mass balance at Urumqi Glacier No. 1, Tian Shan, Central Asia. *Global Planet. Change* 114, 14–22 (2014).truejrn</jrn>truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1017/jog.2016.17" \_id="b37" \_issn="0022-1430"truejrn<jrn1true0false0>true37. Basantes-Serrano, R. et al. Slight mass loss revealed by reanalyzing glacier mass-balance observations on Glaciar Antisana 15 $\alpha$  (inner tropics) during the 1995–2012 period. *J. Glaciol.* 62, 124–136 (2016).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-9-525-2015" \_id="b38"truejrn<jrn1true0false0>true38. Fischer, M., Huss, M. & Hoelzle, M. Surface elevation and mass changes of all Swiss glaciers 1980–2010. *Cryosphere* 9, 525–540 (2015).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3390/rs8121038" \_id="b39" \_issn="1032-9714"truejrn<jrn1true0false0>true39. Vijay, S. & Braun, M. Elevation change rates of glaciers in the Lahaul-Spiti (Western Himalaya, India) during 2000–2012 and 2012–2013. *Remote Sens.* 8, 1038 (2016).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.3189/2015AoG70A227" \_id="b40" \_issn="0260-3055"truejrn<jrn1true0false0>true40. Le Bris, R. & Paul, F. Glacier-specific elevation changes in parts of western Alaska. *Ann. Glaciol.* 56, 184–192 (2015).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1657/1938-4246-45.1.19" \_id="b41" \_issn="1523-0430"truejrn<jrn1true0false0>true41. Falaschi, D., Bravo, C., Masiokas, M., Villalba, R. & Rivera, A. First glacier inventory and recent changes in glacier area in the Monte San Lorenzo region (47 °S), Southern Patagonian Andes, South America. *Arct. Antarct. Alp. Res.* 45, 19–28 (2013).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1029/2006JF000586" \_id="b42" \_issn="0148-0227"truejrn<jrn1true0false0>true42. Larsen, C. F., Motyka, R. J., Arendt, A.

A., Echelmeyer, K. A. & Geissler, P. E. Glacier changes in southeast Alaska and northwest British Columbia and contribution to sea level rise. *J. Geophys. Res.* 112, F01007 (2007).

\_doi="10.3390/rs9070704" \_id="b43" \_issn="1032-9714". Girod, L., Nuth, C., Kääb, A., McNabb, R. W. & Galland, O. MMASTER: improved ASTER DEMs for elevation change monitoring. *Remote Sens.* 9, 704 (2017).

\_doi="10.5194/tc-5-271-2011" \_id="b44". Nuth, C. & Kääb, A. Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change. *Cryosphere* 5, 271–290 (2011).

\_id="b45". Shean, D. High mountain Asia 8-meter DEMs derived from cross-track optical imagery V001 (NASA National Snow and Ice Data Center Distributed Active Archive Center (NSIDS DAAC), 2017). **[Authors: software details (including title) here OK?]**

\_id="b46". McNabb, R., Nuth, C., Kääb, A. & Girod, L. Sensitivity of geodetic glacier mass balance estimation to DEM void interpolation. *Cryosph. Discuss.* <https://doi.org/10.5194/tc-2018-175> (2018). **[Authors: OK?]**

\_doi="10.1038/sdata.2016.32" \_id="b47" \_issn="2052-4463". Korsgaard, N. J. et al. Digital elevation model and orthophotographs of Greenland based on aerial photographs from 1978–1987. *Sci. Data* 3, 160032 (2016).

truejrn<jrn1true0false0>true \_doi="10.3189/2014JoG13J176" \_id="b48" \_issn="0022-1430"truejrn<jrn1true0false0>true48. Pfeffer, W. T. et al. The Randolph Glacier Inventory: a globally complete inventory of glaciers. *J. Glaciol.* 60, 537–552 (2014).truejrn</jrn>true

trueunknown<unknown1true0false0>true \_id="b49"trueunknown<unknown1true0false0>true49. GLIMS glacier database, version 1 (GLIMS and National Snow and Ice Data Center (NSIDC), 2005).trueunknown</unknown>true **[Authors: is this reference OK as edited?]**

truejrn<jrn1true0false0>true \_doi="10.5194/tc-6-1483-2012" \_id="b50"truejrn<jrn1true0false0>true50. Rastner, P. et al. The first complete inventory of the local glaciers and ice caps on Greenland. *Cryosph.* 6, 1483–1495 (2012).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/essd-9-115-2017" \_id="b51" \_issn="1866-3508"truejrn<jrn1true0false0>true51. Huber, J., Cook, A. J., Paul, F. & Zemp, M. A complete glacier inventory of the Antarctic Peninsula based on Landsat 7 images from 2000 to 2002 and other preexisting data sets. *Earth Syst. Sci. Data* 9, 115–131 (2017).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.1017/jog.2016.86" \_id="b52" \_issn="0022-1430"truejrn<jrn1true0false0>true52. Fountain, A. G., Basagic, H. J., IV & Niebuhr, S. Glaciers in equilibrium, McMurdo Dry Valleys, Antarctica. *J. Glaciol.* 62, 976–989 (2016).truejrn</jrn>true

truejrn<jrn1true0false0>true \_id="b53"truejrn<jrn1true0false0>true53. Mernild, S. H. et al. Glacier changes in the circumpolar Arctic and sub-Arctic, mid-1980s to late-

2000s/2011. *Geogr. Tidsskr. J. Geogr.* 115, 39–56

(2015).truejrn</jrn>true

truejrn<jrn1true0false0>true \_doi="10.5194/tc-9-565-2015"

\_id="b54"truejrn<jrn1true0false0>true54. Hannesdóttir, H., Björnsson, H., Pálsson, F., Aðalgeirsdóttir, G. & Guðmundsson, S. Changes in the southeast Vatnajökull ice cap, Iceland, between ~1890 and 2010. *Cryosphere* 9, 565–585

(2015).truejrn</jrn>true

trueunknown<unknown1true0false0>true

\_id="b55"trueunknown<unknown1true0false0>true55. Khromova, T. et al. Impacts of climate change on the mountain glaciers of Russia. *Reg. Environ. Change* 18, xxx–xxx (2018).trueunknown</unknown>true

**[Authors: I can't track this reference down; please could you check the details and add the relevant page numbers?]**

trueunknown<unknown1true0false0>true

\_id="b56"trueunknown<unknown1true0false0>true56. Glacier Regions. *Global Terrestrial Network for Glaciers*. <http://dx.doi.org/10.5904/gtng-glacreg-2017-07> (2017). trueunknown</unknown>true

truejrn<jrn1true0false0>true \_doi="10.1029/2009JF001373" \_id="b57" \_issn="0148-

0227"truejrn<jrn1true0false0>true57. Radić, V. & Hock, R. Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory data. *J. Geophys. Res.* 115, F001373 (2010).truejrn</jrn>true

**[Authors: reference details OK?]**

trueother<other1true0false0>true \_id="b58"trueother<other1true0false0>true58. Dyurgerov,

M. B. Glacier mass balance and regime: data of measurements and analysis.

Occasional Paper No. 55 (Institute of Arctic and Alpine Research, Univ. Colorado, 2002).

[doi="10.2307/1551719"](#) [\\_id="b59"](#) [\\_issn="0004-0851"](#). [Letréguilly, A. & Reynaud, L. Space and time distribution of glacier mass-balance in the Northern Hemisphere. \*Arct. Alp. Res.\* 22, 43–50 \(1990\).](#)

[\\_id="b60"](#). [Cogley, J. G. & Adams, W. P. Mass balance of glaciers other than the ice sheets. \*J. Glaciol.\* 44, 315–325 \(1998\).](#) **[Authors: reference details OK?]**

[\\_doi="10.1038/nmeth.3005"](#) [\\_id="b61"](#) [\\_issn="1548-7091"](#). [Krzywinski, M. & Altman, N. Analysis of variance and blocking. \*Nat. Methods\* 11, 699–700 \(2014\).](#)

[\\_doi="10.3189/002214311795306673"](#) [\\_id="b62"](#) [\\_issn="0022-1430"](#). [Eckert, N., Baya, H., Thibert, E. & Vincent, C. Extracting the temporal signal from a winter and summer mass-balance series: application to a six-decade record at Glacier de Sarennes, French Alps. \*J. Glaciol.\* 57, 134–150 \(2011\).](#) [CrossRef](#)

[\\_doi="10.1038/nmeth.3368"](#) [\\_id="b63"](#) [\\_issn="1548-7091"](#). [Puga, J. L., Krzywinski, M. & Altman, N. Bayesian statistics. \*Nat. Methods\* 12, 377–378 \(2015\); corrigendum 12, 1098 \(2015\).](#)

truejrn<jrn1true0false0>true \_doi="10.1080/01490419.2012.709476" \_id="b64"  
\_issn="0149-0419"truejrn<jrn1true0false0>true64. Cogley, J. G. Area of the ocean.  
*Mar. Geod.* 35, 379–388 (2012).truejrn</jrn>true

trueunknown<unknown1true0false0>true \_id="b65"trueunknown<bok1true0false0>true65.  
GCOS. *The Global Observing System for Climate: Implementation Needs*  
(World Meteorological Organization,  
2016).trueunknown</unknown>true

**Extended Data Fig. 1 | Regional glacier hypsometry and observational coverage. a–s,**  
For each of the 19 first-order regions, glacier hypsometry from the RGI 6.0 (blue) is overlaid  
with glacier hypsometry of both the geodetic (grey) and the glaciological (black) samples  
used here. Values for the total number ( $N$ ) and total area ( $S$ ) of glaciers are given for each  
region, together with the relative coverage of both the glaciological and the geodetic samples.  
Plots are ordered according to the region numbers in RGI 6.0 (see **Table 1**); m a.s.l., metres  
above sea level. **[Authors: Please check all legends carefully]**

**Extended Data Fig. 2 | Cumulative regional glacier changes since the 1960s. a, b,**  
Cumulative mass changes in m w.e. (**a**) and Gt (**b**) are shown for the 19 regions. Specific  
mass changes (**a**) indicate the observed glacier thickness changes. Total glacier mass changes  
(**b**, left  $y$ -axis) correspond to the regional contributions to global mean sea-level rise (**b**, right  
 $y$ -axis). As an example, cumulative specific mass changes were most negative in the Southern  
Andes with an average regional glacier thickness change of approximately  $-40$  m w.e. (**a**),  
resulting in a cumulative mass change of  $-1,200$  Gt (**b**). Glaciers in Alaska experience less  
negative specific mass changes (**a**) but contribute much more to global sea-level rise (**b**)  
because of the larger regional glacier area.

**Extended Data Fig. 3 | Relative annual ice loss for the period from 2006 to 2016.** Annual mass change rates (see Fig. 3b) relative to estimated total ice volumes<sup>2</sup> are plotted as vertical bars (% yr<sup>-1</sup>).

**Extended Data Fig. 4 | Relative error contributions for the period 2006–2016.** Shown are relative contributions (%) of the different sources to the overall regional error bars (Fig. 3b). Taking Alaska as an example, the overall error estimate is dominated by the glaciological and the geodetic errors with contributions of 47% and 37%, respectively, whereas the errors for extrapolation (10%), glacier area (5%), and second-order crossed uncertainties (less than 1%) are of less importance. A special case is Central Europe: the large number of high-quality observations from airborne surveys comes with reported geodetic uncertainties that are one order of magnitude smaller than the spaceborne estimates in other regions. As a result, the overall error bars are much smaller (Fig. 3) and the relative contributions from other error sources become larger. In the Southern Andes, the relative contribution of the geodetic error is reduced by the large sample size, while glaciological and interpolation errors feature large absolute values.

**Extended Data Fig. 5 | Comparison of regional mass changes with results from IPCC AR5.** a, b, Annual specific mass-change rates in m w.e. yr<sup>-1</sup> (a) and in Gt yr<sup>-1</sup> (b), as shown in Fig. 3 but for the period 2003–2009. The estimates and related error bars (corresponding to 95% confidence intervals) found here are shown in blue. The results from IPCC AR5<sup>11,24</sup> are shown in red, differentiating between those based on glaciological and geodetic observations (crosses) and those based on ICESat and/or the Gravity Recovery and Climate Experiment (GRACE; diamonds). Global mass change rates are  $-260 \pm 28$  Gt per year and  $-307 \pm 148$  Gt per year, as estimated by IPCC AR5<sup>11,24</sup> and this study, respectively. **[Authors: Please could you amend the figure to show ‘ref. 11’ instead of ‘Gardner et al.’, and then resubmit the figure?]**

**Extended Data Fig. 6 | Temporal variability in the glaciological mass balance for Alaska and British Columbia, 1961–2016. a, b**, Annual (**a**; m w.e. yr<sup>-1</sup>) and cumulative (**b**; m w.e.) values for the cluster's smooth trend ( $g(t)$ ; blue lines) and annual deviations ( $g(t) + z(t)$ ; orange lines), as reconstructed from the variance decomposition (see Methods, equations (1) and (2)) on the basis of glaciological measurements from 19 glaciers ([Extended Data Table 2](#), cluster 01).

**Extended Data Fig. 7 | Calibration of temporal variability from glaciological sample to geodetic values of individual glaciers.** Schematic representation of the approach to calibrate the cumulative temporal variability (black line; m w.e.), as derived from the variance decomposition (see [Extended Data Fig. 6](#)), to geodetic values of individual glaciers (blue and purple lines; m w.e.). For glacier 1 and glacier 2, the mean annual deviations between the glaciological balance of the cluster and the glacier-individual geodetic balances were 0.1 m w.e. per year and  $-0.2$  m w.e. per year, respectively, over corresponding survey periods between  $t_0$  and  $t_1$  (see Methods, equation (3)).

**Extended Data Table 1 | Overview of new geodetic volume changes**

For each first-order region<sup>56</sup> with new geodetic surveys, the numbers of glaciers and observations are given together with the DEMs used, the range of survey periods (SP), the average length of the survey period (N), and the day of the year (doy) of the average survey date (SD) and of the average reference date (RD). The averages of N, SDdoy and RDdoy are given together with the corresponding standard deviations. The TanDEM-X as used for Greenland is a merged product from surveys between 2010 and 2014 from any month of the year.

**Extended Data Table 2 | Spatial clusters used to analyse temporal variability from glaciological samples**

Spatial clusters and corresponding first- and second-order regions<sup>56</sup> as used for extracting the temporal variability of the glaciological sample.  $N$  indicates the number of available glaciological time series per cluster. We complemented clusters with limited time coverage with long-term mass-balance series from neighbouring regions. For region codes, see [Table 1](#).