

ONE-DIMENSIONAL ACTIONS OF HIGMAN'S GROUP

CRISTÓBAL RIVAS AND MICHELE TRIESTINO

ABSTRACT. We build a faithful action of Higman's group on the line by homeomorphisms, answering a question of Yves de Cornulier. As a by product we obtain many quasimorphisms from the Higman group into the reals. We also show that every action by C^1 -diffeomorphisms of Higman's group on the line or the circle is trivial.

1. INTRODUCTION

The first example of an infinite, finitely generated and simple group was built by Higman [9] as a quotient of the group with the following presentation

$$(1) \quad \langle a_i \ (i \in \mathbb{Z}/4\mathbb{Z}) \mid a_i a_{i+1} a_i^{-1} = a_{i+1}^2 \rangle = H.$$

Indeed, Higman observed that H is infinite, in fact torsion free, and moreover that if four elements in a group satisfy the relations above and each of them has finite order, then they all have to be the group identity. In particular H has no non-trivial finite quotient and the quotient of H by a maximal normal subgroup is infinite, simple and finitely generated. With time, the name *Higman group* was attached to the group with the above presentation.

The fact that H has no non-trivial finite quotient implies that every finite dimensional linear representation $\rho : H \rightarrow \mathrm{GL}(V)$ is trivial, since finitely generated groups of matrices are residually finite by a theorem of Mal'cev [11]. It also implies that every C^1 -action $\rho : H \rightarrow \mathrm{Diff}^1(M)$ having a periodic orbit is trivial if M is a connected manifold. Indeed, since H has no finite quotient other than the trivial, the finite orbit must be a global fixed point and, since H has only trivial linear representation, the derivative of every $h \in H$ must be trivial at that point. It then follows from Thurston Stability Theorem [16] that if $\rho(H)$ is not the trivial group, then $\rho(H)$ (and hence H) must admit a surjective homomorphism onto \mathbb{Z} , which is certainly impossible since H has trivial abelianization. This immediately implies, for instance, that every representation $\rho : H \rightarrow \mathrm{Diff}^1([0, 1])$ is trivial since 0 is always in a finite orbit.

The main purpose of this work is to show that, despite the constraints described above, Higman group can be faithfully represented as a group of orientation-preserving homeomorphisms of the real line¹. This implies, in particular, that H admits faithful representations inside the group of homeomorphisms of any given manifold. Indeed, choose a ball B^d in a manifold M^d , the complement of a point $p \in B^d$ in B^d is homeomorphic to $\mathbb{R} \times \mathbb{S}^{d-1}$; given an action of H on \mathbb{R} , extend it to an action on $\mathbb{R} \times \mathbb{S}^{d-1}$, so that the action on the \mathbb{S}^{d-1} -factor is trivial: this gives an action of H on B^d , which fixes p and the boundary ∂B^d , so that it can be extended as the identity outside B^d .

We start by showing, in Section 2, that H admits a non-trivial action on the real line. This amounts to find four homeomorphisms a, b, c, d of the line satisfying

$$aba^{-1} = b^2, \quad bcb^{-1} = c^2, \quad cdc^{-1} = d^2, \quad dad^{-1} = a^2.$$

In our construction, b and d will be *lifts* to the real line of strong parabolic homeomorphisms of the circle, so that the subgroup $\langle b, d \rangle$ display ping-pong dynamics and acts without global fixed points. The choice of the generators a and c is more subtle and it relies on the fact that the Baumslag-Solitar group $\mathrm{BS}(1, 2) = \langle \alpha, \beta \mid \alpha\beta\alpha^{-1} = \beta^2 \rangle$ admits two different and easy-to-describe actions on the real

¹Note that, since H has no finite index subgroup, every action of H by homeomorphisms of an oriented manifold must preserve the orientation.

line (see the beginning of Section 2). We will build a and c as a limit of homeomorphisms a_n and c_n ($n \geq 0$), where the relations

$$a_n b a_n^{-1} = b^2 \text{ and } c_n d c_n^{-1} = d^2,$$

hold for even n 's, and the relations

$$b c_n b^{-1} = c_n^2 \text{ and } d a_n d^{-1} = a_n^2,$$

hold for odd n 's. At each step, we will perform slight modifications of the homeomorphisms so that, *in the limit*, both sets of relations will be satisfied thus providing a non-trivial representation $\varphi : H \rightarrow \text{Homeo}_+(\mathbb{R})$.

The fact that $\varphi : H \rightarrow \text{Homeo}_+(\mathbb{R})$ is non-trivial entails that $\varphi(H)$ is a *left-orderable* group, that is $\varphi(H)$ admits a total order \preceq such that $\varphi(f) \prec \varphi(g)$ implies that $\varphi(hf) \prec \varphi(hg)$ for every $f, g, h \in H$. In fact, for countable groups being left-orderable is equivalent to admit an injective representation into $\text{Homeo}_+(\mathbb{R})$ (see, for instance, [4, 8]). In our case, we are unable to decide whether φ is injective or not, but, using Kurosh Theorem [15], in Section 3 we will show that the kernel of φ must be a free group, which is one of the basic examples of left-orderable groups [4, 8]. In particular, we can *lexicographically* extend a left-order on $\varphi(H)$ to a left-order on H (see Section 3) so that we can deduce

Theorem A. *Higman group H is left-orderable and hence it admits a faithful action on the real line by homeomorphisms.*

We point out that all the homeomorphisms in our construction of $\varphi : H \rightarrow \text{Homeo}(\mathbb{R})$ actually commute with the translation $T : x \mapsto x + 2$. Thus, we get a faithful action of H on the circle $\mathbb{R}/T(x) \sim x$ without global fixed points. See Proposition 3.2. With the same idea of construction, we are able to describe countably many different (i.e. pairwise non-semi-conjugate²) actions of H on the circle, and uncountably many different actions on the real line, all without global fixed points. See Section 4.

Remark 1.1. Observe that Higman's group H is perfect (that is, its abelianization is trivial), so every homomorphism $\Phi : H \rightarrow \mathbb{R}$ is trivial. A weaker notion is that of *quasimorphism* to \mathbb{R} , that is, a map $\Phi : H \rightarrow \mathbb{R}$ for which there exists $D > 0$ such that for every $g, h \in H$, one has $|\Phi(gh) - \Phi(g) - \Phi(h)| \leq D$. The set of all quasimorphisms $H \rightarrow \mathbb{R}$ defines the real vector space $\mathcal{QM}(H)$. By cohomological considerations, one deduces that the dimension of $\mathcal{QM}(H)$ is the cardinal of continuum: indeed, for perfect groups, the quotient $\mathcal{QM}(H)/\{\text{bounded functions}\}$ is isomorphic to the second bounded cohomology group $H_b^2(H; \mathbb{R})$, whose dimension is the cardinal of continuum since H is a non-trivial amalgam [7]. Similarly, one can consider the second bounded cohomology group $H_b^2(H; \mathbb{Z})$ with integer coefficients. Every action on the circle without global fixed points determines a non-trivial cocycle in $H_b^2(H; \mathbb{Z})$, by the so-called *bounded Euler class*. Moreover, two actions of H are semi-conjugate if and only if their bounded Euler classes in $H_b^2(H; \mathbb{Z})$ are the same [8]. Therefore, the actions constructed in Section 4 provide countably many linearly independent classes in $H_b^2(H; \mathbb{Z})$. See the references [6, 8] for more about bounded cohomology related to group actions on the circle.

In contrast, although our actions have no fixed points neither periodic orbits, we can still show that they cannot be made differentiable. In fact we have:

Theorem B. *Every representation $\rho : H \rightarrow \text{Diff}^1(\mathbb{R})$ is trivial. The same holds for every representation $\rho : H \rightarrow \text{Diff}^1(\mathbb{S}^1)$.*

²Recall that two actions $\psi_1 : G \rightarrow \text{Homeo}_+(\mathbb{R})$ and $\psi_2 : G \rightarrow \text{Homeo}_+(\mathbb{R})$ are semi-conjugate, if there is a proper, non-decreasing map $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ such that $\alpha \circ \psi_1(g) = \psi_2(g) \circ \alpha$ for all $g \in G$. Two actions on the circle are semi-conjugate if they have lifts to the real line which are semi-conjugate. See [2, 8, 12] for further details.

Our proof relies on the knowledge of C^1 -actions of the Baumslag-Solitar group $\text{BS}(1, 2) = \langle \alpha, \beta \mid \alpha\beta\alpha^{-1} = \beta^2 \rangle$ on the line [1] together with an analysis of the possible combinatorics of fixed points of elements of H when acting on the line by homeomorphisms. The proof is given in Section 5 and it is independent of the rest of the paper.

Acknowledgements: C.R. would like to thank the organizers of Groups of Dynamical Origin II where he learned about the question of orderability of Higman's group. In particular, he is grateful to Yves de Cornulier for sharing this question and to Andrés Navas for his constant encouragement. M.T. acknowledges the hospitality of USACH while part of this project was carried on.

C.R. was partially supported by FONDECYT 1181548. M.T. was partially supported MathAmSud RGSd (Rigidity and Geometric Structures in Dynamics) 19-MATH-04 and the project "Jeunes Géomètres" of F. Labourie (financed by the Louis D. Foundation).

2. THE CONSTRUCTION OF φ

Higman's group contains four canonical copies of the Baumslag-Solitar group $\text{BS}(1, 2) = \langle \alpha, \beta \mid \alpha\beta\alpha^{-1} = \beta^2 \rangle$. This latter group acts on the real line via its *standard affine action* $\alpha : x \mapsto 2x$, $\beta : x \mapsto x + 1$ (see Figure 1, left), which is a faithful action since every proper quotient of $\text{BS}(1, 2)$ is abelian. *Non-standard affine* actions of $\text{BS}(1, 2)$ on the line can be obtained from the standard one by blowing up an orbit, following the method of Denjoy (see Figure 1, right). In fact, it is not very hard to see that *if there is* a non-trivial action of H on the line, then a non-standard affine action must appear at least locally (this is, for instance, exploited in the proof of Theorem B, see Section 5). The following example is the one that we will use.

Example 2.1 (Denjoy trick). Consider the affine maps of the real line $f(x) = 2x$ and $g(x) = x + 1$, which generate a group isomorphic to $\text{BS}(1, 2)$. The orbit of 0 coincides with the dyadic rationals $\mathbb{Z}[1/2] = \{n/2^m \mid n \in \mathbb{Z}, m \geq 0\}$. We blow up the orbit of 0 to build a *new* real line, that is, we replace every point $o \in \mathbb{Z}[1/2]$ of the orbit of 0 by an interval J_o in such a way that, for every compact subset K , the sum $\sum_{o \in K} |J_o|$ is bounded (we are denoting by $|J|$ the length of the interval J). For our purpose, if $n/2^m$ is a reduced fraction, we set $|J_{n/2^m}| = \kappa/2^{2^m}$, for some constant $\kappa > 0$ to be fixed later. By convention the reduced fraction of 0 is $0/1$, so $|J_0| = \kappa/4$.

The maps f and g act naturally on the boundary points of the collection of intervals $\{J_o\}$, and this action can be extended to the *new* real line by taking linear interpolation and passing to the closure of the union of intervals. We put the new origin (still denoted by 0) at the midpoint of J_0 . The resulting maps are called \bar{f} and \bar{g} . Observe that since $n/2^m$ is a reduced fraction if and only if $1 + n/2^m = (2^m + n)/2^m$ is a reduced fraction, we have that the homeomorphism \bar{g} preserves the length of the intervals it permutes and hence \bar{g} is still a translation. By adjusting κ , and possibly replacing \bar{g} by some power of it, we can assume that \bar{g} is the translation by 2 and that $|J_0| \leq 1/4$.

Finally, we let $M \subseteq \mathbb{R}$ be the minimal invariant set for $\langle \bar{f}, \bar{g} \rangle$ action on the line. That is, M is closed, $\langle \bar{f}, \bar{g} \rangle$ -invariant, and contains no proper, closed, $\langle \bar{f}, \bar{g} \rangle$ -invariant subsets. In the present case, since J_0 is a wandering interval, and no $\langle \bar{f}, \bar{g} \rangle$ -orbit is discrete, the set M is locally a Cantor set which intersects trivially the $\langle \bar{f}, \bar{g} \rangle$ -orbit of J_0 (see [8]).

Remark 2.2. Observe that \bar{f} from Example 2.1 is the identity on J_0 . For our purpose it will be important to consider \bar{f} not being the identity on J_0 while preserving the action on the minimal set M from the original action (Figure 1, right). See Lemma 2.4 below.

Crucially, $\text{BS}(1, 2)$ admits a different family of actions without fixed points on the real line. These are the so-called *Conradian actions*, which are actions where α acts without fixed points and β fixes a fundamental domain of α ³ (see Figure 2). The following lemma formalizes the idea that for building a Conradian action one just need pick $\alpha \in \text{Homeo}_+(\mathbb{R})$ and *any other* homeomorphism

³In fact, any faithful action of $\text{BS}(1, 2)$ without global fixed points is either of affine type or Conradian, see [14].




FIGURE 1. The standard affine action of $\text{BS}(1,2)$ (left) and an action *à la Denjoy* (right).

defined on a fundamental domain of α which is then *extended* to a homeomorphism of the line using the relation of $\text{BS}(1,2)$. For the statement, we denote by $\text{Homeo}_0(I)$ the group of homeomorphisms of an open subset $I \subset \mathbb{R}$ which are isotopic to the identity. Also, given a homeomorphism $h : I \rightarrow I$, we define its *support* as the open subset $\text{supp}(h) = \{x \in I \mid h(x) \neq x\}$.




FIGURE 2. An example of Conradian action of $\text{BS}(1,2)$ on the real line.

Lemma 2.3. *Given $f \in \text{Homeo}_+(\mathbb{R})$, an open set I such that $f^n(I) \cap I = \emptyset$ for every $n \neq 0$, and $h \in \text{Homeo}_0(I)$, there exists a map $R_1 : (f, I, h) \mapsto g_h \in \text{Homeo}_+(\mathbb{R})$ such that*

- (i) $g_h(x) = h(x)$ for every $x \in I$,
- (ii) g_h is the identity outside $\bigcup_{n \in \mathbb{Z}} f^n(I)$,
- (iii) $f g_h f^{-1}(x) = g_h^2(x)$ for every $x \in \mathbb{R}$.

Proof. We need to define g_h . On $f^{-n}(I)$ ($n > 0$) we define g_h as $f^{-1}h^{2^n}f$ and on $f^n(I)$ we want to define g_h as $f \cdot 2\text{root}^{(n)}(h) \cdot f^{-1}$, where $2\text{root}(h)$ is the square root of h and $2\text{root}^{(n)} = 2\text{root} \circ \dots \circ 2\text{root}$ n -times.

Certainly, every $h \in \text{Homeo}_+(\mathbb{R})$ has a square root in $\text{Homeo}_+(\mathbb{R})$. For instance, if h has no fixed points, then it is conjugate to a translation and $2\text{root}(h)$ can be then defined as half of the translation conjugated back, while for general h one restricts the attention to connected components of the support of h and repeats the argument locally. The problem is that the square root is highly not

unique (for instance, the translation to which a homeomorphism h without fixed points is conjugate to is not unique), so 2root is not well defined (so far).

To give a precise definition of $\text{2root} : \text{Homeo}_0(I) \rightarrow \text{Homeo}_0(I)$ (and hence a precise definition of g_h) we proceed as follows. For every connected component C of the support of $h \in \text{Homeo}_0(I)$, let $x_0 \in C$ be its midpoint (this is well-defined, as from $f(I) \cap I = \emptyset$, every component C is a bounded interval). Assume $h(x_0) > x_0$ (the case $h(x_0) < x_0$ is analogous). Then define a measure μ_0 on $[x_0, h(x_0))$, as the push forward of the Lebesgue measure Leb on $[0, 1)$ by the unique affine map A sending $[0, 1)$ to $[x_0, h(x_0))$ bijectively. So $\mu_0(X) = \text{Leb}(A^{-1}(X))$ for every Borel subset $X \subset [x_0, h(x_0))$. Then there is a unique way of extending μ_0 to an h -invariant measure over C . The resulting measure μ_C is fully supported on C , gives finite mass to compact subsets of C , and has not atoms. We can now use μ_C to conjugate the restriction of h to C to the translation by 1 on the real line. Indeed if we let $F(x) = \text{sign}(x - x_0)\mu([x_0, x))$, then FhF^{-1} is the translation by 1. Then we can take the square root inside the group of translations and conjugate back. This is our preferred 2root . \square

2.1. Choosing b and d : Let J_0 and $\bar{g} : x \mapsto x + 2$ from Example 2.1, and let $\psi : \mathbb{R} \rightarrow (-1, 1)$ be the homeomorphism given by the interpolation of

$$\psi(n) = \begin{cases} 0 & \text{if } n = 0, \\ \sum_{i=1}^n 2^{-i} & \text{if } n > 0, \\ -\sum_{i=1}^{-n} 2^{-i} & \text{if } n < 0, \end{cases}$$

for $n \in \mathbb{Z}$. Let $\beta_0 = \psi \circ \bar{g} \circ \psi^{-1} \in \text{Homeo}_+(-1, 1)$, and call I_0 the image of J_0 under ψ . Call $I_1 = I_0 + 1$ and define $\beta_1 \in \text{Homeo}_+(0, 2)$ by $\beta_1(x) = \beta_0(x - 1) + 1$. Note that I_0 and I_1 are disjoint intervals, one centered at 0, and the other centered at 1, and the maps β_0, β_1 , are piecewise linear. Finally, we extend β_0 and β_1 to homeomorphisms of the real line by imposing that they commute with the translation $T : x \mapsto x + 2$. The resulting homeomorphisms are still denoted β_0 and β_1 . We also let $\mathcal{I}_0 = \bigcup_{n \in \mathbb{Z}} I_0 + 2n$ and $\mathcal{I}_1 = \bigcup_{n \in \mathbb{Z}} I_1 + 2n$, which are disjoint open sets, and choose $N > 0$ such that

$$(2) \quad \beta_0^{\pm N}(\mathbb{R} \setminus \mathcal{I}_1) \subseteq \mathcal{I}_1 \text{ and } \beta_1^{\pm N}(\mathbb{R} \setminus \mathcal{I}_0) \subseteq \mathcal{I}_0.$$

By possibly enlarging N , we can (and will) also assume that for all $n \geq N$ we have

$$(3) \quad (\beta_i^{\pm n})'(x) \leq 1/2 \text{ for all } x \notin \mathcal{I}_{i+1},$$

where the indices are taken in $\mathbb{Z}/2\mathbb{Z}$. Note that by (2) and the ping-pong lemma, the group $\langle \beta_0^N, \beta_1^N \rangle$ is a free group inside $\text{Homeo}_+(\mathbb{R})$. We let $b = \beta_0^N$ and $d = \beta_1^N$.

2.2. Choosing a and c . For f and g homeomorphisms of the line, we let

$$|f - g|_\infty = \sup_{x \in \mathbb{R}} |f(x) - g(x)| + \sup_{x \in \mathbb{R}} |f^{-1}(x) - g^{-1}(x)|.$$

This “norm” may be infinite but in many cases it is bounded, so it becomes a true distance. This happens for instance on

$$\text{Homeo}_{+,T}(\mathbb{R}) = \{f \in \text{Homeo}_+(\mathbb{R}) \mid T \circ f = f \circ T\},$$

where T is a translation (which is our case). Hence, endowed with $|\cdot|_\infty$, $\text{Homeo}_{+,T}(\mathbb{R})$ becomes a complete Polish space. We will build a and c as the limit (in this topology) of homeomorphisms a_n and c_n ($n \geq 0$), where a_n (resp. c_n) will be perturbations of a_{n-1} (resp. c_{n-1}). We keep the notations from §2.1.

Let \bar{f}, \bar{g} be the homeomorphisms from Example 2.1 and M its minimal set. Let $a_{-1} = \psi \circ \bar{f} \circ \psi^{-1} \in \text{Homeo}_+(-1, 1)$, and $c_{-1} \in \text{Homeo}_+(0, 2)$ be defined by $c_{-1}(x) = a_{-1}(x - 1) + 1$. We extend a_{-1} and c_{-1} inside $\text{Homeo}_{+,T}(\mathbb{R})$ by imposing that they commute with the translation $T : x \mapsto x + 2$. We denote the resulting elements by a_0 and c_0 respectively. Observe that $a_0 b a_0^{-1}(x) = b^2(x)$ for

all $x \in \mathbb{R}$ and in the same way $c_0 d c_0^{-1}(x) = d^2(x)$ for all $x \in \mathbb{R}$. Note also that a_0 is supported on $\mathbb{R} \setminus \mathcal{I}_0$ and c_0 is supported on $\mathbb{R} \setminus \mathcal{I}_1$ (because \tilde{f} is the identity on J_0).

We now explain how to obtain a_n and c_n for $n \geq 1$. The construction of the sequences a_n and c_n , as well as the proof of its convergence are independent and completely analogous upon exchanging the roles of b and d (and replacing 0 indices by 1 indices). We only explain the case of a_n . Let $\mathcal{M}_0 = \bigcup_{n \in \mathbb{Z}} \psi(M) + 2n$. We will perform successive perturbations on a_0 on the complement of \mathcal{M}_0 .

Lemma 2.4. *There exists a map $R_2 : h \mapsto a_h \in \text{Homeo}_{+,T}(\mathbb{R})$ defined on elements of $\text{Homeo}_{+,T}(\mathbb{R})$ supported on \mathcal{I}_0 with the following properties.*

- (i) $a_h(x) = h(x)$ for every $x \in \mathcal{I}_0$.
- (ii) $a_h(x) = a_0(x)$ for $x \in \mathcal{M}_0$ and $a_h b a_h^{-1}(x) = b^2(x)$ for every $x \in \mathbb{R}$.
- (iii) R_2 is an isometry with respect to the $\|\cdot\|_\infty$ -norm: for any two $h, h' \in \text{Homeo}_{+,T}(\mathbb{R})$ supported on \mathcal{I}_0 , one has $|a_h - a_{h'}|_\infty = |h - h'|_\infty$.

Proof. Suppose that $h \in \text{Homeo}_{+,T}(\mathbb{R})$ is supported on \mathcal{I}_0 . Since \mathcal{I}_0 is contained in a fundamental domain of b (i.e. $b^n(\mathcal{I}_0) \cap \mathcal{I}_0 = \emptyset$ for $n \neq 0$), the infinite composition $h_1 := \prod_{n \in \mathbb{Z}} b^n h b^{-n}$ is a composition of homeomorphisms with pairwise disjoint support, all commuting with T . Hence h_1 is a well-defined element in $\text{Homeo}_{+,T}(\mathbb{R})$, supported on $\bigcup_{n \in \mathbb{Z}} b^n(\mathcal{I}_0)$, which furthermore commutes with b . We can then define $R_2(h) = a_h = a_0 h_1$ which satisfies item (i).

For (ii) we observe that $h_1(x) = x$ for all $x \in \mathcal{M}_0$ since $b^n \mathcal{I}_0$ and \mathcal{M}_0 are disjoint (see the end of Example 2.1) and h_1 is supported on $\bigcup b^n(\mathcal{I}_0)$. In particular $a_h(x) = a_0(x)$ for all $x \in \mathcal{M}_0$. We can also check that $a_h b(x) = a_0 h_1 b(x) = a_0 b h_1(x) = b^2 a_0 h_1(x) = b^2 a_h(x)$ for all $x \in \mathbb{R}$.

Finally, to show (iii) we note that

$$(4) \quad \sup_{x \in \mathbb{R}} |a_0 h_1(x) - a_0 h'_1(x)| = \sup_{n \in \mathbb{Z}} \sup_{x \in \mathcal{I}_0} |a_0 b^n h(x) - a_0 b^n h'(x)| = \sup_{n \in \mathbb{Z}} \sup_{x \in \mathcal{I}_0} |b^{2n} a_0 h(x) - b^{2n} a_0 h'(x)|.$$

Further, for $x \in \mathcal{I}_0$, the images $h(x), h'(x)$ belong to \mathcal{I}_0 , on which a_0 is the identity and, as \mathcal{I}_0 and \mathcal{I}_1 are disjoint, condition (3) implies that the supremum in the last term is achieved for $n = 0$. After these considerations, equality (4) reduces to

$$\sup_{x \in \mathbb{R}} |a_0 h_1(x) - a_0 h'_1(x)| = \sup_{n \in \mathbb{Z}} \sup_{x \in \mathcal{I}_0} |b^{2n} h(x) - b^{2n} h'(x)| = \sup_{x \in \mathcal{I}_0} |h(x) - h'(x)|.$$

Similarly we control the difference of inverses: we have

$$\sup_{x \in \mathbb{R}} |(a_0 h_1)^{-1}(x) - (a_0 h'_1)^{-1}(x)| = \sup_{x \in \mathbb{R}} |(h_1)^{-1}(x) - (h'_1)^{-1}(x)| = \sup_{n \in \mathbb{Z}} \sup_{x \in \mathcal{I}_0} |b^n(h)^{-1}(x) - b^n(h')^{-1}(x)|$$


which, by (3), equals $\sup_{x \in \mathcal{I}_0} |(h)^{-1}(x) - (h')^{-1}(x)|$. Putting the two pieces together, we obtain the equality $|a_h - a_{h'}|_\infty = |h - h'|_\infty$. \square

The iteration. We will use sequentially Lemma 2.3 and Lemma 2.4 to perturb a_0 outside \mathcal{M}_0 . To this extent we let $\text{Dom}_0 = \{f \in \text{Homeo}_{+,T}(\mathbb{R}) \mid f(x) = a_0(x) \text{ for } x \in \mathcal{M}_0\}$. Observe that Dom_0 is a closed subset of $\text{Homeo}_{+,T}(\mathbb{R})$ and for $f \in \text{Dom}_0$, we have that $f(\mathcal{I}_0) = \mathcal{I}_0$, and $f b f^{-1}(x) = b^2(x)$ for every $x \in \mathcal{M}_0$. In particular, for $f \in \text{Dom}_0$, we can define $\widetilde{R}_2(f) = R_2(f \upharpoonright_{\mathcal{I}_0})$, where $f \upharpoonright_{\mathcal{I}_0}$ denotes the restriction of f to \mathcal{I}_0 , extended as to be the identity outside \mathcal{I}_0 . Certainly \widetilde{R}_2 is an idempotent of Dom_0 (i.e. $\widetilde{R}_2^2 = \widetilde{R}_2$) and, from Lemma 2.4.iii, it also satisfies

$$(5) \quad |\widetilde{R}_2(f) - \widetilde{R}_2(g)|_\infty = |f \upharpoonright_{\mathcal{I}_0} - g \upharpoonright_{\mathcal{I}_0}|_\infty.$$

Note also that by (2), $d^n(\mathbb{R} \setminus \mathcal{I}_0) \subset \mathcal{I}_0$ for all $n \neq 0$ (and hence $d^n(\mathcal{M}_0) \subset \mathcal{I}_0$ for all $n \neq 0$). In particular, for $f \in \text{Dom}_0$ we can apply R_1 from Lemma 2.3 to define $\widetilde{R}_1(f) = R_1(d, \mathbb{R} \setminus \mathcal{I}_0, f \upharpoonright_{\mathbb{R} \setminus \mathcal{I}_0})$. Certainly \widetilde{R}_1 is also an idempotent of Dom_0 which, by (3), satisfies that

$$(6) \quad |\widetilde{R}_1(f) \upharpoonright_{\mathcal{I}_0} - \widetilde{R}_1(g) \upharpoonright_{\mathcal{I}_0}|_\infty \leq \frac{1}{2} |f \upharpoonright_{\mathbb{R} \setminus \mathcal{I}_0} - g \upharpoonright_{\mathbb{R} \setminus \mathcal{I}_0}|_\infty.$$

FIGURE 3. The action of Higman's group on the circle $\mathbb{S}^1 \cong \mathbb{R}/\langle T \rangle$.

Suppose that a_n is defined for $n \leq 2k$ ($k \geq 0$). We define $a_{2k+1} = \widetilde{R}_1(a_{2k})$ and $a_{2k+2} = \widetilde{R}_2(a_{2k+1})$.

Main Claim. *The sequence a_n converges in $\text{Homeo}_{+,T}(\mathbb{R})$.*

Before giving the proof note that if a_n converges to a , then a is both \widetilde{R}_2 and \widetilde{R}_1 invariant so, by Lemma 2.3.iii and Lemma 2.4.ii, it satisfies $aba^{-1} = b^2$ and $dad^{-1} = a^2$ over \mathbb{R} as desired.

Proof of Claim. First observe that, from (5) and (6), it follows that $\widetilde{R}_2\widetilde{R}_1$ and $\widetilde{R}_1\widetilde{R}_2$ are contractions by $1/2$, so each of them have unique (attracting) fixed points in $\text{Dom}_0 \subset \text{Homeo}_{+,T}(\mathbb{R})$, call them $p = \widetilde{R}_2\widetilde{R}_1(p)$ and $q = \widetilde{R}_1\widetilde{R}_2(q)$. Observe that since \widetilde{R}_1 and \widetilde{R}_2 are idempotents, it follows that $\widetilde{R}_1(q) = q$ and $\widetilde{R}_2(p) = p$. We need to show that $p = q$.

By the definition of Dom_0 , we have $p(x) = q(x) = a_0(x)$ for all $x \in \mathcal{M}_0$ and hence

$$|p \upharpoonright_{\mathbb{R} \setminus \mathcal{I}_0} - q \upharpoonright_{\mathbb{R} \setminus \mathcal{I}_0}|_\infty \leq |J|,$$

where J is the largest complementary interval of $\mathcal{M}_0 \cup \mathcal{I}_0$. In particular, applying \widetilde{R}_1 , we get from (6) that

$$|\widetilde{R}_1(p) \upharpoonright_{\mathcal{I}_0} - q \upharpoonright_{\mathcal{I}_0}|_\infty \leq \frac{1}{2}|J|,$$

and then, applying \widetilde{R}_2 , (5) gives

$$|p - \widetilde{R}_2(q)|_\infty \leq \frac{1}{2}|J|.$$

If we now repeat k times this process we get

$$|\widetilde{R}_1(p) \upharpoonright_{\mathcal{I}_0} - q \upharpoonright_{\mathcal{I}_0}|_\infty \leq |J|/2^k \quad \text{and} \quad |p - \widetilde{R}_2(q)|_\infty \leq |J|/2^k.$$

It follows that $p = \widetilde{R}_2(q)$ and hence $\widetilde{R}_1(p) = q$. But, respectively, this implies that $p = q$ over \mathcal{I}_0 (see Lemma 2.4.i) and over $\mathbb{R} \setminus \mathcal{I}_0$ (see Lemma 2.3.i). So $p = q$, and we declare $a = p = q$ as needed. \square

Remark 2.5. Observe that the four homeomorphisms a, b, c, d all commute with the translation T , so they define homeomorphisms of the circle $\mathbb{S}^1 \cong \mathbb{R}/_{T(x) \sim x}$, and hence a representation of H into $\text{Homeo}(\mathbb{S}^1)$. Further, since c and d do not share fixed points, the resulting action on the circle is without global fixed points. See Figure 3.

3. A FAITHFUL ACTION

In the action $\varphi : H \rightarrow \text{Homeo}_{+,T}(\mathbb{R})$ built in Section 2, the subgroup $F = \langle b, d \rangle$ acts as a lift of a ping pong action on the circle (with parabolic elements) so φ is an embedding in restriction to $\langle b, d \rangle$. This remark, together with Kurosh Theorem [15], give the following:

Lemma 3.1. *The kernel of the action φ is a free group and hence is left-orderable.*

Proof. The group H is the amalgamated product

$$(7) \quad H = \langle d, a, b \mid dad^{-1} = a^2, aba^{-1} = b^2 \rangle *_{\langle b, d \rangle} \langle b, c, d \mid bcb^{-1} = c^2, cdc^{-1} = d^2 \rangle.$$

By the preceding remark, $\ker(\varphi)$ has trivial intersection with the edge group $F = \langle b, d \rangle$. Therefore we can apply Kurosh Theorem to $\ker(\varphi)$ in the amalgamated product (7) (in the case of a normal subgroup, it is enough to look at the intersection with the edge group): it gives that $\ker(\varphi)$ is the free product of a free group with intersections of $\ker(\varphi)$ with conjugates of the vertex groups $H_1 = \langle d, a, b \mid dad^{-1} = a^2, aba^{-1} = b^2 \rangle$ and $H_2 = \langle b, c, d \mid bcb^{-1} = c^2, cdc^{-1} = d^2 \rangle$. It is then enough to prove that the intersection of $\ker(\varphi)$ with every conjugate of H_1 and H_2 is also a free group. Observe that by normality of $\ker(\varphi)$, it is enough to prove that the intersections $\ker(\varphi) \cap H_1$ and $\ker(\varphi) \cap H_2$ are free. For this, we repeat nearly the same argument.

Indeed, the group $H_1 = \langle d, a, b \mid dad^{-1} = a^2, aba^{-1} = b^2 \rangle$ is also an amalgamated product

$$(8) \quad H_1 = \langle d, a \mid dad^{-1} = a^2 \rangle *_{\langle a \rangle} \langle a, b \mid aba^{-1} = b^2 \rangle,$$

and the two vertex groups $\langle d, a \mid dad^{-1} = a^2 \rangle, \langle a, b \mid aba^{-1} = b^2 \rangle$ are two copies of $\mathbf{BS}(1, 2)$ which act faithfully by construction of the action (every proper quotient of $\mathbf{BS}(1, 2)$ is abelian, and certainly this is not the case for this action). Therefore the intersection of $\ker(\varphi)$ with the vertex group in H_1 is trivial and this implies that $\ker(\varphi) \cap H_1$ is a free group. The same argument works for the intersection with H_2 . This shows that $\ker(\varphi)$ is a free group (potentially trivial).

Finally, the fact that free groups are left-orderable is well known and can be found in [4, 8]. \square

Proof of Theorem A. As a consequence, the group H fits into the short exact sequence

$$1 \rightarrow \ker(\varphi) \rightarrow H \rightarrow \varphi(H) \rightarrow 1,$$

where $\ker(\varphi)$ and $\varphi(H)$ are both left-orderable after Lemma 3.1. This implies that H is left-orderable as well: denote by \preceq_φ and \preceq_{\ker} the left-invariant orders on $\varphi(H)$ and $\ker(\varphi)$ respectively; then define the order \preceq on H by declaring $id \preceq h$ if and only if

- $h \in \ker(\varphi)$ and $id \preceq_{\ker} h$, or
- $h \notin \ker(\varphi)$ and $\varphi(id) \preceq_\varphi \varphi(h)$,

and extending it so that it is left-invariant (declaring $h \preceq g$ if and only if $id \preceq h^{-1}g$; the fact that \preceq_φ and \preceq_{\ker} are left-invariant implies that the definition is consistent, see [4] for more details). Finally, every countable left-orderable group is isomorphic to a subgroup of $\mathbf{Homeo}_+(\mathbb{R})$ [8], therefore H admits faithful actions on the real line. \square

We can also show the analogous result for the circle.

Proposition 3.2. *H admits a faithful action on the circle without global fixed points.*

Sketch of proof. The proof follows the same lines as for Theorem A:

- (1) For countable groups, acting faithfully on the circle by orientation preserving homeomorphisms is equivalent to admit a circular order invariant under left-multiplications [3, 13].
- (2) If $1 \rightarrow L \rightarrow G \rightarrow C \rightarrow 1$, where C is a circularly orderable group and L is a left-orderable group, then G is circularly orderable [3].

In our case we have that $1 \rightarrow \ker(\varphi) \rightarrow H \rightarrow \bar{\varphi}(H) \rightarrow 1$, where $\bar{\varphi} : H \rightarrow \mathbf{Homeo}(\mathbb{S}^1)$ is the projection of φ to $\mathbb{S}^1 \cong \mathbb{R}_{T(x) \sim x}$, see Remark 2.5. Since $\bar{\varphi}(H)$ acts without global fixed points, the same is true for actions coming from circular orders obtained from the preceding short exact sequence [13]. \square

4. MANY INEQUIVALENT ACTIONS

For the construction of φ , in Section 2, the initial point was the choice of b and d starting from T -periodic extensions of $\beta_0 \in \text{Homeo}_+([-1, 1])$ and $\beta_1 \in \text{Homeo}_+([0, 2])$. It is possible to get an analogous construction of φ by imposing milder periodicity conditions to the extensions of β_0 and β_1 to the real line. Concretely, for arbitrary sequences $(\epsilon_n), (\delta_n) \in \{-1, +1\}^{\mathbb{Z}}$, we can consider extensions of β_0, β_1 verifying

$$\begin{aligned} \beta_0(T^n(x)) &= T^n(\beta_0^{\epsilon_n}(x)) \quad \text{for every } n \in \mathbb{Z} \setminus \{0\} \text{ and every } x \in [-1, 1], \\ \beta_1(T^n(x)) &= T^n(\beta_1^{\delta_n}(x)) \quad \text{for every } n \in \mathbb{Z} \setminus \{0\} \text{ and every } x \in [0, 2]. \end{aligned}$$

The effect in the corresponding choice of the new $b = \beta_0^N$ and $d = \beta_1^N$, is that, locally, they could be switched to the inverse of the old b and d (much in the spirit of [10]). This new b and d (and hence the new a and c) commute with a translation (and hence project to homeomorphisms of the circle) if and only if (ϵ_n) and (δ_n) have a common period. Moreover, any two distinct choices of sequences (which are not shift-equivalent) give actions which are not semi-conjugate one to another.

5. SMOOTH ACTIONS

In this section we give the proof of Theorem B. We will first establish the result for actions on the real line.

Proposition 5.1. *Every representation $\rho : H \rightarrow \text{Diff}^1(\mathbb{R})$ is trivial.*

Our proof relies on the following:

Lemma 5.2. *Let $\psi : H \rightarrow \text{Homeo}(\mathbb{R})$ be a non-trivial representation without global fixed points. Then there is a copy B of the Baumslag-Solitar group $\text{BS}(1, 2)$ inside H such that the restriction of ψ to B is faithful, $\psi(B)$ preserves a compact interval I , and the restriction of the action of $\psi(B)$ to I is not semi-conjugate to the standard affine action.*

Before proving the lemma, let us explain how to deduce the proposition.

Proof of Proposition 5.1. As pointed out before, H contains several copies of $\text{BS}(1, 2) = \langle \alpha, \beta \mid \alpha\beta\alpha^{-1} = \beta^2 \rangle$. After [1] every action of $\text{BS}(1, 2)$ inside $\text{Diff}_+^1(I)$, where I is a compact interval, must be (topologically) conjugate to the standard affine action. By Lemma 5.2 there is a copy of $\text{BS}(1, 2)$, call it $B \leq H$, that preserves a compact interval I and, moreover, the action of B on I is not conjugate to an affine action (it is an action by levels in the sense of Lemma 2.3). In particular, no action of H on the line is conjugate to an action by C^1 -diffeomorphisms. \square

Let us next prove the lemma.

Proof of Lemma 5.2. For the proof it will be convenient to define $H = \langle a_i \mid a_i a_{i+1} a_i^{-1} = a_{i+1}^2 \rangle$ where the indices are taken inside $\mathbb{Z}/4\mathbb{Z}$. Let $\psi : H \rightarrow \text{Homeo}_+(\mathbb{R})$ be a non-trivial representation.

Claim 1. *The set of fixed points of each $\psi(a_i)$ accumulates on $\pm\infty$.*

Proof of Claim. We argue by contradiction. Without loss of generality, we will assume that the fixed points of one generator do not accumulate on $+\infty$. After re-indexation, we may also assume that this generator is a_3 . Note that the relation $a_2 a_3 a_2^{-1} = a_3^2$ implies that the subset $\text{Fix}(\psi(a_3)) = \{x \in \mathbb{R} \mid \psi(a_3)(x) = x\}$ is $\psi(a_2)$ -invariant. Therefore, $\psi(a_2)$ maps any connected component of the complement $\text{supp}(\psi(a_3)) = \mathbb{R} \setminus \text{Fix}(\psi(a_3))$ to a (not necessarily different) connected component of $\text{supp}(\psi(a_3))$.

Let I be an unbounded component of the support of $\psi(a_3)$. Then, by the previous remark, $\psi(a_2)$ must either preserve or move I off itself. Since I is unbounded, we conclude that $\psi(a_2)$ preserves I . It follows from [14] that in this case the action of $\langle \psi(a_2), \psi(a_3) \rangle$ on I is *semi-conjugate* to the

standard affine action of $\mathbf{BS}(1, 2)$ on \mathbb{R} (i.e. where a_3 acts as a translation and a_2 as a dilation by 2), that is there is a continuous and surjective map $\alpha : I \rightarrow \mathbb{R}$ such that

$$\alpha \circ \psi(a_2)(x) = 2\alpha(x), \text{ and } \alpha \circ \psi(a_3)(x) = \alpha(x) + 1.$$

In particular, $\psi(a_2)(x) > \psi(a_3)(x)$ for every x large enough, and hence the set of fixed points of $\psi(a_2)$ does not accumulate on $+\infty$. We can therefore repeat the argument with a_2 and a_1 , then with a_1 and a_0 and then with a_0 and a_3 (this time $\psi(a_3)$ acting as dilation), to get

$$\psi(a_3)(x) < \psi(a_2)(x) < \psi(a_1)(x) < \psi(a_0)(x) < \psi(a_3)(x)$$

for all x large enough. A contradiction. \square

Observe that no $\psi(a_i)$ can be the identity, otherwise the image $\psi(H)$ would be trivial. Therefore every open support $\text{supp}(\psi(a_i)) = \{x \in \mathbb{R} \mid \psi(a_i)(x) \neq x\}$ is non-empty, and by Claim 1 all connected components are bounded intervals.

Claim 2. *There is a pair of consecutive generators a_i, a_{i+1} such that $\psi(a_i)$ moves off itself a connected component of the open support of $\psi(a_{i+1})$.*

Proof of Claim. Let I_3 be a connected component of $\text{supp}(\psi(a_3))$. Arguing as in the proof of Claim 1, we see that if $\psi(a_2)$ does not preserve I_3 , then it must move it off itself, so we are done. So suppose that $\psi(a_2)$ fixes I_3 . Then the action of $\langle \psi(a_2), \psi(a_3) \rangle$ on I_3 is semi-conjugate to the standard affine action of $\mathbf{BS}(1, 2)$ (where $\psi(a_3)$ is acting as the translation), and hence there is a connected component of $\text{supp}(\psi(a_2))$ which is strictly contained in I_3 . If $\psi(a_1)$ does not move I_2 off itself, then we find a connected component $I_1 \subsetneq I_2$ of $\text{supp}(\psi(a_1))$. Again, if $\psi(a_0)$ does not move I_1 off itself, we find a connected component $I_0 \subsetneq I_1 \subsetneq I_2 \subsetneq I_3$ of $\text{supp}(\psi(a_0))$. But then $I_0 \subsetneq \text{supp}(\psi(a_3))$, so $\psi(a_3)$ cannot preserve I_0 , and therefore it must move I_0 off itself. \square

Let i be the index given by Claim 2, and let J be a connected component of $\text{supp}(\psi(a_{i+1}))$ which is moved off itself by $\psi(a_i)$. Let $I \subset \mathbb{R}$ be the smallest interval containing all the images $\psi(a_i)^n(J)$, $n \in \mathbb{Z}$. Claim 1 applied to a_i implies that I is bounded, and moreover its choice gives that it is preserved by both $\psi(a_i)$ and $\psi(a_{i+1})$. Clearly the restriction of the action of $\langle \psi(a_i), \psi(a_{i+1}) \rangle \cong \mathbf{BS}(1, 2)$ to I is faithful, and is not semi-conjugate to the standard affine action. \square

We can finally conclude.

Proof of Theorem B. A direct consequence of Proposition 5.1 is that the analogous result holds for the circle, namely that every representation $\psi : H \rightarrow \text{Diff}^1(\mathbb{S}^1)$ is trivial. This follows from the well-known fact that Higman's group is acyclic [5], in particular $H^2(H, \mathbb{Z}) = 0$ and every (continuous) action on the circle lifts to an action on the real line (the obstruction to such a lift is represented by the Euler class of the action, which is a 2-cocycle [8]).

We can also use a more dynamical argument. For this, suppose such ψ is given.

Claim 3. *For every $i \in \mathbb{Z}/4\mathbb{Z}$, the group $\langle \psi(a_i), \psi(a_{i+1}) \rangle$ has a global fixed point in \mathbb{S}^1 .*

Proof of Claim. Observe that the group $\langle a_i, a_{i+1} \rangle \cong \mathbf{BS}(1, 2)$ is amenable. This implies that there exists a Borel probability measure μ_i on \mathbb{S}^1 which is invariant by the group $\langle \psi(a_i), \psi(a_{i+1}) \rangle$, and moreover, the rotation number restricts to a group homomorphism $\text{rot} : \langle \psi(a_i), \psi(a_{i+1}) \rangle \rightarrow \mathbb{R}/\mathbb{Z}$ (see [8]). We deduce that

$$\text{rot } \psi(a_{i+1}) = \text{rot } \psi(a_i a_{i+1} a_i^{-1}) = \text{rot } \psi(a_{i+1}^2) = 2 \text{rot } \psi(a_{i+1}),$$

so that $\text{rot } \psi(a_{i+1}) = 0$, implying that $\psi(a_{i+1})$ has a fixed point. Cycling over the indices, we get that also $\psi(a_i)$ has a fixed point, and thus the probability measure μ_i must be supported on points that are fixed by both $\psi(a_i)$ and $\psi(a_{i+1})$, and thus by $\langle \psi(a_i), \psi(a_{i+1}) \rangle$. \square

From [1] we know that for every maximal interval $I \subset \mathbb{S}^1$ on which $\langle \psi(a_i), \psi(a_{i+1}) \rangle$ has no global fixed point, we either have

- (1) $\psi(a_{i+1})$ is the identity in restriction to I or
- (2) the restriction to I of the action of $\langle \psi(a_i), \psi(a_{i+1}) \rangle$ is conjugate to the standard affine action, and moreover the derivative of $\psi(a_i)$ at the internal fixed point equals 2.

Continuity of derivatives implies that there are only finitely many intervals of the second type. Therefore $\text{supp}(\psi(a_{i+1}))$ has finitely many connected components. Repeating the argument with the subgroup $\langle \psi(a_{i+1}), \psi(a_{i+2}) \rangle$, we obtain that not only $\text{supp}(\psi(a_{i+2}))$ has finitely many components, but moreover, they are strictly less than those of $\text{supp}(\psi(a_{i+1}))$. Cycling over the indices, one gets a contradiction. \square

REFERENCES

- [1] C. BONATTI, I. MONTEVERDE, A. NAVAS, C. RIVAS. Rigidity for C^1 actions arising from hyperbolicity I: solvable groups. *Math. Z.* **286** (2017), 919–949.
- [2] M. BUCHER, R. FRIGERIO, T. HARTNICK. A note on semi-conjugacy for circle actions, *Enseign. Math.* **62** (2016), 317–360.
- [3] D. CALEGARI, Circular groups, planar groups, and the Euler class. *Proc. of the Casson Fest (Geometry and Topology Monographs)* **7** (2004), 431–491.
- [4] A. CLAY, D. ROLFSEN. Ordered Groups and Topology. *Graduate Studies in Mathematics* 176, AMS (2016).
- [5] E. DYER, A.T. VASQUEZ. Some small aspherical spaces. *J. Austral. Math. Soc.* **16** (1973), 332–352.
- [6] R. FRIGERIO. Bounded cohomology of discrete groups. *Mathematical Surveys and Monographs* 227, AMS (2017).
- [7] K. FUJIWARA. The second bounded cohomology of an amalgamated free product of groups. *Trans. Amer. Math. Soc.* **352** (1999), 1113–1129.
- [8] É. GHYS. Groups acting on the circle. *Enseign. Math.* **47** (2001), 329–407.
- [9] G. HIGMAN. A finitely generated infinite simple group. *J. Lond. Math. Soc.* **26** (1951), 61–64.
- [10] J. HYDE, Y. LODHA. Finitely generated infinite simple groups of homeomorphisms of the real line. *Invent. Math.*, to appear.
- [11] A. I. MAL'CEV. On isomorphic matrix representation of infinite groups of matrices (Russian). *Math. Sb.* **8** (1940), 405–422, & *Amer. Math. Soc. Transl.* **45** (1965), 1–18.
- [12] K. MANN. Rigidity and flexibility of group actions on the circle. *Handbook of group actions, Vol. IV* (2018), 705–752.
- [13] K. MANN, C. RIVAS. Circular orders, dynamics and rigidity. *Ann. Inst. Fourier (Grenoble)* **68** (2018), 1399–1445.
- [14] C. RIVAS. On the space of Conradian group orderings. *J. of Group Theory.* **13** (2010), 337–353.
- [15] J.-P. SERRE. Arbres, amalgames, SL_2 (English summary), rédigé avec la collaboration de H. BASS, *Astérisque* **46** (1977).
- [16] W. THURSTON. A generalization of the Reeb stability theorem. *Topology* **13** (1974), 347–352.

Cristóbal Rivas

Dpto. de Matemáticas y C.C., Universidad de Santiago de Chile
Alameda 3363, Estación Central, Santiago, Chile
`cristobal.rivas@usach.cl`

Michele Triestino

Institut de Mathématiques de Bourgogne (IMB, UMR 5584)
9 av. Alain Savary, 21000 Dijon, France
`michele.triestino@u-bourgogne.fr`