



Beyond mortality: Sterility as a neglected component of parasite virulence

Jessica Lee Abbate, Sarah Kada, Sébastien Lion

► To cite this version:

Jessica Lee Abbate, Sarah Kada, Sébastien Lion. Beyond mortality: Sterility as a neglected component of parasite virulence. PLoS Pathogens, 2015, 11 (12), pp.e1005229. 10.1371/journal.ppat.1005229 . hal-02122395

HAL Id: hal-02122395

<https://hal.science/hal-02122395>

Submitted on 21 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

OPINION

Beyond Mortality: Sterility As a Neglected Component of Parasite Virulence

Jessica L. Abbate^{1,2*}, Sarah Kada¹, Sébastien Lion¹

1 Centre d'Écologie Fonctionnelle et Évolutive (CEFE), CNRS-Université de Montpellier- Université Paul-Valéry Montpellier-EPHE, Montpellier, France, **2** Institute of Ecology and Evolution, University of Bern, Bern, Switzerland

* jessica.abbate@iee.unibe.ch

Abstract

Virulence is generally defined as the reduction in host fitness following infection by a parasite (see [Box 1](#) for glossary) [1]. In general, parasite exploitation of host resources may reduce host survival (mortality virulence), decrease host fecundity (sterility virulence), or even have sub-lethal effects that disturb the way individuals interact within a community (morbidity) [2,3]. In fact, the virulence of many parasites involves a combination of these various effects ([Box 2](#)). In practice, however, virulence is most often defined as disease-induced mortality [1, 4–6]. This is especially true in the theoretical literature, where the evolution of sterility virulence, morbidity, and mixed strategies of host exploitation have received relatively little attention. While the focus on mortality effects has allowed for easy comparison between models and, thus, rapid advancement of the field, we ask whether these theoretical simplifications have led us to inadvertently minimize the evolutionary importance of host sterilization and secondary virulence effects. As explicit theoretical work on morbidity is currently lacking (but see [7]), our aim in this Opinion piece is to discuss what is understood about sterility virulence evolution, its adaptive potential, and the implications for parasites that utilize a combination of host survival and reproductive resources.



OPEN ACCESS

Citation: Abbate JL, Kada S, Lion S (2015) Beyond Mortality: Sterility As a Neglected Component of Parasite Virulence. *PLoS Pathog* 11(12): e1005229. doi:10.1371/journal.ppat.1005229

Editor: Glenn F Rall, The Fox Chase Cancer Center, UNITED STATES

Published: December 3, 2015

Copyright: © 2015 Abbate et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Agence Nationale pour la Recherche grant 10-PDOC-017-01 \textsc{SpatEvolEpid} to SL, The University of Bern for postdoctoral support of JLA, and by a PhD grant from French Ministry of Research to SK. These are all academic institutions providing support to the authors for the time in which the manuscript was completed. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Box 1. Glossary

Fitness: The expected number of adult offspring contributed to the next generation by a focal genotype or phenotype in a given environment.

Virulence: Any reduction in host fitness following infection by a parasite. This includes parasite-induced mortality, sterility, and morbidity (other sub-lethal effects of infection).

Transmission rate: The rate at which susceptible hosts become infected when encountering infected hosts.

Basic reproductive ration (R_0): The expected number of secondary infections produced by an infected host in an otherwise uninfected population. If R_0 is strictly larger than 1, the disease can spread.

Parasite fitness: The expected number of secondary infections produced by a given parasite genotype in a population infected by one or several resident genotypes. As explained in [Box 3](#), there is a simple link between R_0 and parasite fitness only when the epidemiological feedbacks are of a particularly simple kind.

Epidemiological feedback: Any environmental variable, such as the density of susceptible or infected hosts, that affects parasite fitness.

Box 2. Examples of Pleiotropic Virulence Components

Beyond reducing resources allocated to host maintenance and growth (mortality virulence and morbidity), parasite-induced reduction in fecundity (sterility virulence) is a widely-studied and important consequence of host exploitation (reviewed in [\[13\]](#)). Additionally, many parasites have been shown to impact both survival and reproduction of their hosts. For example, commonly sterilizing syphilis and gonorrhea infections in humans [\[57\]](#) caused much mortality and morbidity in the pre-antibiotic era [\[58–60\]](#). Likewise, the bacterium *Pasteuria ramosa* is considered a “sterilizing pathogen” (e.g., [\[61\]](#)) of the small crustacean *Daphnia magna*, as the host produces parasite spores instead of eggs; nevertheless, *P. ramosa* also causes premature death, a process necessary for transmission of the spores [\[62–65\]](#). Other examples include everything from nematode infections of red grouse [\[66\]](#) to viral, crop-destroying infections of their insect vectors [\[67\]](#) and *Puccinia* spp. fungal infections of European weeds [\[68\]](#). While for many pathogens sterilization may only be a consequence of host response to infection, some are able to manipulate such responses [\[69\]](#), and specific targeting of diverse host tissues can also occur. For instance, highly prevalent human herpes simplex viruses (HSV-1 and HSV-2) infecting a wide range of neuronal, ocular, mucosal, and epithelial tissues [\[70\]](#) also localize in sperm and spermatozoa, likely affecting male infertility [\[71\]](#) in addition to juvenile and adult morbidity and mortality [\[72,73\]](#). Similarly, death and morbidity from the human immunodeficiency virus (HIV) infection of blood cells receives a great deal of much-deserved attention, but very little has been done to investigate the potential fertility impacts of elevated sperm abnormalities associated with asymptomatic infection [\[74\]](#).

Diseases in which morbidity effects occur in addition to mortality and/or sterility are far more obvious, if not ubiquitous (e.g. sores, rashes, fever, etc.). Morbidity is often measured in addition to, or as a proxy for, mortality effects, such as in experiments on the evolution of virulence in malaria (e.g., maximum number of red blood cells destroyed [\[75\]](#)), or for “aggressiveness,” as it is termed for phytopathogens (e.g., the number or size of tissue lesions, reviewed in [\[76\]](#)).

Basic Theory

A key quantity in epidemiology is the basic reproductive ratio, R_0 , which is defined as the average number of secondary infections produced by an infected host in an otherwise uninfected population [\[8,9\]](#). In most classical models of horizontally transmitted infections, R_0 is simply a function of parasite transmission and host survival ([Box 3](#)). Early evolutionary models

emphasized a particularly simple result: selection should favor the parasite strain that maximizes R_0 (Box 3). Consequently, the evolution of virulence is solely constrained by the shape of the trade-off between parasite transmission and host survival [6]. Importantly for our purpose, parasite fitness is then predicted to be independent of host fecundity.

Box 3. Parasite Fitness, R_0 Maximization, and Epidemiological Feedbacks

R_0 and parasite fitness. Classical epidemiological theory shows that parasite persistence in the host population depends on the average number of secondary infections produced by a parasite in an uninfected population [6,9]. This defines the basic reproductive ratio, R_0 . In most textbooks, the following expression is found:

$$R_0 = \frac{\beta}{\mu + \alpha + \gamma}. \quad (1)$$

Simply put, R_0 is the product of the transmission rate, β , and of the average lifespan of infected hosts, $1 / (\mu + \alpha + \gamma)$, where μ is intrinsic mortality, α is mortality virulence, and γ is the clearance rate.

In fact, for many classical models, there is often a simple link between parasite fitness and R_0 [6,9], and one may write

$$\mathcal{W} = \frac{\beta_m}{\mu + \alpha_m + \gamma_m} \hat{S} = R_{0,m} \hat{S}, \quad (2)$$

where the m subscript indicates the dependency of the traits on parasite genotype. Eq 2 shows that the fitness of the focal mutant strain, \mathcal{W} , is simply the product of the mutant parasite's basic reproductive ratio, $R_{0,m}$, and of the density, \hat{S} , of susceptible hosts in the population at equilibrium. The mutant parasite is then favored by selection if $\mathcal{W} > 1$. The dependency on \hat{S} illustrates the role of epidemiological feedbacks: parasite fitness depends not only on how well a parasite strain performs in an uninfected population (which is measured by R_0), but also on how many susceptible hosts it encounters in an already infected population.

R_0 maximization. Under the simplest ecological scenarios typically assumed, \hat{S} does not depend on host fecundity and is simply given by $1 / R_{0,w}$, where $R_{0,w}$ is the basic reproductive ratio of the wild-type, resident strain. Then, the parasite fitness can be written as a ratio of basic reproductive ratios,

$$\mathcal{W} = \frac{R_{0,m}}{R_{0,w}}, \quad (3)$$

and the strain with the larger R_0 is selected for. Evolution will then lead to a gradual increase in R_0 , until the maximal R_0 is reached. The precise optimal strategy of the parasite is determined by the trade-off between epidemiological parameters such as transmission, recovery, and virulence [6,9]. For our purpose, an important implication is that host fecundity does not affect the evolutionary outcome, which only depends on parasite transmission and on the survival of infected hosts.

Epidemiological feedbacks: an example. It is important to realize that this simple result only holds true under very restrictive epidemiological assumptions (see main text). Consider multiple infections, for instance. If infected hosts can be infected with

probability σ (superinfection), the fitness of a mutant strain takes the following form [39]:

$$\mathcal{W} = \frac{\beta_m (\hat{S} + \sigma \hat{I})}{\mu + \alpha_m + \gamma_m + \sigma \beta_w \hat{I}}. \quad (4)$$

Now, parasite fitness depends on the densities of both susceptible and infected hosts. Consequently, host fecundity may feed back on the density of infected hosts and affect the evolutionary outcome [19,39,77].

We argue that this textbook result may have played a role in fastening attention on mortality virulence. Within this framework, sterility virulence can only evolve if there is a trade-off between sterilization and the other life-history traits (transmission and survival) [10]. For instance, there is evidence that sterilized, infectious hosts may sometimes have higher contact rates because of increased longevity or promiscuity [11–13], particularly for sexually transmitted infections [14]. If there is a positive linear relationship between transmission and sterilization, parasites are predicted to fully sterilize their host whenever possible [10,15]—a take-home message that has been largely influential in the literature (e.g., [16]). However, the key finding from these theoretical studies was that other ecological assumptions can favor sub-maximal sterility, and empirical evidence suggests that sterilization need not be perfect [17].

As a result of this focalization on mortality virulence, the potential adaptive nature of sterility virulence for the parasite has often been downplayed (but see [18]), and observed changes in host fecundity following infection have generally been interpreted as resulting from various host responses. For instance, hosts may adjust their resource allocation between fecundity and survival following infection [18–22], or suffer reduced fecundity because of immunopathology [23,24]. Likewise, incomplete host sterility following infection by a “sterilizing pathogen” is usually explored as a result of host resistance mechanisms [25] or imperfect specificity [16], rather than the evolution of a less virulent parasite. We think this is not the full picture. In the remainder of this essay, we stress that epidemiological feedbacks may have non-trivial effects on the evolution of two key aspects of parasite virulence: which host life-history traits are affected (e.g., mortality and/or fecundity) and the magnitude of virulence (e.g., to what extent host survival or fecundity is reduced).

Implications of Realistic Epidemiological Feedbacks

Importantly, the simple link between R_0 and parasite fitness (explained in Box 3) only holds true under relatively stringent assumptions about the feedback between epidemiological dynamics and selection. Theoretical studies show that realistic epidemiological feedbacks will cause parasite fitness to depend on additional factors that are likely to be affected by host fecundity. Thus, selective effects due to changes in host fecundity cannot be dismissed a priori. Although this has long been recognized for vertically transmitted parasites, the fitness of which necessarily depends on host reproduction [15], the importance of host fecundity for horizontally transmitted parasites seems to have been underplayed.

Here, we highlight four key ecological factors under which host fecundity is predicted to affect parasite fitness.

Spatial structure

The theory developed in [Box 3](#) considers a well-mixed population where all parasite strains have access to the same amount of susceptible hosts. However, most host–parasite interactions are spatially structured, and this causes different parasite strains to experience different densities of susceptible hosts. As a result, R_0 is no longer maximized by selection because of indirect kin selection effects [\[26,27\]](#). Kin selection can then favor intermediate levels of sterilization [\[10,28\]](#), because higher fecundity in infected hosts will tend to relax local competition for susceptible hosts [\[27\]](#). The key point is that, in spatially structured populations, parasites that differ in their allocation between mortality and sterility virulence will tend to have different opportunities for horizontal transmission.

Epidemiological fluctuations

Many natural host–parasite interactions exhibit fluctuations in host densities, either because of intrinsic nonlinear dynamics or because of external forcing (e.g., seasonality, variation in vector density, etc.). Such non-equilibrium dynamics may, for instance, cause fluctuations in the density of susceptible hosts, thereby affecting short- and long-term selective pressures on parasite traits [\[29–34\]](#). In general, one may expect the non-equilibrium density of susceptible hosts to be affected by the fecundity of infected individuals. By taking this into account, novel theoretical approaches building on quantitative genetics methods [\[30,32\]](#) may help shed light on the origin and maintenance of sterilizing parasites.

Multiple infections

Most natural infections consist of several parasite strains or species [\[35–37\]](#). In hosts infected by multiple parasite strains, the fitness of a focal strain will be reduced by both host mortality and by the rate at which susceptible hosts become infected by other genotypes (see [Box 3](#)). Parasite fitness then depends on the density of infected hosts, and this creates a new route by which an effect of parasites on host fecundity may feed back on parasite evolution [\[38–40\]](#). For instance, the level of sterilization in the population will affect the selective pressures on a mutant parasite with a different virulence strategy.

Variation in host quality

While classical theory considers that all hosts have the same value for the parasite, natural host–parasite interactions often exhibit variation in host quality (e.g., because of natural resistance, treatments, or other forms of class structure such as sex-based differences in immunity). The parasite is then faced with the problem of exploiting a heterogeneous resource. As a result, the optimal strategy for the parasite is determined by the frequencies of each type of host [\[41–43\]](#). In general, these frequencies may be expected to be affected by host fecundity.

These four ecological factors are ubiquitous in natural host–parasite populations and, importantly, are often combined. Thus, under realistic ecological scenarios, we should not expect differences in host fecundity to be irrelevant to the evolution of horizontally transmitted parasites. Epidemiological feedbacks alter the classical predictions of [Box 3](#) and create novel selective pressures that may affect how a parasite should exploit the resources the host allocates to survival and fecundity. So far, we only have a partial theoretical picture (mostly limited to the effect of spatial structure [\[10\]](#)). Combined with genetic constraints (e.g., trade-offs between transmission, sterility, and mortality virulence), epidemiological feedbacks potentially introduce a whole range of possible evolutionary outcomes. In addition, the evolution of parasite

strategies of host exploitation may feed back on population dynamics, potentially affecting host and parasite extinction rates.

From Theoretical to Empirical Studies

We think the focus on classical epidemiological models has contributed to neglecting the potential of sterility virulence as an adaptive trait in the theoretical literature. Taking into account epidemiological feedbacks suggests that the joint evolutionary dynamics of mortality and sterility virulence may be more complex than previously imagined. Although a number of models have investigated the joint effects of mortality and sterility on the population and coevolutionary dynamics of host–parasite interactions (e.g., [25,44–47]), very few have addressed the evolution of parasite allocation between mortality and sterility virulence. For instance, Thrall, et al. [48] have investigated the evolution of diseases with both sexual and non-sexual transmission routes under the assumption that allocation to sexual transmission leads to reduced fecundity and lower mortality, while van Baalen [49] has studied the allocation between mortality and sterility virulence in a semi-spatial model in which the degree of local (parent-to-offspring) and global transmission may vary. Both results show that the allocation between killing or sterilizing a host is very sensitive to epidemiological feedbacks. Given the abundance, diversity, and sometimes long-standing evolutionary history of systems in which sterility virulence occurs [50], the lack of studies on the topic is striking.

We therefore urge that a general theoretical framework for virulence evolution should take into account the multiple effects of parasite exploitation on host traits. Such a generalized framework would have important implications for empirical studies, as there are some potential pitfalls with the current idea that parasite-induced changes in host fecundity do not affect parasite fitness unless there is an obvious trade-off between transmission and fecundity (e.g., for the sexually transmitted and obligately sterilizing anther-smut fungi [51]). First, if one measures only mortality effects of a given disease, other (minor or less obvious) virulence components may go completely unnoticed. Cowpox virus had long been thought to have no apparent “disease effect” in its wild rodent reservoir until the existence of a negative impact on their fecundity was revealed [52]. In contrast, virulence effects in the well-studied *Drosophila*–nematode system are systematically measured in terms of both fertility and longevity [53]. Although secondary effects may not always be important or adaptive (e.g., *Ophreocystis electro-skirra* effects on Monarch butterfly mating success do not necessarily reduce reproductive output [54]), such practice offers a more thorough understanding of the disease’s natural history. Second, tests of theory-derived hypotheses may be biased if the theoretical framework used does not effectively capture the biology of the system. Third, when thorough measurements of lifetime reproductive success are technically impossible, experimental studies often rely on indirect fitness proxies (e.g., [55]). However, the validity of these proxies is not always established [56]. We argue that mathematical models exploring the adaptive potential of complex host exploitation strategies could help identify which traits need to be measured to calculate a meaningful fitness proxy, and could open avenues of new research into the evolution of virulence in both novel and established host–pathogen systems.

Conclusion

Although theoretical studies are largely biased toward mortality virulence, extensions of classical theoretical evolutionary epidemiology reveal that host fecundity may very likely matter for parasite evolution, and that parasite strategies that manipulate both the fecundity and survival of the host may be shaped by natural selection under realistic ecological scenarios. Given the widespread occurrence of parasites that, in contrast to earlier predictions, do not fully sterilize

their host, we urge theoreticians and empiricists to work together toward a better understanding of the adaptive and multifaceted nature of parasite virulence.

Acknowledgments

We thank S. Alizon, M. van Baalen, S. Gandon, T. Giraud, M. Hood, and T. Lefèvre for discussions and comments. Two anonymous reviewers provided helpful comments on a previous version of this manuscript. We also thank C. L. Murall for input on virus examples.

References

1. Read AF. The evolution of virulence. *Trends in Microbiology*. 1994; 2(3):73–76. PMID: [8156274](#)
2. Poulin R, Combes C. The concept of virulence: interpretations and implications. *Parasitology Today*. 1999; 15(12):474–475. PMID: [10557145](#)
3. Bedhomme S, Agnew P, Vital Y, Sidobre C, Michalakakis Y. Prevalence-dependent costs of parasite virulence. *PLoS Biology*. 2005; 3(8):e262. PMID: [16008503](#)
4. May RM, Anderson RM. Epidemiology and genetics in the coevolution of parasites and hosts. *Proceedings of the Royal Society of London B: Biological Sciences*. 1983; 219(1216):281–313. PMID: [6139816](#)
5. Bull JJ. Perspective: virulence. *Evolution*. 1994; 48:1423–1437.
6. Alizon S, Hurford A, Mideo N, Van Baalen M. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *Journal of Evolutionary Biology*. 2009; 22(2):245–259. doi: [10.1111/j.1420-9101.2008.01658.x](#) PMID: [19196383](#)
7. Day T. Parasite transmission modes and the evolution of virulence. *Evolution*. 2001; 55(12):2389–2400. PMID: [11831655](#)
8. Anderson RM, May RM. Population biology of infectious diseases: Part I. *Nature*. 1979; 280(5721):361–367. PMID: [460412](#)
9. Anderson RM, May RM. Coevolution of hosts and parasites. *Parasitology*. 1982; 85:411–426. PMID: [6755367](#)
10. O’Keefe KJ, Antonovics J. Playing by different rules: the evolution of virulence in sterilizing pathogens. *American Naturalist*. 2002; 159(6):597–605. doi: [10.1086/339990](#) PMID: [18707384](#)
11. Baudoin M. Host castration as a parasitic strategy. *Evolution*. 1975; 29:335–352.
12. Obrebski S. Parasite reproductive strategy and evolution of castration of hosts by parasites. *Science*. 1975; 188(4195):1314–1316. PMID: [1145198](#)
13. Lafferty KD, Kuris AM. Parasitic castration: the evolution and ecology of body snatchers. *Trends in Parasitology*. 2009; 25(12):564–572. doi: [10.1016/j.pt.2009.09.003](#) PMID: [19800291](#)
14. Apari P, de Sousa JD, Müller V. Why sexually transmitted infections tend to cause infertility: an evolutionary hypothesis. *PLoS Pathogens*. 2014; 10(8):e1004111. doi: [10.1371/journal.ppat.1004111](#) PMID: [25101790](#)
15. Jaenike J. Suboptimal virulence of an insect-parasitic nematode. *Evolution*. 1996; 50:2241–2247.
16. Sloan DB, Giraud T, Hood ME. Maximized virulence in a sterilizing pathogen: the anther-smut fungus and its co-evolved hosts. *Journal of Evolutionary Biology*. 2008; 21(6):1544–1554. doi: [10.1111/j.1420-9101.2008.01604.x](#) PMID: [18717748](#)
17. Clerc M, Ebert D, Hall MD. Expression of parasite genetic variation changes over the course of infection: implications of within-host dynamics for the evolution of virulence. *Proceedings of the Royal Society of London B: Biological Sciences*. 2015; 282(1804): 20142820.
18. Hurd H. Host fecundity reduction: a strategy for damage limitation? *Trends in Parasitology*. 2001; 17(8):363–368. PMID: [11685895](#)
19. Gandon S, Agnew P, Michalakakis Y. Coevolution between parasite virulence and host life-history traits. *American Naturalist*. 2002; 160(3):374–388. doi: [10.1086/341525](#) PMID: [18707446](#)
20. Bonds MH. Host life-history strategy explains pathogen-induced sterility. *American Naturalist*. 2006; 168(3):281–293. PMID: [16947104](#)
21. Hall SR, Becker C, Caceres CE. Parasitic castration: a perspective from a model of dynamic energy budgets. *Integrative and comparative biology*. 2007; 47(2):295–309. doi: [10.1093/icb/icm057](#) PMID: [21672839](#)
22. Leventhal GE, Dünner RP, Barribeau SM. Delayed Virulence and Limited Costs Promote Fecundity Compensation upon Infection. *American Naturalist*. 2014; 183(4):480–493. doi: [10.1086/675242](#) PMID: [24642493](#)

23. Day T, Burns JG. A consideration of patterns of virulence arising from host-parasite coevolution. *Evolution*. 2003; 57(3):671–676. PMID: [12703956](#)
24. Graham AL, Allen JE, Read AF. Evolutionary causes and consequences of immunopathology. *Annual Review of Ecology, Evolution, and Systematics*. 2005; 36(1):373–397.
25. Best A, White A, Boots M. Resistance is futile but tolerance can explain why parasites do not always castrate their hosts. *Evolution*. 2010; 64(2):348–357. doi: [10.1111/j.1558-5646.2009.00819.x](#) PMID: [19686267](#)
26. Lion S, Boots M. Are parasites “prudent” in space? *Ecology Letters*. 2010; 13(10):1245–1255. doi: [10.1111/j.1461-0248.2010.01516.x](#) PMID: [20727004](#)
27. Lion S, Gandon S. Evolution of spatially structured host–parasite interactions. *Journal of Evolutionary Biology*. 2015; 28(1):10–28. doi: [10.1111/jeb.12551](#) PMID: [25439133](#)
28. O’Keefe KJ. The evolution of virulence in pathogens with frequency-dependent transmission. *Journal of Theoretical Biology*. 2005; 233(1):55–64. PMID: [15615619](#)
29. Lenski RE, May RM. The Evolution of Virulence in Parasites and Pathogens: Reconciliation Between Two Competing Hypotheses. *Journal of Theoretical Biology*. 1994; 169(3):253–265. PMID: [7967617](#)
30. Day T, Proulx SR. A general theory for the evolutionary dynamics of virulence. *American Naturalist*. 2004; 163(4):E40–E63. PMID: [15122509](#)
31. Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latta J, et al. Should we expect population thresholds for wildlife disease? *Trends in Ecology & Evolution*. 2005; 20(9):511–519. PMID: [16701428](#)
32. Day T, Gandon S. Applying population-genetic models in theoretical evolutionary epidemiology. *Ecology Letters*. 2007; 10(10):876–888. PMID: [17845288](#)
33. Bull JJ, Ebert D. Invasion thresholds and the evolution of non-equilibrium virulence. *Evolutionary Applications*. 2008; 1:172–182. doi: [10.1111/j.1752-4571.2007.00003.x](#) PMID: [25567500](#)
34. Hartfield M, Alizon S. Epidemiological feedbacks affect evolutionary emergence of pathogens. *American Naturalist*. 2014; 183(4):E105–117. doi: [10.1086/674795](#) PMID: [24642501](#)
35. Petney TN, Andrews RH. Multiparasite communities in animals and humans: frequency, structure and pathogenic significance. *International journal for parasitology*. 1998; 28(3):377–393. PMID: [9559357](#)
36. Pedersen AB, Fenton A. Emphasizing the ecology in parasite community ecology. *Trends in Ecology & Evolution*. 2007; 22(3):133–139. PMID: [17137676](#)
37. Balmer O, Tanner M. Prevalence and implications of multiple-strain infections. *The Lancet Infectious Diseases*. 2011; 11(11):868–878. doi: [10.1016/S1473-3099\(11\)70241-9](#) PMID: [22035615](#)
38. van Baalen M, Sabelis MW. The dynamics of multiple infection and the evolution of virulence. *American Naturalist*. 1995; 146(520):881–910.
39. Gandon S, Jansen VAA, van Baalen M. Host life history and the evolution of parasite virulence. *Evolution*. 2001; 55(5):1056–1062. PMID: [11430642](#)
40. Lion S. Multiple infections, kin selection and the evolutionary epidemiology of parasite traits. *Journal of Evolutionary Biology*. 2013; 26(10):2107–2122. doi: [10.1111/jeb.12207](#) PMID: [24028471](#)
41. Regoes RR, Nowak MA, Bonhoeffer S. Evolution of virulence in a heterogeneous host population. *Evolution*. 2000; 54(1):64–71. PMID: [10937184](#)
42. Gandon S, Mackinnon M, Nee S, Read A. Imperfect vaccination: some epidemiological and evolutionary consequences. *Proceedings of the Royal Society of London B: Biological Sciences*. 2003; 270(1520):1129–1136.
43. Cousineau SV, Alizon S. Parasite evolution in response to sex-based host heterogeneity in resistance and tolerance. *Journal of Evolutionary Biology*. 2014; 27(12):2753–2766. doi: [10.1111/jeb.12541](#) PMID: [25376168](#)
44. Best A, White A, Boots M. Maintenance of host variation in tolerance to pathogens and parasites. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(52):20786–20791. doi: [10.1073/pnas.0809558105](#) PMID: [19088200](#)
45. Best A, Hoyle A. The evolution of costly acquired immune memory. *Ecology and Evolution*. 2013; 3(7):2223–2232. doi: [10.1002/ece3.611](#) PMID: [23919164](#)
46. Ashby B, Gupta S. Parasitic castration promotes coevolutionary cycling but also imposes a cost on sex. *Evolution*. 2014; 68(8):2234–2244. doi: [10.1111/evo.12425](#) PMID: [24749747](#)
47. Bernhauerová V, Berec L. Role of trade-off between sexual and vertical routes for evolution of pathogen transmission. *Theoretical Ecology*. 2015; 8(1):23–36.
48. Thrall PH, Antonovics J, Wilson WG. Allocation to sexual versus nonsexual disease transmission. *American Naturalist*. 1998; 151(1):29–45. doi: [10.1086/286100](#) PMID: [18811422](#)

49. van Baalen M. Parent-to-offspring infection and the struggle for transmission In: Poulin R, Morand S, Skorping A, editors. Evolutionary biology of host-parasite relationships: theory meets reality. Elsevier, Amsterdam.; 2000. pp. 97–116.
50. Antonovics J, Boots M, Abbate J, Baker C, McFrederick Q, Panjeti V. Biology and evolution of sexual transmission. *Annals of the New York Academy of Sciences*. 2011; 1230:12–24. doi: [10.1111/j.1749-6632.2011.06127.x](https://doi.org/10.1111/j.1749-6632.2011.06127.x) PMID: [21824163](https://pubmed.ncbi.nlm.nih.gov/21824163/)
51. Schäfer AM, Kemler M, Bauer R, Begerow D. The illustrated life cycle of *Microbotryum* on the host plant *Silene latifolia*. *Botany*. 2010; 88(10):875–885.
52. Feore SM, Bennett M, Chantrey J, Jones T, Baxby D, Begon M. The effect of cowpox virus infection on fecundity in bank voles and wood mice. *Proceedings of the Royal Society of London B: Biological Sciences*. 1997; 264(1387):1457–1461.
53. Jaenike J, Perlman SJ. Ecology and evolution of host-parasite associations: mycophagous *Drosophila* and their parasitic nematodes. *American Naturalist*. 2002; 160(S4):S23–S39.
54. de Roode JC, Yates AJ, Altizer S. Virulence-transmission trade-offs and population divergence in virulence in a naturally occurring butterfly parasite. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(21):7489–7494. doi: [10.1073/pnas.0710909105](https://doi.org/10.1073/pnas.0710909105) PMID: [18492806](https://pubmed.ncbi.nlm.nih.gov/18492806/)
55. Refardt D, Ebert D. Inference of parasite local adaptation using two different fitness components. *Journal of Evolutionary Biology*. 2007; 20(3):921–929. PMID: [17465903](https://pubmed.ncbi.nlm.nih.gov/17465903/)
56. Alizon S, Michalakakis Y. Adaptive virulence evolution: the good old fitness-based approach. *Trends in Ecology & Evolution*. 2015; 5(30):248–254.
57. Pellati D, Mylonakis I, Bertoloni G, Fiore C, Andrisani A, Ambrosini G, et al. Genital tract infections and infertility. *European Journal of Obstetrics and Gynecology*. 2008; 140(1):3–11.
58. Sefton AM. The Great Pox that was... syphilis. *Journal of Applied Microbiology*. 2001; 91(4):592–596. PMID: [11576292](https://pubmed.ncbi.nlm.nih.gov/11576292/)
59. Rasnake MS, Conger NG, McAllister K, Holmes KK, Tramont EC. History of U.S. military contributions to the study of sexually transmitted diseases. *Military Medicine*. 2005; 170(4 Suppl):61–65. PMID: [15916284](https://pubmed.ncbi.nlm.nih.gov/15916284/)
60. Lewis DA. Antibiotic-resistant gonococci-past, present and future. *South African Medical Journal*. 2007; 97(11):1146–1150. PMID: [18250926](https://pubmed.ncbi.nlm.nih.gov/18250926/)
61. Vale PF, Little TJ. Fecundity compensation and tolerance to a sterilizing pathogen in *Daphnia*. *Journal of Evolutionary Biology*. 2012; 25(9):1888–1896. doi: [10.1111/j.1420-9101.2012.02579.x](https://doi.org/10.1111/j.1420-9101.2012.02579.x) PMID: [22856460](https://pubmed.ncbi.nlm.nih.gov/22856460/)
62. Ebert D, Carius HJ, Little T, Decaestecker E. The evolution of virulence when parasites cause host castration and gigantism. *American Naturalist*. 2004; 164(S5):S19–S32.
63. Jensen KH, Little T, Skorping A, Ebert D. Empirical Support for Optimal Virulence in a Castrating Parasite. *PLoS Biology*. 2006; 4(7):e197. doi: [10.1371/annotation/6f580f9f-d724-433c-9e12-f402fac28829](https://doi.org/10.1371/annotation/6f580f9f-d724-433c-9e12-f402fac28829) PMID: [16719563](https://pubmed.ncbi.nlm.nih.gov/16719563/)
64. Little TJ, Chadwick W, Watt K. Parasite variation and the evolution of virulence in a *Daphnia*-microparasite system. *Parasitology*. 2008; 135(3):303–308. PMID: [18005474](https://pubmed.ncbi.nlm.nih.gov/18005474/)
65. Izhar R, Ben-Ami F. Host age modulates parasite infectivity, virulence and reproduction. *Journal of Animal Ecology*. 2015; 84(4):1018–1028. doi: [10.1111/1365-2656.12352](https://doi.org/10.1111/1365-2656.12352) PMID: [25661269](https://pubmed.ncbi.nlm.nih.gov/25661269/)
66. Hudson PJ, Newborn D, Dobson AP. Regulation and Stability of a Free-Living Host-Parasite System: *Trichostrongylus tenuis* in Red Grouse. I. Monitoring and Parasite Reduction Experiments. *Journal of Animal Ecology*. 1992; 61(2): 477–486.
67. Rubinstein G, Czosnek H. Long-term association of tomato yellow leaf curl virus with its whitefly vector *Bemisia tabaci*: effect on the insect transmission capacity, longevity and fecundity. *Journal of General Virology*. 1997; 78:2683–2689. PMID: [9349491](https://pubmed.ncbi.nlm.nih.gov/9349491/)
68. Roy BA, Kirchner JW. Evolutionary dynamics of pathogen resistance and tolerance. *Evolution*. 2000; 54(1):51–63. PMID: [10937183](https://pubmed.ncbi.nlm.nih.gov/10937183/)
69. Lefèvre T, Adamo SA, Biron DG, Missé D, Hughes D, Thomas F. Invasion of the body snatchers: the diversity and evolution of manipulative strategies in host-parasite interactions. *Advances in Parasitology*. 2009; 68:45–83. doi: [10.1016/S0065-308X\(08\)00603-9](https://doi.org/10.1016/S0065-308X(08)00603-9) PMID: [19289190](https://pubmed.ncbi.nlm.nih.gov/19289190/)
70. Karasneh G, Shukla D. Herpes simplex virus infects most cell types in vitro: clues to its success. *Virology Journal*. 2011; 8(1):481.
71. Monavari SH, Vaziri MS, Khalili M, Shamsi-Shahrabadi M, Keyvani H, Mollaei H, et al. Asymptomatic seminal infection of herpes simplex virus: impact on male infertility. *Journal of Biomedical Research*. 2013; 27(1):56–61. doi: [10.7555/JBR.27.20110139](https://doi.org/10.7555/JBR.27.20110139) PMID: [23554795](https://pubmed.ncbi.nlm.nih.gov/23554795/)

72. Fatahzadeh M, Schwartz RA. Human herpes simplex virus infections: epidemiology, pathogenesis, symptomatology, diagnosis, and management. *Journal of the American Academy of Dermatology*. 2007; 57:737–763. PMID: [17939933](#)
73. Hjalmarsson A, Blomqvist P, Sköldenberg B. Herpes Simplex Encephalitis in Sweden, 1990–2001: Incidence, Morbidity, and Mortality. *Clinical Infectious Diseases*. 2007; 45(7):875–880. PMID: [17806053](#)
74. Le Tortorec A, Dejucq-Rainsford N. HIV infection of the male genital tract—consequences for sexual transmission and reproduction. *International Journal of Andrology*. 2010; 33(1):e98–108. doi: [10.1111/j.1365-2605.2009.00973.x](#) PMID: [19531082](#)
75. Mackinnon MJ, Read AF. Virulence in malaria: an evolutionary viewpoint. *Philosophical Transactions of the Royal Society of London B Biological Sciences*. 2004; 359(1446):965–986. PMID: [15306410](#)
76. Pariaud B, Ravigne V, Halkett F, Goyeau H, Carlier J, Lannou C. Aggressiveness and its role in the adaptation of plant pathogens. *Plant Pathology*. 2009; 58:409–424.
77. Kada S, Lion S. Superinfection and the coevolution of parasite virulence and host recovery. *Journal of Evolutionary Biology*. 2015. Epub ahead of print. doi: [10.1111/jeb.12753](#) PMID: [26353032](#)