



HAL
open science

Spectral distortions of the CMB as a probe of inflation, recombination, structure formation and particle physics

J. Chluba, A. Kogut, S.P. Patil, M.H. Abitbol, N. Aghanim, Y. Ali-Haïmoud, M.A. Amin, J. Aumont, N. Bartolo, K. Basu, et al.

► To cite this version:

J. Chluba, A. Kogut, S.P. Patil, M.H. Abitbol, N. Aghanim, et al.. Spectral distortions of the CMB as a probe of inflation, recombination, structure formation and particle physics: Astro2020 Science White Paper. Bull.Am.Astron.Soc., 2019, 51 (3), pp.184. hal-02123336

HAL Id: hal-02123336

<https://hal.science/hal-02123336>

Submitted on 9 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Astro2020 Science White Paper

Spectral Distortions of the CMB as a Probe of Inflation, Recombination, Structure Formation and Particle Physics

Primary thematic area: Cosmology and Fundamental Physics

Secondary thematic area: Galaxy Evolution

Corresponding author email: Jens.Chluba@Manchester.ac.uk

J. Chluba¹, A. Kogut², S. P. Patil³, M. H. Abitbol⁴, N. Aghanim⁵, Y. Ali-Haïmoud⁶, M. A. Amin⁷, J. Aumont⁸, N. Bartolo^{9,10,11}, K. Basu¹², E. S. Battistelli¹³, R. Battye¹, D. Baumann¹⁴, I. Ben-Dayan¹⁵, B. Bolliet¹, J. R. Bond¹⁶, F. R. Bouchet¹⁷, C. P. Burgess^{18,19}, C. Burigana^{20,21,22}, C. T. Byrnes²³, G. Cabass²⁴, D. T. Chuss²⁵, S. Clesse^{26,27}, P. S. Cole²³, L. Dai²⁸, P. de Bernardis^{13,29}, J. Delabrouille^{30,31}, V. Desjacques³², G. de Zotti¹¹, J. A. D. Diacoumis³³, E. Dimastrogiovanni^{34,35}, E. Di Valentino¹, J. Dunkley³⁶, R. Durrer³⁷, C. Dvorkin³⁸, J. Ellis³⁹, H. K. Eriksen⁴⁰, M. Fasiello⁴¹, D. Fixsen⁴², F. Finelli⁴³, R. Flauger⁴⁴, S. Galli⁴⁵, J. Garcia-Bellido⁴⁶, M. Gervasi⁴⁷, V. Gluscevic^{36,48}, D. Grin⁴⁹, L. Hart¹, C. Hernández-Monteagudo⁵⁰, J. C. Hill^{28,51}, D. Jeong^{52,53}, B. R. Johnson⁵⁴, G. Lagache⁵⁵, E. Lee¹, A. Lewis²³, M. Liguori^{9,10,11}, M. Kamionkowski⁵⁷, R. Khatri⁵⁸, K. Kohri⁵⁹, E. Komatsu²⁴, K. E. Kunze⁵⁹, A. Mangilli⁶⁰, S. Masi^{13,29}, J. Mather², S. Matarrese^{9,10,11,61}, M. A. Miville-Deschênes⁶², T. Montaruli⁶³, M. Münchmeyer¹⁹, S. Mukherjee^{45,64}, T. Nakama⁶⁵, F. Nati⁴⁷, A. Ota⁶⁶, L. A. Page³⁶, E. Pajer⁶⁷, V. Poulin^{56,68}, A. Ravenni¹, C. Reichardt⁶⁹, M. Remazeilles¹, A. Rotti¹, J. A. Rubiño-Martín^{70,71}, A. Sarkar¹, S. Sarkar⁷², G. Savini⁷³, D. Scott⁷⁴, P. D. Serpico⁷⁵, J. Silk^{56,76}, T. Souradeep⁷⁷, D. N. Spergel^{51,78}, A. A. Starobinsky⁷⁹, R. Subrahmanyam⁸⁰, R. A. Sunyaev²⁴, E. Switzer², A. Tartari⁸¹, H. Tashiro⁸², R. Basu Thakur⁸³, T. Trombetti²⁰, B. Wallisch^{28,44}, B. D. Wandelt⁴⁵, I. K. Wehus⁴⁰, E.J. Wollack², M. Zaldarriaga²⁸, M. Zannoni⁴⁷

Additional endorsers: www.Chluba.de/SDWP-Decadal-2020/Endorsers.pdf

¹ Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, U.K.

² NASA/GSFC, Mail Code: 665, Greenbelt, MD 20771, USA

³ Niels Bohr International Academy and Discovery Center, Blegdamsvej 17, 2100 Copenhagen, Denmark

⁴ University of Oxford, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH, UK

⁵ Institut d'Astrophysique Spatiale (IAS), CNRS (UMR8617), Université Paris-Sud, Batiment 121, 91405 Orsay, France

⁶ Center for Cosmology and Particle Physics, Department of Physics, New York University, New York, NY 10003, USA

⁷ Department of Physics and Astronomy, Rice University, 6100 Main Street, Houston, Texas 77005, USA

⁸ IRAP, Université de Toulouse, CNRS, CNES, UPS, France

⁹ Dipartimento di Fisica e Astronomia 'Galileo Galilei', Università degli Studi di Padova, via Marzolo 8, I-35131, Padova, Italy

¹⁰ INFN, Sezione di Padova, via Marzolo 8, I-35131, Padova, Italy

¹¹ INAF-Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

¹² Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

¹³ Physics department, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185, Rome, Italy

¹⁴ Institute for Theoretical Physics, University of Amsterdam, Science Park 904, Amsterdam, 1098 XH, The Netherlands

¹⁵ Physics Department, Ariel University, Ariel 40700, Israel

¹⁶ Canadian Institute for Theoretical Astrophysics, 60 St. George Street, University of Toronto, Toronto, ON, M5S 3H8, Canada

¹⁷ Institut d'Astrophysique de Paris, UMR7095, CNRS & Sorbonne Université, 98 bis Boulevard Arago, F-75014, Paris, France

¹⁸ McMaster University, 1280 Main St W, Hamilton, ON L8S 4L8, Canada

¹⁹ Perimeter Institute, 31 Caroline Street North, Waterloo, Ontario, N2L 2Y5, Canada

²⁰ INAF, Istituto di Radioastronomia, Via Piero Gobetti 101, I-40129 Bologna, Italy

²¹ Dipartimento di Fisica e Scienze della Terra, Università di Ferrara, Via Giuseppe Saragat 1, I-44122 Ferrara, Italy

²² Istituto Nazionale di Fisica Nucleare, Sezione di Bologna, Via Irnerio 46, I-40126 Bologna, Italy

²³ Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, United Kingdom

²⁴ Max-Planck-Institut für Astrophysik, 85741 Garching, Germany

²⁵ Department of Physics, Villanova University, 800 E. Lancaster Ave., Villanova, PA 19085

²⁶ IRMP, CURL group, Louvain University, 2 Chemin du Cyclotron, 1348 Louvain-la-Neuve, Belgium

²⁷ Namur Institute of Complex Systems, Department of Mathematics, University of Namur, Rempart de la Vierge 8, 5000 Namur, Belgium.

²⁸ Institute for Advanced Study, Princeton, NJ 08540, USA

²⁹ INFN sezione di Roma, P.le A. Moro 2, 00815 Roma, Italy

³⁰ Laboratoire Astroparticule et Cosmologie (APC), CNRS/IN2P3, 10, rue Alice Domon et Léonie Duquet, 75205 Paris Cedex 13, France

³¹ Département d'Astrophysique, CEA Saclay DSM/Irfu, 91191 Gif-sur-Yvette, France

³² Physics Department and Asher Space Science Institute, Technion, Haifa 3200003, Israel

³³ School of Physics, The University of New South Wales, Sydney NSW 2052, Australia

³⁴ Department of Physics/CERCA/Institute for the Science of Origins, Case Western Reserve University, Cleveland, OH 44106-7079, USA

³⁵ School of Physics, The University of New South Wales, Sydney NSW 2052, Australia

³⁶ Joseph Henry Laboratories of Physics, Jadwin Hall, Princeton University, Princeton, NJ 08544, USA

- 37 Department de Physique Theorique, Université de Genève, Quai E. Ansermet 24, 1211 Genève, Switzerland
- 38 Department of Physics, Harvard University, Cambridge, MA, 02138, USA
- 39 Department of Physics, Kings College London, London WC2R 2LS, UK
- 40 Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway
- 41 Institute of Cosmology and Gravitation, University of Portsmouth, Dennis Sciama Building, Burnaby Road, PO1 3FX, U.K.
- 42 Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA
- 43 INAF - Osservatorio di Astrofisica e Scienza dello Spazio, Via Gobetti 101, I-40129 Bologna, Italy
- 44 Department of Physics, University of California San Diego, 9500 Gilman Drive #0319, La Jolla CA 92093, USA
- 45 Sorbonne Université, CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France
- 46 Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain
- 47 Physics Department, University of Milano Bicocca, 20126, Piazza della Scienza 3, Milano
- 48 Department of Physics, University of Florida, Gainesville, FL 32611
- 49 Haverford College, 370 Lancaster Ave, Haverford PA, 19041, USA
- 50 Centro de Estudios de Física del Cosmos de Aragón (CEFCA), Plaza San Juan, 1, planta 2, E-44001, Teruel
- 51 Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, New York, NY 10010, USA
- 52 Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA
- 53 Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA
- 54 Department of Physics, Columbia University, 538 West 120th Street 1122 Pupin Hall, Box 28, MC 5228 New York, NY 10027, USA
- 55 Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France
- 56 Department of Physics and Astronomy, The Johns Hopkins University, 3701 San Martin Drive, Baltimore MD 21218, USA
- 57 Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005 India
- 58 KEK and Sokendai, Tsukuba 305-0801, Japan
- 59 Departamento de Física Fundamental, Universidad de Salamanca, Plaza de la Merced s/n, 37008 Salamanca, Spain
- 60 Institut de Recherche en Astrophysique et Planetologie (IRAP), CNRS, 9 avenue du Colonel Roche 31028 Toulouse, France
- 61 Gran Sasso Science Institute (GSSI) Viale Francesco Crispi 7, I-67100 L'Aquila, Italy
- 62 AIM, CEA, CNRS, Université Paris-Saclay, Université Paris Diderot, Sorbonne Paris Cité, F-91191 Gif-sur-Yvette, France
- 63 Département de physique nucléaire et corpusculaire, Université de Genève, 24 Quai Ernest-Ansermet, 1205 Genève 4, Suisse
- 64 Sorbonne Universités, Institut Lagrange de Paris, 98 bis Boulevard Arago, 75014 Paris, France
- 65 Jockey Club Institute for Advanced Study, The Hong Kong University of Science and Technology, Hong Kong, P.R. China
- 66 Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, CB3 0WA, UK
- 67 DAMTP, Centre for Mathematical Sciences, University of Cambridge, CB3 0WA, UK
- 68 LUPM, CNRS & Université de Montpellier (UMR-5299), Place Eugène Bataillon, F-34095 Montpellier Cedex 05, France
- 69 School of Physics, University of Melbourne, Parkville VIC 3010, Australia
- 70 Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain
- 71 Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain
- 72 University of Oxford, Clarendon Laboratory, Parks Road, Oxford OX1 3PU, UK
- 73 Physics & Astronomy Dept., UCL, London WC1E 6BT, UK
- 74 Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia, 55 Canada
- 75 Université Grenoble Alpes, USMB, CNRS, LAPTh, F-74940 Annecy, France
- 76 Department of Physics and Beecroft Inst. for Particle Astrophysics and Cosmology, University of Oxford, 1 Keble Road, Oxford, OX1 3RH, UK
- 77 IUCAA, Post Bag 4 Ganeshkhind, Pune 411007, India
- 78 Department of Astrophysical Sciences, Peyton Hall, Princeton University, Princeton, NJ 08544, USA
- 79 L. D. Landau Institute for Theoretical Physics RAS, Moscow 119334, Russia
- 80 Raman Research Institute, C V Raman Avenue, Sadashivanagar, Bangalore 560080, India
- 81 INFN Sezione di Pisa Largo B. Pontecorvo 3, Edif. C I-56127 Pisa, Italy
- 82 Division of Particle and Astrophysical Science, Graduate School of Science, Nagoya University, Chikusa, Nagoya 464-8602, Japan
- 83 Department of Physics, California Institute of Technology, Pasadena, California 91125, USA

EXECUTIVE SUMMARY

Following the pioneering observations with COBE in the early 1990s, studies of the cosmic microwave background (CMB) have focused on temperature and polarization anisotropies. CMB spectral distortions – tiny departures of the CMB energy spectrum from that of a perfect blackbody – provide a second, independent probe of fundamental physics, with a reach deep into the primordial Universe. The theoretical foundation of spectral distortions has seen major advances in recent years, which highlight the immense potential of this emerging field. Spectral distortions probe a fundamental property of the Universe – its *thermal history* – thereby providing additional insight into processes within the cosmological standard model¹ (CSM) as well as new physics beyond. Spectral distortions are an important tool for understanding inflation and the nature of dark matter. They shed new light on the physics of recombination and reionization, both prominent stages in the evolution of our Universe, and furnish critical information on baryonic feedback processes, in addition to probing primordial correlation functions at scales inaccessible to other tracers. In principle the range of signals is vast: *many orders of magnitude of discovery space* could be explored by detailed observations of the CMB energy spectrum. Several CSM signals are predicted and provide clear experimental targets, some of which are already observable with present-day technology. Confirmation of these signals would extend the reach of the CSM by orders of magnitude in physical scale as the Universe evolves from the initial stages to its present form. *The absence of these signals would pose a huge theoretical challenge, immediately pointing to new physics.*

I. COSMOLOGY BEYOND THERMAL EQUILIBRIUM

Cosmology is now a precise scientific discipline, with detailed theoretical models that fit a wealth of very accurate measurements. Of the many cosmological data sets, the CMB temperature and polarization anisotropies provide the most stringent and robust constraints on theoretical models, allowing us to determine the key parameters of our Universe (e.g., the total density, expansion rate and baryon content) with unprecedented precision, while simultaneously addressing fundamental questions about inflation and early-universe physics. By studying the statistics of the CMB anisotropies with different experiments over the past decades we have entered the era of precision cosmology, clearly establishing the highly-successful Λ CDM concordance model [1–3].

But the quest continues. Despite its many successes, Λ CDM *is known to be incomplete*. It traces the growth of structure in the Universe from primordial density perturbations to the modern era, but the origin of those perturbations remains poorly understood. In addition, in spite of relentless efforts, the nature of dark matter (DM) and dark energy remains a mystery. Together, these enigmatic components comprise 95% of the energy density of the Universe. Particle and high-energy physics offer candidate solutions of these problems (e.g., inflation and particle dark matter), but these inevitably require new physics beyond the Standard Model of particle physics.

Precision measurements of the CMB energy spectrum open a new window into the physics of the early Universe, constraining models in ways not possible using other techniques. Departures of the CMB energy spectrum from a pure blackbody – commonly referred to as *spectral distortions* – encode unique information about the thermal history of the Universe, from when it was a few months old until today. Since the measurements with COBE/FIRAS in the early '90s, the sky-averaged CMB spectrum is known to be extremely close to a perfect blackbody at a temperature $T_0 = (2.7255 \pm 0.0006)$ K [4, 5], with possible distortions limited to one part in 10^5 . This impressive measurement was awarded the 2006 Nobel Prize in Physics and already rules out cosmologies with extended periods of large energy release.

Spectral distortions are created by processes that drive matter and radiation out of thermal equilibrium after thermalization becomes inefficient at redshift $z \lesssim 2 \times 10^6$. Exam-

ples are energy-releasing mechanisms that heat the baryonic matter, inject photons or other electromagnetically-interacting particles. The associated signals are usually characterized as μ - and y -type distortions, formed by energy exchange between electrons and photons through Compton scattering [6–10]. Compton scattering is inefficient at $z \lesssim 5 \times 10^4$, yielding a y -type distortion, which probes the thermal history during recombination and reionization (Fig. 1).




FIG. 1: Evolution of spectral distortions across time. Distortions probe the thermal history over long periods deep into the primordial Universe that are inaccessible by other means.

phenomenology [16], one prominent example being the distortion created by the cosmological recombination process [17–19] (see Fig. 1). Additional information can be imprinted by non-equilibrium processes in the pre-recombination hydrogen and helium plasma [20–22] or by non-thermal particles in high-energy particle cascades [e.g., 16, 22–25]. Finally, distortion anisotropies are expected and can be correlated against tracers of both primordial density perturbations and large-scale structure to further probe cosmic evolution [26–31].

Spectral distortions thus provide more than just a simple integral constraint for cosmology. They are a unique and powerful probe of a wide range of interactions between particles and CMB photons, reaching back all the way from the present to a few months after the Big Bang and allowing us to access information that cannot be extracted in any other way. Broad overviews of the CMB spectral distortion science case can be found in [11, 18, 32–38].

II. PHYSICS BEYOND Λ CDM AND THE STANDARD MODEL

A central question in modern cosmology is the origin of the observed primordial density perturbations. Measurements from CMB anisotropies and large-scale structure find a nearly scale-invariant power spectrum $\mathcal{P}(k) \simeq k^{n_s-1}$ with spectral index $n_s = 0.965 \pm 0.004$, sampled over a range of spatial scales $k \simeq 10^{-4}$ to 0.1 Mpc^{-1} [39]. Their phase coherence is evidence for their super-Hubble nature, and their near scale-invariance is evidence of a weakly broken shift symmetry in the underlying theory. However, their precise origin is as of yet unknown. Inflation provides a widely accepted framework [40–43], with the simplest models generically predicting a small departure from scale-invariance (with $n_s < 1$) as the inflaton rolls down its potential [44–47]. However, various alternatives to inflation have been proposed [48–57] and no clear theoretical consensus has yet emerged. Searches for a B -mode pattern in CMB polarization could yield compelling evidence

In contrast, a μ -type (or chemical potential) distortion forms at $z \gtrsim 5 \times 10^4$, when Compton scattering is very efficient. A μ -distortion cannot be generated at recent epochs and thus directly probes events in the pre-recombination era.

The simple classical picture has been refined in recent years. We now understand that the transition between μ - and y -type distortions is gradual and that the signal contains additional time-dependent information [11–13]. This extra information is contained in the residual or r -type distortion and can be used to distinguish energy release mechanisms [14, 15]. Distortions created by photon-injection mechanisms exhibit rich spectral phe-

for the simplest inflationary models. CMB polarization measurements so far only provide upper limits [58, 59] with no firm target from theory for a guaranteed detection. However, detection of a tensor to scalar ratio of $r \simeq 10^{-3}$ is a distinguishing benchmark for large-field models, which in certain realizations further manifest the specific relation $r \simeq (1 - n_s)^2$ [e.g., 40, 60].

Spectral distortions provide a unique new probe of primordial density perturbations. Inflation may or may not be a valid description of the early Universe, but density perturbations are known to exist; regardless of their origin, dissipation of these perturbations through photon diffusion (Silk damping) in the early Universe will distort the CMB spectrum at observable levels [61–65]. The signal can be accurately calculated using simple linear physics and depends on the amplitude of primordial perturbations at scales with $k \simeq 1 - 10^4 \text{ Mpc}^{-1}$, some ten e-folds further than what can be probed by CMB anisotropies (Fig. 2). If the near scale-invariance of the power spectrum observed on large scales persists to these much smaller scales, then the predicted distortion, $\mu \simeq (2.3 \pm 0.14) \times 10^{-8}$ [37, 64, 66], could be observed using current technology (§IV). Detecting this signal extends our grasp on primordial density




FIG. 2: Forecast constraints (95 % c.l.) on the primordial power spectrum for features with a k^4 profile that cuts off sharply at someⁱⁱ k_p [see 67, for more details]. μ -distortions constrain perturbations at scales and levels inaccessible to other probes.

from scale-invariance are well-motivated and often produce excess small-scale power (e.g., features [74–77] or inflection points [78–83] in the potential, particle production [84–88], waterfall transitions [89–93], axion inflation [94–96], etc. [97]), implying the presence of new physical scales that can be probed with spectral distortions (Fig. 2). In this respect, spectral distortions could establish a link to a possible primordial origin of the small-scale structure crisis [98, 99]. They could also place limits on primordial black holes [100–102], testing their role in super-massive black-hole formation [103] and as DM candidate [91, 104], the latter seeing renewed interest after the first LIGO/Virgo merger events [105–107]. Spectral distortions are also created by the dissipation of small scale *tensor* perturbations [108, 109] and depend on the perturbation-type (i.e., adiabatic vs. iso-curvature) [110–113], providing ways to test inflation scenarios in uncharted territory. Spectral distortion anisotropies furthermore probe local-type primordial non-Gaussianity at small scales [29, 30, 114–121], an exciting direction that complements other probes and could shed light on multi-field inflation scenarios [122].

Dark matter is another example of how spectral distortions probe new physics. Non-baryonic matter constitutes $\simeq 25\%$ of the energy density of the Universe, but its nature remains unknown.

perturbations by over three orders of magnitude in scale, covering epochs that cannot be probed directly in any other way. A *non-detection at this level would be a serious challenge for Λ CDM, immediately requiring new physics.*

Measurements of μ -distortions are directly sensitive to the power spectrum amplitude and its scale-dependence around $k \simeq 10^3 \text{ Mpc}^{-1}$ [15, 68–70]. Within the slow-roll paradigm, this provides a handle on higher-order slow-roll parameters (i.e., running of the tilt), benefiting from a vastly extended lever arm [71–73]. Outside of standard slow-roll inflation, large departures

The long-favored WIMP-scenario is under increasing pressure [123–128], and emphasis is gradually shifting focus towards alternatives, prominent examples being axions, sterile neutrinos, sub-GeV DM or primordial black holes [100, 129–134]. To solve this puzzle, a multi-tracer approach that combines different particle physics and cosmological probes is needed. Measurements of the CMB anisotropies themselves have clearly helped to establish the presence of DM on cosmological scales and provided tight constraints on DM annihilation and decay [135–143] and DM-SM-interactions [144–149]. However, for DM annihilation and decay CMB anisotropies quickly lose constraining power before recombination ($z \gtrsim 10^3$), being impeded by cosmic variance. Similarly, measurements of light-element abundances [135, 150–152], which are only sensitive to non-thermal energy release above nuclear dissociation thresholds in the pre-recombination era [15, 153], saturated their limits due to astrophysical uncertainties. *This is where CMB spectral distortions offer a valuable complementary probe.* For decaying particle scenarios, distortions are sensitive to particles with lifetimes $t \simeq 10^6 - 10^{12}$ s [11, 15, 154–158], providing direct measurement of particle lifetimes via r -type distortions [14, 15]. Similarly, annihilating particles can be constrained using distortions; μ -distortions are sensitive to light particles ($m \lesssim 100$ keV) and complement γ -ray searches for heavier particles [14, 159]. The rich spectral information added by various non-thermal processes [16, 20–22, 24, 25] will allow us to glean even more information about the nature of dark matter. Significant theoretical work is required, although it is already clear that in addition to the aforementioned examples distortions can meaningfully probe scenarios involving axions [160–162], gravitino decays [155, 163], strings [164, 165], DM-SM-interactions [141, 166, 167], macros [168] and primordial magnetic fields [169–172].

III. PRECISION TESTS OF Λ CDM AND THE STANDARD MODEL

Spectral distortions also enable new tests of the standard cosmological model. The cosmological recombination and reionization eras as well as the cosmic dark ages mark important transitions in the evolution of our Universe. The largest all-sky distortion signal is indeed caused by the reionization and structure-formation processes [26, 173–177]. Energy output from the first stars and shocks heats the baryons and electrons, which then up-scatter CMB photons to create a y -type distortion. The overall distortion is expected to reach $y \simeq \text{few} \times 10^{-6}$ [26, 176, 178, 179], only one order of magnitude below the current upper limit placed by COBE/FIRAS. Such a distortion *must* exist and provides a measurement of the total thermal energy in (ionized) baryons in the Universe. A large part of the signal is due to halos with masses $M \simeq 10^{13} M_\odot$ containing gas with an electron temperature of $kT_e \simeq 2$ keV. This causes a relativistic temperature correction [180–184] to the y -distortion that can directly tell us about feedback mechanisms [179]. The low-redshift y -distortion from reionization is furthermore anisotropic [e.g., 26–28] and thus opens new opportunities for cross-correlation studies (e.g., with CMB and 21 cm tomography).

The cosmological recombination process causes another small but inevitable distortion of the CMB. Line emission from hydrogen and helium injects photons into the CMB, which after redshifting from $z \simeq 10^3$ is visible today as complex frequency structure in the microwave bands (Fig. 1) [17, 185–192]. The cosmological recombination radiation (CRR) has a simple dependence on cosmological parameters and the dynamics of recombination; since it includes not only hydrogen but also two helium recombinations, it probes eras well beyond the last-scattering surface observed by CMB anisotropies [18, 19, 193]. Cosmological recombination is furthermore crucial for understanding and interpreting the CMB temperature and polarization anisotropies [194–197]. It is thus critical to test our physical assumptions during this era [198–201]. *The CRR provides one of the most direct methods to achieve this.* The CRR also enables a pristine measurement of the primordial helium abundance, long before the first stars have formed. It would thus break cosmological parameter degeneracies, e.g., between the primordial helium abundance and neutrino

number [19]. Finally, interactions of CMB photons with atoms can imprint additional frequency-dependent signals through resonance [202–209] and Rayleigh scattering effects [210, 211], both of which provide independent ways for learning about recombination, the dark ages and reionization.

IV. THE PATH FORWARD WITH CMB SPECTRAL DISTORTIONS

The seminal measurements of the CMB blackbody spectrum by COBE/FIRAS in the early '90s cemented the Hot Big Bang model by ruling out any energy release greater than $\Delta U/U \simeq 10^{-5}$ of the energy in CMB photons [4, 5, 212]. Advances since then, in both detector technology and cryogenics, could improve this sensitivity by four orders of magnitude or more (e.g., with experimental concepts like PIXIE [213, 214] or PRISM [33]), opening an enormous discovery space for both predicted distortion signals and those caused by new physics.

COBE/FIRAS was not background limited; its sensitivity was set instead by phonon noise from the 1.4 K detector. Modern detectors, operating at $\simeq 0.1$ K, would have detector (dark) noise well




FIG. 3: Spectral distortions are observable using current technology. The signal amplitudes and required foreground cleaning are comparable to planned *B*-mode searches.

coverage outside the atmospheric windows ultimately requires a space mission to detect $\mu \simeq 10^{-8}$ or the CRR [219–224]. Both the anticipated signal levels (in the range 1–100 Jy sr⁻¹) and level of foreground cleaning are comparable to those encountered for next-generation CMB polarization measurements [e.g., 225]. Therefore, much of the technology and analysis techniques are directly transferable, although a new synergistic approach (combining multiple data sets) and observing strategy (e.g., small-patch vs. all-sky) have yet to be developed.

To conclude, CMB spectral distortions probe many processes throughout the history of the Universe. Precision spectroscopy, possible with existing technology, would provide key tests for processes expected within the CSM and open an enormous discovery space for new physics. This offers unique scientific opportunities for furthering our understanding of inflation, recombination, reionization and particle physics. Many experimental and theoretical challenges have to be overcome before we can fully exploit this new window to early- and late-universe physics. However, the potential gains are immense and the field is entering a phase of accelerated growth after decades of dormancy. With a coordinated approach, the next decade could thus see new precision measurements of one of the fundamental observables of our Universe^{iv}.

below the intrinsic limit set by photon arrival statistics. The sensitivity of a background-limited instrument could be further improved by increasing the instrument's collecting area.

Figure 3 compares several predicted spectral distortions [e.g., 37] and the largest astrophysical foregroundsⁱⁱⁱ to the sensitivity of possible next-generation spectrometers. Pioneering steps towards $y \simeq 10^{-7}$, yielding a clear detection of the expected average distortion caused by groups and clusters (see Sect. III), are possible from the ground and balloons (e.g., using concepts similar to COSMO, OLIMPO [215, 216] and ARCADE [217, 218]). However, because the distortions peak at frequencies above 200 GHz, broad frequency

-
- [1] G. F. Smoot, C. L. Bennett, A. Kogut, E. L. Wright, J. Aymon, N. W. Boggess, E. S. Cheng, G. de Amici, S. Gulkis, M. G. Hauser, et al., *ApJL* **396**, L1 (1992).
- [2] C. L. Bennett, M. Halpern, G. Hinshaw, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, L. Page, D. N. Spergel, G. S. Tucker, et al., *ApJS* **148**, 1 (2003).
- [3] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, and C. Baccigalupi, et al., *A&A* **571**, A16 (2014).
- [4] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright, *ApJ* **473**, 576 (1996), arXiv:astro-ph/9605054.
- [5] D. J. Fixsen, *ApJ* **707**, 916 (2009), 0911.1955.
- [6] Y. B. Zeldovich and R. A. Sunyaev, *ApSS* **4**, 301 (1969).
- [7] R. A. Sunyaev and Y. B. Zeldovich, *ApSS* **7**, 20 (1970).
- [8] A. F. Illarionov and R. A. Sunyaev, *Soviet Astronomy* **18**, 691 (1975).
- [9] C. Burigana, L. Danese, and G. de Zotti, *A&A* **246**, 49 (1991).
- [10] W. Hu and J. Silk, *Phys. Rev. D* **48**, 485 (1993).
- [11] J. Chluba and R. A. Sunyaev, *MNRAS* **419**, 1294 (2012), 1109.6552.
- [12] R. Khatri and R. A. Sunyaev, *JCAP* **9**, 016 (2012), 1207.6654.
- [13] J. Chluba, *MNRAS* **434**, 352 (2013), 1304.6120.
- [14] J. Chluba, *MNRAS* **436**, 2232 (2013), 1304.6121.
- [15] J. Chluba and D. Jeong, *MNRAS* **438**, 2065 (2014), 1306.5751.
- [16] J. Chluba, *MNRAS* **454**, 4182 (2015), 1506.06582.
- [17] V. K. Dubrovich, *Soviet Astronomy Letters* **1**, 196 (1975).
- [18] R. A. Sunyaev and J. Chluba, *Astronomische Nachrichten* **330**, 657 (2009), 0908.0435.
- [19] J. Chluba and Y. Ali-Haïmoud, *MNRAS* **456**, 3494 (2016), 1510.03877.
- [20] Y. E. Lyubarsky and R. A. Sunyaev, *A&A* **123**, 171 (1983).
- [21] J. Chluba and R. A. Sunyaev, *A&A* **501**, 29 (2009), 0803.3584.
- [22] J. Chluba, *MNRAS* **402**, 1195 (2010), 0910.3663.
- [23] T. A. Enßlin and C. R. Kaiser, *A&A* **360**, 417 (2000), arXiv:astro-ph/0001429.
- [24] T. R. Slatyer, *Phys. Rev. D* **93**, 023521 (2016), 1506.03812.
- [25] S. K. Acharya and R. Khatri, *ArXiv e-prints* (2018), 1808.02897.
- [26] A. Refregier, E. Komatsu, D. N. Spergel, and U.-L. Pen, *Phys. Rev. D* **61**, 123001 (2000).
- [27] P. Zhang, U.-L. Pen, and H. Trac, *MNRAS* **355**, 451 (2004), arXiv:astro-ph/0402115.
- [28] C. Pitrou, F. Bernardeau, and J.-P. Uzan, *JCAP* **7**, 19 (2010), 0912.3655.
- [29] E. Pajer and M. Zaldarriaga, *Phys. Rev. Lett* **109**, 021302 (2012), 1201.5375.
- [30] J. Ganc and E. Komatsu, *Phys. Rev. D* **86**, 023518 (2012), 1204.4241.
- [31] J. C. Hill and E. Pajer, *Phys. Rev. D* **88**, 063526 (2013), 1303.4726.
- [32] R. A. Sunyaev and R. Khatri, *IJMPD* **22**, 1330014 (2013), 1302.6553.
- [33] P. André, C. Baccigalupi, A. Banday, D. Barbosa, B. Barreiro, J. Bartlett, N. Bartolo, E. Battistelli, R. Battye, G. Bendo, et al., *JCAP* **2**, 006 (2014), 1310.1554.
- [34] J. Chluba, *ArXiv e-prints* (2014), 1405.6938.
- [35] H. Tashiro, *Prog. of Theo. and Exp. Physics* **2014**, 06B107 (2014).
- [36] G. De Zotti, M. Negrello, G. Castex, A. Lapi, and M. Bonato, *JCAP* **3**, 047 (2016).
- [37] J. Chluba, *MNRAS* **460**, 227 (2016), 1603.02496.
- [38] J. Chluba, *ArXiv e-prints* (2018), 1806.02915.

- [39] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, and R. B. Barreiro, et al., ArXiv e-prints (2018), 1807.06209.
- [40] A. A. Starobinsky, Phys. Lett. **B91**, 99 (1980).
- [41] A. H. Guth, Phys. Rev. **D23**, 347 (1981).
- [42] A. D. Linde, Phys. Lett. **108B**, 389 (1982).
- [43] A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett **48**, 1220 (1982).
- [44] V. F. Mukhanov and G. V. Chibisov, SJETP Letters **33**, 532 (1981).
- [45] S. W. Hawking, Phys. Lett. **115B**, 295 (1982).
- [46] A. A. Starobinsky, JETP Lett. **30**, 682 (1979).
- [47] A. H. Guth and S. Y. Pi, Phys. Rev. Lett **49**, 1110 (1982).
- [48] M. Gasperini and G. Veneziano, Astropart. Phys. **1**, 317 (1993), hep-th/9211021.
- [49] D. Wands, Phys. Rev. **D60**, 023507 (1999), gr-qc/9809062.
- [50] J. Khoury, B. A. Ovrut, N. Seiberg, P. J. Steinhardt, and N. Turok, Phys. Rev. **D65**, 086007 (2002).
- [51] S. Hollands and R. M. Wald, Gen. Rel. Grav. **34**, 2043 (2002), gr-qc/0205058.
- [52] R. H. Brandenberger, A. Nayeri, S. P. Patil, and C. Vafa, Int. J. Mod. Phys. **A22**, 3621 (2007).
- [53] B. Craps, T. Hertog, and N. Turok, Phys. Rev. **D86**, 043513 (2012), 0712.4180.
- [54] P. Creminelli, A. Nicolis, and E. Trincherini, JCAP **1011**, 021 (2010), 1007.0027.
- [55] D. A. Easson, I. Sawicki, and A. Vikman, JCAP **2011**, 021 (2011), 1109.1047.
- [56] A. Ijjas and P. J. Steinhardt, Phys. Rev. Lett **117**, 121304 (2016), 1606.08880.
- [57] D. A. Dobre, A. V. Frolov, J. T. Gálvez Ghersi, S. Ramazanov, and A. Vikman, JCAP **2018**, 020 (2018), 1712.10272.
- [58] N. Aghanim et al. (Planck) (2018), 1807.06209.
- [59] P. A. R. Ade et al. (BICEP2, Keck Array), Submitted to: Phys. Rev. Lett (2018), 1810.05216.
- [60] F. Bezrukov and M. Shaposhnikov, Phys. Lett. **B659**, 703 (2008), 0710.3755.
- [61] R. A. Sunyaev and Y. B. Zeldovich, ApSS **9**, 368 (1970).
- [62] R. A. Daly, ApJ **371**, 14 (1991).
- [63] W. Hu, D. Scott, and J. Silk, ApJL **430**, L5 (1994), arXiv:astro-ph/9402045.
- [64] J. Chluba, R. Khatri, and R. A. Sunyaev, MNRAS **425**, 1129 (2012), 1202.0057.
- [65] R. Khatri, R. A. Sunyaev, and J. Chluba, A&A **543**, A136 (2012), 1205.2871.
- [66] G. Cabass, A. Melchiorri, and E. Pajer, Phys. Rev. **D93**, 083515 (2016), 1602.05578.
- [67] C. T. Byrnes, P. S. Cole, and S. P. Patil (2018), 1811.11158.
- [68] J. Chluba, A. L. Erickcek, and I. Ben-Dayan, ApJ **758**, 76 (2012), 1203.2681.
- [69] E. Pajer and M. Zaldarriaga, JCAP **2**, 036 (2013), 1206.4479.
- [70] R. Khatri and R. A. Sunyaev, JCAP **6**, 026 (2013), 1303.7212.
- [71] B. A. Powell, ArXiv e-prints (2012), 1209.2024.
- [72] S. Clesse, B. Garbrecht, and Y. Zhu, JCAP **2014**, 046 (2014), 1402.2257.
- [73] G. Cabass, E. Di Valentino, A. Melchiorri, E. Pajer, and J. Silk, Phys. Rev. **D94**, 023523 (2016).
- [74] A. A. Starobinsky, JETP Lett. **55**, 489 (1992), [Pisma Zh. Eksp. Teor. Fiz.55,477(1992)].
- [75] J. A. Adams, B. Cresswell, and R. Easther, Phys. Rev. **D64**, 123514 (2001).
- [76] D. K. Hazra, A. Shafieloo, G. F. Smoot, and A. A. Starobinsky, Phys. Rev. Lett **113**, 071301 (2014).
- [77] D. K. Hazra, D. Paoletti, M. Ballardini, F. Finelli, A. Shafieloo, G. F. Smoot, and A. A. Starobinsky, JCAP **1802**, 017 (2018).
- [78] A. G. Polnarev and I. Musco, Class. Quant. Grav. **24**, 1405 (2007), gr-qc/0605122.
- [79] K. Kohri, D. H. Lyth, and A. Melchiorri, JCAP **0804**, 038 (2008), 0711.5006.
- [80] I. Ben-Dayan and R. Brustein, JCAP **2010**, 007 (2010), 0907.2384.
- [81] S. Choudhury and A. Mazumdar, Phys. Lett. **B733**, 270 (2014), 1307.5119.

- [82] S. Clesse and J. Garca-Bellido, *Phys. Rev.* **D92**, 023524 (2015), 1501.07565.
- [83] C. Germani and T. Prokopec, *Phys. Dark Univ.* **18**, 6 (2017), 1706.04226.
- [84] N. Barnaby and Z. Huang, *Phys. Rev.* **D80**, 126018 (2009), 0909.0751.
- [85] J. L. Cook and L. Sorbo, *Phys. Rev.* **D85**, 023534 (2012), 1109.0022.
- [86] D. Battefeld, T. Battefeld, and D. Fiene, *Phys. Rev.* **D89**, 123523 (2014), 1309.4082.
- [87] E. Dimastrogiovanni, M. Fasiello, and T. Fujita, *JCAP* **2017**, 019 (2017), 1608.04216.
- [88] V. Domcke and K. Mukaida, *JCAP* **1811**, 020 (2018), 1806.08769.
- [89] A. D. Linde, *Phys. Rev.* **D49**, 748 (1994), astro-ph/9307002.
- [90] D. H. Lyth and E. D. Stewart, *Phys. Rev.* **D54**, 7186 (1996), hep-ph/9606412.
- [91] J. García-Bellido, A. Linde, and D. Wands, *Phys. Rev.* **D54**, 6040 (1996).
- [92] A. A. Abolhasani, H. Firouzjahi, and M. H. Namjoo, *Class. Quant. Grav.* **28**, 075009 (2011).
- [93] S. Clesse, B. Garbrecht, and Y. Zhu, *JCAP* **1410**, 046 (2014), 1402.2257.
- [94] N. Barnaby and M. Peloso, *Phys. Rev. Lett* **106**, 181301 (2011), 1011.1500.
- [95] N. Barnaby, E. Pajer, and M. Peloso, *Phys. Rev.* **D85**, 023525 (2012), 1110.3327.
- [96] P. D. Meerburg and E. Pajer, *JCAP* **1302**, 017 (2013), 1203.6076.
- [97] J. Chluba, J. Hamann, and S. P. Patil, *IJMP D* **24**, 1530023 (2015), 1505.01834.
- [98] T. Nakama, J. Chluba, and M. Kamionkowski, *Phys. Rev.* **D95**, 121302 (2017), 1703.10559.
- [99] K. Cho, S. E. Hong, E. D. Stewart, and H. Zoc, *JCAP* **2017**, 002 (2017), 1705.02741.
- [100] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, *Phys. Rev.* **D81**, 104019 (2010).
- [101] P. Pani and A. Loeb, *Phys. Rev.* **D88**, 041301 (2013), 1307.5176.
- [102] T. Nakama, B. Carr, and J. Silk, *Phys. Rev.* **D97**, 043525 (2018), 1710.06945.
- [103] K. Kohri, T. Nakama, and T. Suyama, *Phys. Rev.* **D90**, 083514 (2014), 1405.5999.
- [104] S. Clesse and J. García-Bellido, *Phys. Rev.* **D92**, 023524 (2015), 1501.07565.
- [105] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz, A. Raccanelli, and A. G. Riess, *Phys. Rev. Lett* **116**, 201301 (2016), 1603.00464.
- [106] S. Clesse and J. García-Bellido, *Physics of the Dark Universe* **15**, 142 (2017), 1603.05234.
- [107] K. Kohri and T. Terada, *Classical and Quantum Gravity* **35**, 235017 (2018), 1802.06785.
- [108] A. Ota, T. Takahashi, H. Tashiro, and M. Yamaguchi, *JCAP* **10**, 029 (2014), 1406.0451.
- [109] J. Chluba, L. Dai, D. Grin, M. A. Amin, and M. Kamionkowski, *MNRAS* **446**, 2871 (2015).
- [110] W. Hu and N. Sugiyama, *ApJ* **436**, 456 (1994), arXiv:astro-ph/9403031.
- [111] J. B. Dent, D. A. Easson, and H. Tashiro, *Phys. Rev.* **D86**, 023514 (2012), 1202.6066.
- [112] J. Chluba and D. Grin, *MNRAS* **434**, 1619 (2013), 1304.4596.
- [113] T. Haga, K. Inomata, A. Ota, and A. Ravenni, *JCAP* **2018**, 036 (2018).
- [114] M. Biagetti, H. Perrier, A. Riotto, and V. Desjacques, *Phys. Rev.* **D87**, 063521 (2013).
- [115] A. Ota, T. Sekiguchi, Y. Tada, and S. Yokoyama, *JCAP* **2015**, 013 (2015), 1412.4517.
- [116] R. Emami, E. Dimastrogiovanni, J. Chluba, and M. Kamionkowski, *Phys. Rev.* **D91**, 123531 (2015).
- [117] R. Khatri and R. Sunyaev, *JCAP* **2015**, 026 (2015), 1507.05615.
- [118] J. Chluba, E. Dimastrogiovanni, M. A. Amin, and M. Kamionkowski, *MNRAS* **466**, 2390 (2017).
- [119] A. Ota, *Phys. Rev.* **D94**, 103520 (2016).
- [120] A. Ravenni, M. Liguori, N. Bartolo, and M. Shiraishi, *JCAP* **9**, 042 (2017), 1707.04759.
- [121] G. Cabass, E. Pajer, and D. van der Woude, *JCAP* **1808**, 050 (2018), 1805.08775.
- [122] E. Dimastrogiovanni and R. Emami, *JCAP* **12**, 015 (2016), 1606.04286.
- [123] Z. Ahmed et al. (CDMS-II), *Science* **327**, 1619 (2010), 0912.3592.
- [124] E. Aprile et al. (XENON100), *Phys. Rev. Lett* **109**, 181301 (2012), 1207.5988.
- [125] G. Angloher et al. (CRESST), *Eur. Phys. J.* **C76**, 25 (2016), 1509.01515.
- [126] R. Agnese et al. (SuperCDMS), *Phys. Rev. Lett* **116**, 071301 (2016), 1509.02448.

- [127] A. Tan et al. (PandaX-II), Phys. Rev. Lett **117**, 121303 (2016), 1607.07400.
- [128] D. S. Akerib et al. (LUX), Phys. Rev. Lett **118**, 021303 (2017), 1608.07648.
- [129] G. Jungman, M. Kamionkowski, and K. Griest, Phys. Rep. **267**, 195 (1996), hep-ph/9506380.
- [130] J. L. Feng, A. Rajaraman, and F. Takayama, Phys. Rev. Lett **91**, 011302 (2003).
- [131] J. L. Feng, A. Rajaraman, and F. Takayama, Phys. Rev. **D68**, 063504 (2003).
- [132] A. Kusenko, Phys. Rep. **481**, 1 (2009), 0906.2968.
- [133] J. L. Feng, ARA&A **48**, 495 (2010), 1003.0904.
- [134] D. J. E. Marsh, Phys. Rep. **643**, 1 (2016), 1510.07633.
- [135] J. Ellis, G. B. Gelmini, J. L. Lopez, D. V. Nanopoulos, and S. Sarkar, Nu. Phys. B **373**, 399 (1992).
- [136] J. A. Adams, S. Sarkar, and D. W. Sciama, MNRAS **301**, 210 (1998), astro-ph/9805108.
- [137] X. Chen and M. Kamionkowski, Phys. Rev. **D70**, 043502 (2004), arXiv:astro-ph/0310473.
- [138] N. Padmanabhan and D. P. Finkbeiner, Phys. Rev. **D72**, 023508 (2005).
- [139] S. Galli, F. Iocco, G. Bertone, and A. Melchiorri, Phys. Rev. D **80**, 023505 (2009), 0905.0003.
- [140] T. R. Slatyer, N. Padmanabhan, and D. P. Finkbeiner, Phys. Rev. **D80**, 043526 (2009).
- [141] T. R. Slatyer and C.-L. Wu, Phys. Rev. **D95**, 023010 (2017), 1610.06933.
- [142] V. Poulin, J. Lesgourgues, and P. D. Serpico, JCAP **3**, 043 (2017), 1610.10051.
- [143] K. R. Dienes, J. Kumar, P. Stengel, and B. Thomas, ArXiv e-prints (2018), 1810.10587.
- [144] R. J. Wilkinson, C. Boehm, and J. Lesgourgues, JCAP **2014**, 011 (2014), 1401.7597.
- [145] C. Dvorkin, K. Blum, and M. Kamionkowski, Phys. Rev. **D89**, 023519 (2014), 1311.2937.
- [146] R. J. Wilkinson, J. Lesgourgues, and C. é. Boehm, JCAP **2014**, 026 (2014), 1309.7588.
- [147] V. Gluscevic and K. K. Boddy, Phys. Rev. Lett **121**, 081301 (2018), 1712.07133.
- [148] K. K. Boddy and V. Gluscevic, Phys. Rev. **D98**, 083510 (2018), 1801.08609.
- [149] K. K. Boddy, V. Gluscevic, V. Poulin, E. D. Kovetz, M. Kamionkowski, and R. Barkana, Phys. Rev. **D98**, 123506 (2018), 1808.00001.
- [150] M. Kawasaki, K. Kohri, and T. Moroi, Phys. Rev. **D71**, 083502 (2005).
- [151] K. Jedamzik, JCAP **3**, 008 (2008), 0710.5153.
- [152] M. Kawasaki, K. Kohri, T. Moroi, and Y. Takaesu, Phys. Rev. **D97**, 023502 (2018).
- [153] V. Poulin and P. D. Serpico, Phys. Rev. Lett **114**, 091101 (2015), 1502.01250.
- [154] S. Sarkar and A. M. Cooper, Physics Letters B **148**, 347 (1984).
- [155] J. Ellis, D. V. Nanopoulos, and S. Sarkar, Nuclear Physics B **259**, 175 (1985).
- [156] M. Kawasaki and K. Sato, Physics Letters B **169**, 280 (1986).
- [157] W. Hu and J. Silk, Phys. Rev. Lett **70**, 2661 (1993).
- [158] J. L. Aalberts, S. Ando, W. M. Borg, E. Broeils, J. Broeils, S. Broeils, B. J. Kavanagh, G. Leguijt, M. Reemst, D. R. van Arneman, et al., Phys. Rev. **D98**, 023001 (2018).
- [159] P. McDonald, R. J. Scherrer, and T. P. Walker, Phys. Rev. **D63**, 023001 (2001).
- [160] H. Tashiro, J. Silk, and D. J. E. Marsh, Phys. Rev. **D88**, 125024 (2013), 1308.0314.
- [161] D. Ejlli and A. D. Dolgov, Phys. Rev. **D90**, 063514 (2014), 1312.3558.
- [162] S. Mukherjee, R. Khatri, and B. D. Wandelt, JCAP **4**, 045 (2018), 1801.09701.
- [163] E. Dimastrogiovanni, L. M. Krauss, and J. Chluba, Phys. Rev. **D94**, 023518 (2016).
- [164] J. P. Ostriker and C. Thompson, ApJL **323**, L97 (1987).
- [165] H. Tashiro, E. Sabancilar, and T. Vachaspati, JCAP **8**, 035 (2013), 1212.3283.
- [166] Y. Ali-Haïmoud, J. Chluba, and M. Kamionkowski, Phys. Rev. **D115**, 071304 (2015).
- [167] J. A. D. Diacoumis and Y. Y. Y. Wong, JCAP **9**, 011 (2017), 1707.07050.
- [168] S. Kumar, E. Dimastrogiovanni, G. D. Starkman, C. Copi, and B. Lynn, arXiv e-prints (2018), 1804.08601.
- [169] K. Jedamzik, V. Katalinić, and A. V. Olinto, PRL **85**, 700 (2000), arXiv:astro-ph/9911100.

- [170] S. K. Sethi and K. Subramanian, *MNRAS* **356**, 778 (2005), astro-ph/0405413.
- [171] K. E. Kunze and E. Komatsu, *JCAP* **1**, 009 (2014), 1309.7994.
- [172] J. M. Wagstaff and R. Banerjee, *Phys. Rev.* **D92**, 123004 (2015), 1508.01683.
- [173] R. A. Sunyaev and Y. B. Zeldovich, *A&A* **20**, 189 (1972).
- [174] W. Hu, D. Scott, and J. Silk, *Phys. Rev. D* **49**, 648 (1994), arXiv:astro-ph/9305038.
- [175] R. Cen and J. P. Ostriker, *ApJ* **514**, 1 (1999), arXiv:astro-ph/9806281.
- [176] F. Miniati, D. Ryu, H. Kang, T. W. Jones, R. Cen, and J. P. Ostriker, *ApJ* **542**, 608 (2000).
- [177] S. P. Oh, A. Cooray, and M. Kamionkowski, *MNRAS* **342**, L20 (2003).
- [178] R. Khatri and R. Sunyaev, *JCAP* **2015**, 013 (2015), 1505.00781.
- [179] J. C. Hill, N. Battaglia, J. Chluba, S. Ferraro, E. Schaan, and D. N. Spergel, *Phys. Rev. Lett* **115**, 261301 (2015), 1507.01583.
- [180] E. L. Wright, *ApJ* **232**, 348 (1979).
- [181] Y. Rephaeli, *ApJ* **445**, 33 (1995).
- [182] S. Y. Sazonov and R. A. Sunyaev, *ApJ* **508**, 1 (1998).
- [183] N. Itoh, Y. Kohyama, and S. Nozawa, *ApJ* **502**, 7 (1998), arXiv:astro-ph/9712289.
- [184] A. Challinor and A. Lasenby, *ApJ* **499**, 1 (1998), arXiv:astro-ph/9711161.
- [185] G. B. Rybicki and I. P. dell'Antonio, *ApJ* **427**, 603 (1994), astro-ph/9312006.
- [186] V. K. Dubrovich and V. A. Stolyarov, *A&A* **302**, 635 (1995).
- [187] E. E. Kholupenko, A. V. Ivanchik, and D. A. Varshalovich, *Gravitation and Cosmology* **11**, 161 (2005), astro-ph/0509807.
- [188] W. Y. Wong, S. Seager, and D. Scott, *MNRAS* **367**, 1666 (2006), astro-ph/0510634.
- [189] J. A. Rubiño-Martín, J. Chluba, and R. A. Sunyaev, *MNRAS* **371**, 1939 (2006).
- [190] J. Chluba and R. A. Sunyaev, *A&A* **446**, 39 (2006), astro-ph/0508144.
- [191] J. A. Rubiño-Martín, J. Chluba, and R. A. Sunyaev, *A&A* **485**, 377 (2008).
- [192] Y. Ali-Haïmoud, *Phys. Rev. D* **87**, 023526 (2013), 1211.4031.
- [193] J. Chluba and R. A. Sunyaev, *A&A* **478**, L27 (2008), 0707.0188.
- [194] Y. B. Zeldovich, V. G. Kurt, and R. A. Syunyaev, *ZhETF* **55**, 278 (1968).
- [195] P. J. E. Peebles, *ApJ* **153**, 1 (1968).
- [196] R. A. Sunyaev and Y. B. Zeldovich, *ApSS* **7**, 3 (1970).
- [197] P. J. E. Peebles and J. T. Yu, *ApJ* **162**, 815 (1970).
- [198] W. Hu, D. Scott, N. Sugiyama, and M. White, *Phys. Rev. D* **52**, 5498 (1995).
- [199] A. Lewis, J. Weller, and R. Battye, *MNRAS* **373**, 561 (2006), astro-ph/0606552.
- [200] J. A. Rubiño-Martín, J. Chluba, W. A. Fendt, and B. D. Wandelt, *MNRAS* **403**, 439 (2010).
- [201] J. R. Shaw and J. Chluba, *MNRAS* **415**, 1343 (2011), 1102.3683.
- [202] A. Loeb, *ApJL* **555**, L1 (2001), astro-ph/0103505.
- [203] M. Zaldarriaga and A. Loeb, *ApJ* **564**, 52 (2002), astro-ph/0105345.
- [204] K. Basu, C. Hernández-Monteagudo, and R. A. Sunyaev, *A&A* **416**, 447 (2004).
- [205] J. A. Rubiño-Martín, C. Hernández-Monteagudo, and R. A. Sunyaev, *A&A* **438**, 461 (2005).
- [206] C. Hernández-Monteagudo and R. A. Sunyaev, *MNRAS* **359**, 597 (2005), astro-ph/0405487.
- [207] C. Hernández-Monteagudo, L. Verde, and R. Jimenez, *ApJ* **653**, 1 (2006), astro-ph/0604324.
- [208] C. Hernández-Monteagudo, J. A. Rubiño-Martín, and R. A. Sunyaev, *MNRAS* **380**, 1656 (2007).
- [209] C. Hernández-Monteagudo, U. Maio, B. Ciardi, and R. A. Sunyaev, arXiv e-prints (2017), 1707.01910.
- [210] Q. Yu, D. N. Spergel, and J. P. Ostriker, *ApJ* **558**, 23 (2001), astro-ph/0103149.
- [211] A. Lewis, *JCAP* **8**, 053 (2013), 1307.8148.
- [212] J. C. Mather, E. S. Cheng, D. A. Cottingham, R. E. Eplee, Jr., D. J. Fixsen, T. Hewagama, R. B.

- Isaacman, K. A. Jensen, S. S. Meyer, P. D. Noerdlinger, et al., *ApJ* **420**, 439 (1994).
- [213] A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M. Meyer, S. H. Moseley, M. D. Seiffert, et al., *JCAP* **7**, 25 (2011), 1105.2044.
- [214] A. Kogut, J. Chluba, D. J. Fixsen, S. Meyer, and D. Spergel, in *SPIE Conference Series* (2016), vol. 9904 of *Proc.SPIE*, p. 99040W.
- [215] S. Masi, P. Ade, P. de Bernardis, A. Boscaleri, M. De Petris, G. De Troia, M. Fabrini, M. Giacometti, A. Iacoangeli, L. Lamagna, et al., *Memorie della Società Astronomica Italiana* **74**, 96 (2003).
- [216] A. Schillaci, G. D’Alessandro, P. de Bernardis, S. Masi, C. Paiva Novaes, M. Gervasi, and M. Zannoni, *A&A* **565**, A125 (2014).
- [217] A. Kogut, D. Fixsen, S. Fixsen, S. Levin, M. Limon, L. Lowe, P. Mirel, M. Seiffert, J. Singal, P. Lubin, et al., *New Astronomy Reviews* **50**, 925 (2006), arXiv:astro-ph/0609373.
- [218] M. Seiffert, D. J. Fixsen, A. Kogut, S. M. Levin, M. Limon, P. M. Lubin, P. Mirel, J. Singal, T. Villela, E. Wollack, et al., *ApJ* **734**, 6 (2011).
- [219] V. Desjacques, J. Chluba, J. Silk, F. de Bernardis, and O. Doré, *MNRAS* **451**, 4460 (2015).
- [220] M. Sathyanarayana Rao, R. Subrahmanyam, N. Udaya Shankar, and J. Chluba, *ApJ* **810**, 3 (2015).
- [221] P. Serra, O. Doré, and G. Lagache, *ApJ* **833**, 153 (2016), 1608.00585.
- [222] J. Chluba, J. C. Hill, and M. H. Abitbol, *MNRAS* **472**, 1195 (2017), 1701.00274.
- [223] M. H. Abitbol, J. Chluba, J. C. Hill, and B. R. Johnson, *MNRAS* **471**, 1126 (2017).
- [224] M. Remazeilles and J. Chluba, *MNRAS* **478**, 807 (2018), 1802.10101.
- [225] M. Remazeilles, A. J. Banday, C. Baccigalupi, S. Basak, A. Bonaldi, G. De Zotti, J. Delabrouille, C. Dickinson, H. K. Eriksen, J. Errard, et al., *JCAP* **4**, 023 (2018), 1704.04501.
- [226] K. Inomata and T. Nakama, arXiv e-prints (2018), 1812.00674.

Notes

ⁱWhen referring to the cosmological standard model (CSM) we assume the Λ CDM parameterization, supplemented by the Standard Model of particle physics, admitting that the presence of dark matter and dark energy requires physics beyond the latter.

ⁱⁱTo avoid the unrealistic GW spectrum generated by a δ -function scalar power spectrum, we plot all integrated constraints using a k^4 spectrum – see [67] for the reason for this choice. The peak sensitivity for μ -distortions is effectively unchanged were we to instead plot constraints for δ -function features in the power spectrum with the same integrated power (see Fig. 9 therein), and also [226].

ⁱⁱⁱAt high frequencies, these are dominated by dust from the galaxy and the cosmic infrared background, while at low frequencies it is synchrotron and free-free emission.

^{iv}We wish to acknowledge the generosity and support of the CERN Theoretical Physics Department in hosting the workshop “Probing fundamental physics with CMB spectral distortions” from Mar 12–16 2018 where this white paper was first conceived. This work has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No 725456, CMBSPEC). JC is supported by the Royal Society as a Royal Society University Research Fellow at the University of Manchester, UK.