



HAL
open science

Journal of Petroleum Technology and Alternative Fuels
Fatty acid profile and quality parameters of Ceiba
pentandra (L.) seed oil: A potential source of biodiesel

Papin Sourou Montcho, Léopold Tchiakpe, Guévara Nonviho, Fifa Théomaine
Diane Bothon, Assou Sidohounde, Cokou Pascal Agbangnan Dossa, David
Bessieres, Anna Chrostowska, Dominique Codjo Koko Sohounhloue

► **To cite this version:**

Papin Sourou Montcho, Léopold Tchiakpe, Guévara Nonviho, Fifa Théomaine Diane Bothon, Assou Sidohounde, et al.. Journal of Petroleum Technology and Alternative Fuels Fatty acid profile and quality parameters of Ceiba pentandra (L.) seed oil: A potential source of biodiesel. Journal of Petroleum Technology and Alternative Fuels (JPTAF), 2018, 9 (3), pp.14-19. 10.5897/JPTAF2018.0141 . hal-02125246

HAL Id: hal-02125246

<https://hal.science/hal-02125246>

Submitted on 10 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Full Length Research Paper

Fatty acid profile and quality parameters of *Ceiba pentandra* (L.) seed oil: A potential source of biodiesel

Papin Sourou MONTCHO¹, Léopold TCHIAKPE², Guevara NONVIHO¹, Fifa Théomaine Diane BOTHON¹, Assou SIDOHOUNDE¹, Cokou Pascal AGBANGNAN DOSSA¹, David BESSIERES³, Anna CHROSTOWSKA⁴ and Dominique Codjo Koko SOHOUNHLOUE^{1*}

¹Laboratory of Study and Research in Applied Chemistry, Polytechnic School of Abomey-Calavi, University of Abomey Calavi, Benin.

²Unité Mixte de Formation Continue en Santé, Faculty of Pharmacy, University of Aix-Marseille, France.

³CNRS /TOTAL/UNIV PAU & PAYS ADOUR/ E2S UPPA, Laboratory of Complex Fluids and Their Reservoirs-IPRA, UMR5150, 64000, PAU, France.

⁴CNRS/ Univ Pau & Pays Adour/ E2S UPPA, Institute of Analytical Sciences and Physicochemistry for Environment and Materials, UMR5254, 64000, Pau, France.

Received 17 October, 2018; Accepted 29 November, 2018

The probable depletion of fossil energy resources has led the international scientific community to direct research towards biofuels, including vegetable oils. Benin has a rich biodiversity with multiple oilseed species, potential sources of biofuels. Among these, *Ceiba pentandra* was identified and selected for a detailed study of its unconventional vegetable oil. In this order of idea, the harvested seeds were dried in the sun and crushed. It was preserved at 25°C according to the NF T 60 – 201, 1993 standard. The physicochemical parameters and fatty acids content of *C. pentandra* vegetable oil have been determined by standard methods. The biofuel potential of this vegetable oil has also been evaluated. The results revealed that *C. pentandra* vegetable oil is predominant in saturated (40.8%) and polyunsaturated (41.37%) fatty acids. The saturated fatty acids quantified are stearic (20.17%) and palmitic (19.77%), while the most important polyunsaturated fatty acids are linoleic acid (C18:2) (20.95%) and linolelaidic acid (18.28%). Quality indices such as acid (4.52 ± 0.24 mg KOH/g-Oil), peroxide (2.16 ± 0.54 meq O₂/kg-Oil), saponification (152.79 ± 6.07 mg KOH/g-Oil), iodine (129.79 ± 2.81 mg I₂/100 g-Oil) and ester (148.27 ± 5.83 mg KOH/g-Oil) shall comply with the recommended standards for biofuels. These values of quality indices coupled with those of lower calorific value (LCV) (≈ 40002.26 kJ/kg), refractive index (≈ 1.4728 at 30°C) and cetane index (≈ 49.70) make it possible to consider the use of this vegetable oil as a fuel oil.

Key words: *Ceiba pentandra*, vegetable oil, quality indexes, fatty acids, fuel properties, Benin.

INTRODUCTION

The use of biofuels as an alternative to conventional fossil fuels is increasingly discussed in the literature.

*Corresponding author. E-mail: csohoun@gmail.com or ksohoun@bj.refer.org. Tel: +229 21 36 09 93. Fax: +229 21 36 01 99/LERCA.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Among other benefits, this use has the advantage of reducing fuel costs and harmful greenhouse gas emissions (Goldemberg et al., 2008). Among biofuels, vegetable oils used as such or trans-esterified also have a renewed interest. There are generally four sources of vegetable oils with biodiesel potential: edible vegetable oils, recyclable vegetable oils, unconventional vegetable oils and vegetable oils of algal origin. Nowadays, the use of edible vegetable oils as fuels leads to a reduction in food production and indirectly, soaring prices and food insecurity (Djenontin, 2006; Oderinde et al., 2009). If the recycling of used vegetable oils into biofuels is seen as ecologically credible and more ethical than the use of edible oils, their collection and pre-treatment are not less expensive. Similarly, the different algae processing techniques for 3rd generation biofuel production are quite tedious. Besides, these technologies use catalysts that make the cost of vegetable oil barrels of algal origin prohibitive. On the other hand, unconventional vegetable oils as potential sources of biofuels have several advantages, including:

- i) their low (or no) use in food and livestock (Nonviho et al., 2014),
- ii) their high availability due to their production on degraded and marginal soils for food crops,
- iii) their biofuel use that can be done without major treatments when their physicochemical and fuel properties are acknowledged (Sidohoude et al., 2018),
- iv) and an easily reproducible and inexpensive technology (Atabani et al., 2013) through their transformation into biodiesel by transesterification.

Kaimal and Lakshminarayan reported the presence of cyclopropenic fatty acid (malvalic acid) in vegetable oils extracted from the seeds and barks of *Ceiba pentandra* (the kapok tree) (Kaimal et al., 1970). For that purpose, it would be unwise to use the seeds of this plant species for food.

In Benin, this species is widely distributed and also exploited for its wood. Kapok wood (trade name 'fuma') is lightweight, porous, durable wood and useful for making plywood, packaging, carvings and dugout canoes (Orwa et al., 2009; Tessi et al., 2012). *C. pentandra* may disappear if other uses such as seeds are not highlighted. It must be emphasized that the kapok tree under optimal conditions can produce 330 to 400 fruits a year and about 30 kg of seeds (Iko et al., 2015). Moreover, these seeds are high in fiber and the fibers are valued as livestock feed (Chaiarekij et al., 2011; Bationo et al., 2012). It is for this purpose that the present study aims to evaluate the physicochemical characteristics, fuel and fatty acid profile of the unconventional vegetable oil extracted from these seeds.

MATERIALS AND METHODS

Vegetal material

Mature fruits of *C. pentandra* were collected in Kpataba, a village of

Savalou in central Benin.

Seed conditioning and extraction of vegetable oil

The fruits were dried in the laboratory ($27 \pm 2^\circ\text{C}$) for at least seven (7) days. Thereafter, their kernels were collected and all the impurities were removed by manual sorting. These kernels have been milled with a machine and the obtained powders sieved and then stored in an oven (40°C) to constant mass (water and volatile matters elimination).

These powders are then extracted with hexane using the Soxhlet device for 6 h at 69°C respecting the protocol of NF V03-924. The extracted vegetable oil was then stored in opaque dry, flasks, and the extraction yields were evaluated gravimetrically.

$$W = \frac{m_2 - m_1}{m_0} \times 100 \quad (1)$$

where W: extraction yields, m_2 : mass of the balloon containing the oil extracted, m_1 : mass of empty balloon, and m_0 : mass of the test sample.

Determination of the quality indexes from vegetable oil extracted

Water content and acid, iodine and saponification indexes

Water content, volatile matters and the density of the vegetable oil were determined according to DIN EN ISO 12937 and NF T 60-214 methods. The acid (Ia), peroxide (Ip) and saponification (Is) indexes have been determined respecting the French standards: T 60-204; T 60-220 and T 60-206. The iodine (Ii) index has been evaluated using the Winkler method.

Determination of the fuel characteristics of vegetable oils

The ester value (IE) has been calculated on the basis of the analytical data according to the formula (Dahouenon-Ahoussi et al., 2012).

$$Ie = Is - Ia \quad (2)$$

The lower calorific value (LCV) has been calculated using the formula of Batel et al. (1980).

$$LCV = 47645 - 4.15 I_i - 38.31 \quad (3)$$

The refractive index (Ri) has been determined using the formula of Perkins et al. (1995) and Asemave et al. (2012).

$$Ri = 1.45765 + 1.164 \times 10^{-4} I_i \quad (4)$$

The cetane number (CN) was calculated using the formula of Klopfenstein (1982) and Haidara (1996).

$$CN = 46.3 + \frac{5458}{I_s} - 0.225 I_i \quad (5)$$

Determination of fatty acid composition by GC/MS

Oil samples were transesterified as fatty acid methyl esters (FAMES) following a validated method slightly adapted from

Table 1. Quality indices of the vegetable oil of *C. pentandra* seeds.

| Characteristic | <i>Ceiba pentandra</i> |
|-----------------------------------------------|------------------------|
| Moisture content and volatile matters (%) | 6.05 ± 0.36 |
| Extraction yield (%) | 31.62 ± 1.60 |
| Acid number (mg KOH / g-Oil) | 4.52 ± 0.24 |
| Saponification number (mg KOH / g-Oil) | 152.79 ± 6.07 |
| Iodine Number (mg I ₂ / 100 g-Oil) | 129.79 ± 2.81 |
| Peroxide value (meq O ₂ / kg-Oil) | 2.16 ± 0.54 |
| Calculated Ester Index (mg KOH / g-Oil) | 148.27 ± 5.83 |

methods previously described by Lepage and Roy (1984) and by Masood and Stark (2005).

The fatty acid composition of the transesterified unconventional vegetable oil of *C. pentandra* was determined by coupling Gas Chromatography (Thermo Fischer Scientific Ultra Brand) with mass spectrometry (GC/MS).

Chromatographic analyses were performed on Trace GC Ultra equipped with an ASI 3000 autosampler and with Polaris Q spectral mass detector, all from Thermo Fischer Scientific Ultra Brand. The coupling and automatic control of the devices have been done by the software EXCALIBUR 2.0 Thermo Fisher.

The splitless injector was set at 250°C and the ion source temperature set at 250°C. Ultrapure Helium Alpha-gas 2 was the carrier gas set at 1 mL/min constant flow with automatically adjusted pressure. Injections were on split mode. Gas Chromatography was fitted with a fused silica capillary column (DB-FFAP) 30 m (length) × 0.25 mm inner diameter (id) × 0.25 µm film thickness (J & W Scientific, Agilent Technologies).

Initial oven temperature was 130°C. The program temperature was as follows: equilibration time: 0.5 min; linear increase to 178°C at 4°C/min, followed by linear increase to 210°C at 1°C/min, followed by an increase to 245°C at 40°C/min and final 13 min hold. The duration of the analysis was 60 min. The injected volume was 1 µL and the injected amount 10 µg/mL.

Positive ionisation of the FAMES was performed by electronic impact (EI), with 70 eV energy and full scan detection mode. Mass spectra range was 50 to 650 m/z; scan 0.58 s.

Precise identification of the analytes was achieved by their relative retention times and mass spectra on the spectral mass database NIST libraries for fatty acid composition. External fatty acid standards were the 28 FAME compounds NU-CHEK-PREP Inc Elysian USA, (GLC reference standard 462) and the Supelco 37 component FAME mix (CRM 47885).

RESULTS AND DISCUSSION

Moisture content and vegetable oil

The quality indexes of the vegetable oil investigated are shown in Table 1. The water and volatile matter content of *C. pentandra* seeds has been reduced, by heating in the open air to 6.05±0.36% for a vegetable oil extraction yield of 31.62%±1.60%.

Kaimal and Lakshminarayana found a vegetable oil content of *C. pentandra* seeds of 23.6%. This value is much lower than that of the seeds investigated in present study. Researches have shown that the lipid potential of seeds could depend on several parameters such as

those related to seed maturity and edaphic conditions (Atabani et al., 2013).

However, according to the vegetable oil content values obtained (> 30%) in the present study on *C. pentandra* seeds harvested in Benin, we could consider its use as fuel oil (Alabi et al., 2013). For this purpose, we have evaluated the quality indices of this vegetable oil.

Acid index (Ia)

The acid value of a vegetable oil depends on its free fatty acid composition. It characterizes the state of alteration of the oil by hydrolysis (Bettahar et al., 2016). The acid value of the vegetable oil of *C. pentandra* is 4.52 ± 0.24 mg KOH / g-oil. It has been shown that a high acid value of a fuel oil causes the injectors clogging and the metal workpieces corrosion (Stauffer et al., 2005). Khan et al. (2015) found an acid value of *C. pentandra* oil collected in Java (Indonesia) 4.5 times (20.23 mg KOH / g-Oil) higher than our values.

Peroxide index (Ip)

The oxidation stability of this vegetable oil has been evaluated by quantification of its peroxide index. In fact, the high peroxide indexes are at the base of the polymerization of the esters and the formation of the gums and sediments, which clog the filters of the engines (Clark et al., 1984). This index is 2.16 ± 0.54 meq O₂/kg-oil for the vegetable oil of *C. pentandra* of Benin. It is relatively low and indicates that this vegetable oil could have a cetane number that meets the standard. There is indeed a positive correlation between the cetane number of vegetable fuel oils and their peroxide index (Abdul, 1998).

Saponification and ester indexes (Is and Ie)

The saponification and ester index provides a first idea of the lengths of the fatty acids of the vegetable oil studied (Zovi et al., 2011). The vegetable oil of *C. pentandra* has

Table 2. Fatty acids from vegetable oil extracted from *Ceiba pentandra* seeds harvested in Benin compared to literature data.

| Fatty acid | <i>C. pentandra</i> harvested in Benin | <i>C. pentandra</i> (Kaimal et al., 1970; Sivakumar et al., 2013; Silitonga et al., 2013) |
|-----------------------------------|-------------------------------------------|-------------------------------------------------------------------------------------------|
| Lauric acid (C12:0) | - | - |
| Tridecyl acid (C13:0) | 0.14 | - |
| Myristic acid (C14:0) | 0.03 | 0.1-0.25 |
| Palmitic acid (C16:0) | 19.77 | 20.36-24.31 |
| Margaric acid (C17:0) | 0.06 | - |
| Stearic acid (C18:0) | 20.17 | 0.8-2.85 |
| Arachidic acid (C20:0) | 0.3 | 0.7 |
| Behenic acid (C22:0) | 0.16 | 0.44 |
| Tricosylic acid (C23:0) | 0.11 | - |
| Lignocéric acid (C24:0) | 0.06 | - |
| Total saturated fatty acids | 40.8 | 21.26-28.55 |
| Pentadecanoic acid (C15:1) | - | - |
| Palmitoleic acid (C16:1) | 0.16 | 0.4 |
| margaroleic acid (C17:1) | 0.21 | - |
| Oleic acid (C18:1) | 18.82 | 17.95-30.0 |
| Gondoic acid (C20 :1) | 0.08 | - |
| Total monounsaturated fatty acids | 19.27 | 17.95-30.4 |
| Linolelaidic acid (C18:2) | 18.28 | - |
| Linoleic acid (C18:2) | 20.95 | 32.9-41.43 |
| α -Linolenic acid (C18:3) | 1.14 | - |
| γ -Linolenic acid (C18:3) | 0.92 | - |
| Eicosapentaenoic acid (C20:5) | 0.08 | - |
| Total polyunsaturated fatty acids | 41.37 | 32.9-41.43 |
| Malvalic acid | - | 7.18-18.5 |
| Dihydromalvalic acid | - | 0.7 |
| Sterculic acid | - | 2.96-3.4 |
| Total cyclopropenic fatty acids | - | 0.7-22.6 |

-: Unquantified fatty acids.

respectively, saponification and ester indexes of 152.79 ± 6.07 mg KOH / g-oil and 148.27 ± 5.83 mg KOH / g-oil. Berry (1979) found a higher saponification index (183 mg KOH / g-oil) for vegetable oil extracted from *C. pentandra* seeds harvested in Malaysia (Berry et al., 1979). This index variation depends on the fatty acid profiles of the vegetable oils and shows that *C. pentandra* vegetable oil from Benin could have a different fatty acid profile from that of Malaysia.

Iodine index (Ii)

A classification of the vegetable oils fuels, according to their index of iodine, has been made by Vaitilingom et al. (1983). The vegetable oil of *C. pentandra* had an iodine value higher than the value of 110 mg I₂/100 g oil. According to Vaitiligom's classification, the oil of *C. pentandra* can be classified among linoleic semi-drying

oils (Vaitilingom et al., 1983).

Fatty acid profiles of vegetable oils

There is evidence that the fatty acid composition of a vegetable oil can have a great influence on its fuel characteristics (Ferhat et al., 2014). The quantified fatty acids of the oil of *C. pentandra* are listed in Table 2.

In accordance with its relatively high iodine value, the vegetable oil of *C. pentandra* is rich in polyunsaturated fatty acid (41.37%) including linoleic acids (cis and trans C18: 39.23%). Our chromatographic analyses were able to quantify the cis (all-cis- $\Delta^{9,12}$: 20.95%) and trans (all-trans- $\Delta^{9,12}$: 18.28%) isomers of the linoleic acid of this vegetable oil. Previous work has revealed relatively higher proportions of linoleic acids combined in *C. pentandra* (Atabani et al., 2013; Abdul, 1998).

A good cetane index has been associated with

Table 3. Fuel characteristics of the vegetable oil of *C. pentandra* seeds.

| Characteristics | <i>Ceiba pentandra</i> |
|-------------------------------------|------------------------|
| Density at 30°C | 0.92 |
| Refractive index at 30°C | 1.47 ± 0.01 |
| Cetane index | 52.85 ± 1.45 |
| Lower calorific value (LCV) (kJ/kg) | 41248.17 ± 244.19 |

vegetable fuels having a good saturated fatty acid composition such as palmitic acid and stearic acid (Gerhard et al., 2003). The proportion of saturated fatty acid (40.8%) of *C. pentandra* vegetable oil harvested in Benin is different from the values reported in the literature (Atabani et al., 2013; Sivakumar et al., 2013).

Palmitic acid and stearic acid (palmitic: 19.77 and stearic: 20.17%) are the most important saturated fatty acids of *C. pentandra* vegetable oil from Benin. As indicated above, this is in conformity with its saponification index, in negative correlation. This oil is composed of oleic acid, palmitic acid, followed by linoleic acid. These components are able to improve not only certain important fuel properties like cetane number, heat of combustion, oxidative stability, and kinematic viscosity (C18:1, C16:0), but also the cold flow properties of biodiesel (C18:2) as shown in the works of Knothe et al. (2008).

Based on this profile of fatty acid composition, it is clearly assumed that *C. pentandra* oil is suitable for biodiesel production.

Against all expectations, *C. pentandra* vegetable oil from Benin is found to be free of malvalic acid; usually quantified in this oil by Halphen tests or other similar methods (Pawlowski et al., 1972). However, the presence of oleic acid (18.82%), fatty acid at the origin of the biosynthesis of malvalic acid, is observed. This could be explained by the genotypic differences related to soil, climate and other parameters not studied here.

Fuel characteristics

An estimation of the potential fuels of this vegetable oil has been made (Table 3). The refractive index of the oil varies according to their degree of unsaturation. The refractive index and the density have been evaluated at a temperature of 30°C. Thus, the density and refractive index values of *C. pentandra* oil are 0.92 and 1.47, respectively. This obtained value of refractive index reveals the need to purify these oils before any use.

The lower calorific value (LCV) is the amount of energy released when one kilogram of fuel is burned. The LCV of *C. pentandra* oil from Benin is 41248.17 ± 244.19 kJ/kg. This value is of the same order of magnitude as that of *C. pentandra* (40493 kJ/kg), *Jatropha curcas* (40224 kJ/kg) and that of palm oil (40151 kJ/kg) found by Sivakumar et

al. (2013), and then Yunus et al. (2014). It is higher than the value of 35000 kJ/kg recommended for pure vegetable fuels (Ong et al., 2013; Sidohoude et al., 2018).

The cetane index measures a fuel ability to ignite itself (Aligrot, 1994). This characteristic is particularly important for diesel fuel where the fuel must "ignite" under the effect of the compression of the air enclosed in the cylinder. The higher the cetane number, the shorter the ignition delay and the better the combustion quality (Haidara et al., 1996). The cetane index of *C. pentandra* vegetable oil is 52.85 ± 1.45 (Table 3). This value was close to that of petrol diesel recommended by ASTM D975 (40-55) (Sidohoude et al., 2018).

Conclusion

This study evaluated the physicochemical properties, fuel potentials and fatty acid composition of unconventional vegetable oil extracted from *C. pentandra* seeds harvested from Benin. The vegetable oil possessed quality indexes and a fatty acid composition that indicated its possible adaptation to use as energy oils.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Abdul M (1998). The effect of biodiesel oxidation on engine performance and emissions, Retrospective Theses and Dissertations. Iowa State University Capstones. 11950. <https://lib.dr.iastate.edu/rtd/11950>
- Alabi K A (2013). Analysis of fatty acid composition of Thevetia peruviana and Hura crepitans seed oils using GC-FID. Fountain Journal of Natural and Applied Sciences, 2(2).
- Asemave K, Ubwa S T, Anhwange B A, Gbaamende A G (2012). Comparative evaluation of some metals in palm oil, groundnut oil and soybean oil from Nigeria. International Journal of Modern Chemistry, 1(1), 28-35.
- Atabani AE, Silitonga AS, Ong HC, Mahlia TMI, Masjuki HH, Badruddin IA, Fayaz H (2013). Non-edible vegetable oils: a critical evaluation of oil extraction, fatty acid compositions, biodiesel production, characteristics, engine performance and emissions production. Renewable and Sustainable Energy Reviews 18:211-245.
- Batel W, Graef M, Mejer G J, Möller R, Schoedder F (1980). Pflanzenöle für die Kraftstoff-und Energieversorgung. Grundlagen der landtechnik 30(2).

- Bationo BA, Kalinganire A, Bayala J (2012). Potentials of woody species in the practice of conservation agriculture in arid and semi-arid zones of West Africa: Overview of some candidate systems. ICRAF Technical Manual (17).
- Berry S K (1979). The characteristics of the kapok (*Ceiba pentandra*, Gaertn.) seed oil. *Pertanika* 2(1):1-4.
- Bettahar Z, Cheknane B, Boutemak K (2016). Study of transesterification of a mixture of waste oil for biodiesel production. *Renewable and Sustainable Energy Reviews* 19(4):605-615.
- Chaiarekij S, Apirakchaiskul A, Suvarnakich K, Kiatkamjornwong S (2011). Kapok I: characteristics of Kapok fiber as a potential pulp source for papermaking. *Bio Resources* 7(1):0475-0488.
- Clark SJ, Wagner L, Schrock MD, Piennaar PG (1984). Methyl and ethyl soybean esters as renewable fuels for diesel engines. *Journal of the American Oil Chemists Society* 61(10):1632-1638.
- Dahouenon-Ahoussi E, Djenontin TS, Codjia DR, Tchobo FP, Alitonou AG, Dangou J, Sohounhloue CD (2012). Fruit morphology and some physical and chemical characteristics of oil and oilcake from *Irvingia gabonensis* (Irvingiaceae). *International Journal of Biological and Chemical Sciences* 6(5):2263-2273.
- Djenontin TS (2006). Benin Oilseed Study: Chemical Characteristics, Fractionation and Biocidal Activity; thesis defended on December 15, 2006.
- Gerhard K, Andrew C, Matheaus T, Ryan W (2003). Cetane numbers of branched and straight-chain fatty esters determined in an ignition quality tester. *Fuel* 82:971-975.
- Goldemberg J (2008). Environmental and ecological dimensions of biofuels. In *Proceedings of the conference on the ecological dimensions of biofuels*, Washington, DC Vol. 10.
- Haidara H, Vonna L, Schultz J (1996). Kinetics and thermodynamics of surfactant adsorption at model interfaces: evidence of structural transitions in the adsorbed films. *Langmuir* 12(13):3351-3355.
- Iko W, Eze S, Oscar O (2015). Gas Chromatography Mass Spectrometry of *Quassia undulata* Seed Oil. *Nigerian Journal of Biotechnology* 30(1):53-58.
- Kaimal TNB, Lakshminarayana G (1970). Fatty acid compositions of lipids isolated from different parts of *Ceiba pentandra*, *Sterculia foetida* and *Hydnocarpus wightiana*. *Phytochemistry* 9(10):2225-2229.
- Khan TY, Atabani AE, Badruddin IA, Ankalgi RF, Khan TM, Badarudin A (2015). *Ceiba pentandra*, *Nigella sativa* and their blend as prospective feedstocks for biodiesel. *Industrial Crops and Products* 65:367-373.
- Klopfenstein WE (1982). Estimation of Cetane Index for Esters of Fatty Acids. *Journal of the American Oil Chemists Society* 59(12):531-533.
- Knothe G (2008). "Designer" Biodiesel: Fatty Ester Composition to Improve Fuel Properties. *Energy and Fuels* 22:1358-1364.
- Lepage G, Roy CC (1984). Improved recovery of fatty acid through direct transesterification without prior extraction or purification. *Journal of Lipid Research* 25:1391-1396.
- Masood A, Stark KD (2005). Salem N, A simplified and efficient method for the analysis of fatty acid methyl esters suitable for large clinical studies. *Journal of Lipid Research* 46:2299-2305.
- Nonviho G, Paris C, Muniglia L, Sessou P, Agbangnan DCP, Brosse N, Sohounhloue D (2014). Chemical characterization of *Lophira lanceolata* and *Carapa procera* seed oils: Analysis of fatty acids, sterols, tocopherols and tocotrienols. *Research Journal of Chemical Sciences* 4(9):57-62.
- Oderinde RA, Ajayi IA, Adewuyi A (2009). Characterization of seed and seed oil of *Hura crepitans* and the kinetics of degradation of the oil during heating. *Electron. Journal Environmental Agricultural Food Chemistry* 8(3):201-208.
- Ong LK, Effendi C, Kurniawan A, Lin CX, Zhao XS, Ismadi S (2013). Optimization of catalyst-free production of biodiesel from *Ceiba pentandra* (kapok) oil with high free fatty acid contents. *Energy* 57:615-623.
- Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A (2009). *Agroforestry database: a tree species reference and selection guide version 4.0*. World Agroforestry Centre ICRAF, Nairobi, KE.
- Pawlowski NE, Nixon JE, Sinnhuber RO (1972). Assay of cyclopropanoid lipids by nuclear magnetic resonance. *Journal of the American Oil Chemists Society* 49(6):387-392.
- Perkins EG (1995). Physical Properties of Soybeans and Soybean Products. In *Practical Handbook of Soybean Processing and Utilization* pp. 29-38.
- Sidhounde A, Agbangnan DCP, Nonviho G, Montcho PS, Sohounhloue CKD (2018). Biodiesel potentials of two phenotypes of *Cyperus esculentus* unconventional oils: *Journal of Petroleum Technology and Alternative Fuels* 9(1):1-6.
- Sivakumar P, Sindhanaiselvan S, Gandhi NN, Devi SS, Renganathan S (2013). Optimization and kinetic studies on biodiesel production from underutilized *Ceiba pentandra* oil. *Fuel* 103:693-698.
- Stauffer E (2005). A review of the analysis of vegetable oil residues from fire debris samples: spontaneous ignition, vegetable oils, and the forensic approach. *Journal of Forensic Science* 50(5):JFS2004510-10.
- Tessi DY, Akouhou GS, Ganglo JC (2012). Structural and ecological characteristics of the populations of *Antiaris toxicaria* (Pers.) Lesch and *Ceiba pentandra* (L.) Gaertn in the relict forests of South Benin. *International Journal of Biological and Chemical Sciences* 6(6):5056-5067.
- Vaitilingom G (1983). Use of vegetable oils as fuel for diesel engines. *Journal of Tropical Agricultural Machinery* 82:58-65.
- Yunus KTM, Atabani AE, Badruddin IA, Badarudin A, Khayoond MS, Tri-wahyond S (2014). Recent scenario and technologies to utilize non edible oils for biodiesel production. *Renewable and Sustainable Energy Reviews* 37(0):840-851.
- Zovi O, Lecamp L, Loutelier-Bourhis C, Lange CM, Bunel C (2011). Stand reaction of linseed oil. *European Journal of Lipid Science and Technology* 113(5):616-626.