

Modeling Trap-Awareness and Related Phenomena in Capture-Recapture Studies

Roger Pradel, Ana Sanz-Aguilar

► To cite this version:

Roger Pradel, Ana Sanz-Aguilar. Modeling Trap-Awareness and Related Phenomena in Capture-Recapture Studies. PLoS ONE, 2012, 7 (3), pp.e32666. 10.1371/journal.pone.0032666 . hal-02126307

HAL Id: hal-02126307 https://hal.science/hal-02126307

Submitted on 11 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modeling Trap-Awareness and Related Phenomena in Capture-Recapture Studies

Roger Pradel*, Ana Sanz-Aguilar

Biostatistics and Population Biology Group, Centre d'Ecologie Fonctionnelle et Evolutive, Centre National de la Recherche Scientifique, Montpellier, France

Abstract

Trap-awareness and related phenomena whereby successive capture events are not independent is a feature of the majority of capture-recapture studies. This phenomenon was up to now difficult to incorporate in open population models and most authors have chosen to neglect it although this may have damaging consequences. Focusing on the situation where animals exhibit a trap response at the occasion immediately following one where they have been trapped but revert to their original naïve state if they are missed once, we show that trap-dependence is more naturally viewed as a state transition and is amenable to the current models of capture-recapture. This approach has the potential to accommodate lasting or progressively waning trap effects.

Citation: Pradel R, Sanz-Aguilar A (2012) Modeling Trap-Awareness and Related Phenomena in Capture-Recapture Studies. PLoS ONE 7(3): e32666. doi:10.1371/journal.pone.0032666

Editor: Mark S. Boyce, University of Alberta, Canada

Received September 5, 2011; Accepted January 31, 2012; Published March 2, 2012

Copyright: © 2012 Pradel, Sanz-Aguilar. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Dr. Sanz-Aguilar was funded by the Marie Curie Fellowship MATERGLOBE (grant #251809). The Spanish Ministry of Science funded the study through project CGL 2009-08298. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: roger.pradel@cefe.cnrs.fr

Introduction

Live-trapping is a fundamental tool in the study of wildlife species and populations. When different trapping methods are used, empirical studies have found that different devices tend to catch different individuals [1-3]. While trappability with a particular device can sometimes be related to an identifiable feature (sex, age, weight [1], temperament [4]), this is not always possible. There is also evidence that knowledge of the trap fades with the passing of time [2]. The trap response issue is thus particularly acute when intervals between trapping occasions are short as is the case in closed population studies aiming at estimating population size. In these studies, it is generally considered that once an individual has been captured, its trappability changes for the rest of the study. Because trap response in this context is strong and because population size tends to be largely underestimated when the phenomenon is ignored, most work has been devoted to correcting for it in closed population models [5–7]. On the other hand, trap response in open population studies where occasions are generally separated by long intervals, typically a year, is much less considered. Yet, although the phenomenon is probably less intense, underestimation of survival is a true risk [8]. In this paper, we focus on short-time trap response in open populations, namely response affecting trappability solely at the occasion following one when the animal was trapped. This situation results in successive capture events being correlated and can be detected by appropriate tests-'Test 2.CT' for data from a single site [9] and 'Test M.ITEC' for multisite or multistate data [10]. However, the reciprocal is not true. With the above tests, trap dependence between successive occasions has been found when animals are captured in baited traps (trap-dependence stricto sensu) (e.g. [11-13]) but also in studies where individuals are not physically captured (trap-dependence lato sensu). Some situations where trap-dependence lato sensu occurs are: 1) When observers tend to visit some parts of the study area more often when marked individuals have been detected [14-15]; 2) When some patches of a heterogeneous habitat are more accessible so that individuals stationed there have higher resighting probabilities [16-17]; 3) When age, sex or social status are unknown, but determine individual movements or activity patterns so that the susceptibility to be recaptured or resighted varies [18-19]; 4) Or when non random temporary emigration occurs [8], often in relation to skipped reproduction [20-22]. For simplicity, we speak hereafter of 'trap dependence' to designate any correlation between capture events whatever its nature, as it is difficult to know for sure what type of trap dependence is at play in a particular study.

A survey of the literature shows that trap-dependence is a frequent phenomenon (Appendix S1). However, although the corresponding tests are largely available (program U-CARE, [23]), not all studies examine trap-dependence and it is not always clear whether this has been done in a particular study. Taking as a vardstick the papers citing Pradel (1993) where details of the way to detect and model trap-dependence in open populations were first expounded, the prevalence of trap dependence can be estimated at 71% (94/133) and touches several animal groups: birds, mammals, reptiles, amphibians, fish and insects (see Appendix S1). As for its nature, 32 papers put forward no interpretation, 26 evoke temporary emigration, 16 trap response, 8 individual heterogeneity, 7 the sampling protocol (biased sampling of known nests [14-15], unequal nest accessibility [3,24]) and 5 some behavioral feature not directly related to the trap such as dominance. For some, in particular those evoking individual heterogeneity, the restriction of sighting dependence to one

occasion may be too crude an approximation; specific models would be more appropriate [25]. Similarly, there exist specific models for temporary emigration [26]. Remarkably, only 76 of the 94 studies went on to incorporate trap-dependence at the data analysis stage. The method originally proposed to model trapdependence [9] is indeed cumbersome and unnatural as a single individual has to be represented by several capture histories. In particular, it is difficult to combine with age-dependency. Another approach using individual covariates to code for capture at the preceding occasion [27–28] is probably more natural but still uneasy to put in practice. We propose here a new approach where trap-dependence is modeled as a change of state allowing it to be naturally incorporated in the current capture-recapture models [29].

Methods

Immediate Trap Effect seen as an animal state

Here, we describe the implementation of the basic Immediate Trap Effect on Capture model (ITEC, [9]) using trappability states. This model assumes that, when an animal is caught, it becomes aware of the trap and, depending on the case, will seek it or try to avoid it at the next occasion. However, if it is not caught, it reverts immediately to the 'trap unaware' state. This model is best described by examining the state of the animal at the end of each recapture session (denoted t⁺ when the precise timing need be specified) and how this state changes from one session to the next (alternatively, it is possible to consider the state at the beginning of each recapture session, but this approach would cause difficulties in the treatment of censored individuals, a situation frequently encountered). The individual is actually moving back and forth in a Markovian way between the state 'trap aware' (A) -its original state when it is first released after marking- and the state 'trap unaware' (U) which follows any occasion where it is not captured. At one point, the animal may also enter the state 'dead' (†), never to leave it again. To describe the capture histories under this model, we need three kinds of parameters: survival probabilities between capture sessions (ϕ), capture probabilities of trap aware individuals (p'), and capture probabilities of trap unaware individuals (p). Several kinds of dependency may be considered on these parameters (e.g., constancy, time or age dependency or individual characteristics, etc.) but the treatment of trap-dependence remains the same. Hence, for simplicity, we present the model as if parameters were constant.

The transition matrix, $\mathbf{\Phi}_{\cdot t}$. from the state at t^{\dagger} (in line) to the state at $t+I^{\dagger}$ (in column) can be written as

But it may be useful to separate the survival process (S), which takes place between times t^+ and $t+1^-$ (i.e. the instant just before occasion t+1) from the trap awareness process (P) assumed to take place between $t+1^-$ and $t+1^+$. Below, the time is specified as an index.

$$\Phi_t = S_t P_{t+1}$$
 with

$$A_{t+1} - U_{t+1} - \dagger_{t+1} - A_{t} \begin{pmatrix} \phi & 0 & 1 - \phi \\ 0 & \phi & 1 - \phi \\ 0 & 0 & 1 \end{pmatrix} \text{ and}$$

$$S_{t} = U_{t} \begin{pmatrix} A_{t+1} + U_{t+1} + \dagger_{t+1} + A_{t+1} - V_{t+1} + U_{t+1} + \delta_{t+1} + V_{t+1} + \delta_{t+1} - V_{t+1} + V_{t+1} + \delta_{t+1} - V_{t+1} + \delta_{t+1} - V_{t+1} + \delta_{t+1} - V_{t+1} + \delta_{t+1} + \delta_{t+1} - \delta_{t+1} + \delta_{t+1$$

This model can be implemented as a multievent model [29] in program E-SURGE [30] or as a state-space model [31–32]. We detail here the first approach. Besides the transitions between states, the multievent formulation, which has a hidden Markov model structure, requires that probabilities of initial states be specified along with probabilities of the two events ('encountered', 'not encountered') conditional on the underlying state. However, initial state, assessed at the time of initial release, is necessarily 'trap aware' (A). As for the event probabilities, they are also trivial. If an animal is trap-aware at t⁺, that means that it has just been captured (conventional code '1'). If it is trap unaware or dead, it has not been captured during this session (conventional code '0'). This is summarized in the following matrix of event probabilities (E) with states in row and events in column.

$$E_{t} = \begin{array}{c} & & & & \\ & & & \\ & & & \\ &$$

Using this approach, we were able to reproduce an analysis of survival of Cory's shearwaters (*Calonectris diomedea*) in presence of temporary emigration [20]. The new multievent approach proved strictly equivalent to the old approach where capture histories had to be split after each capture (Table 1). Table 1 also shows that ignoring trap-dependence would have led to an underestimation of survival. The practical implementation in program E-SURGE of model 5 of Table 2 in Sanz-Aguilar et al. (2011) is given in Appendix S2.

For more complex situations where there are several types of observations, probabilities associated to each type of observation appear in the event matrix [29]. Appendix S2 contains such an example.

Immediate Trap Effect with several sites or states

Most often, an analysis will involve state considerations, such as the breeding status or the geographical location. We treat here the multistate version of the ITEC model where, further to being 'trap aware' or 'trap unaware', individuals support another state classification. Without loss of generality, we assume that there are only two 'other' states. When combined with 'aware' and 'unaware', this leads to 4 (live) operational states: 'aware' and 'unaware' in state 1 (A1 and U1 respectively); 'aware' and 'unaware' in state 2 (A2 and U2 respectively). To which we add the state 'dead' (†). In what follows, we reserve the term 'state' for the states of 'interest'. In addition to survival and capture

PLoS ONE | www.plosone.org

Model	$(\phi_1, \phi_2, \mathbf{p}_t)$	(φ ₁ ,φ ₂ , p _{t+m})	(ϕ_1,ϕ_2, p_{t+m})		
	no treatment	new approach	traditional approach		
	of trap dependence	(trap-awareness states)	(split capture histories)		
φ1	0.75 (0.69–0.80)	0.77 (0.70–0.82)	0.77 (0.70–0.82)		
φ ₂	0.84 (0.80–0.87)	0.87 (0.82–0.90)	0.87 (0.82–0.90)		

The current approach to modelling trap-dependence is compared to the traditional approach and to the model that ignores trap-dependence in a survival analysis of Cory's shearwaters (from [10]). Because there are transient individuals in this data set, two survival values are estimated: ϕ_1 , the apparent survival of newly-marked individuals, which is affected by the presence of transients, and ϕ_2 , the survival of previously marked individuals. Capture probability p is time-dependent-only in model (ϕ_1, ϕ_2, p_t) and time- and trap-dependent in model $(\phi_1, \phi_2, p_{t+m})$. In this last model, trap and time dependencies are additive. This model was fitted with the current approach, which considers trap-awareness states and with the traditional approach as in ([10] Model 5, Table 2), which involves the special preparation of the data detailed in [12]. The 95% confidence intervals are in parentheses.

probabilities as in the previous section, we consider now transition probabilities \boldsymbol{o} between the states of interest 1 and 2. In what follows, all parameters are explicitly shown as state-specific. Like in the single-site case, the model can be summarized through a transition matrix.

For instance, an individual which is in state 1 and is trap aware at t (operational state A1) may survive, remain in state 1, and not be caught at occasion t+1. In which case, it reverts to being trap unaware at t+1⁺. Its operational state at this moment changes to U1. The associated probability $\phi_1\psi_{11}(1-p'_1)$ is found in row 1, column 2 of matrix $\mathbf{\Phi}_t$. Now, it may be more illuminating to consider 3 steps: the survival process (S), which takes place between times t and t+1⁻, the state transition process (T), assumed to take place by the end of the interval at t+1⁻, and eventually the trap awareness process (P) assumed to take place between t+1⁻ and t+1⁺. Again, the time is specified through an index.

$$\Phi_t = S_t T_t P_{t+1}$$
 with

$$\begin{array}{rcl} & A1_{t+1} & U1_{t+1} - A2_{t+1} - U2_{t+1} - \dagger_{t+1} - \\ & A1_{t+1} - \\ & U1_{t+1} - \\ & U1_{t+1} - \\ & U2_{t+1} - \\ & \dagger_{t+1} - \end{array} \begin{pmatrix} \psi_{11} & 0 & \psi_{12} & 0 & 0 \\ 0 & \psi_{11} & 0 & \psi_{12} & 0 \\ \psi_{21} & 0 & \psi_{22} & 0 & 0 \\ 0 & \psi_{21} & 0 & \psi_{22} & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \text{ and}$$

$$\begin{aligned} \mathbf{A1}_{t+1} + \mathbf{U1}_{t+1} + \mathbf{A2}_{t+1} + \mathbf{U2}_{t+1} + \dagger_{t+1} + \\ \mathbf{A1}_{t+1} - & P_1 & 0 & 0 & 0 \\ \mathbf{U1}_{t+1} - & p_1' & 1 - p_1' & 0 & 0 & 0 \\ p_1 & 1 - p_1 & 0 & 0 & 0 \\ 0 & 0 & p_2' & 1 - p_2' & 0 \\ 0 & 0 & p_2 & 1 - p_2 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \end{aligned} \right).$$

When an individual is initially released, it is trap aware but may be in state 1 or 2. There is thus a probability π_1 that it is in state 1 (operational state A1) and the complementary probability $1 - \pi_1$ that it is in state 2 (operational state A2). This is summarized by the following vector of initial state probabilities.

A1 U1 A2 U2 †

$$\Pi_t = (\pi_1 \ 0 \ 1 - \pi_1 \ 0 \ 0)$$

Like in the single state case, the specification of event probabilities is trivial. Like in the previous section, the code will necessarily be '0' (not encountered) for trap-unaware and dead individuals. As for trapaware individuals, we assume here that their state is recognized without error: '1' for individuals in state 1, '2' for individuals in state 2.

	'0'	'1'	'2'
$A1_t$	(0	1	0)
$U1_t$	1	0	0
$E_t = A2_t$	0	0	1
$U2_t$	1	0	0
† _t	1	0	0)

For a practical implementation of this model with program E-SURGE, see Appendix S2.

Discussion

In the above models, unlike in traditional multistate capturerecapture models, capture probabilities appear among the transitions. This may be surprising to those used to the traditional models but is perfectly understandable when one realizes that, in presence of trap-dependence *stricto sensu*, the capture process does effect a change of state: after being captured, the animal knows of the trap and will adapt its behavior; the capture probability is thus a legitimate transition probability. In cases of trap-dependence lato sensu (overlap of survey area with territory, dominant individual with a conspicuous behavior, reproductive skipping, etc.), the capture event does not truly effect a change of state, but rather unveils a preexisting state (e.g. [20]). In these cases, dependence among sighting probabilities may well extend beyond one occasion, the extreme being intrinsic individual heterogeneity where the same individuals are always the highly catchable. For this last case, mixture models [27] are clearly more appropriate. One-step dependence and fixed heterogeneity represent actually two extremes of a gradient where the correlation lasts a more or less long time and may weaken progressively. With genuine trap response, this can be related to fading memory. When correlation is due to the overlap of the survey area with the individual territories, it may also be lost over time if territories and sampling protocol evolve progressively. The proposed approach could be extended to treat such cases by introducing appropriate holding times in the trap-aware state (semi-Markov process). At the moment, we recommend that in the absence of a clear understanding of the situation in a particular study where the tests for trap-dependence are significant, both immediate trap dependence and mixture models be tried. Temporary emigration

References

- Bisi F, Newey S, Nodari M, Wauters LA, Harrison A, et al. (2011) The strong and the hungry: bias in capture methods for mountain hares Lepus timidus. Wildlife Biology 17: 311–316.
- Gilbert DJ, McKenzie JR, Davies NM (2001) Evidence from tag recapture experiments that fish learn to avoid fishing gear. Journal of Agricultural Biological and Environmental Statistics 6: 281–291.
- Sanz-Aguilar A, Tavecchia G, Minguez E, Massa B, Lo Valvo F, et al. (2010) Recapture processes and biological inference in monitoring burrow-nesting seabirds. Journal of Ornithology 151: 133–146.
- Reale D, Gallant BY, Leblanc M, Festa-Bianchet M (2000) Consistency of temperament in bighorn ewes and correlates with behaviour and life history. Animal Behaviour 60: 589–597.
- Otis DL, Burnham KP, White GC, Anderson DR (1978) Statistical inference from capture data on closed animal populations. Wildlife Monographs 62: 1–135.
- Chao A, Chu WT, Hsu CH (2000) Capture-recapture when time and behavioral response affect capture probabilities. Biometrics 56: 427–433.
- Yang HC, Chao A (2005) Modeling animals' behavioral response by Markov chain models for capture-recapture experiments. Biometrics 61: 1010–1017.
- Sandland RL, Kirkwood GP (1981) Estimation of survival in marked populations with possibly dependent sighting probabilities. Biometrika 68: 531–541.
- Pradel R (1993) Flexibility in survival analysis from recapture data: handling trap-dependence. In: Lebreton JD, North PM, eds. Marked individuals in the study of bird population. Basel: Birkhauser Verlag. pp 29–37.
- Pradel R, Wintrebert CMA, Gimenez O (2003) A proposal for a goodness-of-fit test to the Arnason–Schwarz multisite capture–recapture model. Biometrics 59: 43–53.
- Pradel R, Rioux N, Tamisier A, Lebreton JD (1997) Individual turnover among wintering teal in Camargue: a mark-recapture study. J Wildl Manage 61: 816–821.
- Wormald CL, Steele MA (2008) Testing assumptions of mark-recapture theory in the coral reef fish *Lutjanus apodus*. J Fish Biol 73: 498–509.
- Kraus C, Eberle M, Kappeler PM (2008) The costs of risky male behaviour: sex differences in seasonal survival in a small sexually monomorphic primate. Proc Roy Soc Lond B 275: 1635–1644.
- Schaub M, Kania W, Koppen U (2005) Variation of primary production during winter induces synchrony in survival rates in migratory white storks *Ciconia ciconia*. J Anim Ecol 74: 656–666.
- Kaufman MJ, Frick WF, Linthicum J (2003) Estimation of habitat-specific demography and population growth for peregrine falcons in California. Ecol Appl 13: 1802–1816.
- Cam E, Oro D, Pradel R, Jimenez J (2004) Assessment of hypotheses about dispersal in a long-lived scabird using multistate capture-recapture models. J Anim Ecol 73: 723–736.

models may accommodate intermediate situations even when transitions do not correspond to geographical movements.

Supporting Information

Appendix S1 Studies investigating trap-dependence. Studies citing Pradel (1993) in which a trap-dependence effect has been found (research on ISI Web of Knowledge). (DOCX)

Appendix S2 Practical implementation of multievent trap-dependence models with program E-SURGE: a medium-term monitoring program on Cory's Shearwater (*Calonectris diomedea*) as a case example. (DOCX)

Acknowledgments

We thank the Population Ecology Group (IMEDEA, Esporles, Spain) for making the Cory's shearwater data available to us.

Author Contributions

Conceived and designed the experiments: RP AS. Analyzed the data: AS RP. Wrote the paper: RP AS.

- Peron G, Crochet PA, Choquet R, Pradel R, Lebreton JD, et al. (2010) Capturerecapture models with heterogeneity to study survival senescence in the wild. Oikos 119: 524–532.
- Crespin L, Choquet R, Lima M, Merritt J, Pradel R (2008) Is heterogeneity of catchability in capture-recapture studies a mere sampling artefact or a relevant feature of the population? Pop Ecol 50: 247–256.
- Cubaynes S, Pradel R, Choquet R, Duchamp C, Gaillard JM, et al. (2010) Importance of accounting for detection heterogeneity when estimating abundance: the case of french wolves. Cons Biol 24: 621–626.
- Sanz-Aguilar A, Tavecchia G, Genovart M, Igual JM, Oro D, et al. (2011) Studying the reproductive skipping behavior in long-lived birds by adding nestinspection to individual-based data. Ecol Appl 21: 555–564.
- Schmidt BR, Schaub M, Anholt BR (2002) Why you should use capturerecapture methods when estimating survival and breeding probabilities: on bias, temporary emigration, overdispersion, and common toads. Amph-Rept 23: 375–388.
- Sendor T, Simon M (2003) Population dynamics of the pipistrelle bat: effects of sex, age and winter weather on seasonal survival. J Anim Ecol 72: 308–320.
- Choquet R, Lebreton JD, Gimenez O, Reboulet AM, Pradel R (2009) U-CARE: Utilities for performing goodness of fit tests and manipulating CApture-REcapture data. Ecography 32: 1071–1074.
- Sandvik H, Erikstad KE, Barrett RT, Yoccoz NG (2005) The effect of climate on adult survival in five species of North Atlantic seabirds. Journal of Animal Ecology 74: 817–831.
- Pledger S, Pollock KH, Norris JL (2003) Open capture-recapture models with heterogeneity: I. Cormack-Jolly-Seber model. Biometrics 59: 786–794.
- Fujiwara M, Caswell H (2002) Estimating population projection matrices from multi-stage mark-recapture data. Ecology 83: 3257–3265.
- Huggins RM (1989) On the statistical analysis of capture experiments. Biometrika 76: 133–140.
- Ratcliffe N, Newton S, Morrison P, Merne O, Cadwallender T, et al. (2008) Adult Survival and Breeding Dispersal of Roseate Terns Within the Northwest European Metapopulation. Waterbirds 31: 320–329.
- 29. Pradel R (2005) Multievent: An extension of multistate capture-recapture models to uncertain states. Biometrics 61:442-447.
- Choquet R, Rouan L, Pradel R (2009) Program E–SURGE: a software application for fitting multievent models. In: Thomson DL, Cooch EG, Conroy MJ, eds. Modeling Demographic Processes in Marked Populations. Berlin: Springer. pp 845–865.
- Gimenez O, Rossi V, Choquet R, Dehais C, Doris B, et al. (2007) State-space modelling of data on marked individuals. Ecol Model 206: 431–438.
- Royle JA (2008) Modeling individual effects in the Cormack-Jolly-Seber Model: A state-space formulation. Biometrics 64: 364–370.

Appendix S1

Studies investigating trap-dependence

Studies citing Pradel (1993) in which a trap-dependence effect has been found (research on ISI Web of Knowledge).

Species group	Number of studies treating	Number of studies not
	trap-dependence	treating trap-dependence
Birds	52 (83.87%)	10 (16.13%)
Mammals	18 (82.82%)	4 (18.18%)
Reptiles	2 (40%)	3 (60%)
Amphibians	2 (100%)	0
Fish	1 (50%)	1 (50%)
Insects	1 (100%)	0
TOTAL	76 (80.85 %)	18 (19.14 %)

Bird studies correcting for trap-dependence

- Bachler E, Schaub M (2007) The effects of permanent local emigration and encounter technique on stopover duration estimates as revealed by telemetry and mark-recapture. Condor 109: 142-154.
- Balanca G, Schaub M (2005) Post-breeding migration ecology of Reed Acrocephalus scirpaceus, Moustached A-melanopogon and Cetti's Warblers *Cettia cetti* at a Mediterranean stopover site. Ardea 93: 245-257.
- Barbraud C, Weimerskirch H (2003) Climate and density shape population dynamics of a marine top predator. Proc Roy Soc Lond B 270: 2111-2116.
- Barbraud C, Weimerskirch H, Guinet C, Jouventin P (2000) Effect of sea-ice extent on adult survival of an Antarctic top predator: the snow petrel *Pagodroma nivea*. Oecol 125: 483-488.
- Belda EJ, Barba E, Monros JS (2007) Resident and transient dynamics, site fidelity and survival in wintering Blackcaps *Sylvia atricapilla*: evidence from capture-recapture analyses. Ibis 149: 396-404.
- Bokony V, Liker A, Lendvai AZ, Kulcsár A (2008) Risk-taking and survival in the House Sparrow *Passer domesticus*: are plumage ornaments costly? Ibis 150: 139-151.
- Cam E, Oro D, Pradel R, Jimenez J (2004) Assessment of hypotheses about dispersal in a long-lived seabird using multistate capture-recapture models. J Anim Ecol 73: 723-736.
- Clobert J (1995) Capture-recapture and evolutionary ecology: A difficult wedding? J Appl Stat 22: 989-1008.

- Clucas RJ, Fletcher DJ, Moller H (2008) Estimates of adult survival rate for three colonies of Sooty Shearwater (*Puffinus griseus*) in New Zealand. Emu 108: 237-250.
- Crespin L, Harris MP, Lebreton JD, Wanless S (2006) Increased adult mortality and reduced breeding success with age in a population of common guillemot *Uria aalge* using marked birds of unknown age. J Avian Biol 37: 273-282.
- Erikstad KE, Sandvik H, Fauchald P, Tveraa T (2009) Short- and long-term consequences of reproductive decisions: an experimental study in the puffin. Ecology 90: 3197-3208.
- Evans RJ, Wilson JD, Amar A, Douse A, Maclennan A, et al. (2009) Growth and demography of a re-introduced population of White-tailed Eagles *Haliaeetus albicilla*. Ibis 151: 244-254.
- Forero MG, Tella JL, Oro D (2001) Annual survival rates of adult Red-necked Nightjars *Caprimulgus ruficollis*. Ibis 143: 273-277.
- Frederiksen M, Wanless S, Harris MP, Rothery P, Wilson LJ (2004) The role of industrial fisheries and oceanographic change in the decline of North Sea black-legged kittiwakes. J Appl Ecol 41: 1129-1139.
- Gauthier, G, Pradel R, Menu S, Lebreton JD (2001) Seasonal survival of Greater Snow Geese and effect of hunting under dependence in sighting probability. Ecology 82: 3105-3119.
- Grosbois V, Harris MP, Anker-Nilssen T, McCleery RH, Shaw DN, et al. (2009) Modeling survival at multi-population scales using mark-recapture data. Ecology 90: 2922-2932.
- Grosbois V, Thompson PM (2005) North Atlantic climate variation influences survival in adult fulmars. Oikos 109: 273-290.
- Hario M, Mazerolle MJ, Saurola PS (2009) Survival of female common eiders *Somateria m. mollissima* in a declining population of the northern Baltic Sea. Oecol 159: 747-756.
- Harris, M. P, Anker-Nilssen T, Mccleery RH, Erikstad KE, Shaw DN, et al. (2005) Effect of wintering area and climate on the survival of adult Atlantic puffins *Fratercula arctica* in the eastern Atlantic. Mar Ecol Progr Ser 297: 283-296.
- Harris MP, Freeman SN, Wanless S, Morgan BJT, Wernham CV (1997) Factors influencing the survival of Puffins *Fratercula arctica* at a North Sea colony over a 20-year period. J Avian Biol 28: 287-295.
- Jenouvrier S, Barbraud C, Weimerskirch H (2003) Effects of climate variability on the temporal population dynamics of southern fulmars. J Anim Ecol 72: 576-587.
- Jenouvrier S, Barbraud C, Weimerskirch H (2005) Long-term contrasted responses to climate of two Antarctic seabird species. Ecology 86: 2889-2903.
- Jenouvrier S, Thibault J C, Viallefont A, Vidals P, Ristow D et al. (2009) Global climate patterns explain range-wide synchronicity in survival of a migratory seabird. Glob Ch Biol 15: 268-279.

- Kaiser A (1995) Estimating turnover, movements and capture parameters of resting passerines in standardized capture-recapture studies. J Appl Stat 22: 1039-1047.
- Kauffman MJ, Frick WF, Linthicum J (2003) Estimation of habitat-specific demography and population growth for peregrine falcons in California. Ecol Appl 13: 1802-1816.
- Kendall WL, Nichols JD (1995) On the use of secondary capture-recapture samples to estimate temporary emigration and breeding proportions. J Appl Stat 22: 751-762.
- Kery M, Madsen J, Lebreton JD (2006) Survival of Svalbard pink-footed geese Anser brachyrhynchus in relation to winter climate, density and land-use. J Anim Ecol 75: 1172-1181.
- Madsen J, Frederiksen M, Ganter B (2002) Trends in annual and seasonal survival of Pinkfooted Geese *Anser brachyrhynchus*. Ibis 144: 218-226.
- Monticelli D, Ramos JA, Guerreiro-Milheiras SA, Doucet JL (2008) Adult survival of Tropical Roseate Terns breeding on Aride Island, Seychelles, Western Indian Ocean. Waterbirds 31: 330-337.
- Nevoux M, Barbraud C (2006) Relationships between sea ice concentration, sea surface temperature and demographic traits of thin-billed prions. Polar Biol 29: 445-453.
- Nevoux M, Weimerskirch H, Barbraud C (2010) Long- and short-term influence of environment on recruitment in a species with highly delayed maturity. Oecol 162: 383-392.
- Oro D, De Leon A, Minguez E, Furness RW (2005) Estimating predation on breeding European storm-petrels (*Hydrobates pelagicus*) by yellow-legged gulls (*Larus Michahellis*). J Zool 265: 421-429.
- Peron G, Crochet PA, Choquet R, Pradel R, Lebreton JD, et al. (2010) Capture-recapture models with heterogeneity to study survival senescence in the wild. Oikos 119: 524-532.
- Pons J, Migot M (1995) Life-history strategy of The Herring Gull changes in survival and fecundity in a population subjected to various feeding conditions. J Anim Ecol 64: 592-599.
- Pradel R, Rioux N, Tamisier A, Lebreton JD (1997) Individual turnover among wintering teal in Camargue: a mark-recapture study. J Wildl Manage 61: 816-821.
- Pugesek BH, Nations C, Diem KL, Pradel R (1995) Mark-resighting analysis of a California gull population. J Appl Stat 22: 625-639.
- Ratcliffe N, Newton S, Morrison P, Merne O, Cadwallender T, et al. (2008) Adult survival and breeding dispersal of Roseate Terns within the Northwest European metapopulation. Waterbirds 31: 320-329.
- Rolland V, Barbraud C, Weimerskirch H (2008) Combined effects of fisheries and climate on a migratory long-lived marine predator. J Appl Ecol 45: 4-13.

- Rolland V, Barbraud C, Weimerskirch H (2009) Assessing the impact of fisheries, climate and disease on the dynamics of the Indian yellow-nosed Albatross. Biol Cons 142: 1084-1095.
- Rolland V, Nevoux M, Barbraud C, Weimerskirch H (2009) Respective impact of climate and fisheries on the growth of an albatross population. Ecol App 19: 1336-1346.
- Sandvik, H, Erikstad K E, Barrett R T, Yoccoz N G (2005) The effect of climate on adult survival in five species of North Atlantic seabirds. J Anim Ecol 74: 817-831.
- Sandvik H, Erikstad KE, Fauchald P, Tveraa T (2008) High survival of immatures in a longlived seabird: Insights from a long-term study of the Atlantic Puffin (*Fratercula arctica*). Auk 125: 723-730.
- Sanz-Aguilar A, Tavecchia G, Mínguez E, Massa B, Lo Valvo F, et al. (2010) Recapture processes and biological inference in monitoring burrowing nesting seabirds. J Ornithol 151: 133-146.
- Sanz-Aguilar A, Tavecchia G, Genovart M, Igual JM, Oro D, et al. (2011) Studying the reproductive skipping behavior in long-lived birds by adding nest-inspection to individual-based data. Ecol Appl 21: 555-564.
- Schaub M, Kania W, Koppen U (2005) Variation of primary production during winter induces synchrony in survival rates in migratory white storks *Ciconia ciconia*. J Anim Ecol 74: 656-666.
- Spendelow JA, Nichols JD, Hines JE, Lebreton JD, Pradel R (2002) Modelling postfledging survival and age-specific c breeding probabilities in species with delayed maturity: a case study of Roseate Terns at Falkner Island, Connecticut. J Appl Stat, 29: 385-405.
- Szep T (1999) Effects of age- and sex-biased dispersal on the estimation of survival rates of the Sand Martin *Riparia riparia* population in Hungary. Bird Stu 46: 169-177.
- Tavecchia G, Minguez E, De León A, Louzao M, Oro D (2008) Living close, doing differently: Small-scale asynchrony in demography of two species of seabirds. Ecology 89: 77-85.
- Tavecchia G, Viedma C, Martínez-Abraín A, Bartolomé MA, Gómez JA, et al. (2009) Maximizing re-introduction success: Assessing the immediate cost of release in a threatened waterfowl. Biological Conservation, 142, 3005-3012.
- Viallefont A, Cooch EG, Cooke F (1995) Estimation of trade-offs with capture-recapture models: A case study on the lesser snow goose. J Appl Stat 22: 847-861.
- Viallefont A, Cooke F, Lebreton JD (1995) Age-specific costs of first-time breeding. Auk 112: 67-76.
- Votier SC, Hatchwell BJ, Beckerman A, McCleery RH, Hunter FM, et al. (2005) Oil pollution and climate have wide-scale impacts on seabird demographics. Ecol Lett 8: 1157-1164.
- Bird studies not correcting for trap-dependence

- Bearhop S, Ward RM, Evans PR (2003) Long-term survival rates in colour-ringed shorebirds - practical considerations in the application of mark-recapture models. Bird Stu 50: 271-279.
- Cooch, E. G, Blank DB, Rockwell RF, Cooke F (1999) Body size and age of recruitment in Snow Geese *Anser c. caerulescens*. Bird Stu 46: 112-119.
- Dugger, K. M, Ainley DG, Lyver POB, Barton K, Ballardef G (2010) Survival differences and the effect of environmental instability on breeding dispersal in an Adelie penguin meta-population. Proc Natl Acad Sci USA 107: 12375-12380.
- Faustino, C.R, Jennelle CS, Connolly V, Davis AK, Swarthout EC, et al. (2004) *Mycoplasma gallisepticum* infection dynamics in a house finch population: seasonal variation in survival, encounter and transmission rate. J Anim Ecol 73: 651-669.
- Frederiksen M, Bregnballe T (2000) Evidence for density-dependent survival in adult cormorants from a combined analysis of recoveries and resightings. J Anim Ecol 69: 737-752.
- Frederiksen M, Bregnballe T (2001) Conspecific reproductive success affects age of recruitment in a great cormorant, *Phalacrocorax carbo sinensis*, colony. Proc Roy Soc Lond B 268: 1519-1526.
- Lebreton, J. D, Hines JE, Pradel R, Nichols JD, Spendelow JA (2003) Estimation by capturerecapture of recruitment and dispersal over several sites. Oikos 101: 253-264.
- Peach WJ, Hanmer DB, Oatley TB (2001) Do southern African songbirds live longer than their European counterparts? Oikos 93: 235-249.
- Schaub M, Jenni L (2001) Stopover durations of three warbler species along their autumn migration route. Oecol 128: 217-227.
- Votier SC, Birkhead TR, Oro D, Trinder M, Grantham MJ, et al. (2008) Recruitment and survival of immature seabirds in relation to oil spills and climate variability. J Anim Ecol 77: 974-983.

Mammal studies correcting for trap-dependence

- Crespin L, Choquet R, Lima M, Merritt J, Pradel R (2008) Is heterogeneity of catchability in capture-recapture studies a mere sampling artefact or a relevant feature of the population? Pop Ecol 50: 247-256.
- Crespin L, Papillon Y, Abdoulaye D, Granjon L, Sicard B (2008) Annual flooding, survival and recruitment in a rodent population from the Niger River plain in Mali. J Trop Ecol 24: 375-386.
- Guitton JS, Devillard S, Guénézan M, Fouchet D, Pontier D, et al. (2008) Vaccination of free-living juvenile wild rabbits (*Oryctolagus cuniculus*) against myxomatosis improved their survival. Prev Vet Med 84: 1-10.
- Hoyle SD, Pople AR, Toop GJ (2001) Mark-recapture may reveal more about ecology than about population trends: Demography of a threatened ghost bat (*Macroderma gigas*)

population. Austral Ecol 26: 80-92.

- Johannesen E, Aars J, Andreassen HP, Ims RA (2003) A demographic analysis of vole population responses to fragmentation and destruction of habitat. Pop Ecol 45: 47-58.
- Julliard R, Leirs H, Stenseth NC, Yoccoz NG, Prevot-Julliard AC, et al. (1999) Survivalvariation within and between functional categories of the African multimammate rat. J Anim Ecol 68: 550-561.
- 1Kraus C, Eberle M, Kappeler PM (2008) The costs of risky male behaviour: sex differences in seasonal survival in a small sexually monomorphic primate. Proc Roy Soc Lond B 275: 1635-1644.
- Langtimm CA, O'Shea TJ, Pradel R, Beck CA (1998) Estimates of annual survival probabilities for adult Florida manatees (*Trichechus manatus latirostris*). Ecology 79: 981-997.
- Largo E, Gaillard JM, Festa-Bianchet M Toïgo C, Bassano B, et al. (2008) Can ground counts reliably monitor ibex *Capra ibex* populations? Wildl Biol 14: 489-499.
- Letty J, Ubineau JA, Marchandeau S, Lobert JC (2003) Effect of translocation on survival in wild rabbit (*Oryctolagus cuniculus*). Mammal Biol 68: 250-255.
- Lima M, Merritt JF, Bozinovic F (2002) Numerical fluctuations in the northern short-tailed shrew: evidence of non-linear feedback signatures on population dynamics and demography. J Anim Ecol 71: 159-172.
- Pendleton GW, Pitcher KW, Fritz L, Raum-Suryan KL, Loughlin T, et al. (2006) Survival of Steller sea lions in Alaska: a comparison of increasing and decreasing populations. Cn J Zool 84: 1163-1172.
- Pradel R, Choquet R, Lima MA, Merritt J, Crespin L (2010) Estimating population growth rate from capture-recapture data in presence of capture heterogeneity. J Agrt Biol Environ Stat 15: 248-258.
- Ramp, C. Bérubé M, Hagen W, Sears R (2006) Survival of adult blue whales *Balaenoptera musculus* in the Gulf of St. Lawrence, Canada. Mar Ecol Progr Ser 319: 287-295.
- Ramp C, Bérubé M, Palsbøll P, Hagen W, Sears R (2010) Sex-specific survival in the humpback whale *Megaptera novaeangliae* in the Gulf of St. Lawrence, Canada. Mar Ecol Progr Ser 400: 267-276.
- Regehr EV, Regehr EV, Lunn N, Amstrup SC, Stirling I (2007) Effects of earlier sea ice breakup on survival and population size of polar bears in western Hudson bay. J Wildl Manage 71: 2673-2683.
- Sendor T, Simon M (2003) Population dynamics of the pipistrelle bat: effects of sex, age and winter weather on seasonal survival. J Anim Ecol 72: 308-320.
- Telfer S, Bennett M, Bown K, Cavanagh R, Crespin Laurent, et al. (2002) The effects of cowpox virus on survival in natural rodent populations: increases and decreases. J Anim Ecol 71: 558-568.

Mammal studies not correcting for trap-dependence

- Beauplet G, Barbraud C, Dabin W, Küssener C, Guinet C (2006) Age-specific survival and reproductive performances in fur seals: evidence of senescence and individual quality. Oikos 112/430-441.
- Graham IM, Lambin X (2002) The impact of weasel predation on cyclic field-vole survival: the specialist predator hypothesis contradicted. J Anim Ecol 71: 946-956.
- Schwarz CJ, Stobo WT (2000) Estimation of juvenile survival, adult survival, and agespecific pupping probabilities for the female grey seal (*Halichoerus gryprus*) on Sable Island from capture-recapture data. Can J Fish Aqu Sci 57: 247-253.
- Sluydts V, Crespin L, Davis S, Lima M, Leirs H (2007) Survival and maturation rates of the African rodent, Mastomys natalensis: density-dependence and rainfall. Integr Zool 2/ 220-232.

Reptile studies correcting for trap-dependence

- Rivalan P, Pradel R, Choquet R, Girondot M, Prévot-Julliard AC (2006) Estimating clutch frequency in the sea turtle Dermochelys coriacea using stopover duration. Mar Ecol Progr Ser 317: 285-295.
- Rivalan P, Prévot-Julliard AC, Choquet R, Pradel R, Jacquemin B, et al. (2005) Trade-off between current reproductive effort and delay to next reproduction in the leatherback sea turtle. Oecol 145: 564-574.
- Reptile studies not correcting for trap-dependence
- Bjorndal KA, Bolten AB, Chaloupka MY (2003) Survival probability estimates for immature green turtles *Chelonia mydas* in the Bahamas. Mar Ecol Progr Ser 252: 273-281.
- Bjorndal KA, Bolten AB, Chaloupka MY (2005) Evaluating trends in abundance of immature green turtles, *Chelonia mydas*, in the Greater Caribbean. Ecol Appl 15: 304-314.
- Chaloupka MY, Limpus CJ (2002) Survival probability estimates for the endangered loggerhead sea turtle resident in southern Great Barrier Reef waters. Mar Biol 140: 267-277.

Amphibian studies correcting for trap-dependence

- Fretey T, Cam E, Le Garff B, Monnat JY (2004) Adult survival and temporary emigration in the common toad. Can J Zool 82: 859-872.
- Schmidt BR, Schaub M, Anholt BR (2002) Why you should use capture-recapture methods when estimating survival and breeding probabilities: on bias, temporary emigration, overdispersion, and common toads. Amph-Rept 23: 375-388.

Fish studies correcting for trap-dependence

Wormald CL, Steele MA (2008) Testing assumptions of mark-recapture theory in the coral reef fish *Lutjanus apodus*. J Fish Biol 73: 498-509.

Fish studies not correcting for trap-dependence

Holmberg J, Norman B, Arzoumanian Z (2008) Robust, comparable population metrics through collaborative photo-monitoring of whale sharks *Rhincodon Typus*. Ecol Appl 18: 222-233.

Insect studies correcting for trap-dependence

Stoks R (2001) What causes male-biased sex ratios in mature damselfly populations? Ecol Entomol 26: 188-197.

Appendix S2

Practical implementation of multievent trap-dependence models with program <u>E-</u> <u>SURGE</u>: a medium-term monitoring program on Cory's Shearwater (*Calonectris diomedea*) as a case example.

The Cory's shearwater (*Calonectris diomedea*) is a long-lived burrow nesting seabird. Skipping reproduction is typical among them and usually confounded with a recapture failure. Because skippers in one year have a higher probability of skipping the following year, trapdependence (*lato sensu*) is usually found when analyzing Cory's Shearwater capture histories. Sanz-Aguilar et al. (2011) combined individual-based information with nest-based information collected as part of the Cory's Shearwater monitoring program of the <u>Population</u> <u>Ecology Group</u> (IMEDEA, Esporles, Spain) carried out at Pantaleu Island (Balearic Islands, Spain). Using multievent models, they estimated simultaneously recapture, survival, reproductive skipping and within-colony breeding dispersal probabilities. Here, for the sake of illustrating how to treat trap-dependence in different contexts, we consider simpler and more typical situations which use only part of the available information.

- I) We start with the situation described in the first section of the methods, i.e. we keep only the information about whether the individual is 'seen' (code 1) or 'not seen' (code 0), and we show how to fit the single state model with trap-dependence allowing the estimation of survival and capture probabilities.
- II) We next illustrate the second section of the methods by considering 2 observable states: the bird occupies its 'last known burrow' (code 1) or the bird occupies a 'new burrow' (code 2). We show how to fit a multistate model with trap-dependence allowing the estimation of survival, capture probabilities and the probability of burrow change.

III) Finally, we consider a situation with two ambiguous events relative to the underlying state: the classical event 'not seen with no further information' (code 0) and the more informative but still imperfect event 'not seen but known to be absent from its previous burrow' (previous burrow is empty or occupied by others) (code 2). Code 1 is kept as usual for the event 'seen'. We show how to fit this pure multievent model with trap dependence to estimate survival, capture probabilities and the probability of knowing that the bird is absent from its previous burrow. This model has no more interest than showing the basics of how trap-dependence and ambiguous events can be combined. For a fuller exploitation of burrow information, we refer the reader to Sanz-Aguilar et al. (2011).

Specification of the multievent modelling approach in program E-SURGE (extracted from Sanz-Aguilar et al. 2011)

Multievent models are built in several stages using program E-SURGE (Choquet et al. 2009). Each step represents one type of the different parameters to estimate. This is done by means of row-stochastic matrices, i.e. each row corresponds to a multinomial. Consequently, the total of cell probabilities in the same row is 1. Because of this constraint, one and only one cell probability in each row will be calculated as the complement to 1 of the others. This particular cell is denoted with a '*' symbol. Inactive cells, i.e. cells whose associated probability is structurally 0 are denoted with a '-' symbol. An active cell receives an arbitrary letter. Note that the same letter in two cells *does not mean* that the two values should be equal.

Case I: Estimating survival and recapture probabilities.

The individual states considered are:

A, "trap-aware"

U, "trap-unaware"

D, dead

The possible events are:

0, not recaptured

1, captured or recaptured

The symbols for parameters are:

φ, survival probability

p, capture probability

Initial State probabilities

A	U
*	-

Transition probabilities, step 1: Survival

	A_{t+1}	U_{t+1}	D_{t+1}
A_t	φ	-	*
U_t	-	ф	*

	Transition	probabilities,	step	o 2:	Trap	awareness	process
--	------------	----------------	------	------	------	-----------	---------

	A _{t+1}	<i>U</i> _{<i>t</i>+1}	D _{t+1}
A_{t+1}	р	*	-
U_{t+1}	р	*	-
D_{t+1}	-	-	*

Event probabilities:

	0	1
A_t	-	*
\boldsymbol{U}_{t}	*	-
\boldsymbol{D}_{t}	*	-

Detailed example of fitting model 5, Table 2 of Sanz-aguilar et al. (2011) with program E-SURGE

After reading the data into program E-SURGE, the number of states is changed to 3 and the number of age-classes is changed to 2 (2 age-classes are needed to account for the presence of transients in this data set, see Pradel et al. 1997).



Then, enter the GEPAT interface to specify the patterns as we have seen above: first, for the initial state probabilities,

4 Gepat interface									
Input-Output for patterns Parameters Pre-defined Patterns									
Diagonal Matrix Empty Matrix Full Matrix	Initial state Number of steps Current step & Label	1	IS >>						
	Matrix Pattern								
->1	1 2								
			# of Rows 1						
			# of Columns 2						
			Update Now						
	EXIT								



then for the transitions for which 2 steps must be specified.



and eventually for the events.



After that, click EXIT and, back on the main window, GEMACO to enter the GEMACO interface where effects are specified on each type of parameter in turn.

For the initial state probability, there is no active parameter. The keyword 'i' will do.



The first step of transitions corresponds to survival probabilities which depend only on age: keyword 'a'.

🚺 Gema	ico interface					_ D X		
Input-Ou	Input-Output for constraint matrix Parameters Gemaco							
	Transition							
	Î	Current st	ep & Label	<< 1	Т	>>		
	Constra	int matrix		SHO	ORTCUTS List			
		^	1	not yet defined	not ye	t defined		
				<	0 >			
				Mo	odel definition			
		-			а	*		
	Transitio	ns pattern	L					
	ψ-* -ψ* *	Í				.		
			No	external variables		-		
	•			File selected for ex	ternal variables :	defaultfile		
		c	Call Gen	naco		EXIT		

The second step involves capture probabilities, which in this model depend on the trapawareness status and on time, the two effects being additive: phrase 'f+t'. ('f' short for 'from' means that there is a row effect, which here is the trap-awareness status effect).

E	🤼 Gema	co inte	rface								-	X	
	Input-Output for constraint matrix Parameters Gemaco												
				Transition									
				Currer	nt step (& Label	<<	2	Т	-	>>		
			Constrai	nt matrix				SHO	RTCUTS	List			
					^	1	not yet de	fined	nc	ot yet (defined		
								<	0	>			
								Mod	del definit	ion			
					Ŧ				f+t			^	
			Transitio	ns pattern									
		π*- π*- *			•							•	
		•			+ +	No	external va File selecte	riables ed for exte	ernal varial	bles: de	efaultfile		•
					Cal	ll Gen	naco				EXIT		



Event probabilities are not active. The keyword 'i' will do the job.

After clicking EXIT, then back in the main windows, IVFV and EXIT again, you can run the model by clicking RUN.

Case II: Estimating survival, nest dispersal and recapture probabilities.

The individual states considered are:

- A1, "trap-aware" and breeding in the same nest as in the previous year
- U1, "trap-unaware" and breeding in the same nest as in the previous year
- A2, "trap-aware" and breeding in another nest
- U2, "trap-unaware" and breeding in another nest
- D, dead

The possible events are:

- 0, not recaptured
- 1, captured for the first time or recaptured breeding in the last known nest
- 2, recaptured breeding in a different nest

The symbols for parameters are:

- ϕ , survival probability
- ψ , nest dispersal probability, conditional on survival
- **p**, capture probability

Initial State probabilities

A1	U1	A2	U2
π	-	*	-

	A1 _{t+1}	<i>U1</i> _{t+1}	$A2_{t+1}$	$U2_{t+1}$	D_{t+1}
A1 _t	¢	-	-	-	*
$U1_t$	-	ф	-	-	*
A2 t	-	-	ø	-	*
$U2_t$	-	-	-	ф	*
\boldsymbol{D}_t	-	-	-	-	*

Transition probabilities, step 1: Survival

Transition probabilities, step 2: Nest dispersal

	$A1_{t+1}$	$U1_{t+1}$	$A2_{t+1}$	$U2_{t+1}$	D_{t+1}
	ale				
AI_{t+1}	*	-	Ψ	-	-
<i>U1</i> _{t+1} ⁻	-	*	-	Ψ	-
A2 t+1	*	-	ψ	-	-
$U2_{t+1}$	-	*	-	Ψ	-
D_{t+1}	-	-	-	-	*

Transition probabilities, step 3: Trap awareness process

	A1 t+1	U1 t+1	A2 t+1	$U2_{t+1}$	D _{t+1}
A.T		*			
AI $t+1$	р	*	-	-	-
$U1_{t+1}$	р	*	-	-	-
A2 t+1	-	-	р	*	-
$U2_{t+1}$	-	-	р	*	-
D_{t+1}	-	-	-	-	*

Event probabilities:

	0	1	2
$A1_t$	-	*	-
U1 _t	*	-	-
A2 t	-	-	*
U2 t	*	-	-
D_t	*	-	-

Detailed example of fitting a multistate model with trap-dependence with program E-SURGE

After reading the data into program E-SURGE, the number of states is changed to 5 and the number of age-classes to 2 (this is to deal with transients). Then we go through GEPAT to specify the patterns as indicated above.









In GEMACO, we specify that the initial state probabilities will be constant using the keyword 'i'. This will suffice because all Cory's shearwaters are arbitrarily assumed to be in the same nest as the year before when first encountered, i.e. initial state A1. (Alternatively, for this particular data set, we could have entered (* - -) in GEPAT)





Survival as above is assumed to depend on age to account for the presence of transients.

Burrow transition is assumed independent of a previous transition, trap-awareness, etc.: keyword 'i'.

Gemaco interface					
Input-Output for constraint matrix Parameters Gemaco					
Transition					
Current step & La	^{abel} << 2 T >>				
Constraint matrix	SHORTCUTS List				
	not yet defined not yet defined				
	< 0 >				
	Model definition				
	i *				
Transitions pattern					
-ψ --ψ- *-ψ					
*					
	No external variables				
	File selected for external variables : defaultfile				
Call G	Semaco EXIT				

Capture probability depends as before on time and trap-awareness status additively. Because trap-aware individuals are those in the operative states A1 (row 1) and A2 (row 3), the phrase is 't+f(1 3)'.



🛃 Gemaco interface						
Input-Output for constraint matrix Parameters Ge	Input-Output for constraint matrix Parameters Gemaco					
EV	/ENT					
Current step & La	^{abel} << 1 ^C >>					
Constraint matrix	SHORTCUTS List					
^	not yet defined not yet defined					
	< 0 >					
	Model definition					
	i ^					
•						
Transitions pattern						
*- *- *						
*						
*	•					
4	No external variables					
Call G	Semaco EXIT					

There is no active event probability. This step is as above.

The rest is pretty much unchanged, except that in IVFV it may help to fix the initial state probability to 1 as we know that all individuals start in state A1.

🚺 Initial Valu	🥂 Initial Value or Fixed Value for beta (IVFV).					
FILE Selected	FILE Selected Beta Value Space					
			Initial stat	e	[0,1]	
Coordinates of the const F=Departure To=Arrival T=Time A=Age G=Group S=Step	beta value raint matrix ;	es are with non with	zero elements			
		FTOTAC	⇒ s	Fixed Value	Initial Value ? of Beta	
	Beta #1	1111		Fix. or	✓ 1	
E	Beta # 2		*			
E	Beta # 3		÷.			
E	Beta # 4		*			
E	Beta # 5		<u></u>			
E	Beta # 6		÷.			
E	Beta # 7		÷			
E	Beta # 8		÷.			
E	Beta # 9		\$			
В	eta # 10		^			
~	< Previo	us	EXIT	Next >>	Last >	

Case III: Estimating survival and recapture probabilities, and the probability of knowing that a bird is absent from its previous burrow.

The individual states considered are:

A, "trap-aware"

U, "trap-unaware"

D, dead

The possible events are:

0, not recaptured without further information

1, captured or recaptured

2, not recaptured and known to be absent from its previous burrow (previous burrow is found

empty or occupied by others)

The symbols for parameters are:

φ, survival probability

- **p**, capture probability
- $\boldsymbol{\epsilon}$, probability of knowledge of absence from previous burrow

Initial State probabilities

A	U
*	-

Transition probabilities, step 1: Survival

	A_{t+1}	U_{t+1}	D_{t+1}
A_t	ф	-	*
U_t	-	¢	*

Transition probabilities, step 2: Trap awareness process

	A _{t+1}	U_{t+1}	D _{t+1}
A_{t+1}	р	*	-
U_{t+1}	р	*	-
D_{t+1}	-	-	*

Event probabilities:

	0	1	2
A_t	-	*	-
U_t	*	-	3
D_t	*	-	3

Detailed example of fitting a pure multievent model with trap-dependence with program E-SURGE

After reading the data into program E-SURGE, the number of states is changed to 3 and the number of age-classes to 2 (this last point is to account for the presence of transients). Then we go through GEPAT to specify the patterns as indicated above.









In GEMACO, as there is no active parameter in the initial state probabilities, we can use the keyword 'i'.

🚺 Gemaco interface					
Input-Output for constraint matrix Parameters Gemaco					
Init s	state				
Current step & Lab	^{el} << 1 IS >>				
Constraint matrix	SHORTCUTS List				
not defined	not yet defined not yet defined				
	< 0 >				
	Model definition				
	i ^				
Transitions pattern					
* <u>-</u>					
▼ N	lo external variables 🔹				
	File selected for external variables : defaultfile				
Call Ge	EXIT				



For the survival probabilities, we need age to account for transients.

For capture probabilities, we specify an additive effect of time and trap-awareness status. Here, trap-aware individuals are those in state A1, the first state.

Gemaco interface								
Input-Output for constraint matrix Parameters Gemaco								
Transition								
Current step) & Label	<< 2	Т	>>		
	Constraint matrix			SHORTCUTS List				
		•	r	not yet defined	not yet	: defined		
				<	0 >			
				Mod	el definition			
		Ŧ		t·	+f(1)	*		
	Transitio	ns pattern						
	π*- π*- *	^				Ţ		
		-	No	external variables				
	4	► F		File selected for exte	rnal variables :	defaultfile		
		Cá	all Gen	naco		EXIT		

The event probabilities step is where a new parameter appears: the probability of knowing that an individual is absent from its previous burrow. We assume that this probability is unique over time, etc.: keyword 'i'.

	Gemaco interface								
Input-Output for constraint matrix Parameters Gemaco									
	EV	'ENT							
	Current step & La	^{bel} << 1 C >>							
	Constraint matrix	SHORTCUTS List							
		not yet defined not yet defined							
		< 0 >							
		Model definition							
	-	i ^							
	Transitions pattern								
	- * - ^ ^								
	* - v								
		No external variables 🗸							
	4 ▶	File selected for external variables : defaultfile							
	Call G	emaco							

References:

Choquet, R., L. Rouan, and R. Pradel. 2009. Program E–SURGE: a software application for fitting multievent models. Pages 845–865 *in* D. L. Thomson, E. G. Cooch, and M. J. Conroy, editors. Modeling Demographic Processes in Marked Populations. Springer, Berlin, Germany.

- Pradel, R., J.E. Hines, J.D. Lebreton, and J.D. Nichols. 1997. Capture-recapture survival models taking account of transients. Biometrics **53**:60–72.
- Sanz-Aguilar, A., G. Tavecchia, M. Genovart, J. M. Igual, D. Oro, L. Rouan, and R. Pradel. 2011. Studying the reproductive skipping behavior in long-lived birds by adding nestinspection to individual-based data. Ecological Applications 21:555–564.