



# Reduction in baseline corticosterone secretion correlates with climate warming and drying across wild lizard populations

Andréaz Dupoué, Alexis Rutschmann, Jean-François Le Galliard, Jean Clobert, Pauline Blaimont, Barry Sinervo, Donald B. Miles, Claudy Haussy, Sandrine Meylan

## ► To cite this version:

Andréaz Dupoué, Alexis Rutschmann, Jean-François Le Galliard, Jean Clobert, Pauline Blaimont, et al.. Reduction in baseline corticosterone secretion correlates with climate warming and drying across wild lizard populations. *Journal of Animal Ecology*, 2018, 87 (5), pp.1331-1341. 10.1111/1365-2656.12843 . hal-02129407

**HAL Id: hal-02129407**

**<https://hal.science/hal-02129407>**

Submitted on 23 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Reduction of baseline corticosterone secretion correlates with  
climate warming and drying across wild lizard populations**

Andréaz Dupoué <sup>1\*</sup>, Alexis Rutschmann <sup>2</sup>, Jean François Le Galliard <sup>1,3</sup>, Jean Clobert <sup>2</sup>,  
Pauline Blaimont <sup>4</sup>, Barry Sinervo <sup>4</sup>, Donald B Miles <sup>2,5</sup>, Claudy Haussy <sup>1</sup>, Sandrine Meylan  
<sup>1,6</sup>

<sup>1</sup> Sorbonne Université, CNRS, iEES Paris, UMR 7618, 4 place Jussieu, F-75005 Paris, France

<sup>2</sup> Station d'Ecologie Théorique et Expérimentale du CNRS à Moulis, UMR 5321, 2 route du  
CNRS, 09200 Saint Giron, France

<sup>3</sup> Centre de recherche en écologie expérimentale et prédictive (CEREPEP-Ecotron  
IleDeFrance), Ecole normale supérieure, CNRS, PSL Research University, UMS 3194, 78 rue  
du château, 77140 Saint-Pierre-lès-Nemours, France

<sup>4</sup> Department of Ecology & Evolutionary Biology, University of California, Santa Cruz, 1156  
High Street, Santa Cruz, CA 95064, USA

<sup>5</sup> Department of Biological Sciences, Ohio University, Athens, OH, 45701, USA

<sup>6</sup> ESPE de Paris, Sorbonne Université, 10 Rue Molitor, 75016 Paris, France

\*Correspondence: andreaz.dupoue@gmail.com

19   **Abstract**

- 20   1. Climate change should lead to massive loss of biodiversity in most taxa but the detailed  
21       physiological mechanisms underlying population extinction remain largely elusive so far.  
22       In vertebrates, baseline levels of hormones such as glucocorticoids (GCs) may be  
23       indicators of population state since their secretion to chronic stress can impair survival  
24       and reproduction. However, the relationship between GC secretion, climate change and  
25       population extinction risk remains unclear.
- 26   2. In this study we investigated whether levels of baseline corticosterone (the main GCs in  
27       reptiles) correlate with environmental conditions and associated extinction risk across  
28       wild populations of the common lizard *Zootoca vivipara*.
- 29   3. First, we performed a cross-sectional comparison of baseline corticosterone levels along  
30       an altitudinal gradient among 14 populations. Then, we used a longitudinal study in 8  
31       populations to examine the changes in corticosterone levels following the exposure to a  
32       heat wave period.
- 33   4. Unexpectedly, baseline corticosterone decreased with increasing thermal conditions at  
34       rest in females, and was not correlated with extinction risk. In addition, baseline  
35       corticosterone levels decreased after exposure to an extreme heat wave period. This  
36       seasonal corticosterone decrease was more pronounced in populations without access to  
37       standing water.
- 38   5. We suggest that low basal secretion of corticosterone may entail down-regulating activity  
39       levels and limit exposure to adverse climatic conditions, especially to reduce water loss.  
40       These new insights suggest that rapid population decline might be preceded by a down-  
41       regulation of the corticosterone secretion.

42   **Keywords.** Altitude, corticosterone, ectotherm, population decline, temperature, water  
43   availability.

44     **Introduction**

45     An increasing number of living organisms are on the verge of extinction, mostly due to  
46     human-related factors such as land use, exploitation or climate change (Pereira et al., 2010;  
47     Thomas et al., 2004). For instance, nearly half of all plant and animal species have already  
48     faced population extinctions over the last 50 years (Wiens, 2016), and the pace of extinction is  
49     greater than expected in vertebrates (Ceballos, Ehrlich, & Dirzo, 2017). This Earth "sixth  
50     mass extinction crisis" has led to an intense effort to understand the liabilities of global  
51     change on biodiversity loss, and the search of reliable physiological mechanisms underlying  
52     population declines and of physiological determinants of extinction risk (Cooke et al., 2013;  
53     Wikelski & Cooke, 2006). Indeed, physiological studies are useful because they could unravel  
54     individual-level mechanisms underlying population declines and the identification of relevant  
55     biomarkers of the extinction risk could ease conservation status assessment and prioritization  
56     of management actions.

57             In vertebrates, hormones such as glucocorticoids (GCs) modulate daily and seasonal  
58     routines, such as reproductive cycles, as well as behavioural and life history responses to cope  
59     with both predictable and unpredictable events (Angelier & Wingfield, 2013; Landys,  
60     Ramenofsky, & Wingfield, 2006; Wingfield et al., 1998). The baseline secretion of GCs by  
61     the hypothalamic–pituitary–adrenal axis help individuals to mobilise energy associated with  
62     the demands of specific homeostatic states, where homeostasis refers to the tendency of the  
63     individual organism to maintain a state of physiological equilibrium on the long run (Landys  
64     et al., 2006; Romero, Dickens, & Cyr, 2009). However, individuals chronically exposed to  
65     stressors may show increased levels of baseline GCs, changes in the magnitude of the GCs  
66     response to an acute stress, and/or a reduction in the capacity to recover from perturbations,  
67     all of which may impair survival and/or reproduction on the long term (Angelier, Wingfield,  
68     Weimerskirch, & Chastel, 2010; Breuner, Patterson, & Hahn, 2008; Meylan, Miles, &

69 Clobert, 2012; Romero & Wikelski, 2010; Wingfield, 2013). Therefore, the regulation of GCs  
70 secretion may trigger adaptive responses to environmental changes, although empirical  
71 studies have shown that the fitness outcomes of the stress response are often complex and  
72 species-specific (Angelier et al., 2010; Bonier, Martin, Moore, & Wingfield, 2009; Bonier,  
73 Moore, & Robertson, 2011).

74 In wild populations, a range of chronic stressors from anthropogenic disturbance,  
75 including exposure of animals to unregulated ecotourism (Ellenberg, Setiawan, Cree,  
76 Houston, & Seddon, 2007; French, DeNardo, Greives, Strand, & Demas, 2010), habitat  
77 pollution (Crino, Klassen Van Oorschot, Johnson, Malisch, & Breuner, 2011; Meillère et al.,  
78 2016; Wikelski, Romero, & Snell, 2001), or landscape fragmentation (Janin, Léna, & Joly,  
79 2011; Martínez-Mota, Valdespino, Sánchez-Ramos, & Serio-Silva, 2007) may result in  
80 chronic elevation of baseline GC level, population declines and increased risks of rapid  
81 population extinction. There is also some direct evidence that lower corticosterone level may  
82 improve fitness of individuals exposed to desiccation and thermal stress in terrestrial  
83 ectotherms (Jessop, Letnic, Webb, & Dempster, 2013). And yet, a demonstration that  
84 secretion of GCs consistently correlate with the status and future fate of populations exposed  
85 to climate warming has not emerged yet, because of a wide variation in the sensitivity of GCs  
86 to anthropogenic disturbances across individuals and variable pathways between patterns of  
87 GCs secretion and individual fitness (Dantzer, Fletcher, Boonstra, & Sheriff, 2014; Dickens &  
88 Romero, 2013). These results thus question the GCs as a pertinent bio-marker of population  
89 extinction status.

90 Climate change represents an ubiquitous environmental challenge for living organisms  
91 that may interact with other environmental stressors and accelerate population declines  
92 (Bellard, Bertelsmeier, Leadley, Thuiller, & Courchamp, 2012; Cahill et al., 2012; Flesch,  
93 Rosen, & Holm, 2017). Ectothermic vertebrates are expected to be particularly vulnerable due

to their behavioural and physiological sensitivity to environmental temperature (Deutsch et al., 2008; Frishkoff, Hadly, & Daily, 2015; Kingsolver, Diamond, & Buckley, 2013; Le Galliard, Massot, Baron, & Clobert, 2012; Telemeco et al., 2017). In these species, flexible secretion GCs may therefore determine the ability of individuals and populations to cope with acute or chronic changes in thermal conditions (Jessop et al., 2016). For instance, the baseline secretion of GCs may be up-regulated to help individuals adjust their physiology (e.g., metabolic rate) and/or behaviour (e.g., flight response, behavioural thermoregulation) when they are exposed to non-optimal environmental temperatures (Dupoué, Brischoux, Lourdaïs, & Angelier, 2013; Telemeco & Addis, 2014). Still, although body and environmental temperatures generally correlate with secretion of GCs across and within species in ectothermic vertebrates (Jessop et al., 2016), the relationships between baseline levels of GCs, climate change and extinction risk in wild populations have not been investigated.

In this study, we examined the variation of baseline plasma corticosterone (the primary GCs in birds and reptiles) across 14 populations of the European common lizard (*Zootoca vivipara*) distributed across an altitudinal gradient. Our study populations occupy habitats that differ in thermal microclimates and access to free standing water (Dupoué, Rutschmann, Le Galliard, Miles, et al., 2017; Rutschmann et al., 2016), and are distributed along an extinction risk gradient including rapidly declining populations at the lowest altitudes and steady populations at mountaintops (Chamaillé-Jammes, Massot, Aragon, & Clobert, 2006; Sinervo et al., 2010). So far, population extinction has been directly related to warmer conditions in lowland populations although the proximate mechanisms remain unknown (Bestion, Teyssier, Richard, Clobert, & Cote, 2015). It is noteworthy that natural populations also differ in other parameters including vegetation cover (Lorenzon, Clobert, Oppliger, & John-Alder, 1999; Rutschmann et al., 2016), or slope orientation and local wind speed conditions (pers. obs.), which may influence corticosterone levels. However, we

focused here on the environmental covariates that are associated with population decline (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017). In each population, we sampled the representative demographic structure by capturing females and males from two age classes, including reproductive adults and non-reproductive yearlings. In Summer 2015, we combined two complementary studies to compare and clarify the factors responsible for variation in baseline corticosterone levels along the altitudinal gradient. First, we designed a cross-sectional study in which we compared baseline corticosterone levels across the 14 populations at a single time point in the early summer, which corresponds to the mid-gestation period in females. We then used a longitudinal study on lizards from 8 out of these 14 populations to investigate seasonal changes in corticosterone after three weeks in the middle of summer. During this specific time period, all populations experienced an extreme heat wave which ranked 6<sup>th</sup> in severity of European heat waves since 1950 according to meteorological reports (Russo, Sillmann, & Fischer, 2015). Our general hypothesis is that baseline corticosterone levels should be higher in populations where lizards are chronically exposed to extreme environmental conditions, such as consistently higher temperatures at the lower altitudes or extended periods of high temperatures (e.g., heat waves), ultimately leading to population collapse.

## **Material and methods**

### *Studied species and populations*

The European common lizard, *Zootoca vivipara*, is a small (adult size ~ 50 - 75 mm), widespread species (Family: Lacertidae) that inhabits peat bogs and heathlands across northern Eurasia. In 2015, we studied 14 populations found in the Massif Central Mountains in south-central France, which corresponds to the southern range limits for the viviparous form of the species in France (Rutschmann et al., 2016). In our focal populations, males emerge in mid-April while females emerge in early May. Males copulate with females shortly

after their emergence with fertilization occurring in mid to late May (Bleu et al., 2013). Gestation has a duration of 2 to 3 months, with parturition occurring between late-June and early August.

In each population, we characterized local environmental conditions, including altitude, permanent or periodic access to free standing water, and thermal microclimate (Table S1) (Dupoué, Rutschmann, Le Galliard, Miles, et al., 2017). We also calculated the relative changes in lizard abundance observed during the past decade and the extinction status according to IUCN indexes (IUCN, 2017). We distinguished the populations with permanent access to water in peat bogs or humid meadow habitats from those with periodic access in dry meadows with no water during summer except precipitations and morning dew. This objective score correlates with physiological regulation of the water balance in these populations such that lizards from populations without permanent access to water down-regulate their water loss rates to remain normosmotic (Dupoué, Rutschmann, Le Galliard, Miles, et al., 2017). We used two to three temperature data loggers (iButtons, Maxim Integrated Products, Sunnyvale, CA, USA,  $\pm 0.0625^{\circ}\text{C}$ ) per population. Loggers were placed within vegetation in the shade and protected in a polyvinyl cylinder pipe (diameter=5 cm, length=15 cm) to measure microclimatic temperature every hour from 29 June to 17 July between our two sampling sessions. This three weeks sampling period was chosen to reflect accurately the differences in microclimatic conditions during the active season among populations. We extracted the mean daily minimum and mean daily maximum temperatures ( $T_{\min}$ , and  $T_{\max}$ , respectively) to assess the thermal microclimate of each population. Population indices calculated with this method were correlated between years (2015-2017) either for  $T_{\min}$  (Pearson's  $r = 0.75$ ) or  $T_{\max}$  ( $r = 0.42$ ).

These populations have been monitored since 2005, so we could estimate relative changes in lizard abundance over a decade. Lizards are captured during days in active seasons

when the weather conditions and external temperature are optimal. During the searches, people randomly scan the populations to find lizards that are then captured by hand. Lizard abundance was calculated in summers 2005, 2015 and 2017 based on the number of lizards captured divided by the total time spent searching, the number of people capturing and the area of the study site (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017). We used the relative changes in abundance to determine the IUCN status of each population, and we identified two populations extremely at risk, one population at risk and 11 non-threatened populations (Table S1). This classification is consistent with the predictive model of climate change effect on population trends derived from a previous field experiment (Bestion et al., 2015). We managed to sample 135 individuals from those populations in a critical state, and it is worth noting that studies including as much individuals and replicates of collapsing populations are rare and urgently needed. Besides, many populations from this Mountain range have been extirpated during the 2-3 last decades following higher exposure to warm spells and summer droughts (Sinervo et al., under review) suggesting that population decline precedes inevitable extinction.

#### *Sampling procedures*

##### Cross-sectional study

Between the 19 and 26 of June 2015 (early summer session), we caught a total of 312 adult pregnant females (mean  $\pm$  SE, body mass (BM) =  $4.77 \pm 0.07$  g, snout-vent length (SVL) =  $61.24 \pm 0.25$  mm), 132 adult males (BM =  $3.54 \pm 0.06$  g, SVL =  $54.75 \pm 0.30$  mm), 87 yearling females (BM =  $1.85 \pm 0.06$  g, SVL =  $47.02 \pm 0.49$  mm), and 131 yearling males (BM =  $2.01 \pm 0.05$  g, SVL =  $46.10 \pm 0.32$  mm) from the 14 populations. Within 5 min of capture, we bled all individuals using a standard protocol (Meylan, Dufty, & Clobert, 2003). Blood samples (40-60  $\mu$ l whole blood) were collected from the post-orbital sinus using 2-3 20  $\mu$ l microcapillary tubes. Samples were kept fresh in a cooler on ice and brought back to a field

laboratory the same day. In the laboratory, blood samples were centrifuged for 5 min at 11,000 rpm, plasma and blood cells were separated and kept frozen in airtight tubes until subsequent analyses.

Adult females and males were transferred to the laboratory and housed in individual terraria (18 x 12 x 12 cm) with sterilized soil, a shelter, and basking opportunities to record parturition date and measurement of reproductive effort (litter size and mass) (Rutschmann et al., 2016). We calculated the reproductive timing (i.e., the embryonic stage at the sampling) as the difference between parturition and sampling dates. Each lizard was provided a 20-30°C thermal gradient for 6 hours per day (09:00-12:00 and 14:00-17:00) using a 25 W incandescent light bulb placed over one end of each terrarium. We also provided water 3 times per day and fed lizards with 2 crickets (*Acheta domesticus*) every two days. In 2017, we measured the critical maximal temperature limits (CT<sub>max</sub>) in a subsample of 57 individuals from the monitored populations, using a standard protocol (Gilbert & Miles, 2017). Individuals were continuously heated (~1°C per minute) under a 60 W incandescent light bulb, and flipped on the back every minute. Once they stop responding to this stimuli by righting themselves within 5s, we measured their cloacal temperature with a digital thermometer, considering as the individual CT<sub>max</sub>. We released each male and female together with its offspring at its exact capture location within three days after parturition.

#### Longitudinal study

In 8 out of the 14 populations, we repeated the sampling procedures as described above between 19 and 23 July (mid-summer session). We focused on yearlings to avoid confounding effects of seasonal changes in reproductive state with those of seasonal changes in environmental conditions during a heat wave period. We sampled 76 yearling females (BM =  $2.50 \pm 0.06$  g, SVL =  $53.36 \pm 0.42$  mm) and 77 males (BM =  $2.81 \pm 0.06$  g, SVL =  $51.29 \pm 0.30$  mm). Although we repeated sampling from the same populations than the cross-sectional

study, the probability to recapture the very same individuals was pretty low relative to the sample size (Meylan, Clobert, & Sinervo, 2007), so all measurements were considered as independent.

#### *Plasma corticosterone levels*

Plasma corticosterone assays were performed with a competitive enzyme-immunoassay method (IDS Corticosterone EIA kit, ref AC-14F1, Immunodiagnostic Systems Ltd, France) after 1:10 dilution of all samples. This method quantifies total plasmatic corticosterone using a polyclonal corticosterone antibody and is based on a highly repeatable colorimetric assay of absorbance at 450 nm. The reported sensitivity of the kit is  $0.55 \text{ ng mL}^{-1}$ , and our estimates of corticosterone levels were indeed highly repeatable [12 plates with 4 repeats of a standard per plate: intra-plate repeatability:  $r = 0.98$ ,  $F_{1,34} = 168.7$ ,  $p < 0.001$ ; inter-plate repeatability:  $r = 0.81$ ,  $F_{11,34} = 9.6$ ,  $p < 0.001$  (Lessells & Boag, 1987)]. Such repeatability, the coefficient of variation we obtained (intra-plate: 14%; inter-plate: 23%) and the similar levels of corticosterone assessed either by ELISA or by RIA methods (Meylan et al., 2003) suggest reliable results of corticosterone assays.

#### *Statistical analyses*

Plasma corticosterone were  $\log_{10}$  transformed to achieve normal distribution and analysed with linear models in the R software (R Development Core Team, version 3.2.0, <http://cran.r-project.org/>). In the cross-sectional study, the initial model included the quadratic effect of time of day to test for non-linear daily variation in corticosterone (Dauphin-Villemant & Xavier, 1987), and the fixed effects of population, sex, age class, and first- and second order interaction terms. In the longitudinal study, the initial model included the quadratic effect of time of day, fixed effects of population, sex, and sampling session, and first- and second order interaction terms. In the studied populations, lizard morphology strongly varies among and between populations, sex and age class as previously demonstrated (e.g., Chamaillé-Jammes

et al., 2006; Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017; Massot, Clobert, Pilorge, Lecomte, & Barbault, 1992). Therefore to avoid multicollinearity, lizard morphological indexes were not included in the model selection since we aimed to specifically test the influence of environmental conditions on baseline corticosterone levels independently from variation in morphology. We compared all models to a null model including the intercept only (cross-sectional study: Table S2; longitudinal study: Table S3).

We found significant variation in corticosterone levels among populations in the final model of each study, and therefore tested the influence of environmental covariates on corticosterone levels in a second step. To do so, we used an approach using the Akaike information criterion corrected for small sample size (AICc, package AICcmodavg, Mazerolle 2016). We compared mixed-effect linear models [package nlme, (Pinheiro, Bates, DebRoy, Sarkar, & R, 2016)] in which population identity was included as a random factor to account for non-independence within the same population. In all selected models, the random effects of populations had significant contribution to explain variation in baseline corticosterone (all  $p < 0.001$ ). Environmental covariates were treated one by one. Water access was treated as a categorical factor while the relative changes in abundance, temperature metrics (i.e.,  $T_{\min}$  and  $T_{\max}$ ), and altitude, were treated as linear covariates. Given that several variables were highly correlated and related to population collapse, we computed the relative changes in abundance,  $T_{\min}$  and the altitude together in a principal component analysis [package ade4, (Dray & Dufour, 2007)]. We used the first axis (PC<sub>1</sub>) as a composite score of extinction risk since it was mainly determined by the relative changes in lizard abundance (Table S4), since this integrative score of population decline has been demonstrated to correlate with molecular markers of physiological stress (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017). We compared all models, including additive effects of each environmental covariate or the PC<sub>1</sub> score (6 variables), as well as two-way interactions with age and sex (cross-sectional study,

Table S5) and two-way interactions with sex and sampling session (longitudinal study, Table S6).

In adult pregnant females, we also investigated the relationships between baseline corticosterone levels and reproduction. We compared mixed effects models (population as random factor), with reproductive timing and reproductive effort as linear covariates and compared to a null model with the intercept only (Table S7). It is indeed predicted that GC secretion may increase during gestation and be higher in females with a higher reproductive effort (Dauphin-Villemant & Xavier, 1986; Lorion, Angelier, & Lourdais, 2016). For all model comparisons, the best model was then chosen as the one with the lowest AICc and models that have a difference of AICc lower than 2 comparably support the data (see Tables S2, S3, S5, S6 & S7 for model comparisons).

## Results

### *Cross-sectional study*

Baseline corticosterone levels differed across the 14 populations ( $F_{13,619} = 3.86$ ,  $p < 0.001$ ), between sexes ( $F_{1,619} = 38.42$ ,  $p < 0.001$ ), but did not show daily variation (time of day:  $F_{13,617} = 1.48$ ,  $p = 0.225$ , time of day<sup>2</sup>:  $F_{13,617} = 0.92$ ,  $p = 0.338$ ). Baseline corticosterone levels were significantly impacted by the two-way interactions between age and sex ( $F_{1,619} = 31.46$ ,  $p < 0.001$ ), so that corticosterone levels differed between age classes in females but not in males (mean  $\pm$  SE; pregnant females:  $36.35 \pm 1.34$  ng.ml<sup>-1</sup>, yearling females:  $23.44 \pm 1.83$  ng.ml<sup>-1</sup>, adult males:  $17.07 \pm 1.33$  ng.ml<sup>-1</sup>, and yearling males:  $17.26 \pm 1.18$  ng.ml<sup>-1</sup>).

Baseline corticosterone levels were also significantly impacted by the two-way interactions between age and population ( $F_{13,619} = 2.33$ ,  $p = 0.005$ ) and between sex and population ( $F_{13,619} = 3.53$ ,  $p < 0.001$ ). According to our model selection procedure, most of the inter-population variation in baseline corticosterone levels was explained by  $T_{\min}$  and the two-way interactions between  $T_{\min}$  and age class and between  $T_{\min}$  and sex (model likelihood

$w_i = 0.74$ , see Table S5). Specifically, baseline corticosterone levels decreased with  $T_{\min}$  in females (adults:  $t_{1,12} = -2.31$ ,  $p = 0.039$ , Fig. 1a; yearlings:  $t_{1,12} = -4.01$ ,  $p = 0.002$ , Fig. 1b) but not in males (adults:  $t_{1,12} = 0.98$ ,  $p = 0.347$ , Fig. 1c; yearlings:  $t_{1,12} = -1.33$ ,  $p = 0.209$ , Fig. 1d). Other models had very low relative statistical support ( $w_i \leq 0.09$ ) indicating that the relationships with other covariates were negligible (Table S5). In addition, in adult pregnant females, we did not find any significant relationship between baseline corticosterone and reproductive timing or reproductive effort (Table S7).

### *Longitudinal study*

During the heat wave period in 2015, temperatures were higher and overpassed the lizard CTmax 42.5% of time during normal activity period (10:00 to 19:00), while remaining lower and non-constraining (0% of CTmax overpassing) in 2017 (Fig. 2). Overall, baseline corticosterone levels were lower after the heat wave period than before (early summer session:  $20.29 \pm 1.40 \text{ ng.ml}^{-1}$ , mid-summer session:  $13.68 \pm 0.83 \text{ ng.ml}^{-1}$ ;  $F_{1,244} = 9.82$ ,  $p = 0.002$ ), and this seasonal variation between sampling sessions differed among populations (population x sampling session:  $F_{7,244} = 3.45$ ,  $p = 0.002$ ). In addition, corticosterone levels showed non-linear daily variation (time of day:  $F_{1,244} = 4.51$ ,  $p = 0.035$ , time of day<sup>2</sup>:  $F_{1,244} = 4.09$ ,  $p = 0.044$ , Fig. 3a), and were overall higher in yearling females compared to males ( $F_{1,244} = 7.66$ ,  $p = 0.006$ ).

Differences among populations in their seasonal changes in baseline corticosterone levels were best explained by the access to free standing water since the two best models for baseline corticosterone levels included a two-way interaction between this covariate and the measurement session (combined likelihood  $w_i = 0.35$ , see Table S6). Specifically, according to the top ranking model, baseline corticosterone levels decreased during the heat wave in yearlings from populations characterized by periodic access to water ( $t_{1,257} = -4.05$ ,  $p < 0.001$ , Fig. 3b), while corticosterone levels remained similar in populations with permanent access to

water ( $t_{1,257} = -0.81$ ,  $p = 0.417$ , Fig. 3b). Other models had lower degree of relative support ( $w_i \leq 0.08$ ) suggesting that the relationships with other covariates were not important (Table S6).

## Discussion

We designed two complementary field studies to examine geographic variation in baseline corticosterone levels among wild populations of lizards across a gradient of environmental conditions and extinction risk. We found strong geographic variation in baseline corticosterone levels, interactively or additively with lizard sex and age class. Contrary to our expectations, these differences were poorly related with extinction risks as indicated by an integrative score (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017) and IUCN conservation status. Instead, baseline corticosterone levels of females were lower in local populations with higher minimum daily temperature, and decreased on average in yearlings after a heat wave, suggesting that chronic heat stress was correlated with a lower corticosterone secretion. Importantly, our results also revealed that access to free standing water in each population may determine the sensitivity of corticosterone secretion during the exposure to an intense heat wave.

In this study, we found greater plasma corticosterone levels in adult females compared to males or yearling which is consistent with previous findings showing temporal increase in corticosterone over pregnancy in squamate reptiles (Dauphin-Villemant & Xavier, 1987; Lorigou et al., 2016; Taylor, DeNardo, & Jennings, 2004). Surprisingly in pregnant females, the higher baseline corticosterone levels was not correlated with any index of reproductive performance, thus questioning the proximate causes of variations in hormonal profile during gestation. Besides these predictable variations in baseline corticosterone across life stages, we found sex-specific relationships between baseline corticosterone levels and local thermal conditions during the night (i.e., minimum daily temperatures), which describe local climate conditions when individuals are at rest. Baseline corticosterone levels decreased with  $T_{\min}$  in

females but not in males. This sex-specific pattern may result from a differential sensitivity to temperature, which is likely related to sex-specific life history strategies in this species (Massot et al., 1992). In support of this, it is noteworthy that baseline corticosterone concentration strongly depends on body temperature in ectotherms (Dupoué et al., 2013; Jessop et al., 2016). Further studies would help to clarify if thermoregulation differs within (sex and age specific response) and between populations, and whether differences across populations are a direct consequence of changes in body temperature or due to functional plasticity or local adaptations.

These relationships could mirror the negative correlation between baseline GCs and environmental temperatures observed across species in reptiles (Jessop et al., 2016). One explanation for the higher baseline corticosterone levels in reptile species inhabiting colder habitats is that a high baseline GCs level helps in supporting the faster energy demand and mobilisation needed during activity periods when thermoregulatory opportunities are fewer (Dupoué et al., 2013; Jessop et al., 2016; Telemeco & Addis, 2014). If this hypothesis holds true, we would expect a stronger correlation between baseline GCs levels and thermal conditions during the activity period of the day than during the resting period. However, we did not find any influence of daily maximal temperature ( $T_{\max}$ ), suggesting that geographic differences in thermoregulation opportunities during the day did not explain the changes in corticosterone levels. Alternatively, lower minimum temperatures may trigger higher level of circulating corticosterone to compensate and/or maintain a level of physiological performances that are classical decreased at lower body temperature (Angilletta, 2009; Qualls & Andrews, 1999). Experiments where lizards are exposed to separate changes in nocturnal and daily minimum and maximum environmental temperatures are needed to test if low minimum daily temperatures can indeed induce higher GC secretion and therefore represent "more stressful" conditions than high temperatures.

We suggest instead that individuals responded to abnormally high thermal conditions at rest (i.e., during night-time) because lowering plasma corticosterone may help down regulate maintenance costs at rest and energy expenditure during activity (e.g., locomotion or the proportion of time basking). Low secretion of GC in the warmest populations may thus be an adaptive physiological response in order to save energy and water (Cote, Clobert, Meylan, & Fitze, 2006; Cote, Clobert, Poloni, Haussy, & Meylan, 2010; Preest & Cree, 2008). Indeed, high metabolism at rest and strong behavioural activity in warmer environments could induce diverse physiological costs such as higher energy expenditure, higher risk of dehydration, and eventually negative impacts on survival and reproduction (Bestion et al., 2015; Dillon, Wang, & Huey, 2010; Huey et al., 2012; Kearney, Shine, & Porter, 2009; McKechnie & Wolf, 2010). Alternatively, the studied populations can associate micro-adaptations (e.g., significant genetic differentiation among populations), different trajectory in life history strategies (Chamaillé-Jammes et al., 2006; Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017; Rutschmann et al., 2016), which might explain some variations in corticosterone levels between populations. Besides, we must acknowledge that baseline corticosterone alone may not always correlate with individual stress and animal welfare (Otovic & Hutchinson, 2015). Instead, the acute GC response to a stressor following a standard capture-restrain process or the capacity to recover from such stress response represent complementary indicators of GC secretion that might better predict reproduction and/or survival rate than baseline levels (Blas, Bortolotti, Tella, Baos, & Marchant, 2007; Romero & Wikelski, 2010). However, analyses of the GC secretion in response to acute stressors would require repeated blood sampling, which could not be done for ethical reasons and logistic in this large scale study with small bodied lizards (adult body mass ~3 - 5g) involving many populations. Other physiological indicators of stress could also provide crucial insights on the responses to abnormally high temperatures and desiccation. For instance, the heat shock proteins represent particularly promising

components of the stress response that are likely impacted by global warming and should correlate with population extinction status (Sørensen, 2010). Besides, we recently documented in these populations that telomere length, another integrative indicator of aging and thermal stresses has been recently shown to non-linearly shorten in populations facing high risk of extinction when compared to non-threatened ones (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017). Therefore, further comparative studies are required to depict the specific causes of variation in baseline GC level, GC stress response, GC recovery, or other pertinent physiological indexes of stress and the further effects on individual fitness.

Following a similar tendency, we found that baseline corticosterone decreases seasonally during an intense heat wave period, especially in the driest (i.e., with no access to free standing water) populations. The heat wave was particularly pronounced in 2015 since it ranked 6th in severity in Europe since 1950 (Russo et al., 2015) with abnormally high temperatures and no precipitation (Dong, Sutton, Shaffrey, & Wilcox, 2016). Instead, according to our initial hypothesis, a seasonal increase in baseline corticosterone was expected as individuals were most likely exposed to chronic heat stress and summer droughts, especially in the populations without access to water (Wingfield, 2013). Seasonal differences in mean corticosterone levels could come from intra-individual changes (i.e., physiological flexibility) but also from inter-individual viability selection (i.e., selective removal of individuals with different initial baseline levels) or from a combination of both processes. We cannot exclude that individuals with higher corticosterone levels at the beginning of the summer were selected against through lower survival rate during the heat wave period, which could have been determined with a repeating sampling all over the exposure to the heat waves. However, this hypothesis would involve a strong selection and a high repeatability of corticosterone levels over time, which is very unlikely according to current estimates of inter-individual repeatability of GC levels (Holtmann, Lagisz, & Nakagawa, 2017). Our results thus

alternatively suggest that individual down regulated corticosterone secretion and decreased their resting metabolism and behavioural activity to limit the costs associated with warm and dry environments (see above). Since this seasonal decrease was restricted to populations without access to free standing water, regulation of the water balance may be an important target in the behavioural regulation of activity level in wild lizards (Lorenzon et al., 1999). Water balance is the sum of water entrance (i.e., drinking behaviour) and water exit (i.e., faeces, evaporative water losses through ventilation and transpiration). The rate of evaporative water loss and eventually the rate of dehydration will depend on temperature, humidity and activity levels. When facing extreme heat waves, the behavioural regulation of the water balance through lower corticosterone secretion could represent an adaptive response to avoid overheating, dehydration and premature death (Jessop et al., 2013). Further investigations should now clarify the role of corticosterone in sheltering, microhabitat selection, and the following outcomes on the regulation of water balance, and on individual fitness and general demographic trends in the context of climate change.

Heat wave frequency, intensity and duration will considerably increase in the next decades (Easterling et al., 2000; Meehl & Tebaldi, 2004). Although the responses to extreme temperatures are complex and various among taxa (Buckley & Huey, 2016), the exposure to such climatic events is predicted to lead to massive changes in demographic trends (McKechnie & Wolf, 2010). Within individuals, the allostasis concept posits that hormones such as corticosterone constitute the crucial interfaces between the energy demands associated with predictable life-history stages and unpredictable environmental changes, such as heat waves and dry episodes (McEwen & Wingfield, 2010; Romero et al., 2009). To avoid the exposure to abnormally high temperature and water deprivation, lowering activity and sheltering may thus be crucial to ensure immediate survival in many ectothermic species, and our results suggest that baseline corticosterone may contribute to mediate this adaptive

response in the common lizard. Yet, even if baseline corticosterone might be essential to mitigate the immediate costs of being active in harmful conditions, it was uncorrelated with population decline, thus questioning the pertinence of using high baseline corticosterone as a relevant biomarker of population extinction risk. In the long run, lower activity may even alter foraging or reproductive opportunities and eventually lead to population extinction in our study populations (Dupoué, Rutschmann, Le Galliard, Clobert, et al., 2017; Sinervo et al., 2010). Together, our results therefore suggest that climate change is associated with a down-regulation of corticosterone GC secretion preceding the population decline.

**Acknowledgments**

We thank Pauline Dufour, Laurène Duhalde, Amélie Faure, Julia Rense, and Qiang Wu for their help with fieldwork. We also thank Clotilde Biard for lending us some of the loggers. We are grateful to the ‘Office Nationale des Forêts’, the ‘Parc National des Cévennes’, and the regions Auvergne, Rhône Alpes and Languedoc Roussillon for allowing us to sample lizards.

**Funding**

This study was funded by the Centre National de la Recherche Scientifique (CNRS) the Agence Nationale de la Recherche (ANR-13-JSV7-0011-01 to SM).

**Authors' Contributions**

AD, AR, JFLG, JC, and SM conceived the ideas and design methodology, and all authors contributed to data collection; AD analysed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

**Ethics of Experiment**

All methods were performed in accordance with laws relative to capture, transport and experiments on *Zootoca vivipara* (DREAL Languedoc Roussillon permit #2013-274-0002,

DREAL Midi-Pyrénées permit #81-2013-05, and DREAL Auvergne, permit  
2013/DREAL/259).

### Conflicts of interest

We declare no competing financial interest.

### References

Angelier, F., & Wingfield, J. C. (2013). Importance of the glucocorticoid stress response in a  
changing world: theory, hypotheses and perspectives. *General and Comparative  
Endocrinology*, 190, 118–128. doi:10.1016/j.ygcen.2013.05.022

Angelier, F., Wingfield, J. C., Weimerskirch, H., & Chastel, O. (2010). Hormonal correlates  
of individual quality in a long-lived bird: a test of the “corticosterone – fitness  
hypothesis.” *Biology Letters*, 6(6), 846–849. doi:10.1098/rsbl.2010.0376

Angilletta, M. J. (2009). *Thermal adaptation: a theoretical and empirical synthesis*. Oxford:  
Oxford University Press.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of  
climate change on the future of biodiversity. *Ecology Letters*, 15(4), 365–377.  
doi:10.1111/j.1461-0248.2011.01736.x

Bestion, E., Teyssier, A., Richard, M., Clobert, J., & Cote, J. (2015). Live Fast, Die Young:  
Experimental Evidence of Population Extinction Risk due to Climate Change. *PLoS  
Biology*, 13(10), e1002281. doi:10.1371/journal.pbio.1002281

Blas, J., Bortolotti, G. R., Tella, J. L., Baos, R., & Marchant, T. A. (2007). Stress response  
during development predicts fitness in a wild, long lived vertebrate. *Proceedings of the*

- 491       *National Academy of Sciences of the United States of America*, 104(21), 8880–8884.  
492       doi:10.1073/pnas.0700232104
- 493   Bleu, J., Le Galliard, J. F., Fitze, P. S., Meylan, S., Clobert, J., & Massot, M. (2013).  
494       Reproductive allocation strategies: a long-term study on proximate factors and temporal  
495       adjustments in a viviparous lizard. *Oecologia*, 171(1), 141–151. doi:10.1007/s00442-  
496       012-2401-1
- 497   Bonier, F., Martin, P. R., Moore, I. T., & Wingfield, J. C. (2009). Do baseline glucocorticoids  
498       predict fitness? *Trends in Ecology & Evolution*, 24(11), 634–642.  
499       doi:10.1016/j.tree.2009.04.013
- 500   Bonier, F., Moore, I. T., & Robertson, R. J. (2011). The stress of parenthood? Increased  
501       glucocorticoids in birds with experimentally enlarged broods. *Biology Letters*, 7(May),  
502       944–946. doi:10.1098/rsbl.2011.0391
- 503   Breuner, C. W., Patterson, S. H., & Hahn, T. P. (2008). In search of relationships between the  
504       acute adrenocortical response and fitness. *General and Comparative Endocrinology*,  
505       157(3), 288–295. doi:10.1016/j.ygcen.2008.05.017
- 506   Buckley, L. B., & Huey, R. B. (2016). Temperature extremes: geographic patterns, recent  
507       changes, and implications for organismal vulnerabilities. *Global Change Biology*,  
508       22(12), 3829–3842. doi:10.1111/gcb.13313
- 509   Cahill, A. E., Aiello-Lammens, M. E., Fisher-Reid, M. C., Hua, X., Karanewsky, C. J., Yeong  
510       Ryu, H., ... Wiens, J. J. (2012). How does climate change cause extinction? *Proceedings*  
511       *of the Royal Society B: Biological Sciences*, 280, 20121890. doi:10.1098/rspb.2012.1890
- 512   Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth  
513       mass extinction signaled by vertebrate population losses and declines. *Proceedings of the*

- 514 *National Academy of Sciences of the United States of America*, 114(30), E6089–E6096.  
515 doi:10.1073/pnas.1704949114
- 516 Chamaillé-Jammes, S., Massot, M., Aragon, P., & Clobert, J. (2006). Global warming and  
517 positive fitness response in mountain populations of common lizards *Lacerta vivipara*.  
518 *Global Change Biology*, 12, 392–402. doi:10.1111/j.1365-2486.2005.01088.x
- 519 Cooke, S. J., Sack, L., Franklin, C. E., Farrell, A. P., Beardall, J., Wikelski, M., & Chown, S.  
520 L. (2013). What is conservation physiology? Perspectives on an increasingly integrated  
521 and essential science. *Conservation Physiology*, 1(January), 1–23.  
522 doi:10.1093/conphys/cot001
- 523 Cote, J., Clobert, J., Meylan, S., & Fitze, P. S. (2006). Experimental enhancement of  
524 corticosterone levels positively affects subsequent male survival. *Hormones and*  
525 *Behavior*, 49(3), 320–327. doi:10.1016/j.yhbeh.2005.08.004
- 526 Cote, J., Clobert, J., Poloni, L. M., Haussy, C., & Meylan, S. (2010). Food deprivation  
527 modifies corticosterone-dependent behavioural shifts in the common lizard. *General and*  
528 *Comparative Endocrinology*, 166(1), 142–151. doi:10.1016/j.ygcen.2009.11.008
- 529 Crino, O. L., Klassen Van Oorschot, B., Johnson, E. E., Malisch, J. L., & Breuner, C. W.  
530 (2011). Proximity to a high traffic road: glucocorticoid and life history consequences for  
531 nestling white-crowned sparrows. *General and Comparative Endocrinology*, 173(2),  
532 323–332. doi:10.1016/j.ygcen.2011.06.001
- 533 Dantzer, B., Fletcher, Q. E., Boonstra, R., & Sheriff, M. J. (2014). Measures of physiological  
534 stress: a transparent or opaque window into the status , management and conservation of  
535 species? *Conservation Physiology*, 2(May), 1–18. doi:10.1093/conphys/cou023.
- 536 Dauphin-Villemant, C., & Xavier, F. (1986). Adrenal activity in the females *Lacerta vivipara*

- 537       Jacquin: possible involvement in the success of gestation. In I. Assemacher & J. Boissin  
538       (Eds.), *Endocrine regulation as adaptive mechanism to environment* (pp. 241–250).  
539       Paris: CNRS.
- 540       Dauphin-Villemant, C., & Xavier, F. (1987). Nycthemeral variations of plasma  
541       corticosteroids in captive female *Lacerta vivipara* Jacquin: influence of stress and  
542       reproductive state. *General and Comparative Endocrinology*, 67, 292–302.  
543       doi:10.1016/0016-6480(87)90183-3
- 544       Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C.,  
545       & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across  
546       latitude. *Proceedings of the National Academy of Sciences of the United States of*  
547       *America*, 105(18), 6668–6672. doi:10.1073/pnas.0709472105
- 548       Dickens, M. J., & Romero, L. M. (2013). A consensus endocrine profile for chronically  
549       stressed wild animals does not exist. *General and Comparative Endocrinology*, 191,  
550       177–189. doi:10.1016/j.ygcen.2013.06.014
- 551       Dillon, M. E., Wang, G., & Huey, R. B. (2010). Global metabolic impacts of recent climate  
552       warming. *Nature*, 467(7316), 704–706. doi:10.1038/nature09407
- 553       Dong, B., Sutton, R., Shaffrey, L., & Wilcox, L. (2016). The 2015 European heat wave.  
554       *Bulletin of the American Meteorological Society*, 97, S57–S62. doi:10.1175/BAMS-D-  
555       16-0140.1
- 556       Dray, S., & Dufour, A. B. (2007). The ade4 package: implementing the duality diagram for  
557       ecologists. *Journal of Statistical Software*, 22(4), 1–20.
- 558       Dupoué, A., Brischoux, F., Lourdais, O., & Angelier, F. (2013). Influence of temperature on  
559       the corticosterone stress-response: an experiment in the Children's python (*Antaresia*

- 560 childreni). *General and Comparative Endocrinology*, 193, 178–184.  
561 doi:10.1016/j.ygcen.2013.08.004
- 562 Dupoué, A., Rutschmann, A., Le Galliard, J. F., Clobert, J., Angelier, F., Marciau, C., ...  
563 Meylan, S. (2017). Shorter telomeres precede population extinction in wild lizards.  
564 *Scientific Reports*, 7, 16976. doi:10.1038/s41598-017-17323-z
- 565 Dupoué, A., Rutschmann, A., Le Galliard, J. F., Miles, D. B., Clobert, J., DeNardo, D. F., ...  
566 Meylan, S. (2017). Water availability and environmental temperature correlate with  
567 geographic variation in water balance in common lizards. *Oecologia*, 185, 561–571.  
568 doi:10.1007/s00442-017-3973-6
- 569 Easterling, D. R., Meehl, G. A., Parmesan, C., Changnon, S. A., Karl, T. R., & Mearns, L. O.  
570 (2000). Climate extremes: Observations, modeling, and impacts. *Science*, 289, 2068–  
571 2074. doi:10.1126/science.289.5487.2068
- 572 Ellenberg, U., Setiawan, A. N., Cree, A., Houston, D. M., & Seddon, P. J. (2007). Elevated  
573 hormonal stress response and reduced reproductive output in Yellow-eyed penguins  
574 exposed to unregulated tourism. *General and Comparative Endocrinology*, 152(1), 54–  
575 63. doi:10.1016/j.ygcen.2007.02.022
- 576 Flesch, A. D., Rosen, P. C., & Holm, P. (2017). Long-term changes in abundances of Sonoran  
577 Desert lizards reveal complex responses to climatic variation. *Global Change Biology*,  
578 (June), 1–17. doi:10.1111/gcb.13813
- 579 French, S. S., DeNardo, D. F., Greives, T. J., Strand, C. R., & Demas, G. E. (2010). Human  
580 disturbance alters endocrine and immun responses in the Galapagos marine iguana  
581 (*Amblyrhynchus cristatus*). *Hormones and Behavior*, 58(5), 792–799.  
582 doi:10.1016/j.yhbeh.2010.08.001.Human

- 583 Frishkoff, L. O., Hadly, E. A., & Daily, G. C. (2015). Thermal niche predicts tolerance to  
584 habitat conversion in tropical amphibians and reptiles. *Global Change Biology*, 21(11),  
585 3901–3916. doi:10.1111/gcb.13016
- 586 Gilbert, A. L., & Miles, D. B. (2017). Natural selection on thermal preference, critical thermal  
587 maxima and locomotor performance. *Proceedings of the Royal Society B: Biological*  
588 *Sciences*, 284(1860), 20170536. doi:10.1098/rspb.2017.0536
- 589 Holtmann, B., Lagisz, M., & Nakagawa, S. (2017). Metabolic rates, and not hormone levels,  
590 are a likely mediator of between-individual differences in behaviour: a meta-analysis.  
591 *Functional Ecology*, 31, 685–696. doi:10.1111/1365-2435.12779
- 592 Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., & Williams, S.  
593 E. (2012). Predicting organismal vulnerability to climate warming: roles of behaviour ,  
594 physiology and adaptation. *Philosophical Transactions of the Royal Society of London.*  
595 *Series B, Biological Sciences*, 367, 1665–1679. doi:10.1098/rstb.2012.0005
- 596 IUCN. (2017). IUCN Red List Categories and Criteria and Guidelines for Using the IUCN  
597 Red List Categories and Criteria. Scenario A2.
- 598 Janin, A., Léna, J. P., & Joly, P. (2011). Beyond occurrence: Body condition and stress  
599 hormone as integrative indicators of habitat availability and fragmentation in the  
600 common toad. *Biological Conservation*, 144(3), 1008–1016.  
601 doi:10.1016/j.biocon.2010.12.009
- 602 Jessop, T. S., Lane, M. L., Teasdale, L., Stuart-Fox, D., Wilson, R. S., Careau, V., & Moore,  
603 I. T. (2016). Multiscale evaluation of thermal dependence in the glucocorticoid response  
604 of vertebrates. *The American Naturalist*, 188(3), 342–356. doi:10.1086/687588
- 605 Jessop, T. S., Letnic, M., Webb, J. K., & Dempster, T. (2013). Adrenocortical stress responses

- 606 influence an invasive vertebrate's fitness in an extreme environment. *Proceedings of the*  
607 *Royal Society B: Biological Sciences*, 280(1768), 20131444. doi:10.1098/rspb.2013.1444
- 608 Kearney, M., Shine, R., & Porter, W. P. (2009). The potential for behavioral thermoregulation  
609 to buffer “cold-blooded” animals against climate warming. *Proceedings of the National*  
610 *Academy of Sciences of the United States of America*, 106(10), 3835–3840.  
611 doi:10.1073/pnas.0808913106
- 612 Kingsolver, J. G., Diamond, S. E., & Buckley, L. B. (2013). Heat stress and the fitness  
613 consequences of climate change for terrestrial ectotherms. *Functional Ecology*, 27(6),  
614 1415–1423. doi:10.1111/1365-2435.12145
- 615 Landys, M. M., Ramenofsky, M., & Wingfield, J. C. (2006). Actions of glucocorticoids at a  
616 seasonal baseline as compared to stress-related levels in the regulation of periodic life  
617 processes. *General and Comparative Endocrinology*, 148(2), 132–149.  
618 doi:10.1016/j.ygcen.2006.02.013
- 619 Le Galliard, J. F., Massot, M., Baron, J. P., & Clobert, J. (2012). Ecological effects of climate  
620 change on European reptiles. In J. Brodie, E. Post, & D. Doak (Eds.), *Wildlife*  
621 *conservation in a changing climate* (pp. 179–203). Chicago: University of Chicago  
622 Press.
- 623 Lessells, C. M., & Boag, P. T. (1987). Unrepeatable repeatabilities: A common mistake. *The*  
624 *Auk*, 104(1), 116–121. doi:10.2307/4087240
- 625 Lorenzon, P., Clobert, J., Oppliger, A., & John-Alder, H. (1999). Effect of water constraint on  
626 growth rate , activity and body temperature of yearling common lizard (*Lacerta*  
627 *vivipara*). *Oecologia*, 118, 423–430. doi:10.1007/s004420050744
- 628 Lorient, S., Angelier, F., & Lourdais, O. (2016). Are glucocorticoids good indicators of

- 629 pregnancy constraints in a capital breeder? *General and Comparative Endocrinology*,  
630 232, 125–133. doi:10.1016/j.ygcen.2016.04.007
- 631 Martínez-Mota, R., Valdespino, C., Sánchez-Ramos, M. A., & Serio-Silva, J. C. (2007).  
632 Effects of forest fragmentation on the physiological stress response of black howler  
633 monkeys. *Animal Conservation*, 10(3), 374–379. doi:10.1111/j.1469-1795.2007.00122.x
- 634 Massot, M., Clobert, J., Pilorge, T., Lecomte, J., & Barbault, R. (1992). Density dependence  
635 in the common lizard: demographic consequences of a density manipulation. *Ecology*,  
636 73(5), 1742–1756. doi:10.2307/1940026
- 637 Mazerolle, M. J. (2016). AICcmodavg: Model selection and multimodel inference based on  
638 (Q)AIC(c). Retrieved from <http://cran.r-project.org/package=AICcmodavg>
- 639 McEwen, B. S., & Wingfield, J. C. (2010). What's in a name? Integrating homeostasis,  
640 allostasis and stress. *Hormones and Behavior*, 57(2), 105–111.  
641 doi:10.1016/j.yhbeh.2009.09.011
- 642 McKechnie, A. E., & Wolf, B. O. (2010). Climate change increases the likelihood of  
643 catastrophic avian mortality events during extreme heat waves. *Biology Letters*, 6(2),  
644 253–256. doi:10.1098/rsbl.2009.0702
- 645 Meehl, G. A., & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat  
646 waves in the 21st century. *Science*, 305(5686), 994–997. doi:10.1126/science.1098704
- 647 Meillère, A., Brischoux, F., Bustamante, P., Michaud, B., Parenteau, C., Marciau, C., &  
648 Angelier, F. (2016). Corticosterone levels in relation to trace element contamination  
649 along an urbanization gradient in the common blackbird (*Turdus merula*). *Science of the*  
650 *Total Environment*, 566–567, 93–101. doi:10.1016/j.scitotenv.2016.05.014
- 651 Meylan, S., Clobert, J., & Sinervo, B. (2007). Adaptive significance of maternal induction of

- density-dependent phenotypes. *Oikos*, 116, 650–661. doi:10.1111/j.2007.0030-1299.15432.x
- Meylan, S., Dufty, A. M., & Clobert, J. (2003). The effect of transdermal corticosterone application on plasma corticosterone levels in pregnant *Lacerta vivipara*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 134, 497–503. doi:10.1016/S1095-6433(02)00343-4
- Meylan, S., Miles, D. B., & Clobert, J. (2012). Hormonally mediated maternal effects, individual strategy and global change. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 367, 1647–1664. doi:10.1098/rstb.2012.0020
- Otovic, P., & Hutchinson, E. (2015). Limits to using HPA axis activity as an indication of animal welfare. *Altex*, 32(1), 41–50. doi:10.14573/altex.1406161
- Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P. W., Araújo, M. B., ... Robert, J. (2010). Scenarios for Global Biodiversity in the 21st Century. *Science*, 330, 1496–1501. doi:10.1126/science.1196624
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R, C. T. (2016). nlme: Linear and Nonlinear Mixed Effects Models. Retrieved from <http://cran.r-project.org/package=nlme>
- Preest, M. R., & Cree, A. (2008). Corticosterone treatment has subtle effects on thermoregulatory behavior and raises metabolic rate in the New Zealand common gecko, *Hoplodactylus maculatus*. *Physiological and Biochemical Zoology*, 81(5), 641–650. doi:10.1086/590371
- Qualls, C. P., & Andrews, R. M. (1999). Cold climates and the evolution of viviparity in reptiles: cold incubation temperatures produce poor-quality offspring in the lizard, *Sceloporus virgatus*. *Biological Journal of the Linnean Society*, 67, 353–376.

- 675 Romero, L. M., Dickens, M. J., & Cyr, N. E. (2009). The reactive scope model - a new model  
676 integrating homeostasis, allostasis, and stress. *Hormones and Behavior*, 55(3), 375–389.  
677 doi:10.1016/j.yhbeh.2008.12.009
- 678 Romero, L. M., & Wikelski, M. (2010). Stress physiology as a predictor of survival in  
679 Galapagos marine iguanas. *Proceedings of the Royal Society B: Biological Sciences*,  
680 277(1697), 3157–3162. doi:10.1098/rspb.2010.0678
- 681 Russo, S., Sillmann, J., & Fischer, E. M. (2015). Top ten European heatwaves since 1950 and  
682 their occurrence in the coming decades. *Environmental Research Letters*, 10(12),  
683 124003. doi:10.1088/1748-9326/10/12/124003
- 684 Rutschmann, A., Miles, D. B., Le Galliard, J. F., Richard, M., Moulherat, S., Sinervo, B., &  
685 Clobert, J. (2016). Climate and habitat interact to shape the thermal reaction norms of  
686 breeding phenology across lizard populations. *Journal of Animal Ecology*, 85(2), 457–  
687 466. doi:10.1111/1365-2656.12473
- 688 Sinervo, B., Mendez-de-la-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa  
689 Cruz, M., ... Sites, J. W. (2010). Erosion of lizard diversity by climate change and  
690 altered thermal niches. *Science*, 328, 894–899. doi:10.1126/science.1184695
- 691 Sørensen, J. G. (2010). Application of heat shock protein expression for detecting natural  
692 adaptation and exposure to stress in natural populations. *Current Zoology*, 56(6), 703–  
693 713.
- 694 Taylor, E. N., DeNardo, D. F., & Jennings, D. H. (2004). Seasonal steroid hormone levels and  
695 their relation to reproduction in the western diamond-backed rattlesnake, *Crotalus atrox*  
696 (Serpentes: Viperidae). *General and Comparative Endocrinology*, 136(3), 328–337.  
697 doi:10.1016/j.ygcen.2004.01.008

- 698 Telemeco, R. S., & Addis, E. A. (2014). Temperature has species-specific effects on  
699 corticosterone in alligator lizards. *General and Comparative Endocrinology*, 206(July),  
700 184–192. doi:10.1016/j.ygcen.2014.07.004
- 701 Telemeco, R. S., Fletcher, B., Levy, O., Riley, A., Rodriguez-Sanchez, Y., Smith, C., ...  
702 Buckley, L. B. (2017). Lizards fail to plastically adjust nesting behavior or thermal  
703 tolerance as needed to buffer populations from climate warming. *Global Change*  
704 *Biology*, 23(3), 1075–1084. doi:10.1111/gcb.13476
- 705 Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y.  
706 C., ... Williams, S. E. (2004). Extinction risk from climate change. *Nature*, 427(6970),  
707 145–148. doi:10.1038/nature02121
- 708 Wiens, J. J. (2016). Climate-related local extinctions are already widespread among plant and  
709 animal species. *PLoS Biology*, 14(12), e2001104. doi:10.1371/journal.pbio.2001104
- 710 Wikelski, M., & Cooke, S. J. (2006). Conservation physiology. *Trends in Ecology &*  
711 *Evolution*, 21(2), 38–46. doi:10.1016/j.tree.2005.10.018
- 712 Wikelski, M., Romero, L. M., & Snell, H. L. (2001). Marine iguanas oiled in the Galápagos.  
713 *Science*, 292(5516), 437–438. doi:10.1126/science.292.5516.437c
- 714 Wingfield, J. C. (2013). Ecological processes and the ecology of stress: the impacts of abiotic  
715 environmental factors. *Functional Ecology*, 27(1), 37–44. doi:10.1111/1365-2435.12039
- 716 Wingfield, J. C., Maney, D. L., Breuner, C. W., Jacobs, J. D., Lynn, S., Ramenofsky, M., &  
717 Richardson, R. D. (1998). Ecological bases of hormone-behavior interactions: the  
718 “emergency life history stage.” *American Zoologist*, 38, 191–206.  
719 doi:10.1093/icb/38.1.191
- 720

**Figure captions**

**Figure 1.** Relationships between baseline corticosterone levels and minimal daily ambient temperature ( $T_{\min}$ ) across 14 populations of common lizards. Baseline corticosterone decreases slightly with  $T_{\min}$  in a) adult females ( $n = 312$ ), b) yearling females ( $n = 87$ ), but not in c) adult males ( $n = 132$ ), and d) yearling males ( $n = 131$ ). For significant relationships, the predictions of the best model from Table S3 were fitted on the data (solid line) together with the 95% confidence interval (dashed lines). Note the logarithmic scale of corticosterone levels and wide range of variation within each population.

**Figure 2.** Thermal profile of the 8 populations from the longitudinal study during the heat waves (2015) and during the same period of a year with normal climatic conditions (2017). During the heat wave period, air temperatures measured in shade within vegetation approached or overpassed the maximal critical thermal limit (CT<sub>max</sub>) ranging from 37.7 to 42.6°C, thus highlighting a strong constraint for activity to avoid overhear (Sinervo et al., 2010).

**Figure 3.** Daily and seasonal variation in baseline corticosterone during a summer heat wave in yearling common lizards inside eight populations with periodic (closed circles) or permanent (filled circles) access to water. Baseline corticosterone levels a) non-linearly decreased within the day (time of day:  $F_{1,244} = 4.51$ ,  $p = 0.035$ , time of day<sup>2</sup>:  $F_{1,244} = 4.09$ ,  $p = 0.044$ ) and b) decreased during the heat wave in yearlings from populations with periodic access to water ( $t_{1,257} = -4.05$ ,  $p < 0.001$ ), while remaining similar in populations with permanent access to water ( $t_{1,257} = -0.81$ ,  $p = 0.417$ ).




Figure 1. Relationships between baseline corticosterone levels and minimal daily ambient temperature ( $T_{min}$ ) across 14 populations of common lizards. Baseline corticosterone decreases slightly with  $T_{min}$  in a) adult females ( $n = 312$ ), b) yearling females ( $n = 87$ ), but not in c) adult males ( $n = 132$ ), and d) yearling males ( $n = 131$ ). For significant relationships, the predictions of the best model from Table S3 were fitted on the data (solid line) together with the 95% confidence interval (dashed lines). Note the logarithmic scale of corticosterone levels and wide range of variation within each population.

279x361mm (300 x 300 DPI)




Figure 2. Thermal profile of the 8 populations from the longitudinal study during the heat waves (2015) and during the same period of a year with normal climatic conditions (2017). During the heat wave period, air temperatures measured in shade within vegetation approached or overpassed the maximal critical thermal limit (CTmax) ranging from 37.7 to 42.6°C, thus highlighting a strong constraint for activity to avoid overheat (Sinervo et al., 2010).

279x361mm (300 x 300 DPI)




Figure 3. Daily and seasonal variation in baseline corticosterone during a summer heat wave in yearling common lizards inside eight populations with periodic (closed circles) or permanent (filled circles) access to water. Baseline corticosterone levels a) non-linearly decreased within the day (time of day:  $F_{1,244} = 4.51$ ,  $p = 0.035$ , time of day<sup>2</sup>:  $F_{1,244} = 4.09$ ,  $p = 0.044$ ) and b) decreased during the heat wave in yearlings from populations with periodic access to water ( $t_{1,257} = -4.05$ ,  $p < 0.001$ ), while remaining similar in populations with permanent access to water ( $t_{1,257} = -0.81$ ,  $p = 0.417$ ).

279x361mm (300 x 300 DPI)

**Table S1.** Localisation (GPS coordinates), elevation, climatic conditions and water access in the focal populations of the common lizard (*Z. vivipara*). Values of environmental temperature (minimal:  $T_{\min}$  and maximal:  $T_{\max}$ ) are the mean  $\pm$  SE, which were recorded between June 29 and July 17, 2015. We used the relative change in abundance ( $r$ ) to assess the index of IUCN conservation status of each population [ $r = (\text{Abundance}^{2015} - \text{Abundance}^{2005}) / \text{Abundance}^{2005}$ ]<sup>1</sup>. See text for details.

<sup>1</sup> *IUCN Red List Categories and Criteria and Guidelines for Using the IUCN Red List Categories and Criteria. Scenario A2.* Endangered (EN): very high risk of extinction in the wild; Vulnerable (VU): high risk of extinction in the wild; Least Concern (LC): non-threatened populations.

Population		Coordinates	Altitude (m)	$T_{\min}$ (°C)	$T_{\max}$ (°C)	Water access	Abundance <sup>2005</sup>	Abundance <sup>2015</sup>	$r$	IUCN status
Barnesac	(BAS)	44°25'59.48"N 3°45'20.07"E	1515	9.04 ± 0.62	38.50 ± 0.62	Permanent	1,30	2,53	1,0	LC
Belair	(BEL)	44°40'20.79"N 4° 1'29.84"E	1418	8.80 ± 0.56	33.96 ± 0.49	Periodic	1,09	1,55	0,4	LC
M <sup>t</sup> Caroux	(CAR)	43°36'08.75"N 2°58'54.25"E	1047	11.39 ± 0.43	27.31 ± 0.50	Permanent	0,17	0,06	-0,6	EN
Carmantran	(CARM)	45°09'22.05"N 2°50'16.30"E	1267	9.68 ± 0.92	32.97 ± 0.79	Permanent	3,13	3,18	0,0	LC
Chalet du M <sup>t</sup> Lozère	(CHA)	44°26'56.72"N 3°44'51.79"E	1429	10.49 ± 0.77	35.63 ± 0.68	Periodic	0,38	0,39	0,0	LC
Col du cheval mort	(COM)	44°40'1.29"N 3°31'57.98"E	1405	10.20 ± 0.61	29.93 ± 0.68	Permanent	0,42	0,41	0,0	LC
Col du pendu	(COP)	44°39'18.38"N 4°01'49.79"E	1421	7.16 ± 0.43	44.74 ± 0.77	Permanent	0,53	0,75	0,4	LC
Source de la Loire	(JOC)	44°50'6.66"N 4°12'39.65"E	1296	8.77 ± 0.53	35.78 ± 0.61	Permanent	0,68	1,81	1,6	LC
Gerbier de jonc	(JON)	44°50'30.31"N 4°12'54.30"E	1398	10.06 ± 0.43	33.73 ± 0.68	Periodic	0,60	0,74	0,2	LC
Lajo	(LAJ)	44°50'41.10"N 3°25'51.01"E	1330	8.79 ± 0.46	39.04 ± 0.66	Periodic	1,55	1,64	0,1	LC
Montselgues	(MON)	44°30'40.26"N 4°00'29.36"E	1049	11.42 ± 0.42	33.13 ± 0.32	Permanent	0,27	0,09	-0,7	EN
Pejouzou	(PEJ)	45°09'52.24"N 2°50'37.52"E	1254	8.00 ± 0.95	39.19 ± 1.04	Permanent	0,92	1,02	0,1	LC
Puy Mary	(PUY)	45°6'25.77"N 2°41'6.10"E	1434	12.63 ± 0.60	32.55 ± 0.84	Permanent	0,07	0,05	-0,3	VU
Viala	(VIA)	44°20'17.43"N 3°46'04.76"E	1191	8.53 ± 0.52	35.45 ± 0.58	Permanent	0,50	0,66	0,3	LC

**Table S2.** Initial AICc based model selection comparing the effect of age, sex, population and their interactions on the plasma corticosterone levels in common lizards (n = 662). Models are compared to a null model (model 12) including the intercept only.

Model number	Specification of fixed effects	k	AICc	ΔAICc	$w_i$	Log likelihood	$r^2_m$
1	Age + Sex + Population + Age : Sex + Age : Population + Sex : Population	44	168.01	0.00	0.93	-36.80	39.48
2	Age + Sex + Population + Age : Sex + Age : Population + Sex : Population + Age : Sex : Population	57	173.18	5.17	0.07	-24.12	41.15
3	Age + Sex + Population + Age : Sex	18	184.23	16.22	0.00	-73.58	33.58
4	Age + Sex + Population + Sex : Population	30	203.93	35.92	0.00	-70.49	33.77
5	Age + Sex + Age : Sex	5	212.40	44.38	0.00	-101.15	28.31
6	Sex + Age + Population + Age : Population	30	214.57	46.56	0.00	-75.81	32.72
7	Sex + Population + Sex : Population	29	220.19	52.17	0.00	-79.72	31.98
8	Sex	3	257.85	89.84	0.00	-125.91	22.82
9	Age + Population + Age : Population	29	357.43	189.41	0.00	-148.34	16.85
10	Age	3	373.01	204.99	0.00	-183.49	8.18
11	Population	15	398.98	230.97	0.00	-184.12	7.87
12	<i>Null</i>	2	427.54	259.53	0.00	-211.76	0.00

k : number of parameters, ΔAICc: difference with AICc of the best model,  $w_i$ : model likelihood,  $r^2_m$  : marginal R-squared

**Table S3.** Initial AICc based model selection comparing the effects of sampling session, sex, population and their interaction in yearling common lizards (n = 371). Models are compared to a null model (model 12) including the intercept only.

Model number	Specification of fixed effects	k	AICc	$\Delta$ AICc	$w_i$	Log likelihood	$r^2_m$
1	Sex + Session + Session:Population	20	106.62	0.00	0.70	-31.63	25.80
2	Session + Sex + Population + Session:Sex + Session:Population + Sex:Population	28	108.40	1.78	0.29	-22.85	29.56
3	Session + Sex + Population + Sex:Population	20	115.46	8.84	0.01	-36.05	23.46
4	Session + Population + Session:Population	19	116.08	9.46	0.01	-37.53	22.74
5	Session + Sex + Population + Session:Sex + Session:Population + Sex:Population + Session:Sex:Population	35	119.52	12.90	0.00	-19.40	30.59
6	Session + Sex + Population + Session:Sex	14	119.63	13.01	0.00	-44.99	18.88
7	Sex + Population + Sex:Population	19	123.95	17.33	0.00	-41.46	20.57
8	Session + Sex + Session:Sex	7	128.09	21.47	0.00	-56.83	12.05
9	Session	5	132.09	25.47	0.00	-60.93	9.45
10	Population	11	132.76	26.15	0.00	-54.87	13.11
11	Sex	5	134.54	27.92	0.00	-62.16	8.64
12	Null	4	139.02	32.40	0.00	-65.43	6.45

k : number of parameters,  $\Delta$ AICc: difference with AICc of the best model,  $w_i$ : model likelihood,  $r^2_m$ : marginal R-squared

**Table S4. Principal component analysis (PCA) including the main determinants of population collapse.** Variables included in the PCA are the relative change in abundance ( $\Delta$ abundance), the minimal temperatures ( $T_{\min}$ ), and the altitude. Table shows the eigenvalue of the 3 axes (PC<sub>1</sub> to PC<sub>3</sub>), percentage of variance explained, and the inertia of each variables on the first two axes.

		PCA Axes		
		PC <sub>1</sub>	PC <sub>2</sub>	PC <sub>3</sub>
Eigenvalue		1.89	0.78	0.33
Variance		63.01	25.94	11.04
Inertia	$\Delta$ abundance	4273	54	-
	$T_{\min}$	3219	3733	-
	Altitude	2508	6213	-

**Table S5.** AICc based model selection comparing the effect of environmental conditions in early summer session on the plasma corticosterone levels in adult or yearling and female or male common lizards ( $n = 662$ ). Each environmental covariate was treated one by one (5 models per covariate). Environmental covariates were also computed in a principal component analysis to extract an integrative index of population extinction risk ( $PC_1$  – first axis of a principal component analysis including relative change in abundance,  $T_{\min}$ , and altitude, see Table S2). Population was treated as a random factor to account for non-independence.

Model number	Specification of fixed effects	k	AICc	$\Delta AICc$	$w_i$	Log likelihood	$r^2_m$	$r^2_c$
1	Age + Sex + $T_{\min}$ + Age : Sex + Age : $T_{\min}$ + Sex : $T_{\min}$	9	180.87	0.00	0.74	-81.30	30.79	33.77
2	Age + Sex + $PC_1$ + Age : Sex + Age : $PC_1$ + Sex : $PC_1$	9	185.02	4.16	0.09	-83.37	29.29	33.30
3	Age + Sex + $T_{\min}$ + Age : Sex + Sex : $T_{\min}$	8	185.14	4.27	0.09	-84.46	30.10	33.20
4	Age + Sex + water access + Age : Sex + Age : water access + Sex : water access	9	188.13	7.26	0.02	-84.93	28.68	33.01
5	Age + Sex + water access + Age : Sex + Age : water access	8	188.18	7.31	0.02	-85.98	28.44	32.83
6	Age + Sex + $T_{\min}$ + Age : Sex + Age : $T_{\min}$	8	189.72	8.85	0.01	-86.75	29.76	32.54
7	Age + Sex + $T_{\min}$ + Age : Sex	7	190.73	9.86	0.01	-88.28	29.39	32.28
8	Age + Sex + $PC_1$ + Age : Sex + Age : $PC_1$	8	191.24	10.37	0.00	-87.51	28.43	32.42
9	Age + Sex + $\Delta$ abundance + Age : Sex + Age : $\Delta$ abundance + Sex : $\Delta$ abundance	9	191.94	11.07	0.00	-86.83	28.35	32.42
10	Age + Sex + $T_{\max}$ + Age : Sex + Age : $T_{\max}$ + Sex : $T_{\max}$	9	191.94	11.08	0.00	-86.83	28.82	32.61
11	Age + Sex + Age : Sex	6	192.84	11.97	0.00	-90.35	27.53	31.90
12	Age + Sex + $\Delta$ abundance + Age : Sex + Age : $\Delta$ abundance	8	192.95	12.08	0.00	-88.37	28.05	32.06
13	Age + Sex + altitude + Age : Sex + Age : altitude	8	193.02	12.15	0.00	-88.40	28.04	32.06
14	Age + Sex + $T_{\max}$ + Age : Sex + Sex : $T_{\max}$	8	193.11	12.25	0.00	-88.45	28.49	32.26

15	Age + Sex + PC <sub>1</sub> + Age : Sex + Sex : PC <sub>1</sub>	8	193.19	12.33	0.00	-88.49	28.13	32.47
16	Age + Sex + T <sub>max</sub> + Age : Sex + Age : T <sub>max</sub>	8	193.21	12.35	0.00	-88.50	28.50	32.25
17	Age + Sex + T <sub>max</sub> + Age : Sex	7	193.28	12.41	0.00	-89.55	28.27	32.02
18	Age + Sex + altitude + Age : Sex + Age : altitude + Sex : altitude	9	193.37	12.50	0.00	-87.55	28.24	32.21
19	Age + Sex + PC <sub>1</sub> + Age : Sex	7	194.55	13.69	0.00	-90.19	27.79	32.04
20	Age + Sex + altitude + Age : Sex	7	194.80	13.93	0.00	-90.31	27.53	31.85
21	Age + Sex + Δabundance + Age : Sex	7	194.83	13.96	0.00	-90.33	27.51	31.84
22	Age + Sex + water access + Age : Sex	7	194.87	14.00	0.00	-90.35	27.52	32.88
23	Age + Sex + Δabundance + Age : Sex + Sex : Δabundance	8	196.01	15.14	0.00	-89.90	27.57	32.00
24	Age + Sex + altitude + Age : Sex + Sex : altitude	8	196.59	15.73	0.00	-90.19	27.56	31.88
25	Age + Sex + water access + Age : Sex + Sex : water access	8	196.60	15.74	0.00	-90.19	27.56	31.90
26	Age + Sex	5	221.76	40.89	0.00	-105.83	24.36	28.59
27	Sex	4	238.51	57.65	0.00	-115.23	22.08	26.74
28	Age	4	357.56	176.70	0.00	-174.75	7.46	12.38
29	T <sub>min</sub>	4	405.79	224.92	0.00	-198.86	2.19	6.17
30	Null	3	408.20	227.33	0.00	-201.08	0.00	6.18
31	T <sub>max</sub>	4	408.41	227.54	0.00	-200.17	0.87	6.07
32	PC <sub>1</sub>	4	409.35	228.48	0.00	-200.64	0.47	6.18
33	Δabundance	4	410.17	229.30	0.00	-201.05	0.03	6.19
34	water access	4	410.17	229.31	0.00	-201.06	0.03	6.18

---

35	altitude	4	410.22	229.35	0.00	-201.08	0.00	6.18
----	----------	---	--------	--------	------	---------	------	------

---

k : number of parameters,  $\Delta\text{AICc}$ : difference with AICc of the best model,  $w_i$ : model likelihood,  
 $r^2_{\text{m}}$  : marginal R-squared,  $r^2_{\text{c}}$  : conditional R-squared

**Table S6.** AICc based model selection comparing the effects of environmental conditions during the heat wave period on plasma corticosterone levels in yearling common lizards (n = 371). Each environmental covariate was treated one by one (5 models per covariate). Environmental covariates were also computed in a principal component analysis to extract an integrative index of population extinction risk (PC<sub>1</sub> – first axis of a principal component analysis including relative change in abundance,  $T_{\min}$ , and altitude, see Table S2). Population was treated as a random factor to account for non-independence.

Model number	Specification of fixed effects	k	AICc	$\Delta$ AICc	$w_i$	Log likelihood	$r^2_m$	$r^2_c$
1	Session + Sex + water access + Session : water access + Sex : water access	10	117.95	0.00	0.24	-48.55	16.43	21.82
2	Session + Sex + water access + Session : water access	9	119.55	1.61	0.11	-50.43	15.31	20.57
3	Session + Sex + $T_{\min}$	8	120.26	2.31	0.08	-51.85	15.94	17.89
4	Session + Sex + $\Delta$ abundance + Sex : $\Delta$ abundance	9	120.45	2.50	0.07	-50.88	15.62	19.68
5	Session + Sex + PC <sub>1</sub> + Sex : PC <sub>1</sub>	9	120.91	2.96	0.06	-51.11	15.71	18.82
6	Session + Sex + $\Delta$ abundance + Session : $\Delta$ abundance + Sex : $\Delta$ abundance	10	121.03	3.08	0.05	-50.09	15.86	19.90
7	Session + Sex + PC <sub>1</sub> + Session : PC <sub>1</sub> + Sex : PC <sub>1</sub>	10	121.07	3.12	0.05	-50.11	16.12	19.19
8	Session + Sex + $T_{\min}$ + Sex : $T_{\min}$	9	121.47	3.52	0.04	-51.39	16.20	18.13
9	Session + Sex + $T_{\min}$ + Session : $T_{\min}$	9	121.65	3.70	0.04	-51.48	16.07	17.96
10	Session + Sex + PC <sub>1</sub> + Session : PC <sub>1</sub>	9	121.97	4.02	0.03	-51.64	15.18	18.41
11	Session + Sex + PC <sub>1</sub>	8	122.17	4.22	0.03	-52.81	14.69	17.98
12	Session + Sex + $T_{\max}$	8	122.27	4.32	0.03	-52.86	14.74	18.05
13	Session + Sex	7	122.29	4.34	0.03	-53.93	13.55	18.68
14	Session + Sex + water access + Sex : water access	9	122.51	4.56	0.02	-51.91	14.82	20.08
15	Session + Sex + $T_{\min}$ + Session : $T_{\min}$ + Sex : $T_{\min}$	10	122.66	4.71	0.02	-50.90	16.39	18.26
16	Session + Sex + $\Delta$ abundance + Session : $\Delta$ abundance	9	123.14	5.19	0.02	-52.22	14.52	18.72

17	Session + Sex + $\Delta$ abundance	8	123.36	5.42	0.02	-53.41	14.08	18.32
18	Session + Sex + altitude	8	123.70	5.75	0.01	-53.57	14.44	18.74
19	Session + Sex + $T_{\max}$ + Session : $T_{\max}$	9	124.35	6.40	0.01	-52.83	14.79	18.12
20	Session + Sex + water access	8	124.39	6.44	0.01	-53.92	13.58	18.70
21	Session + Sex + $T_{\max}$ + Sex : $T_{\max}$	9	124.41	6.46	0.01	-52.86	14.74	18.05
22	Session + Sex + altitude + Sex : altitude	9	124.63	6.68	0.01	-52.97	14.81	19.12
23	Session + Sex + altitude + Session : altitude	9	125.75	7.80	0.00	-53.53	14.39	18.57
24	Session + Sex + $T_{\max}$ + Session : $T_{\max}$ + Sex : $T_{\max}$	10	126.50	8.56	0.00	-52.83	14.79	18.11
25	Session + Sex + altitude + Session : altitude + Sex : altitude	10	126.77	8.83	0.00	-52.96	14.79	19.04
26	Session	6	129.21	11.26	0.00	-58.45	10.46	14.91
27	Sex	6	130.54	12.59	0.00	-59.11	11.09	16.32
28	$T_{\min}$	6	135.22	17.27	0.00	-61.45	10.00	11.85
29	<i>Null</i>	5	136.48	18.53	0.00	-63.13	8.06	12.51
30	PC <sub>1</sub>	6	136.75	18.80	0.00	-62.22	9.02	12.04
31	$T_{\max}$	6	136.91	18.97	0.00	-62.30	9.05	12.12
32	$\Delta$ abundance	6	137.79	19.84	0.00	-62.74	8.52	12.34
33	altitude	6	138.12	20.18	0.00	-62.90	8.74	12.63
34	water access	6	138.57	20.62	0.00	-63.12	8.04	12.48

k : number of parameters,  $\Delta$ AICc: difference with AICc of the best model,  $w_i$ : model likelihood,  $r^2_m$ : marginal R-squared,  $r^2_c$ : conditional R-squared