



HAL
open science

Complete Genome Sequence of the *Silicimonas algicola* Type Strain, a Representative of the Marine Roseobacter Group Isolated from the Cell Surface of the Marine Diatom *Thalassiosira delicatula*

Klervi Crenn, Boyke Bunk, Cathrin Spröer, Jörg Overmann, Christian Jeanthon

► To cite this version:

Klervi Crenn, Boyke Bunk, Cathrin Spröer, Jörg Overmann, Christian Jeanthon. Complete Genome Sequence of the *Silicimonas algicola* Type Strain, a Representative of the Marine Roseobacter Group Isolated from the Cell Surface of the Marine Diatom *Thalassiosira delicatula*. *Microbiology Resource Announcements*, 2019, 8 (9), 10.1128/MRA.00108-19 . hal-02130564

HAL Id: hal-02130564

<https://hal.science/hal-02130564>

Submitted on 15 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Complete Genome Sequence of the *Silicimonas algicola* Type Strain, a Representative of the Marine *Roseobacter* Group Isolated from the Cell Surface of the Marine Diatom *Thalassiosira delicatula*

Klervi Crenn,^{a*} Boyke Bunk,^b Cathrin Spröer,^b Jörg Overmann,^{b,c} Christian Jeanthon^a

^aCNRS & Sorbonne Université, Station Biologique de Roscoff, Adaptation et Diversité en Milieu Marin, Team ECOMAP-Ecology of Marine Plankton, Roscoff, France

^bLeibniz Institute DSMZ—German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany

^cMicrobiology, Braunschweig University of Technology, Braunschweig, Germany

ABSTRACT *Silicimonas algicola* strain KC90B^T is an alphaproteobacterium of the *Roseobacter* clade that was isolated from a culture of the marine diatom *Thalassiosira delicatula*. Here, we report the complete genome sequence of this type strain, which is 4,351,658 bp in size with 4,272 coding sequences and an average G+C content of 65.2%.

Members of the *Roseobacter* clade (*Alphaproteobacteria*) are often dominant in natural assemblages with marine algae and are often found in laboratory cultures of marine phytoplankton (1–3). *Silicimonas algicola* strain KC90B^T (= DSM 103371T = RCC 4681^T) was isolated from the cell surface of the marine diatom *Thalassiosira delicatula* RCC 2560 (4) in July 2013. This microalgal culture isolated from the coastal long-term monitoring station SOMLIT-Astan (north of Roscoff, France, in the western English Channel) has been maintained in the RCC since its isolation in January 2011.

Cells were grown in modified marine broth (2.5 g peptone, 0.5 g yeast extract, and 35 g sea salts dissolved in 1 liter of Milli-Q water) at 20°C, and 500 mg was harvested after 15 days. Genomic DNA was isolated using Qiagen Genomic-tip 100/G (Qiagen, Hilden, Germany) according to the manufacturer's instructions. For long read sequencing, a SMRTbell template library was prepared following the procedure and checklist—greater than 10 kb template preparation (Pacific Biosciences, Menlo Park, USA). Briefly, 8 μg genomic DNA was sheared for preparation of 15-kb libraries. DNA was end-repaired and ligated overnight to hairpin adapters applying components from the DNA/polymerase binding kit P6 (Pacific Biosciences). BluePippin size selection to greater than 4 kb was performed (Sage Science, Beverly, USA). Conditions for annealing of sequencing primers and binding of polymerase to a purified SMRTbell template were assessed with the calculator in RS Remote (Pacific Biosciences). Single-molecule real-time (SMRT) sequencing was carried out on the PacBio RS II system (Pacific Biosciences). SMRT sequencing revealed a total of 69,135 reads with a mean read length of 14,050 bp and an N_{50} value of 17,953 bp. From the same batch of DNA, a short insert library was created using the NEBNext ultra DNA library prep kit for Illumina (NEB, Ipswich, USA). Short-read sequencing was carried out on an Illumina HiSeq 2500 platform resulting in 4,006,700 paired-end reads of 2×101 bp.

Genome assembly was performed by applying the Hierarchical Genome Assembly Process version 3 (RS_HGAP3) protocol included in SMRT Portal 2.3.0 using default parameters. The assembly revealed a single circular chromosome with a coverage of 146×. The chromosome was circularized, and artificial redundancies at the ends of the contigs were removed and adjusted to *dnaA* as the first gene. Error correction was

Citation Crenn K, Bunk B, Spröer C, Overmann J, Jeanthon C. 2019. Complete genome sequence of the *Silicimonas algicola* type strain, a representative of the marine *Roseobacter* group isolated from the cell surface of the marine diatom *Thalassiosira delicatula*. *Microbiol Resour Announc* 8:e00108-19. <https://doi.org/10.1128/MRA.00108-19>.

Editor J. Cameron Thrash, University of Southern California

Copyright © 2019 Crenn et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christian Jeanthon, jeanthon@sb-roscoff.fr.

* Present address: Klervi Crenn, Immunrise Biocontrol, Pessac, France.

Received 29 January 2019

Accepted 8 February 2019

Published 28 February 2019

performed by a mapping of Illumina short reads onto the finished genome sequence using Burrows-Wheeler Alignment (BWA) 0.6.2 in paired-end (sample) mode using default settings (5) with subsequent variant and consensus calling using VarScan 2.3.6 (6). A consensus concordance of quality value (QV) 60 was reached. Automated genome annotation was carried out using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (7).

The complete genome sequence of *Silicimonas algicola* strain KC90B^T consists of a single circular chromosome with 4,351,658 bp and a G+C content of 65.2%. The NCBI PGAP predicted 4,272 coding sequences, 42 tRNA genes, and 1 rRNA operon. The genome contains biogeochemically relevant pathways previously reported for roseobacters, such as carbon monoxide oxidation (*cox*), sulfur oxidation (*sox*), and phosphonate utilization (*phn*) genes (8). It also shows a prevalence of genes involved in environmental stress response and detoxification, tripartite ATP-independent periplasmic (TRAP) and ABC transporters, and carbohydrate- and electron-accepting reactions. The genome annotation also revealed the presence of quorum-sensing genes, such as an *N*-acyl-L-homoserine lactone synthetase (*luxI*) and a regulator protein belonging to the *luxR* family on the same operon (9).

Data availability. The complete genome sequence of *S. algicola* strain KC90^T (= DSM 103371T = RCC 4681^T) has been deposited in the NCBI GenBank under the accession no. CP034588 (BioProject no. PRJNA504651, BioSample no. SAMN10396653). The version described in this paper is the first version, CP034588.1. Raw sequence reads have been submitted to the NCBI SRA under the accession no. SRR8529664 (PacBio) and SRR8529665 (Illumina).

ACKNOWLEDGMENTS

This work was supported by the French national program EC2CO-Microbien (project MICROMAR) and the MaCuMBA project funded by the European Union's Seventh Framework Programme (grant agreement no. 311975). Klervi Crenn received a doctoral grant funded by Région Bretagne and CNRS.

We thank Simone Severitt and Nicole Heyer for excellent technical assistance.

REFERENCES

- Buchan A, González JM, Moran MA. 2005. Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* 71:5665–5677. <https://doi.org/10.1128/AEM.71.10.5665-5677.2005>.
- Luo H, Moran MA. 2014. Evolutionary ecology of the marine *Roseobacter* clade. *Microbiol Mol Biol Rev* 78:573–587. <https://doi.org/10.1128/MMBR.00020-14>.
- Amin SA, Parker MS, Armbrust EV. 2012. Interactions between diatoms and bacteria. *Microbiol Mol Biol Rev* 76:667–684. <https://doi.org/10.1128/MMBR.00007-12>.
- Crenn K, Serpin D, Lepleux C, Overmann J, Jeanthon C. 2016. *Silicimonas algicola* gen. nov., sp. nov., a novel member of the *Roseobacter* clade isolated from the cell surface of the marine diatom *Thalassiosira delicatula*. *Int J Syst Evol Microbiol* 66:4580–4588. <https://doi.org/10.1099/ijsem.0.001394>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
- Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* 22:568–576. <https://doi.org/10.1101/gr.129684.111>.
- Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Moran MA, Belas R, Schell MA, González JM, Sun F, Sun S, Binder BJ, Edmonds J, Ye W, Orcutt B, Howard EC, Meile C, Palefsky W, Goesmann A, Ren Q, Paulsen I, Ulrich LE, Thompson LS, Saunders E, Buchan A. 2007. Ecological genomics of marine roseobacters. *Appl Environ Microbiol* 73:4559–4569. <https://doi.org/10.1128/AEM.02580-06>.
- Fuqua WC, Winans SC, Greenberg EP. 1994. Quorum sensing in bacteria: the luxR-luxI family of cell density-responsive transcriptional regulators. *J Bacteriol* 176:269–275. <https://doi.org/10.1128/jb.176.2.269-275.1994>.