



HAL
open science

Koszul calculus of preprojective algebras

Roland Berger, Rachel Taillefer

► **To cite this version:**

Roland Berger, Rachel Taillefer. Koszul calculus of preprojective algebras. Journal of the London Mathematical Society, 2020, Journal of the London Mathematical Society, 102 (3), pp.1241-1292. 10.1112/jlms.12362 . hal-02132927v2

HAL Id: hal-02132927

<https://hal.science/hal-02132927v2>

Submitted on 17 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Koszul calculus of preprojective algebras

Roland Berger and Rachel Taillefer

ABSTRACT

We show that the Koszul calculus of a preprojective algebra, whose graph is distinct from A_1 and A_2 , vanishes in any (co)homological degree $p > 2$. Moreover, its (higher) cohomological calculus is isomorphic as a bimodule to its (higher) homological calculus, by exchanging degrees p and $2 - p$, and we prove a generalised version of the 2-Calabi-Yau property. For the ADE Dynkin graphs, the preprojective algebras are not Koszul and they are not Calabi-Yau in the sense of Ginzburg's definition, but they satisfy our generalised Calabi-Yau property and we say that they are Koszul complex Calabi-Yau (Kc-Calabi-Yau) of dimension 2. For Kc-Calabi-Yau (quadratic) algebras of any dimension, defined in terms of derived categories, we prove a Poincaré Van den Bergh duality theorem. We compute explicitly the Koszul calculus of preprojective algebras for the ADE Dynkin graphs.

1. Introduction

Preprojective algebras are quiver algebras with quadratic relations, that play an important role in the representation theory of quiver algebras [27, 17, 1, 10], with various applications [15, 16] and many developments [14, 26, 8, 29]. In [29], the reader will find an introduction to the various aspects of the preprojective algebras in representation theory, with an extended bibliography. In our paper, we are interested in some homological properties linked to Hochschild cohomology.

In the last two decades, the Hochschild cohomology of preprojective algebras, as well as some extra algebraic structures, have been computed in several steps, as follows.

- 1) Erdmann and Snashall [18, 19] determined the Hochschild cohomology and its cup-product in type A.
- 2) Crawley-Boevey, Etingof and Ginzburg [14] determined the Hochschild cohomology for all preprojective algebras of non-Dynkin type (which are Koszul in this case [33, 30, 10]).
- 3) In type DE and characteristic zero, Etingof and Eu [21] determined the Hochschild cohomology and Eu [22] the cup product. The cyclic homology was computed in type ADE in [21].
- 4) Assembling and completing the previous results in characteristic zero, Eu gave an explicit description of the Tamarkin-Tsygan calculus [39] of the preprojective algebras in type ADE, that is, the homology and the cohomology, the cup product, the contraction map and the Lie derivative, the Connes differential and the Gerstenhaber bracket [23].
- 5) Eu and Schedler extended the ADE results to the case where the base ring is \mathbb{Z} , and obtained the corresponding ADE results in any characteristic [24].

In [7], a Koszul calculus was associated with any quadratic algebra over a field, in order to produce new homological invariants for non-Koszul quadratic algebras. We begin this paper by extending the Koszul calculus to quadratic quiver algebras. We shall compute the Koszul calculus of the preprojective algebras whose graphs are Dynkin of type ADE (the preprojective

algebras are then finite dimensional). Except for types A_1 and A_2 , these quadratic quiver algebras are not Koszul [33, 30, 10], so that the Koszul calculus and the Hochschild calculus provide different information.

Before presenting our computations, we state and develop a Poincaré Van den Bergh duality theorem [41] for the Koszul homology/cohomology of *any preprojective algebra* whose graph is different from A_1 and A_2 . This theorem is formulated as follows and constitutes the first main result of the present paper. The duality is precisely part (ii) in this theorem.

THEOREM 1.1. *Let A be the preprojective algebra of a (non-labelled) connected graph Δ distinct from A_1 and A_2 , over a field \mathbb{F} . Let M be an A -bimodule.*

- (i) *The Koszul bimodule complex $K(A)$ of A has length 2. In particular, $\mathrm{HK}^p(A, M) = \mathrm{HK}_p(A, M) = 0$ for any $p > 2$.*
- (ii) *The $\mathrm{HK}^\bullet(A)$ -bimodules $\mathrm{HK}^\bullet(A, M)$ and $\mathrm{HK}_{2-\bullet}(A, M)$ are isomorphic.*
- (iii) *The $\mathrm{HK}_{hi}^\bullet(A)$ -bimodules $\mathrm{HK}_{hi}^\bullet(A, M)$ and $\mathrm{HK}_{2-\bullet}^{hi}(A, M)$ are isomorphic.*

In this statement, following [7], $\mathrm{HK}_p(A, M)$ and $\mathrm{HK}^p(A, M)$ denote the Koszul homology and cohomology spaces with coefficients in M , while $\mathrm{HK}_p^{hi}(A, M)$ and $\mathrm{HK}_{hi}^p(A, M)$ denote the higher Koszul homology and cohomology spaces. When $M = A$, these notations are simplified into $\mathrm{HK}_p(A)$, $\mathrm{HK}^p(A)$, $\mathrm{HK}_p^{hi}(A)$ and $\mathrm{HK}_{hi}^p(A)$.

In the general setting [7], the Koszul calculus of a quadratic algebra A consists of the graded associative algebra $\mathrm{HK}^\bullet(A)$ endowed with the Koszul cup product and, for all A -bimodules M , of the graded $\mathrm{HK}^\bullet(A)$ -bimodules $\mathrm{HK}^\bullet(A, M)$ and $\mathrm{HK}_\bullet(A, M)$, with actions respectively defined by the Koszul cup and cap products. The higher Koszul calculus of A is given by the analogous data, adding the subscript and superscript *hi*. Sometimes (as will be the case with our computations in ADE types), these calculi are *restricted*, meaning that the data is limited to $M = A$, so that the restricted Koszul calculus consists of the graded associative algebra $\mathrm{HK}^\bullet(A)$ and of the graded $\mathrm{HK}^\bullet(A)$ -bimodule $\mathrm{HK}_\bullet(A)$ – similarly for the higher version.

Using Theorem 1.1 for Δ Dynkin of type ADE, we shall deduce the (higher) homological restricted Koszul calculus from the computation of the (higher) cohomological restricted Koszul calculus.

Part (ii) in Theorem 1.1 comes from an explicit isomorphism from the complex C_1 of Koszul cochains with coefficients in M , whose p th cohomology is $\mathrm{HK}^p(A, M)$, to the complex C_2 of Koszul chains with coefficients in M , whose p th homology is $\mathrm{HK}_{2-p}(A, M)$, described as follows.

PROPOSITION 1.2. *Let A be the preprojective algebra of a connected graph Δ distinct from A_1 and A_2 . Let M be an A -bimodule. The Koszul cup and cap products are denoted by $\underset{K}{\smile}$ and $\underset{K}{\frown}$. Define $\omega_0 = \sum_i e_i \otimes \sigma_i$, where the sum runs over the vertices i of Δ and, for each vertex i , e_i is the idempotent and σ_i is the quadratic relation in A associated with i .*

For each Koszul p -cochain f with coefficients in M , we define the Koszul $(2 - p)$ -chain $\theta_M(f)$ with coefficients in M by

$$\theta_M(f) = \omega_0 \underset{K}{\frown} f.$$

Then $\theta_M : C_1 \rightarrow C_2$ is an isomorphism of complexes. Moreover, the equalities

$$\theta_{M \otimes_A N}(f \underset{K}{\smile} g) = \theta_M(f) \underset{K}{\frown} g = f \underset{K}{\frown} \theta_N(g)$$

hold for any Koszul cochains f and g with coefficients in bimodules M and N respectively.

The proof of Proposition 1.2 relies on some manipulations of the defining formula of θ_M with fundamental formulas of Koszul calculus [7], using actions involving $\underset{K}{\smile}$ and $\underset{K}{\frown}$. The fundamental formulas of Koszul calculus express the differential b_K of C_1 and the differential b^K of C_2 respectively as a cup bracket and a cap bracket, namely

$$b_K = -[\epsilon_A, -]_{\underset{K}{\smile}}, \quad b^K = -[\epsilon_A, -]_{\underset{K}{\frown}},$$

where $\epsilon_A : V \rightarrow A$ is a fundamental Koszul 1-cocycle defined on the arrow space V by $\epsilon_A(x) = x$ for all $x \in V$.

In order to extract a generalised version of the 2-Calabi-Yau property from our Poincaré Van den Bergh duality (Theorem 1.1) for quadratic algebras, we apply this theorem to the left A^e -module $M = A^e := A \otimes A^{op}$ viewed as an A -bimodule. We show that the complex of Koszul chains with coefficients in the left A^e -module A^e is naturally isomorphic, as a right A^e -module, to the Koszul bimodule complex $K(A)$. Using the fact that the homology of $K(A)$ is isomorphic to A in degree 0, and to 0 in degree 1, we obtain a generalisation of the 2-Calabi-Yau property, formulated as follows.

THEOREM 1.3. *Let A be the preprojective algebra of a connected graph Δ distinct from A_1 and A_2 , over a field \mathbb{F} . Let us denote by $K(A)$ the Koszul bimodule complex of A . Then the A -bimodule $\text{HK}^p(A, A^e)$ is isomorphic to the A -bimodule $H_{2-p}(K(A))$ for $0 \leq p \leq 2$. In particular, we have the following.*

- (i) *The A -bimodule $\text{HK}^2(A, A^e)$ is isomorphic to the A -bimodule A .*
- (ii) *$\text{HK}^1(A, A^e) = 0$.*
- (iii) *The A -bimodule $\text{HK}^0(A, A^e)$ is isomorphic to the A -bimodule $H_2(K(A))$, which is always non-zero when Δ is Dynkin of type ADE.*

We then say that the preprojective algebra A is a *Koszul complex Calabi-Yau algebra of dimension 2*. We generalise this definition to any quadratic algebra and any dimension n in Definition 1.4 below, better formulated in terms of derived categories. Since there is an \mathbb{F} -linear isomorphism

$$H(\theta_M) = \bar{\omega}_0 \underset{K}{\frown} - : \text{HK}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}(A, M),$$

we say that the class $\bar{\omega}_0 \in \text{HK}_2(A)$ is the *fundamental class* of the Koszul complex Calabi-Yau algebra A , by analogy with Poincaré’s duality in singular homology/cohomology [31]. In Definition 1.7, we give a stronger version of Definition 1.4 in order to obtain a Poincaré-like duality, that is, a duality isomorphism expressed as a cap action by a suitably defined fundamental class.

Let us remark that the A -bimodule structures in Theorem 1.3 are compatible with the Koszul cup and cap actions of $\text{HK}^\bullet(A)$ on $\text{HK}^\bullet(A, A^e)$ and $H(K(A))$. These actions can be viewed as graded actions of left $\text{HK}^\bullet(A)^e$ -modules, while the A -bimodules can be viewed as compatible right A^e -modules. So the isomorphism $\text{HK}^\bullet(A, A^e) \cong H_{2-\bullet}(K(A))$ in Theorem 1.3 is an isomorphism of graded $\text{HK}^\bullet(A)^e$ - A^e -bimodules. This enriched isomorphism is the expression of the stronger version of the Koszul complex Calabi-Yau property, as we shall see in Definition 1.7.

Note that if Δ is not Dynkin ADE, then A is Koszul, so Theorem 1.3 enables us to recover the well-known result that A is 2-Calabi-Yau in the sense of Ginzburg [14, 9]. However, if Δ is Dynkin ADE, then A is not homologically smooth since its minimal projective resolution has infinite length, so that Ginzburg’s definition of Calabi-Yau algebras cannot be applied in this case [28]. Moreover, the restricted Hochschild calculus is drastically different from the

restricted Koszul calculus, because by [20] there is a cohomological Hochschild periodicity

$$\mathrm{HH}^{p+6}(A) \cong \mathrm{HH}^p(A), \quad p > 0$$

and, consequently, there are non-zero spaces $\mathrm{HH}^p(A)$ for infinitely many values of p . Even taking into account this 6-periodicity, the list of cohomological Koszul invariants consists only of $\mathrm{HK}^0(A)$, $\mathrm{HK}^1(A)$ and $\mathrm{HK}^2(A)$ and is therefore shorter than the list of Hochschild invariants.

In [24], Eu and Schedler define periodic Calabi-Yau Frobenius algebras, for finite dimensional algebras only. Their main example is given by the preprojective algebras of Dynkin ADE graphs [24, Example 2.3.10]. Then the above cohomological Hochschild periodicity is a part of remarkable isomorphisms in Hochschild calculus for any periodic Calabi-Yau Frobenius algebra [24, Theorem 2.3.27 and Theorem 2.3.47].

From Theorem 1.3, we are led to introduce a general definition.

DEFINITION 1.4. Let $\mathcal{Q} = (\mathcal{Q}_0, \mathcal{Q}_1)$ be a finite quiver, and let \mathbb{F} be a field. Let A be an \mathbb{F} -algebra defined on the path algebra $\mathbb{F}\mathcal{Q}$ of \mathcal{Q} by homogeneous quadratic relations. Define the ring $k = \mathbb{F}\mathcal{Q}_0$, so that A is regarded as a quadratic k -algebra. We say that A is Koszul complex Calabi-Yau (Kc-Calabi-Yau) of dimension n , for an integer $n \geq 0$, if

- (i) the bimodule Koszul complex $K(A)$ of A has length n , and
- (ii) $\mathrm{RHom}_{A^e}(K(A), A^e) \cong K(A)[-n]$ in the bounded derived category of A -bimodules.

In our context (that of quadratic algebras), Definition 1.4 is a definition of a new Calabi-Yau property, valid whether A is finite dimensional or not. In this definition, we do not impose that $K(A)$ be a resolution of A , that is, A is not necessarily Koszul, meaning that the bimodules $\mathrm{HK}^p(A, A^e)$ for $0 \leq p \leq n - 2$ may be non-zero. Under the assumptions of Definition 1.4, we verify that, if A is Koszul, Definition 1.4 is equivalent to Ginzburg's definition of n -Calabi-Yau algebras [28, 43]. We then prove a new Poincaré Van den Bergh duality for Kc-Calabi-Yau algebras, adapted to Koszul (co)homologies.

THEOREM 1.5. *Let A be a Kc-Calabi-Yau algebra of dimension n . Then for any A -bimodule M , the \mathbb{F} -vector spaces $\mathrm{HK}^p(A, M)$ and $\mathrm{HK}_{n-p}(A, M)$ are isomorphic.*

DEFINITION 1.6. Let A be a Kc-Calabi-Yau algebra of dimension n . The image $c \in \mathrm{HK}_n(A)$ of the unit 1 of the algebra A under the isomorphism $\mathrm{HK}^0(A) \cong \mathrm{HK}_n(A)$ in Theorem 1.5 is called the fundamental class of the Kc-Calabi-Yau algebra A .

In order to describe the duality isomorphism of Theorem 1.5 explicitly as a cap-product by the fundamental class for strong Kc-Calabi-Yau algebras, we shall use derived categories in the general context of DG algebras, as presented and detailed in the preprint book by Yekutieli [45]. Let us present briefly what we need in this general context.

We introduce the DG algebra $\tilde{A} = \mathrm{Hom}_{A^e}(K(A), A)$. The complexes $K(A)$ and $\mathrm{Hom}_{\tilde{A}^e}(K(A), A^e)$ of A -bimodules have an enriched structure since they can be viewed as DG \tilde{A} -bimodules in the abelian category $A\text{-Bimod}$ of A -bimodules, in the sense of [45].

Denote by $\mathcal{C}(\tilde{A}, A\text{-Bimod})$ the category of DG \tilde{A} -bimodules in $A\text{-Bimod}$ [45]. Let M be an A -bimodule. For any chain DG \tilde{A} -bimodule C in $A\text{-Bimod}$, $\mathrm{Hom}_{A^e}(C, M)$ is a cochain DG \tilde{A} -bimodule in the abelian category $\mathrm{Vect}_{\mathbb{F}}$ of \mathbb{F} -vector spaces (in $A\text{-Bimod}$ when $M = A^e$). For any cochain DG \tilde{A} -bimodule C' in $A\text{-Bimod}$, $M \otimes_{A^e} C'$ is a cochain DG \tilde{A} -bimodule in $\mathrm{Vect}_{\mathbb{F}}$. The bounded derived categories $\mathcal{D}^b(\tilde{A}, A\text{-Bimod})$ and $\mathcal{D}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ are defined in [45]. However we

do not know if the functors $\text{Hom}_{A^e}(-, M)$ and $M \otimes_{A^e} -$ from $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ to $\mathcal{C}^b(\tilde{A}, \text{Vect}_{\mathbb{F}})$ are derivable.

DEFINITION 1.7. Let A be a Kc-Calabi-Yau algebra of dimension n . Then A is said to be strong Kc-Calabi-Yau if the derived functor of the endofunctor $\text{Hom}_{A^e}(-, A^e)$ of $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ exists and if $\text{RHom}_{A^e}(K(A), A^e) \cong K(A)[-n]$ in the bounded derived category $\mathcal{D}^b(\tilde{A}, A\text{-Bimod})$.

THEOREM 1.8. Let A be a Kc-Calabi-Yau algebra of dimension n and let c be its fundamental class. We assume that A is strong Kc-Calabi-Yau and that the derived functors of the functors $\text{Hom}_{A^e}(-, A)$ and $A \otimes_{A^e} -$ from $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ to $\mathcal{C}^b(\tilde{A}, \text{Vect}_{\mathbb{F}})$ exist. Then

$$c \underset{K}{\frown} - : \text{HK}^{\bullet}(A) \rightarrow \text{HK}_{n-\bullet}(A)$$

is an isomorphism of $\text{HK}^{\bullet}(A)$ -bimodules, inducing an isomorphism of $\text{HK}_{hi}^{\bullet}(A)$ -bimodules from $\text{HK}_{hi}^{\bullet}(A)$ to $\text{HK}_{n-\bullet}^{hi}(A)$. For all $\alpha \in \text{HK}^P(A)$, we have $c \underset{K}{\frown} \alpha = (-1)^{np} \alpha \underset{K}{\frown} c$.

Let us describe the contents of the paper. In Section 2, we extend the general formalism – including some results – of Koszul calculus [7] to quadratic quiver algebras. In Section 3, we introduce a right action which is an important tool in order to adapt the definition of Calabi-Yau algebras to quadratic quiver algebras endowed with the Koszul calculus instead of the Hochschild calculus. The Poincaré Van den Bergh duality for preprojective algebras is presented in Section 4, where Theorem 1.1, Proposition 1.2 and Theorem 1.3 of our introduction are proved. In Section 5, we define our generalisations of Calabi-Yau algebras and we thoroughly explain the new objects and remaining results outlined in the introduction. Section 6 is devoted to the computations of the Koszul calculus in ADE Dynkin types. As an application of the computations, we prove that the spaces $\text{HK}_{hi}^0(A)$, $\text{HK}_{hi}^1(A)$ and $\text{HK}_{hi}^2(A)$ form a minimal complete list of cohomological invariants for the ADE preprojective algebras.

Acknowledgement. The authors are grateful to the anonymous referee, whose useful and detailed comments helped to improve this manuscript.

2. Koszul calculus for quiver algebras with quadratic relations

2.1. Setup

Let \mathcal{Q} be a *finite* quiver, meaning that the vertex set \mathcal{Q}_0 and the arrow set \mathcal{Q}_1 are finite. Let \mathbb{F} be a field. The vertex space $k = \mathbb{F}\mathcal{Q}_0$ becomes a commutative ring by associating with \mathcal{Q}_0 a complete set of orthogonal idempotents $\{e_i; i \in \mathcal{Q}_0\}$. The ring k is isomorphic to $\mathbb{F}^{|\mathcal{Q}_0|}$, where $|\mathcal{Q}_0|$ is the cardinal of \mathcal{Q}_0 . Throughout the paper, the case $|\mathcal{Q}_0| = 1$ will be called *the one vertex case*, which is equivalent to saying that k is a field. Koszul calculus over a field k is treated in [7].

For each arrow $\alpha \in \mathcal{Q}_1$, denote its source vertex by $\mathfrak{s}(\alpha)$ and its target vertex by $\mathfrak{t}(\alpha)$. The arrow space $V = \mathbb{F}\mathcal{Q}_1$ is a k -bimodule for the following actions: $e_j \alpha e_i$ is equal to zero if $i \neq \mathfrak{s}(\alpha)$ or $j \neq \mathfrak{t}(\alpha)$, and is equal to α if $i = \mathfrak{s}(\alpha)$ and $j = \mathfrak{t}(\alpha)$.

Via the ring morphism $\mathbb{F} \rightarrow k$ that maps 1 to $\sum_{i \in \mathcal{Q}_0} e_i$, the tensor k -algebra $T_k(V)$ of the k -bimodule V is an \mathbb{F} -algebra isomorphic to the path algebra $\mathbb{F}\mathcal{Q}$, so that $V^{\otimes_k m}$ is identified with $\mathbb{F}\mathcal{Q}_m$, where \mathcal{Q}_m is the set of paths of length m . For two arrows α and β , note that

$\alpha \otimes_k \beta$ is zero if $\mathfrak{t}(\beta) \neq \mathfrak{s}(\alpha)$, and otherwise $\alpha \otimes_k \beta$ is identified with the composition $\alpha\beta$ of paths (where paths are written from right to left, as in [4]).

Let R be a sub- k -bimodule of $V \otimes_k V \cong \mathbb{F}\mathcal{Q}_2$. The unital associative k -algebra $A = T_k(V)/(R)$, where (R) denotes the two-sided ideal of $T_k(V)$ generated by R , is called a *quadratic k -algebra over the finite quiver \mathcal{Q}* . The degree induced on A by the path length is called the *weight*, so that A is a graded algebra for the weight grading. The component of weight m of A is denoted by A_m . Clearly, $A_0 \cong k$ and $A_1 \cong V$. The algebra A is \mathbb{F} -central, meaning that the left action of $\lambda \in \mathbb{F}$ on A is the same as its right action. However if there is an arrow α joining two distinct vertices i and j , A is not k -central since the left and right actions of e_i on α are different. The A -bimodules considered in this paper are not necessarily k -symmetric, meaning that the left and right actions of an element of k are not necessarily equal, but they are always assumed to be \mathbb{F} -symmetric. Setting $A^e = A \otimes_{\mathbb{F}} A^{op}$, any A -bimodule can be viewed as a left (or right) A^e -module, as usual.

For brevity, the notation $\otimes_{\mathbb{F}}$ is replaced by the unadorned tensor product \otimes . Similarly for the notations $\text{Hom}_{\mathbb{F}}$ and $\dim_{\mathbb{F}}$ abbreviated to Hom and \dim . If unspecified, a vector space is an \mathbb{F} -vector space and a linear map is \mathbb{F} -linear.

The tensor product \otimes_k is different from the unadorned tensor product \otimes . However, if M is a right A -module and N is a left A -module, then the natural linear map $Me_i \otimes e_i N \rightarrow Me_i \otimes_k e_i N$ is an isomorphism, so that for $a \in Me_i$ and $b \in e_i N$, we can identify $a \otimes_k b = a \otimes b$. Similarly, if M and N are A -bimodules, $e_j Me_i$ and $e_i Ne_j$ are k -bimodules, that may be viewed as left and right k^e -modules, where $k^e = k \otimes_k k$. The natural linear map $e_j Me_i \otimes e_i Ne_j \rightarrow e_j Me_i \otimes_{k^e} e_i Ne_j$ is an isomorphism, so for $a \in e_j Me_i$ and $b \in e_i Ne_j$, we can identify $a \otimes_{k^e} b = a \otimes b$. We shall freely use these identifications, without explicitly mentioning them.

Although the algebra A is not k -central, we define its bar resolution $B(A)$ following the standard text [44] by $(A \otimes_k A^{\otimes_k \bullet} \otimes_k A, d)$ with

$$\begin{aligned} d(a \otimes_k a_1 \dots a_p \otimes_k a') &= aa_1 \otimes_k a_2 \dots a_p \otimes_k a' + \sum_{1 \leq i \leq p-1} (-1)^i a \otimes_k a_1 \dots (a_i a_{i+1}) \dots a_p \otimes_k a' \\ &\quad + (-1)^p a \otimes_k a_1 \dots a_{p-1} \otimes_k a_p a', \end{aligned}$$

for a, a' and a_1, \dots, a_p in A . When A is not k -central, the extra degeneracy is defined and is still a contracting homotopy, hence $B(A)$ is a resolution of A by projective A -bimodules. See Lemma 2.1 below for the fact that the A -bimodules $A \otimes_k A^{\otimes_k p} \otimes_k A$ are projective.

For any A -bimodule M , Hochschild homology and cohomology are defined by

$$\text{HH}_{\bullet}(A, M) = \text{Tor}_{\bullet}^{A^e}(M, A) = H_{\bullet}(M \otimes_{A^e} B(A), M \otimes_{A^e} d),$$

$$\text{HH}^{\bullet}(A, M) = \text{Ext}_{A^e}^{\bullet}(A, M) = H^{\bullet}(\text{Hom}_{A^e}(B(A), M), \text{Hom}_{A^e}(d, M)).$$

Given any k -bimodule E , there are well known vector space isomorphisms

$$\begin{aligned} M \otimes_{A^e} (A \otimes_k E \otimes_k A) &\rightarrow M \otimes_{k^e} E \\ m \otimes_{A^e} (a \otimes_k x \otimes_k a') &\mapsto a' m a \otimes_{k^e} x \end{aligned} \tag{2.1}$$

$$\begin{aligned} \text{and} \quad \text{Hom}_{k^e}(E, M) &\rightarrow \text{Hom}_{A^e}(A \otimes_k E \otimes_k A, M) \\ f &\mapsto (a \otimes_k x \otimes_k a' \mapsto a f(x) a') \end{aligned}$$

Taking $E = A^{\otimes_k p}$ and transporting $M \otimes_{A^e} d$ and $\text{Hom}_{A^e}(d, M)$ via these isomorphisms, we obtain the Hochschild differentials b^H and b_H , so that

$$\text{HH}_{\bullet}(A, M) \cong H_{\bullet}(M \otimes_{k^e} A^{\otimes_k \bullet}, b^H),$$

$$\text{HH}^{\bullet}(A, M) \cong H^{\bullet}(\text{Hom}_{k^e}(A^{\otimes_k \bullet}, M), b_H).$$

The Hochschild homology differential is then defined, for $m \in M$ and a_1, \dots, a_p in A , by

$$b_p^H(m \otimes_{k^e} (a_1 \dots a_p)) = ma_1 \otimes_{k^e} (a_2 \dots a_p) + \sum_{1 \leq i \leq p-1} (-1)^i m \otimes_{k^e} (a_1 \dots (a_i a_{i+1}) \dots a_p) + (-1)^p a_p m \otimes_{k^e} (a_1 \dots a_{p-1}).$$

The Hochschild cohomology differential (including a Koszul sign in $\text{Hom}_{A^e}(d, M)$) is defined, for $f \in \text{Hom}_{k^e}(A^{\otimes_k p}, M)$ and a_1, \dots, a_{p+1} in A , by

$$b_H^{p+1}(f)(a_1 \dots a_{p+1}) = f(a_1 \dots a_p) a_{p+1} - \sum_{1 \leq i \leq p} (-1)^{p+i} f(a_1 \dots (a_i a_{i+1}) \dots a_{p+1}) - (-1)^p a_1 f(a_2 \dots a_{p+1}).$$

2.2. Koszul homology and cohomology

Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Following [36, 40, 2, 7], the Koszul complex $K(A)$ is the subcomplex of the bar resolution $B(A)$ defined by the sub- A -bimodules $A \otimes_k W_p \otimes_k A$ of $A \otimes_k A^{\otimes_k p} \otimes_k A$, where $W_0 = k$, $W_1 = V$ and, for $p \geq 2$,

$$W_p = \bigcap_{i+2+j=p} V^{\otimes_k i} \otimes_k R \otimes_k V^{\otimes_k j}. \tag{2.2}$$

Here W_p is considered as a sub- k -bimodule of $V^{\otimes_k p} \subseteq A^{\otimes_k p}$. It is immediate that the differential d of $K(A)$ is defined on $A \otimes_k W_p \otimes_k A$ by

$$d(a \otimes_k x_1 \dots x_p \otimes_k a') = ax_1 \otimes_k x_2 \dots x_p \otimes_k a' + (-1)^p a \otimes_k x_1 \dots x_{p-1} \otimes_k x_p a', \tag{2.3}$$

for a, a' in A and $x_1 \dots x_p$ in W_p .

In this paper, we systematically follow [7] for the notation of elements of W_p . Let us recall this notation. As in (2.3), an arbitrary element of W_p is denoted by a product $x_1 \dots x_p$ thought of as a sum of such products, where x_1, \dots, x_p are in V . Moreover, regarding W_p as a subspace of $V^{\otimes_k q} \otimes_k W_r \otimes_k V^{\otimes_k s}$ with $q + r + s = p$, the element $x_1 \dots x_p$ viewed in $V^{\otimes_k q} \otimes_k W_r \otimes_k V^{\otimes_k s}$ will be denoted by the *same* notation, meaning that the product $x_{q+1} \dots x_{q+r}$ represents an element of W_r and the other x_i are arbitrary in V .

We pursue along the same lines as [7]. We present the different objects with their fundamental results more quickly. We keep the same notations as in [7] and we leave the details to the reader when they are the same as in the one vertex case.

The homology of $K(A)$ is equal to A in degree 0, and to 0 in degree 1. The quadratic algebra A is said to be *Koszul* if the homology of $K(A)$ is 0 in any degree > 1 . Denote by $\mu : A \otimes_k A \rightarrow A$ the multiplication of A . Then A is Koszul if and only if $\mu : K(A) \rightarrow A$ is a resolution of A . If $R = 0$ and if $R = V \otimes_k V$, then A is Koszul. Besides these extreme examples, many Koszul algebras occur in the literature, see for instance [35, 34] among many others, and it is well-known that preprojective algebras are Koszul when the graph is not Dynkin of type ADE (see Proposition 4.1 and the references in its proof).

The A -bimodules $A \otimes_k W_p \otimes_k A$ forming $K(A)$ are projective and finitely generated. Indeed, W_p is a sub- k -bimodule of $V^{\otimes_k p}$, so that this fact is an immediate consequence of the following well known lemma (see for instance [12, Proof of Lemma 2.1]). We give an elementary proof here.

LEMMA 2.1. *Let E be a k -bimodule.*

- (i) *The A -bimodule $A \otimes_k E \otimes_k A$ is projective.*
- (ii) *If E is finite dimensional, then the A -bimodule $A \otimes_k E \otimes_k A$ is finitely generated.*

Proof. Clearly $E = \bigoplus_{i,j \in \mathcal{Q}_0} e_j E e_i$. From $A = \bigoplus_{i \in \mathcal{Q}_0} A e_i = \bigoplus_{j \in \mathcal{Q}_0} e_j A$, we deduce that the A -bimodule $A \otimes_k E \otimes_k A$ is isomorphic to the A -bimodule

$$F_1 = \bigoplus_{i,j \in \mathcal{Q}_0} A e_j \otimes e_j E e_i \otimes e_i A.$$

Considering F_1 as a sub- A -bimodule of $F = A \otimes E \otimes A$, we see that $F = F_1 \oplus F_2$, where

$$F_2 = \bigoplus_{i_1, i_2, i_3, i_4 \in \mathcal{Q}_0} A e_{i_1} \otimes e_{i_2} E e_{i_3} \otimes e_{i_4} A$$

in which the sum is taken over the set of indices with $i_1 \neq i_2$ and $i_3 \neq i_4$. As the A -bimodule F is free, we conclude that F_1 is projective. Part (ii) follows from the fact that E is finite dimensional if and only if all the $(e_j E e_i)$ are finite dimensional and is left to the reader. \square

DEFINITION 2.2. For any A -bimodule M , Koszul homology and cohomology are defined by

$$\mathrm{HK}_\bullet(A, M) = H_\bullet(M \otimes_{A^e} K(A)) \text{ and } \mathrm{HK}^\bullet(A, M) = H^\bullet(\mathrm{Hom}_{A^e}(K(A), M)).$$

We set $\mathrm{HK}_\bullet(A) = \mathrm{HK}_\bullet(A, A)$ and $\mathrm{HK}^\bullet(A) = \mathrm{HK}^\bullet(A, A)$.

Since $K(A)$ is a complex of projective A -bimodules, $M \mapsto \mathrm{HK}_\bullet(A, M)$ and $M \mapsto \mathrm{HK}^\bullet(A, M)$ define δ -functors from the category of A -bimodules to the category of vector spaces, that is, a short exact sequence of A -bimodules naturally gives rise to a long exact sequence in Koszul homology and in Koszul cohomology [44, Chapter 2]. As in [7], $\mathrm{HK}_p(A, M)$ (respectively $\mathrm{HK}^p(A, M)$) is isomorphic to a Hochschild hyperhomology (respectively hypercohomology) space.

The inclusion $\chi : K(A) \rightarrow B(A)$ is a morphism of complexes that induces the following morphisms of complexes

$$\tilde{\chi} = M \otimes_{A^e} \chi : M \otimes_{A^e} K(A) \rightarrow M \otimes_{A^e} B(A),$$

$$\chi^* = \mathrm{Hom}_{A^e}(\chi, M) : \mathrm{Hom}_{A^e}(B(A), M) \rightarrow \mathrm{Hom}_{A^e}(K(A), M).$$

The linear maps $H(\tilde{\chi}) : \mathrm{HK}_p(A, M) \rightarrow \mathrm{HH}_p(A, M)$ and $H(\chi^*) : \mathrm{HH}^p(A, M) \rightarrow \mathrm{HK}^p(A, M)$ are always isomorphisms for $p = 0$ and $p = 1$, and if A is Koszul they are isomorphisms for any p .

Taking $E = W_p$ in the isomorphisms (2.1), we get isomorphisms

$$\begin{aligned} \mathrm{HK}_\bullet(A, M) &\cong H_\bullet(M \otimes_{k^e} W_\bullet, b^K), \\ \mathrm{HK}^\bullet(A, M) &\cong H^\bullet(\mathrm{Hom}_{k^e}(W_\bullet, M), b_K) \end{aligned}$$

with differentials $b^K : M \otimes_{k^e} W_p \rightarrow M \otimes_{k^e} W_{p-1}$ and $b_K : \mathrm{Hom}_{k^e}(W_p, M) \rightarrow \mathrm{Hom}_{k^e}(W_{p+1}, M)$ given by

$$b_p^K(m \otimes_{k^e} x_1 \dots x_p) = m.x_1 \otimes_{k^e} x_2 \dots x_p + (-1)^p x_p.m \otimes_{k^e} x_1 \dots x_{p-1}, \quad (2.4)$$

$$b_K^{p+1}(f)(x_1 \dots x_{p+1}) = f(x_1 \dots x_p).x_{p+1} - (-1)^p x_1.f(x_2 \dots x_{p+1}), \quad (2.5)$$

where $m \in M$ and $x_1 \dots x_p \in W_p$, respectively $f \in \mathrm{Hom}_{k^e}(W_p, M)$ and $x_1 \dots x_{p+1} \in W_{p+1}$.

Note that the k -algebra A is augmented by the natural projection $\epsilon_A : A \rightarrow A_0 \cong k$. Let us examine now the particular case $M = k$, where k is the A -bimodule defined by ϵ_A . The action on k of an element of A_p with $p > 0$ is zero, so that the Koszul differentials vanish when $M = k$. Consequently, we have the linear isomorphisms

$$\mathrm{HK}_p(A, k) \cong k \otimes_{k^e} W_p \cong \bigoplus_{i \in \mathcal{Q}_0} e_i W_p e_i,$$

$$\mathrm{HK}^p(A, k) \cong \mathrm{Hom}_{k^e}(W_p, k) \cong \bigoplus_{i \in \mathcal{Q}_0} \mathrm{Hom}(e_i W_p e_i, \mathbb{F}).$$

In particular $\mathrm{HK}^p(A, k) \cong \mathrm{Hom}(\mathrm{HK}_p(A, k), \mathbb{F})$, generalising [7, Proposition 2.8].

2.3. Koszul cup and cap products

Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . As in [7], the usual cup and cap products \smile and \frown in Hochschild cohomology and homology provide, by restriction from $B(A)$ to $K(A)$, the Koszul cup and cap products \smile_K and \frown_K in Koszul cohomology and homology. Let us give these products, expressed on Koszul cochains and chains. Let P, Q and M be A -bimodules. For $f \in \mathrm{Hom}_{k^e}(W_p, P)$, $g \in \mathrm{Hom}_{k^e}(W_q, Q)$ and $z = m \otimes_{k^e} x_1 \dots x_q \in M \otimes_{k^e} W_q$, we define $f \smile_K g \in \mathrm{Hom}_{k^e}(W_{p+q}, P \otimes_A Q)$, $f \frown_K z \in (P \otimes_A M) \otimes_{k^e} W_{q-p}$ and $z \frown_K f \in (M \otimes_A P) \otimes_{k^e} W_{q-p}$ by

$$(f \smile_K g)(x_1 \dots x_{p+q}) = (-1)^{pq} f(x_1 \dots x_p) \otimes_A g(x_{p+1} \dots x_{p+q}), \quad (2.6)$$

$$f \frown_K z = (-1)^{(q-p)p} (f(x_{q-p+1} \dots x_q) \otimes_A m) \otimes_{k^e} x_1 \dots x_{q-p}, \quad (2.7)$$

$$z \frown_K f = (-1)^{pq} (m \otimes_A f(x_1 \dots x_p)) \otimes_{k^e} x_{p+1} \dots x_q. \quad (2.8)$$

For any Koszul cochains f, g, h and any Koszul chain z , we have the *associativity relations*

$$\begin{aligned} (f \smile_K g) \smile_K h &= f \smile_K (g \smile_K h), \\ f \frown_K (g \frown_K z) &= (f \smile_K g) \frown_K z, \\ (z \frown_K g) \frown_K f &= z \frown_K (g \smile_K f), \\ f \frown_K (z \frown_K g) &= (f \frown_K z) \frown_K g, \end{aligned}$$

inducing the same relations on Koszul classes.

As in [7, Subsection 3.1], for any Koszul cochains $f \in \mathrm{Hom}_{k^e}(W_p, P)$ and $g \in \mathrm{Hom}_{k^e}(W_q, Q)$, we have the identity

$$b_K(f \smile_K g) = b_K(f) \smile_K g + (-1)^p f \smile_K b_K(g), \quad (2.9)$$

so that $(\mathrm{Hom}_{k^e}(W_\bullet, A), b_K, \smile_K)$ is a DG algebra. This DG algebra will play an essential role in Section 3 and will be denoted by \tilde{A} . Note that $H(\tilde{A}) = \mathrm{HK}^\bullet(A)$ is a graded algebra for \smile_K . Moreover, formula (2.9) and analogous formulas for $b^K(f \frown_K z)$ and $b^K(z \frown_K f)$ show that for any A -bimodule M , $\mathrm{Hom}_{k^e}(W_\bullet, M)$ and $M \otimes_{k^e} W_\bullet$ are DG bimodules over \tilde{A} for the actions of \smile_K and \frown_K respectively, so that $\mathrm{HK}^\bullet(A, M)$ and $\mathrm{HK}_\bullet(A, M)$ are graded $\mathrm{HK}^\bullet(A)$ -bimodules.

DEFINITION 2.3. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over a finite quiver \mathcal{Q} .

- (i) The general Koszul calculus of A is the datum of all the spaces $\mathrm{HK}^\bullet(A, P)$ and $\mathrm{HK}_\bullet(A, M)$ endowed with \smile_K and \frown_K , when the A -bimodules P and M vary.
- (ii) The Koszul calculus of A consists of the graded associative algebra $\mathrm{HK}^\bullet(A)$ and of all the graded $\mathrm{HK}^\bullet(A)$ -bimodules $\mathrm{HK}^\bullet(A, M)$ and $\mathrm{HK}_\bullet(A, M)$, when the A bimodule M vary.
- (iii) The restricted Koszul calculus of A consists of the graded associative algebra $\mathrm{HK}^\bullet(A)$ and of the graded $\mathrm{HK}^\bullet(A)$ -bimodule $\mathrm{HK}_\bullet(A)$.
- (iv) The scalar Koszul calculus of A consists of the graded associative algebra $\mathrm{HK}^\bullet(A, k)$ and of the graded $\mathrm{HK}^\bullet(A, k)$ -bimodule $\mathrm{HK}_\bullet(A, k)$.

Since $\mathrm{HK}^0(A) = Z(A)$ is the centre of the algebra A , the spaces $\mathrm{HK}^p(A, M)$ and $\mathrm{HK}_p(A, M)$ are symmetric $Z(A)$ -bimodules (left and right actions coincide). However $\mathrm{HK}^p(A, M)$ and $\mathrm{HK}_p(A, M)$ are not k -bimodules in general. Indeed, $\mathrm{HK}^0(A) = Z(A)$ itself is not a k -bimodule whenever there is an arrow joining two different vertices i and j , since in this case $e_i 1 = e_i$ is not in $Z(A)$.

EXAMPLE 2.4. If $\mathcal{Q}_1 = \emptyset$, then $V = 0$ and A is reduced to k . The (general) Koszul calculus of k coincides with the (tensor) category of k -bimodules.

EXAMPLE 2.5. In order to illustrate the notation and the forthcoming results in the paper, we present the case of the preprojective algebra of type A_3 . This algebra is not Koszul and its Koszul calculus differs from its Hochschild calculus (see Subsection 6.6). This is a special case of the more general examples detailed in Section 6.

Let A be the preprojective algebra of type A_3 over $\mathbb{F} = \mathbb{C}$, that is, the \mathbb{C} -algebra defined by the quiver

$$\overline{\mathcal{Q}} \quad 0 \begin{array}{c} \xrightarrow{a_0} \\ \xleftarrow{a_0^*} \end{array} 1 \begin{array}{c} \xrightarrow{a_1} \\ \xleftarrow{a_1^*} \end{array} 2$$

subject to the relations

$$\sigma_0 = -a_0^* a_0, \quad \sigma_1 = a_0 a_0^* - a_1^* a_1, \quad \sigma_2 = a_1 a_1^*.$$

The algebra A has dimension 10 and a basis of A over \mathbb{C} is given by the elements e_i for $0 \leq i \leq 2$, a_i and a_i^* for $0 \leq i \leq 1$, $a_1^* a_1$, $a_1 a_0$ and $a_0^* a_1^*$. We then have $W_0 = \mathbb{C}\overline{\mathcal{Q}}_0 = \mathbb{C}\langle e_1, e_1, e_2 \rangle = k$, $W_1 = \mathbb{C}\overline{\mathcal{Q}}_1 = \mathbb{C}\langle a_1, a_1, a_0^*, a_1^* \rangle = V$ and $W_2 = \mathbb{C}\langle \sigma_0, \sigma_1, \sigma_2 \rangle = R$.

Now consider $W_3 = (V \otimes_k R) \cap (R \otimes_k V)$, viewed inside $\mathbb{C}\overline{\mathcal{Q}}_3$. An element u in W_3 can therefore be written as a path in $\mathbb{C}\overline{\mathcal{Q}}_3$ in two ways:

$$u = \sum_{i=0}^1 (\lambda_i a_i \sigma_i + \lambda_i^* a_i^* \sigma_{i+1}) = \sum_{i=0}^1 (\mu_i \sigma_{i+1} a_i + \mu_i^* \sigma_i a_i^*)$$

with $\lambda_i, \lambda_i^*, \mu_i, \mu_i^*$ in \mathbb{C} . Then, in $\mathbb{C}\overline{\mathcal{Q}}_3$, we have

$$\begin{aligned} & (-\lambda_0 - \mu_0) a_0 a_0^* a_0 + (-\lambda_1 - \mu_1) a_1 a_1^* a_1 + (\lambda_0^* + \mu_0^*) a_0^* a_0 a_0^* + (\lambda_1^* + \mu_1^*) a_1^* a_1 a_1^* \\ & + \lambda_1 a_1 a_0 a_0^* - \lambda_0^* a_0^* a_1^* a_1 + \mu_0 a_1^* a_1 a_0 - \mu_1^* a_0^* a_0^* a_1^* = 0 \end{aligned}$$

so that all the coefficients $\lambda_i, \lambda_i^*, \mu_i$ and μ_i^* must be zero, hence $u = 0$ and $W_3 = 0$.

Since $W_p = (W_{p-1} \otimes_k V) \cap (V \otimes_k W_{p-1})$ for all $p \geq 3$, it follows that $W_p = 0$ for all $p \geq 3$. This is true for any preprojective algebra of type Δ with Δ different from A_1 and A_2 , see Theorem 4.2.

The Koszul complex $K(A)$ is therefore

$$\cdots \rightarrow 0 \rightarrow A \otimes_k R \otimes_k A \xrightarrow{d_2} A \otimes_k V \otimes_k A \xrightarrow{d_1} A \otimes_k A \rightarrow 0$$

with

$$\begin{aligned} d_1(a \otimes_k x \otimes_k a') &= ax \otimes_k a' - a \otimes_k xa' \\ d_2(a \otimes_k \sum_{i=1}^n \lambda_i x_i y_i \otimes_k a') &= \sum_{i=1}^n \lambda_i (ax_i \otimes_k y_i \otimes_k a'_a \otimes_k x_i \otimes_k y_i a'). \end{aligned}$$

Applying $\mathrm{Hom}_{A^e}(-, A)$ and using the natural isomorphism $\mathrm{Hom}_{A^e}(A \otimes_k E \otimes_k A, A) \cong \mathrm{Hom}_{k^e}(E, A)$ for any k -bimodule E (2.1), we get the complex

$$0 \rightarrow \mathrm{Hom}_{k^e}(k, A) \xrightarrow{b_K^1} \mathrm{Hom}_{k^e}(V, A) \xrightarrow{b_K^2} \mathrm{Hom}_{k^e}(R, A) \rightarrow 0 \rightarrow \cdots$$

Before we describe the maps, let us note that $\text{Hom}_{k^e}(k, A) \cong \bigoplus_{i=0}^2 e_i A e_i$ has basis $\{e_0, e_1, e_2, a_1^* a_1\}$, that a general element f in $\text{Hom}_{k^e}(V, A)$ is defined by $f(a_i) = \lambda_i a_i$ and $f(a_i^*) = \lambda_i^* a_i^*$ for $i = 0, 1$ with λ_i, λ_i^* in \mathbb{C} , and that a general element g in $\text{Hom}_{k^e}(R, A)$ is defined by $g(\sigma_i) = \alpha_i e_i$ for $i = 0, 2$ and $g(\sigma_2) = \alpha_2 e_1 + \beta a_1^* a_1$ for some scalars α_i and β . Then

$$b_K^1\left(\sum_{i=0}^2 u_i e_i + v a_1^* a_1\right) \text{ is defined by } a_i \mapsto (u_{i+1} - u_i) a_i \text{ and } a_i^* \mapsto (u_i - u_{i+1}) a_i^* \text{ for } i = 0, 1$$

$$b_K^2(f) \text{ is defined by } \sigma_0 \mapsto 0, \sigma_1 \mapsto (\lambda_0 + \lambda_0^* - \lambda_1 - \lambda_1^*) a_1^* a_1 \text{ and } \sigma_2 \mapsto 0.$$

It is then easy to see that $\text{HK}^0(A) = \mathbb{C}\langle z_0 = 1, z_1 = a_1^* a_1 \rangle$, that $\text{HK}^1(A) = \mathbb{C}\langle \bar{\zeta}_0 \rangle$ with $\zeta_0 \in \text{Hom}_{k^e}(V, A)$ defined by $\zeta_0(a_i) = a_i$ and $\zeta_0(a_i^*) = 0$, and that $\text{HK}^2(A) = \mathbb{C}\langle \bar{h}_0, \bar{h}_1, \bar{h}_2 \rangle$ with $h_i \in \text{Hom}_{k^e}(R, A)$ defined by $h_i(\sigma_j) = \delta_{ij} e_i$.

Moreover, the fundamental 1-cocycle ϵ_A is equal to $2\zeta_0 + b_K^1(2e_0 + e_1)$.

The Koszul cup products can easily be found using the formula (2.6). It follows that $\underset{K}{\smile}$ is graded commutative, that $1 \underset{K}{\smile} x = x$ for any $x \in \text{HK}^\bullet(A)$ and that all other cup products are 0 in $\text{HK}^\bullet(A)$. For instance, $z_1 \underset{K}{\smile} h_1$ is the coboundary $b_K^2(f)$ where f sends a_0 to a_0 and all other arrows to 0. In particular, $\epsilon_A \underset{K}{\smile} \epsilon_A = 0$.

In order to determine the Koszul homology of A , we could also apply the functor $A \otimes_{A^e} -$ to the complex $K(A)$ and compute the homology of the complex obtained. However, we can also use our duality result, Theorem 4.4. Set $\omega_0 = \sum_{i=0}^2 e_i \otimes \sigma_i \in A \otimes_{k^e} R$. There is an isomorphism $\theta_A : \text{HK}^\bullet(A) \rightarrow \text{HK}_{2-\bullet}(A)$ given by $\bar{f} \mapsto \bar{\omega}_0 \underset{K}{\frown} f$. Explicitly in our example,

$$\begin{aligned} \theta_A(z_0) &= \bar{\omega}_0 \text{ and } \theta_A(z_1) = \overline{z_1 \otimes \sigma_1} \text{ form a basis of } \text{HK}_2(A); \\ \theta_A(\bar{\zeta}_0) &= \overline{a_0 \otimes a_0^* + a_1 \otimes a_1^*} \text{ forms a basis of } \text{HK}_1(A); \\ \theta_A(\bar{h}_i) &= \overline{e_i \otimes e_i} \text{ for } 0 \leq i \leq 2 \text{ form a basis of } \text{HK}_0(A). \end{aligned}$$

The Koszul cap products can also be obtained using duality and they all vanish except the cap products $z_0 \underset{K}{\frown} x = x = x \underset{K}{\frown} z_0$ for all $x \in \text{HK}_\bullet(A)$.

2.4. Fundamental formulas of Koszul calculus

Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . We continue to follow the one vertex case [7]. First, we define the Koszul cup and cap brackets. Let P, Q and M be A -bimodules, and take $f \in \text{Hom}_{k^e}(W_p, P)$, $g \in \text{Hom}_{k^e}(W_q, Q)$, $z \in M \otimes_{k^e} W_q$. When P or Q is equal to A , we set

$$[f, g]_{\underset{K}{\smile}} = f \underset{K}{\smile} g - (-1)^{pq} g \underset{K}{\smile} f. \tag{2.10}$$

When P or M is equal to A , we set

$$[f, z]_{\underset{K}{\frown}} = f \underset{K}{\frown} z - (-1)^{pq} z \underset{K}{\frown} f. \tag{2.11}$$

These brackets induce brackets on the Koszul classes.

The Koszul 1-cocycles $f : V \rightarrow M$ are called Koszul derivations with coefficients in M . Such an f extends to a unique derivation from the k -algebra A to the A -bimodule M , realising an isomorphism from the space of Koszul derivations with coefficients in M to the space of derivations from A to M . In particular, the Koszul 1-cocycle from V to A coinciding with the identity map on V , is sent to the Euler derivation D_A of the graded algebra A . This Koszul 1-cocycle is denoted by ϵ_A and is called the *fundamental 1-cocycle*. Its Koszul class is denoted by $\bar{\epsilon}_A$ and is called the *fundamental 1-class*. In the one vertex case, ϵ_A is not a coboundary if $V \neq 0$ [7], but this property does not hold in general.

LEMMA 2.6. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} with $\mathcal{Q}_1 \neq \emptyset$. If the underlying graph of \mathcal{Q} is simple, that is, it contains neither loops nor multiple edges, then ϵ_A is a coboundary.*

Proof. The 1-cocycle ϵ_A is a coboundary if and only if there exists a k^e -linear map $c : k \rightarrow k$ such that $\epsilon_A = b_K(c)$. Such a map is of the form $c(e_i) = \lambda_i e_i$ with $\lambda_i \in \mathbb{F}$, for all $i \in \mathcal{Q}_0$. Then $\epsilon_A = b_K(c)$ if and only if $\lambda_{t(\alpha)} - \lambda_{s(\alpha)} = 1$ for any $\alpha \in \mathcal{Q}_1$. The assumption on the graph means that \mathcal{Q} has no loop and that given two distinct vertices, there is at most one arrow joining them. Then we can choose $\lambda_{t(\alpha)} = 1$ and $\lambda_{s(\alpha)} = 0$. \square

This proof shows that if the quiver \mathcal{Q} has a loop, ϵ_A is not a coboundary. The same conclusion holds if $\text{char } \mathbb{F} \neq 2$ and \mathcal{Q} contains an oriented 2-cycle.

The following propositions are proved as [7, Theorem 3.7, Theorem 4.4, Corollary 3.10, Corollary 4.6] of the one vertex case. Formulas (2.12) and (2.13) are the fundamental formulas of Koszul calculus.

PROPOSITION 2.7. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . For any Koszul cochain f and any Koszul chain z with coefficients in an A -bimodule M , we have*

$$b_K(f) = -[\epsilon_A, f]_{\underset{K}{\smile}}, \quad (2.12)$$

$$b^K(z) = -[\epsilon_A, z]_{\underset{K}{\frown}}. \quad (2.13)$$

PROPOSITION 2.8. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} and let M be an A -bimodule. For any $\alpha \in \text{HK}^p(A, M)$ with $p = 0$ or $p = 1$, $\beta \in \text{HK}^q(A)$ and $\gamma \in \text{HK}_q(A)$, we have the identities*

$$[\alpha, \beta]_{\underset{K}{\smile}} = 0, \quad (2.14)$$

$$[\alpha, \gamma]_{\underset{K}{\frown}} = 0. \quad (2.15)$$

Identity (2.15) also holds if $p = q \notin \{0, 1\}$.

2.5. Higher Koszul calculus

Higher Koszul homology is the homology of the Koszul homology, and similarly for cohomology. Precisely, let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Formula (2.6) shows that the map $\epsilon_A \underset{K}{\smile} \epsilon_A : W_2 = R \rightarrow A$ is zero. Therefore, $\epsilon_A \underset{K}{\smile} -$ is a cochain differential on $\text{Hom}_{k^e}(W_\bullet, M)$, and $\bar{\epsilon}_A \underset{K}{\smile} -$ is a cochain differential on $\text{HK}^\bullet(A, M)$. Similarly, $\epsilon_A \underset{K}{\frown} -$ is a chain differential on $M \otimes_{k^e} W_\bullet$, and $\bar{\epsilon}_A \underset{K}{\frown} -$ is a chain differential on $\text{HK}_\bullet(A, M)$. For a p -cocycle $f : W_p \rightarrow M$ and $x_1 \dots x_{p+1}$ in W_{p+1} , we have

$$(\epsilon_A \underset{K}{\smile} f)(x_1 \dots x_{p+1}) = f(x_1 \dots x_p) \cdot x_{p+1}.$$

For a p -cycle $z = m \otimes_{k^e} x_1 \dots x_p$ in $M \otimes_{k^e} W_p$, we have

$$\epsilon_A \underset{K}{\frown} z = mx_1 \otimes_{k^e} x_2 \dots x_p.$$

DEFINITION 2.9. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over a finite quiver \mathcal{Q} and let M be an A -bimodule. The differentials $\bar{\epsilon}_A \underset{K}{\smile} -$ and $\bar{\epsilon}_A \underset{K}{\frown} -$ are denoted by ∂_{\smile} and ∂_{\frown} .

The homologies of the complexes $(\mathrm{HK}^\bullet(A, M), \partial_-)$ and $(\mathrm{HK}_\bullet(A, M), \partial_-)$ are called the higher Koszul cohomology and homology of A with coefficients in M and are denoted by $\mathrm{HK}_{hi}^\bullet(A, M)$ and $\mathrm{HK}_\bullet^{hi}(A, M)$. We set $\mathrm{HK}_{hi}^\bullet(A) = \mathrm{HK}_{hi}^\bullet(A, A)$ and $\mathrm{HK}_\bullet^{hi}(A) = \mathrm{HK}_\bullet^{hi}(A, A)$.

The higher classes of Koszul classes will be denoted between square brackets. For example, the unit 1 of A is still the unit of $\mathrm{HK}^\bullet(A)$, and $\partial_-(1) = \bar{\epsilon}_A$ implies that $[\bar{\epsilon}_A] = 0$. If $\bar{\epsilon}_A \neq 0$, the unit of $\mathrm{HK}^\bullet(A)$ does not survive in higher Koszul cohomology.

As in the one vertex case, the actions of the Koszul cup and cap products of $\mathrm{HK}^\bullet(A)$ on $\mathrm{HK}^\bullet(A, M)$ and $\mathrm{HK}_\bullet(A, M)$ induce actions on higher cohomology and homology. Thus $\mathrm{HK}_{hi}^\bullet(A)$ is a graded algebra, and $\mathrm{HK}_{hi}^\bullet(A, M)$, $\mathrm{HK}_\bullet^{hi}(A, M)$ are graded $\mathrm{HK}_{hi}^\bullet(A)$ -bimodules, constituting *the higher Koszul calculus of A* . If $\bar{\epsilon}_A = 0$, the higher Koszul calculus coincides with the Koszul calculus. It is the case when $A = k$ as in Example 2.4.

For $M = k$, $\epsilon_A \underset{K}{\smile} -$ and $\epsilon_A \underset{K}{\frown} -$ vanish, so that the higher scalar Koszul calculus coincides with the scalar Koszul calculus. Proposition 3.12 in [7] generalises immediately as follows.

PROPOSITION 2.10. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} and let M be an A -bimodule. Then $\mathrm{HK}_{hi}^0(A, M)$ is the space of elements u in $Z(M)$ such that there exists $v \in M$ satisfying $u.a = v.a - a.v$ for any a in \mathcal{Q}_1 .*

2.6. Grading the restricted Koszul calculus by the weight

A Koszul p -cochain $f : W_p \rightarrow A_m$ is said to be homogeneous of weight m . Since \mathcal{Q}_1 is finite, the spaces W_p are finite dimensional, thus the space of Koszul cochains $\mathrm{Hom}_{k^e}(W_\bullet, A)$ is $\mathbb{N} \times \mathbb{N}$ -graded by the *biweight* (p, m) , where p is called the *homological weight* and m is called the *coefficient weight*. If $f : W_p \rightarrow A_m$ and $g : W_q \rightarrow A_n$ are homogeneous of biweights (p, m) and (q, n) respectively, then $f \underset{K}{\smile} g : W_{p+q} \rightarrow A_{m+n}$ is homogeneous of biweight $(p+q, m+n)$. Moreover b_K is homogeneous of biweight $(1, 1)$ and the algebra $\mathrm{HK}^\bullet(A)$ is $\mathbb{N} \times \mathbb{N}$ -graded by the biweight. The homogeneous component of biweight (p, m) of $\mathrm{HK}^\bullet(A)$ is denoted by $\mathrm{HK}^p(A)_m$. Since

$$\partial_- : \mathrm{HK}^p(A)_m \rightarrow \mathrm{HK}^{p+1}(A)_{m+1},$$

the algebra $\mathrm{HK}_{hi}^\bullet(A)$ is $\mathbb{N} \times \mathbb{N}$ -graded by the biweight, and its (p, m) -component is denoted by $\mathrm{HK}_{hi}^p(A)_m$. From Proposition 2.10, we deduce the following.

PROPOSITION 2.11. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Assume that A is finite dimensional. Let \max be the highest m such that $A_m \neq 0$. Then $\mathrm{HK}_{hi}^0(A)_{\max}$ is isomorphic to the space spanned by the cycles of \mathcal{Q} of length \max .*

Similarly, a Koszul q -chain z in $A_n \otimes_{k^e} W_q$ is said to be homogeneous of weight n . The space of Koszul chains $A \otimes_{k^e} W_\bullet$ is $\mathbb{N} \times \mathbb{N}$ -graded by the *biweight* (q, n) , where q is called the *homological weight* and n is called the *coefficient weight*. Moreover b^K is homogeneous of biweight $(-1, 1)$ and the space $\mathrm{HK}_\bullet(A)$ is $\mathbb{N} \times \mathbb{N}$ -graded by the biweight. The homogeneous component of biweight (q, n) of $\mathrm{HK}_\bullet(A)$ is denoted by $\mathrm{HK}_q(A)_n$. Since

$$\partial_- : \mathrm{HK}_q(A)_n \rightarrow \mathrm{HK}_{q-1}(A)_{n+1},$$

the space $\mathrm{HK}_\bullet^{hi}(A)$ is $\mathbb{N} \times \mathbb{N}$ -graded by the biweight, and its (q, n) -component is denoted by $\mathrm{HK}_q^{hi}(A)_n$.

If $f : W_p \rightarrow A_m$ and $z \in A_n \otimes_{k^e} W_q$ are homogeneous of biweights (p, m) and (q, n) respectively, then $f \underset{K}{\frown} z$ and $z \underset{K}{\frown} f$ are homogeneous of biweight $(q - p, m + n)$ where

$$f \underset{K}{\frown} z = (-1)^{(q-p)p} f(x_{q-p+1} \dots x_q) a \otimes_{k^e} x_1 \dots x_{q-p}, \quad (2.16)$$

$$z \underset{K}{\frown} f = (-1)^{pq} a f(x_1 \dots x_p) \otimes_{k^e} x_{p+1} \dots x_q, \quad (2.17)$$

and $z = a \otimes_{k^e} x_1 \dots x_q$. The $\text{Hom}_{k^e}(W_\bullet, A)$ -bimodule $A \otimes_{k^e} W_\bullet$, the $\text{HK}^\bullet(A)$ -bimodule $\text{HK}_\bullet(A)$ and the $\text{HK}_{hi}^\bullet(A)$ -bimodule $\text{HK}^{hi}(A)$ are thus $\mathbb{N} \times \mathbb{N}$ -graded by the biweight. The proof of the following is left to the reader.

PROPOSITION 2.12. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . We have*

$$\text{HK}_0^{hi}(A)_0 \cong \text{HK}_0(A)_0 \cong k.$$

Moreover $\text{HK}_0(A)_1 \cong \text{HK}_1(A)_0$ is isomorphic to the space spanned by the loops of \mathcal{Q} , and $\partial_\frown : \text{HK}_1(A)_0 \rightarrow \text{HK}_0(A)_1$ identifies with the identity map on this space. As a consequence,

$$\text{HK}_0^{hi}(A)_1 \cong \text{HK}_1^{hi}(A)_0 = 0.$$

2.7. Invariance of Koszul calculus

In [6], the first author proved that the Koszul calculus of an N -homogeneous algebra A over a field k only depends on the structure of associative algebra of A , independently of any presentation $A = T_k(V)/(R)$ of A as an N -homogeneous algebra. This result was based on an isomorphism lemma due to Bell and Zhang [3]. In the quadratic case $N = 2$, we are going to extend this Koszul calculus invariance to any quadratic quiver algebra. For that, we shall use an extension of the isomorphism lemma to quiver algebras with homogeneous relations, due to Gaddis [25].

Let \mathcal{Q} and \mathcal{Q}' be finite quivers, and \mathbb{F} be a field. We introduce the commutative rings $k = \mathbb{F}\mathcal{Q}_0$ and $k' = \mathbb{F}\mathcal{Q}'_0$, the k -bimodule $V = \mathbb{F}\mathcal{Q}_1$ and the k' -bimodule $V' = \mathbb{F}\mathcal{Q}'_1$. As explained in Subsection 2.1, we make the identifications of graded algebras $T_k(V) \cong \mathbb{F}\mathcal{Q}$ and $T_{k'}(V') \cong \mathbb{F}\mathcal{Q}'$. We are interested in the graded \mathbb{F} -algebra isomorphisms $u : T_k(V) \rightarrow T_{k'}(V')$ given by a ring isomorphism $u_0 : k \rightarrow k'$ and by a k -bimodule isomorphism $u_1 : V \rightarrow V'$, where V' is a k -bimodule via u_0 . By [25, Lemma 4], this implies that u_0 maps \mathcal{Q}_0 to \mathcal{Q}'_0 , and the bijection $\mathcal{Q}_0 \rightarrow \mathcal{Q}'_0$ induced by u_0 transforms the adjacency matrix of \mathcal{Q} into the adjacency matrix of \mathcal{Q}' .

Let us fix a sub- k -bimodule R of $V \otimes_k V$ and a sub- k' -bimodule R' of $V' \otimes_{k'} V'$. We define the graded k -algebra $A = T_k(V)/(R)$ and the graded k' -algebra $A' = T_{k'}(V')/(R')$. Following the terminology of the one vertex case, a graded \mathbb{F} -algebra isomorphism $u : A \rightarrow A'$ is called a *Manin isomorphism* if u is defined by a ring isomorphism $u_0 : k \rightarrow k'$ (so V' is a k -bimodule via u_0), and by a k -bimodule isomorphism $u_1 : V \rightarrow V'$, such that the k -bimodule isomorphism $u_1^{\otimes_{k^e} 2} : V^{\otimes_{k^e} 2} \rightarrow V'^{\otimes_{k^e} 2}$ satisfies $u_1^{\otimes_{k^e} 2}(R) = R'$. In particular, u is an isomorphism of the augmented k -algebra A to the augmented k' -algebra A' , the augmentations being the projections $A \rightarrow A_0 \cong k$ and $A' \rightarrow A'_0 \cong k'$.

As in [6], for any A -bimodule M , the Manin isomorphism u naturally defines an isomorphism of complexes from $(M \otimes_{k^e} W_\bullet, b^K)$ to $(M \otimes_{k^e} W'_\bullet, b^K)$, where M is an A' -bimodule via u , inducing natural isomorphisms $\text{HK}_\bullet(A, M) \cong \text{HK}_\bullet(A', M)$. Similarly, u induces natural isomorphisms $\text{HK}^\bullet(A', M) \cong \text{HK}^\bullet(A, M)$. It is clear from the definitions in Subsection 2.3 that these isomorphisms respect the Koszul cup and cap products. To summarise all these properties, we say that a Manin isomorphism induces isomorphic (general) Koszul calculi. Since $u_1(\epsilon_A) = \epsilon_{A'}$ by functoriality, it also induces isomorphic higher Koszul calculi.

Using Gaddis’s theorem [25, Theorem 5], we can now prove ungraded invariance. Let C be a commutative ring. Let A be an augmented associative C -algebra (not necessarily C -central) having a quadratic quiver algebra presentation B , meaning that the augmented C -algebra A is isomorphic to a quadratic k -algebra $B = T_k(V)/(R)$ over a finite quiver \mathcal{Q} , naturally augmented over k by the projection $B \rightarrow B_0 \cong k$. This implies that the ring C is isomorphic to $k \cong \mathbb{F}\mathcal{Q}_0$. Then we can define the (general) Koszul calculus of A as being the (general) Koszul calculus of B . Indeed, if $B' = T_{k'}(V')/(R')$ over a finite quiver \mathcal{Q}' is another quadratic quiver algebra presentation of A , the ungraded augmented k -algebra B is isomorphic to the ungraded augmented k' -algebra B' . By Gaddis’s theorem, there exists a Manin isomorphism from B to B' , thus the (general) Koszul calculi of B and B' are isomorphic by Manin invariance. The higher Koszul calculus of A is also defined as being the higher Koszul calculus of B .

2.8. Comparing Koszul (co)homology with Hochschild (co)homology in degree 2

Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Recall that, for $p = 0$ and $p = 1$, we have linear isomorphisms $\mathrm{HK}_p(A, M) \cong \mathrm{HH}_p(A, M)$ and $\mathrm{HH}^p(A, M) \cong \mathrm{HK}^p(A, M)$ (Subsection 2.2). It is no longer true if $p \geq 2$ and if A is an arbitrary non-Koszul algebra. Preprojective algebras of Dynkin type will give infinitely many counterexamples when $p = 2$. However, in general, we can compare the Koszul and Hochschild spaces when $p = 2$, by providing a surjection $\mathrm{HK}_2(A, M) \rightarrow \mathrm{HH}_2(A, M)$ and an injection $\mathrm{HH}^2(A, M) \rightarrow \mathrm{HK}^2(A, M)$. To prove that, we use a minimal projective resolution of the graded k -algebra A , described as follows.

As in the one vertex case [7], we know that, in the category of graded A -bimodules, A has a minimal projective resolution $P(A)$ whose component of homological degree p can be written as $A \otimes_k E_p \otimes_k A$, where E_p is a weight-graded k -bimodule. Then $P_l(A) = P(A) \otimes_A k$ (respectively $P_r(A) = k \otimes_A P(A)$) is a minimal projective resolution of the graded left (respectively right) A -module k (see for instance [5]) and the differential δ of $P(A)$ is the graded sum of the differentials $\delta \otimes_A \mathrm{id}_k$ and $\mathrm{id}_k \otimes_A \delta$ naturally extended to $P(A)$.

Define the left (respectively right) Koszul complex $K_l(A) = K(A) \otimes_A k$ (respectively $K_r(A) = k \otimes_A K(A)$). Now using [2, Subsections 2.7 and 2.8], [35, Chapter 1, Proposition 3.1] adapted to the case where k is a semisimple ring (rather than a field), and the construction of $\mathrm{Ext}_A(k, k)$ from the resolutions $P_l(A)$ and $P_r(A)$, we can show that $K_l(A)$ (respectively $K_r(A)$) is isomorphic as a left (respectively right) A -module to the diagonal part of the graded resolution $P_l(A)$ (respectively $P_r(A)$). We know (see for instance [40, Section 3]) that the differential d of $K(A)$ is the graded sum of the differentials $d \otimes_A \mathrm{id}_k$ and $\mathrm{id}_k \otimes_A d$ naturally extended to $K(A)$. Thus the inclusions $A \otimes_k W_p \otimes_k A \hookrightarrow A \otimes_k E_p \otimes_k A$ constitute an inclusion map $\iota : K(A) \hookrightarrow P(A)$ of weight-graded A -bimodule complexes. So we can view the complex $K(A)$ as the diagonal part of the weight-graded resolution $P(A)$, and A is Koszul if and only if $P(A) = K(A)$. The beginning of $P(A)$ coincides with $K(A)$, that is, $E_0 = k$, $E_1 = V$, $E_2 = R$, and the differential δ of $P(A)$ coincides with the differential d of $K(A)$ in degrees 1 and 2.

For any A -bimodule M , ι induces $\tilde{\iota} = M \otimes_{A^e} \iota$ and $\iota^* = \mathrm{Hom}_{A^e}(\iota, M)$ decomposed in $\tilde{\iota}_p : M \otimes_{k^e} W_p \rightarrow M \otimes_{k^e} E_p$ and $\iota_p^* : \mathrm{Hom}_{k^e}(E_p, M) \rightarrow \mathrm{Hom}_{k^e}(W_p, M)$. The linear maps

$$H(\tilde{\iota})_p : \mathrm{HK}_p(A, M) \rightarrow \mathrm{HH}_p(A, M) \text{ and } H(\iota^*)_p : \mathrm{HH}^p(A, M) \rightarrow \mathrm{HK}^p(A, M) \tag{2.18}$$

are isomorphisms for $p = 0$ and $p = 1$, and for any p if A is Koszul. Since ι_p is an identity map for $p = 0, 1, 2$, $\tilde{\iota}_p$ and ι_p^* are also identity maps for the same p 's. Therefore, setting $\tilde{\delta} = M \otimes_{A^e} \delta$ and $\delta^* = \mathrm{Hom}_{A^e}(\delta, M)$, we have the commutative diagrams

$$\begin{array}{ccccc} M \otimes_{k^e} W_3 & \xrightarrow{b_3^K} & M \otimes_{k^e} R & \xrightarrow{b_2^K} & M \otimes_{k^e} V \\ \tilde{\iota}_3 \downarrow & & id \downarrow & & id \downarrow \\ M \otimes_{k^e} E_3 & \xrightarrow{\tilde{\delta}_3} & M \otimes_{k^e} R & \xrightarrow{\tilde{\delta}_2} & M \otimes_{k^e} V \end{array} \tag{2.19}$$

$$\begin{array}{ccccc}
\mathrm{Hom}_{k^e}(V, M) & \xrightarrow{\delta_2^*} & \mathrm{Hom}_{k^e}(R, M) & \xrightarrow{\delta_3^*} & \mathrm{Hom}_{k^e}(E_3, M) \\
id \downarrow & & id \downarrow & & \tilde{\iota}_3^* \downarrow \\
\mathrm{Hom}_{k^e}(V, M) & \xrightarrow{b_K^2} & \mathrm{Hom}_{k^e}(R, M) & \xrightarrow{b_K^3} & \mathrm{Hom}_{k^e}(W_3, M)
\end{array} \tag{2.20}$$

where $\tilde{\iota}_3$ is injective and $\tilde{\iota}_3^*$ is surjective (the ring $k^e \cong \mathbb{F}^{|\mathcal{Q}_0|^2}$ is semisimple), so that we obtain the following.

PROPOSITION 2.13. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . For any M ,*

- (i) $H(\tilde{\iota})_2 : \mathrm{HK}_2(A, M) \rightarrow \mathrm{HH}_2(A, M)$ is surjective with kernel isomorphic to $\mathrm{Im}(\tilde{\delta}_3)/\mathrm{Im}(b_3^K)$,
- (ii) $H(\tilde{\iota}^*)_2 : \mathrm{HH}^2(A, M) \rightarrow \mathrm{HK}^2(A, M)$ is injective with image isomorphic to $\mathrm{Ker}(\delta_3^*)/\mathrm{Im}(b_K^2)$.

We can be more specific when $M = A$, by using the weight grading (Subsection 2.6). Unlike the Koszul differentials b^K and b_K , the Hochschild differentials b^H and b_H are not homogeneous for the coefficient weight, but only for the total weight. The grading of $\mathrm{HH}_p(A)$ and $\mathrm{HH}^p(A)$ for the total weight t is denoted by $\mathrm{HH}_p(A)_t$ and $\mathrm{HH}^p(A)_t$. Denote the weight of a homogeneous element a of A by $|a|$. Recall that the total weight of a homogeneous p -chain $z = a \otimes_{k^e} (a_1 \dots a_p)$ is equal to $t = |a| + |a_1| + \dots + |a_p|$, and the total weight of a homogeneous p -cochain f mapping $a_1 \dots a_p$ to an element of A_m is equal to $t = m - |a_1| - \dots - |a_p|$. Then $H(\tilde{\iota})_2$ is homogeneous from the coefficient weight r to the total weight $r + 2$, while $H(\tilde{\iota}^*)_2$ is homogeneous from the total weight $r - 2$ to the coefficient weight r .

COROLLARY 2.14. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} .*

- (i) $H(\tilde{\iota})_2$ is an isomorphism from $\mathrm{HK}_2(A)_r$ to $\mathrm{HH}_2(A)_{r+2}$ if $r = 0$ and $r = 1$.
- (ii) Assume that A is finite dimensional. Let \max be the highest m such that $A_m \neq 0$. Then $H(\tilde{\iota}^*)_2$ is an isomorphism from $\mathrm{HH}^2(A)_{r-2}$ to $\mathrm{HK}^2(A)_r$ if $r = \max$ and $r = \max - 1$.

Proof. Denote by $E_{p,m}$ the homogeneous component of weight m of E_p . Since $E_{3,2} = 0$ and $E_{3,3} = W_3$, both maps δ_3^* and b_3^K vanish on the component of total weight 2 of $A \otimes_{k^e} E_3$, while on that of total weight 3, they coincide with the inclusion map of W_3 into $V \otimes_{k^e} R$. Then we deduce (i) from (i) of the proposition.

Under the assumptions of (ii), if $f : R \rightarrow A_{\max}$, then $b_K^3(f) = 0$. Moreover, any other component of $\delta_3^*(f)$ mapping $E_{3,m}$ to $A_{\max+m-2} = 0$ vanishes as well. Thus $\delta_3^*(f) = 0$, and we conclude by (ii) of the proposition. The same proof works if $f : R \rightarrow A_{\max-1}$ since $\delta_3^*(f)$ is then reduced to a map $W_3 \rightarrow A_{\max}$ coinciding with $b_K^3(f)$. \square

3. A right action on the Koszul calculus

This section presents an important tool which we use in Section 5 to adapt the known definition of Calabi-Yau algebras due to Ginzburg to the context of quadratic quiver algebras endowed with the Koszul calculus. The idea is to put together two compatible bimodule actions on Koszul chains and cochains : the action of the quadratic quiver algebra A and the action of the associated DG algebra \tilde{A} defined just before Definition 2.3.

3.1. Compatibility

LEMMA 3.1. *Let A and B be unital associative \mathbb{F} -algebras. Let M be an A -bimodule (hence the induced \mathbb{F} -bimodule is symmetric). Assume that M is a right B -module such that the actions of \mathbb{F} induced on M by A and by B are the same. Let $A^e = A \otimes_{\mathbb{F}} A^{op}$ be the enveloping algebra. The following are equivalent.*

- (i) *Viewing M as a left A^e -module, M is an A^e - B -bimodule.*
- (ii) *Viewing M as a right A^e -module, the right actions of A^e and B on M commute.*
- (iii) *M is an A - B -bimodule and the right actions of A and B on M commute.*

The proof is straightforward. Under the assumptions of the lemma and if the equivalent assertions hold, we say that the right action of B on M is *compatible* with the A -bimodule M .

EXAMPLE 3.2. With $B = M = A^e$, A^e is a natural A^e - A^e -bimodule for the multiplication of the \mathbb{F} -algebra A^e . Recall that the left A^e -module A^e is isomorphic to the A -bimodule $A \overset{\circ}{\otimes} A$ for the outer action $(a \otimes b).(\alpha \otimes \beta) = (a\alpha) \otimes (b\beta)$, while the right A^e -module A^e is isomorphic to the A -bimodule $A \overset{\circ}{\otimes} A$ for the inner action $(\alpha \otimes \beta).(a \otimes b) = (\alpha a) \otimes (b\beta)$.

3.2. DG bimodules over the DG algebra \tilde{A}

Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Fix a unital associative \mathbb{F} -algebra B and an A -bimodule M . We assume that M is a right B -module compatible with the A -bimodule structure. Then the space $M \otimes_{k^e} W_q$ is a right B -module for the action of $b \in B$ on $z = m \otimes_{k^e} x_1 \dots x_q \in M \otimes_{k^e} W_q$ defined by

$$z.b = (m.b) \otimes_{k^e} x_1 \dots x_q.$$

It is well-defined since $(\lambda m \mu).b = \lambda(m.b)\mu$ for any λ and μ in k . From (2.4) and (2.7), we check that b^K and $\epsilon_A \underset{K}{\frown} -$ are B -linear. Thus $\text{HK}_{\bullet}(A, M)$ and $\text{HK}_{\bullet}^{hi}(A, M)$ are graded right B -modules.

Just before Definition 2.3, we have associated to A the \mathbb{F} -central DG algebra

$$\tilde{A} = \text{Hom}_{A^e}(K(A), A) \cong \text{Hom}_{k^e}(W_{\bullet}, A)$$

whose grading is given by the cohomological degree of cochains, whose differential is b_K and whose multiplication is $\underset{K}{\smile}$. We have also mentioned that $\text{Hom}_{k^e}(W_{\bullet}, M)$ and $M \otimes_{k^e} W_{\bullet}$ are DG bimodules over \tilde{A} for the actions of $\underset{K}{\smile}$ and $\underset{K}{\frown}$ respectively, so that $\text{HK}^{\bullet}(A, M)$ and $\text{HK}_{\bullet}(A, M)$ are graded $\text{HK}^{\bullet}(A)$ -bimodules.

For any k -bimodule morphism $f : W_p \rightarrow A$, we verify that

$$f \underset{K}{\frown} (z.b) = (f \underset{K}{\frown} z).b \text{ and } (z.b) \underset{K}{\frown} f = (z \underset{K}{\frown} f).b,$$

so that the right action of B on $M \otimes_{k^e} W_{\bullet}$ is compatible with the \tilde{A} -bimodule structure. Therefore the right action of B on $\text{HK}_{\bullet}(A, M)$ and on $\text{HK}_{\bullet}^{hi}(A, M)$ is compatible with the structure of $\text{HK}^{\bullet}(A)$ -bimodule and of $\text{HK}_{hi}^{\bullet}(A)$ -bimodule respectively. Let us sum up what we have obtained at the level of complexes.

PROPOSITION 3.3. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} , let B be a unital associative \mathbb{F} -algebra and let M be an A -bimodule. We assume that M is a right B -module and that this structure is compatible with the A -bimodule structure. Denote by $\text{Mod-}B$ the category of right B -modules. Then*

- (i) the complex $(M \otimes_{k^e} W_\bullet, b^K)$ is a complex in $\text{Mod-}B$,
 - (ii) the complex $(M \otimes_{k^e} W_\bullet, b^K)$ is a DG-bimodule over the \mathbb{F} -central DG-algebra \tilde{A} ,
 - (iii) the right action of B and the bimodule action of \tilde{A} on $M \otimes_{k^e} W_\bullet$ are compatible.
- In this situation, following Yekutieli [45, Definition 3.8.1], we say that $M \otimes_{k^e} W_\bullet$ is a DG \tilde{A} -bimodule in the abelian category $\text{Mod-}B$.

In order to reflect the fact that the right action of B and the bimodule actions of $\text{HK}^\bullet(A)$ and $\text{HK}_{hi}^\bullet(A)$ on $\text{HK}_\bullet(A, M)$ and $\text{HK}_\bullet^{hi}(A, M)$ respectively are compatible, we shall also say that $\text{HK}_\bullet(A, M)$ is a graded $\text{HK}^\bullet(A)$ -bimodule in $\text{Mod-}B$, and that $\text{HK}_\bullet^{hi}(A, M)$ is a graded $\text{HK}_{hi}^\bullet(A)$ -bimodule in $\text{Mod-}B$. By Lemma 3.1, it is equivalent to saying that $\text{HK}_\bullet(A, M)$ is a graded $\text{HK}^\bullet(A)^e$ - B -bimodule, similarly for $\text{HK}_\bullet^{hi}(A, M)$.

Similarly, $\text{Hom}_{k^e}(W_\bullet, M)$ is a right B -module for the action of b on $f : W_p \rightarrow M$ defined by

$$(f.b)(x_1 \dots x_p) = f(x_1 \dots x_p).b.$$

Then b_K and $\epsilon_A \underset{K}{\smile} -$ are B -linear, so that $\text{HK}^\bullet(A, M)$ and $\text{HK}_{hi}^\bullet(A, M)$ are graded right B -modules. For $g : W_q \rightarrow A$, we have

$$g \underset{K}{\smile} (f.b) = (g \underset{K}{\smile} f).b \text{ and } (f.b) \underset{K}{\smile} g = (f \underset{K}{\smile} g).b.$$

We obtain an analogue of Proposition 3.3, that is, $\text{Hom}_{k^e}(W_\bullet, M)$ is a DG \tilde{A} -bimodule in $\text{Mod-}B$.

PROPOSITION 3.4. *We keep the notation and assumptions of the previous proposition. Then*

- (i) the complex $\text{Hom}_{k^e}(W_\bullet, M)$ is a complex in $\text{Mod-}B$,
- (ii) the complex $\text{Hom}_{k^e}(W_\bullet, M)$ is a DG \tilde{A} -bimodule,
- (iii) the right action of B and the bimodule action of \tilde{A} are compatible on $\text{Hom}_{k^e}(W_\bullet, M)$.

Therefore $\text{HK}^\bullet(A, M)$ is a graded $\text{HK}^\bullet(A)$ -bimodule in $\text{Mod-}B$, and $\text{HK}_{hi}^\bullet(A, M)$ is a graded $\text{HK}_{hi}^\bullet(A)$ -bimodule in $\text{Mod-}B$.

3.3. Application to the Koszul complex $K(A)$

Let us specialise to $B = M = A^e$ as in Example 3.2. Then $M = A \overset{\circ}{\otimes} A$ is a left A^e -module for the outer structure, and a right A^e -module for the inner structure. Our aim is to identify the A -bimodule complex $K(A)$ with the complex $((A \overset{\circ}{\otimes} A) \otimes_{k^e} W_\bullet, b^K)$ endowed with the right action of A^e . The statement is the following.

PROPOSITION 3.5. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} .*

- (i) For any $q \geq 0$, the bilinear map $\varphi_q : (A \overset{\circ}{\otimes} A) \times W_q \rightarrow A \otimes_k W_q \otimes_k A$ defined by

$$\varphi_q(\alpha \otimes \beta, x_1 \dots x_q) = \beta \otimes_k (x_1 \dots x_q) \otimes_k \alpha$$

induces an isomorphism $\tilde{\varphi}_q : (A \overset{\circ}{\otimes} A) \otimes_{k^e} W_q \rightarrow A \otimes_k W_q \otimes_k A$.

- (ii) The direct sum $\tilde{\varphi}$ of the maps $\tilde{\varphi}_q$ is an isomorphism from the complex $((A \overset{\circ}{\otimes} A) \otimes_{k^e} W_\bullet, b^K)$ to the Koszul complex $(K(A), d)$.
- (iii) The isomorphism $\tilde{\varphi}$ is right A^e -linear.

Proof. The A -bimodule $A \overset{\circ}{\otimes} A$ is a k -bimodule for the actions $\lambda(\alpha \otimes \beta)\mu = \lambda\alpha \otimes \beta\mu$, with α and β in A , λ and μ in k , thus it is a right k^e -module for $(\alpha \otimes \beta)(\lambda \otimes \mu) = \mu\alpha \otimes \beta\lambda$. Then it is easy to check that

$$\varphi_q(\mu\alpha \otimes \beta\lambda, x_1 \dots x_q) = \varphi_q(\alpha \otimes \beta, \lambda x_1 \dots x_q \mu),$$

proving the existence of $\tilde{\varphi}_q$. We define similarly an inverse linear map, therefore $\tilde{\varphi}_q$ is an isomorphism, which gives (i).

Let us show that $\tilde{\varphi}$ is a morphism of complexes. From

$$b^K((\alpha \otimes \beta) \otimes_{k^e} x_1 \dots x_q) = (\alpha \otimes (\beta x_1)) \otimes_{k^e} x_2 \dots x_q + (-1)^q((x_q \alpha) \otimes \beta) \otimes_{k^e} x_1 \dots x_{q-1},$$

we get

$$\tilde{\varphi} \circ b^K((\alpha \otimes \beta) \otimes_{k^e} x_1 \dots x_q) = \beta x_1 \otimes_k x_2 \dots x_q \otimes_k \alpha + (-1)^q \beta \otimes_k x_1 \dots x_{q-1} \otimes_k x_q \alpha$$

whose right-hand side is equal to $d(\beta \otimes_k x_1 \dots x_p \otimes_k \alpha)$, as expected.

Let us prove (iii). Here the A -bimodule $A \otimes_k W_q \otimes_k A$ is seen as a right A^e -module. For $z = (\alpha \otimes \beta) \otimes_{k^e} x_1 \dots x_q$ and a, b in A , we have

$$\begin{aligned} \tilde{\varphi}_q(z.(a \otimes b)) &= \tilde{\varphi}_q((\alpha a \otimes b \beta) \otimes_{k^e} x_1 \dots x_q) \\ &= b \beta \otimes_k (x_1 \dots x_q) \otimes_k \alpha a \\ &= \tilde{\varphi}_q(z).(a \otimes b), \end{aligned}$$

therefore $\tilde{\varphi}_q$ is A^e -linear. \square

So $\tilde{\varphi}$ is an isomorphism from the A -bimodule complex $((A \overset{\circ}{\otimes} A) \otimes_{k^e} W_\bullet, b^K)$ whose A -bimodule structure is the inner one, to the A -bimodule complex $K(A)$. Denote by $A\text{-Bimod}$ the category of A -bimodules. According to Proposition 3.3, $(A \overset{\circ}{\otimes} A) \otimes_{k^e} W_\bullet$ is a DG \tilde{A} -bimodule in $A\text{-Bimod}$. We transport this structure via $\tilde{\varphi}$ and we obtain.

PROPOSITION 3.6. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Then the Koszul complex $K(A)$ is a DG \tilde{A} -bimodule in $A\text{-Bimod}$.*

This DG bimodule will play an essential role in the generalisations of Calabi-Yau algebras (Sections 4 and 5). Moreover, $H(K(A))$ is a graded $\text{HK}^\bullet(A)$ -bimodule in $A\text{-Bimod}$, so that $H(\tilde{\varphi}) : \text{HK}_\bullet(A, A \overset{\circ}{\otimes} A) \rightarrow H(K(A))$ is an isomorphism of graded $\text{HK}^\bullet(A)$ -bimodules in $A\text{-Bimod}$.

Let us give explicitly the underlying \tilde{A} -bimodule structure of the DG \tilde{A} -bimodule $K(A)$. Consider $z = (\alpha \otimes \beta) \otimes_{k^e} x_1 \dots x_q$ in $(A \overset{\circ}{\otimes} A) \otimes_{k^e} W_q$ and f in $\text{Hom}_{k^e}(W_p, A)$, we easily derive from (2.7) that the left action of f on $K(A)$ is defined by

$$f \underset{K}{\frown} (\beta \otimes_k x_1 \dots x_q \otimes_k \alpha) = (-1)^{(q-p)p} \beta \otimes_k x_1 \dots x_{q-p} \otimes_k f(x_{q-p+1} \dots x_q) \alpha. \quad (3.1)$$

Analogously, using (2.8), we define the right action of f on $K(A)$ by

$$(\beta \otimes_k x_1 \dots x_q \otimes_k \alpha) \underset{K}{\frown} f = (-1)^{pq} \beta f(x_1 \dots x_p) \otimes_k x_{p+1} \dots x_q \otimes_k \alpha. \quad (3.2)$$

The fundamental formula (2.13) reduces to

$$d(z') = -\epsilon_A \underset{K}{\frown} z' + (-1)^q z' \underset{K}{\frown} \epsilon_A \quad (3.3)$$

on $K(A)$, where $z' = \beta \otimes_k x_1 \dots x_q \otimes_k \alpha$, and

$$\epsilon_A \underset{K}{\frown} z' = (-1)^{q-1} \beta \otimes_k x_1 \dots x_{q-1} \otimes_k x_q \alpha, \quad (3.4)$$

$$z' \underset{K}{\frown} \mathbf{e}_A = (-1)^q \beta x_1 \otimes_k x_2 \dots x_q \otimes_k \alpha. \tag{3.5}$$

The differential $\mathbf{e}_A \underset{K}{\frown} -$ induces a differential, still denoted by ∂_{\frown} , on $H(K(A))$. The homology of $(H(K(A)), \partial_{\frown})$ is denoted by $H^{hi}(K(A))$ and is called the higher homology of $K(A)$. Then $H^{hi}(K(A))$ is a graded $\mathrm{HK}_{hi}^\bullet(A)$ -bimodule in $A\text{-Bimod}$ and $H(H(\tilde{\varphi})) : \mathrm{HK}_{hi}^\bullet(A, A \overset{\circ}{\otimes} A) \rightarrow H^{hi}(K(A))$ is an isomorphism of graded $\mathrm{HK}_{hi}^\bullet(A)$ -bimodules in $A\text{-Bimod}$.

4. Poincaré Van den Bergh duality of preprojective algebras

4.1. Preprojective algebras

Throughout this section, Δ is a connected graph whose vertex set and edge set are finite. Following a usual presupposition in the papers devoted to Hochschild (co)homology of preprojective algebras, we assume that the graph Δ is not labelled, that is, the labels of the edges are all equal to $(1, 1)$ [4, Definition 4.1.9]. In particular, the Dynkin graphs are limited to types ADE, and the Euclidean (or extended) Dynkin graphs are limited to types $\tilde{A}\tilde{D}\tilde{E}$ [4, Definition 4.5.1].

Let Q be a quiver whose underlying graph is Δ . Define a quiver Q^* whose vertex set is Q_0 and whose arrow set is $Q_1^* = \{a^*; a \in Q_1\}$ where $\mathfrak{s}(a^*) = \mathfrak{t}(a)$ and $\mathfrak{t}(a^*) = \mathfrak{s}(a)$. Let \overline{Q} be the double quiver of Q , that is, the quiver whose vertex set is $\overline{Q}_0 = Q_0$ and whose arrow set is the disjoint union $\overline{Q}_1 = Q_1 \cup Q_1^*$. We shall view $(-)^*$ as an involution of \overline{Q}_1 .

Let \mathbb{F} be a field. As before, we denote the ring $\mathbb{F}Q_0$ by k and the k -bimodule $\mathbb{F}\overline{Q}_1$ by V and we identify the graded k -algebras $T_k(V) \cong \mathbb{F}\overline{Q}$ (see Subsection 2.1).

The preprojective algebra associated with the graph Δ over the field \mathbb{F} is the quadratic k -algebra $A(\Delta)$ over \overline{Q} defined by $A(\Delta) = \mathbb{F}\overline{Q}/(R)$, where the sub- k -bimodule R of $\mathbb{F}\overline{Q}_2$ is generated by

$$\sigma_i := \sum_{\substack{a \in Q_1 \\ \mathfrak{t}(a)=i}} aa^* - \sum_{\substack{a \in Q_1 \\ \mathfrak{s}(a)=i}} a^*a = \sum_{\substack{a \in \overline{Q}_1 \\ \mathfrak{t}(a)=i}} \varepsilon(a)aa^* \quad \text{for all } i \in Q_0,$$

where $\varepsilon(a) = 1$ if $a \in Q_1$, $\varepsilon(a) = -1$ if $a \in Q_1^*$.

If Q' is another quiver whose underlying graph is Δ , and R' is the sub- k -bimodule of $\mathbb{F}\overline{Q}'_2$ generated by the relations $\sigma'_i = \sum_{\substack{a \in \overline{Q}'_1 \\ \mathfrak{t}(a)=i}} \varepsilon(a)aa^*$ for all $i \in Q'_0 = Q_0$, then the preprojective

algebras $\mathbb{F}\overline{Q}/(R)$ and $\mathbb{F}\overline{Q}'/(R')$ are isomorphic, the isomorphism being given by exchanging pairs of arrows a and a^* and changing the sign of one arrow in each pair (see [13, Remark 2.2(3)], or [38, Lemma 1.3.7] for a complete proof in the case of generalised preprojective algebras). Therefore, according to Subsection 2.7, the quadratic k -algebra $A(\Delta)$ and the (general, higher) Koszul calculus of $A(\Delta)$ depend only on the graph Δ and not on Q , justifying the notation $A(\Delta)$. If Δ is a tree, $A(\Delta)$ is isomorphic to the preprojective algebra defined without signs (that is, $\varepsilon(a) = 1$ for all $a \in \overline{Q}_1$, as in [18, 19, 20]).

If $\Delta = A_1$, then $A(\Delta) = k$. If $\Delta = A_2$, then $R = \mathbb{F}\overline{Q}_2$ and $A(\Delta) = \mathbb{F}Q_0 \oplus \mathbb{F}\overline{Q}_1$. These quadratic k -algebras are Koszul, but they are the only exceptions among the Dynkin graphs. More precisely, the following standard result holds, for which we just give proof references (see also [10, Corollary 4.3]).

PROPOSITION 4.1. *Assume that the graph Δ is distinct from A_1 and A_2 . The following are equivalent.*

- (i) Δ is Dynkin of type ADE.

- (ii) $A(\Delta)$ is not Koszul.
 (iii) $A(\Delta)$ is finite dimensional.

Proof. The equivalence (i) \Leftrightarrow (ii) is treated in [33] if Δ is a tree, in [30] otherwise. The equivalence (i) \Leftrightarrow (iii) for any Dynkin graph is cited in [34] as a result by Gelfand and Ponomarev [27]. \square

Sections 2 and 3 can be applied to preprojective algebras. For example, according to the remark following Lemma 2.6, the fundamental 1-cocycle $\epsilon_{A(\Delta)}$ is not a coboundary if Δ has a loop or if $\text{char } \mathbb{F} \neq 2$ and $\Delta \neq A_1$. In the remainder of this section, we often abbreviate $A(\Delta)$ to A and we freely use notations and results from Sections 2 and 3.

4.2. The Koszul complex $K(A)$ has length 2

If $\Delta = A_1$, then $K(A)$ has length 0. If $\Delta = A_2$, then $K(A)$ has infinite length. However, when Δ is not Dynkin ADE, the algebra $A = A(\Delta)$ has global dimension 2 (this is a consequence of [10, Proposition 4.2], inspired by manuscript notes of Crawley-Boevey), and since $k \cong \mathbb{F}^{|Q_0|}$ is separable, it follows that the minimal projective A -bimodule resolution of A , which is $K(A)$ because A is Koszul, has length 2 (see for instance [37, Proposition 3.18]).

Actually, the fact that the length of $K(A)$ is 2 is true for all graphs Δ other than A_1 and A_2 , and we now give a unified proof of this.

THEOREM 4.2. *Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. Then the Koszul complex $K(A)$ of A has length 2. Consequently, $\text{HK}^p(A, M) \cong \text{HK}_p(A, M) = 0$ for all A -bimodules M and all $p \geq 3$.*

Proof. From the defining equality (2.2) of W_p , we have $W_p = (W_{p-1} \otimes_k V) \cap (V \otimes_k W_{p-1})$ for all $p \geq 3$. Moreover $R \neq 0$, therefore it is enough to prove that $W_3 = 0$, that is, $(R \otimes_k V) \cap (V \otimes_k R) = 0$. For that, we only assume that $\Delta \neq A_1$. Our goal is to prove that $W_3 \neq 0$ implies $\Delta = A_2$.

Let u be a non-zero element in W_3 , viewed as an element in $\mathbb{F}\overline{Q}_3$. There exist vertices e, f in Q_0 such that $eu f \neq 0$, therefore we may assume that u is in $eW_3 f$. Then u can be written uniquely as

$$u = \sum_{\substack{a \in e\overline{Q}_1 \\ \alpha \in e\overline{Q}_1 f}} \lambda_\alpha \epsilon(a) a a^* \alpha = \sum_{\substack{b \in f\overline{Q}_1 \\ \beta \in f\overline{Q}_1 e}} \mu_\beta \epsilon(b) \beta b b^*.$$

We now use the fact that \overline{Q}_1 is the disjoint union of Q_1 and Q_1^* and the definition of ϵ to write

$$\begin{aligned} u &= \sum_{\substack{\alpha \in eQ_1 f \\ a \in eQ_1}} \lambda_\alpha a a^* \alpha - \sum_{\substack{\alpha \in eQ_1 f \\ a \in Q_1 e}} \lambda_\alpha a^* a \alpha + \sum_{\substack{\alpha \in fQ_1 e \\ a \in eQ_1}} \lambda_{\alpha^*} a a^* \alpha^* - \sum_{\substack{\alpha \in fQ_1 e \\ a \in Q_1 e}} \lambda_{\alpha^*} a^* a \alpha^* \\ &= \sum_{\substack{\beta \in eQ_1 f \\ b \in fQ_1}} \mu_\beta \beta b b^* - \sum_{\substack{\beta \in eQ_1 f \\ b \in Q_1 f}} \mu_\beta \beta b^* b + \sum_{\substack{\beta \in fQ_1 e \\ b \in fQ_1}} \mu_{\beta^*} \beta^* b b^* - \sum_{\substack{\beta \in fQ_1 e \\ b \in Q_1 f}} \mu_{\beta^*} \beta^* b^* b. \end{aligned}$$

From these expressions, we obtain the following identities in the path algebra $\mathbb{F}\overline{Q}$:

$$\sum_{\substack{\alpha \in fQ_1 e \\ a \in eQ_1}} \lambda_\alpha a a^* \alpha^* = 0 \quad (4.1)$$

$$\sum_{\substack{\alpha \in eQ_1f \\ a \in Q_1e}} \lambda_\alpha a^* a \alpha = 0 \quad (4.2)$$

$$\sum_{\substack{\beta \in eQ_1f \\ b \in fQ_1}} \mu_\beta \beta b b^* = 0 \quad (4.3)$$

$$\sum_{\substack{\beta \in fQ_1e \\ b \in Q_1f}} \mu_{\beta^*} \beta^* b^* b = 0 \quad (4.4)$$

$$\sum_{\substack{\alpha \in eQ_1f \\ a \in eQ_1}} \lambda_\alpha a a^* \alpha = - \sum_{\substack{\beta \in eQ_1f \\ b \in Q_1f}} \mu_\beta \beta b^* b \quad (4.5)$$

$$\sum_{\substack{\alpha \in fQ_1e \\ a \in Q_1e}} \lambda_\alpha a^* a \alpha^* = - \sum_{\substack{\beta \in fQ_1e \\ b \in fQ_1}} \mu_\beta \beta^* b b^* \quad (4.6)$$

Indeed, identity (4.1) follows from the fact that no other path that occurs in the expressions of u ends with two arrows in Q^* , and the other identities are obtained from similar arguments.

In the path algebra $\mathbb{F}\overline{Q}$, where there are no relations between paths, the identities (4.1) to (4.4) above are equivalent to

$$\forall \alpha \in fQ_1e, \forall a \in eQ_1, \lambda_{\alpha^*} = 0 \quad (4.7)$$

$$\forall \alpha \in eQ_1f, \forall a \in Q_1e, \lambda_\alpha = 0 \quad (4.8)$$

$$\forall \beta \in eQ_1f, \forall a \in fQ_1, \lambda_\beta = 0 \quad (4.9)$$

$$\forall \beta \in fQ_1e, \forall a \in Q_1f, \lambda_{\beta^*} = 0. \quad (4.10)$$

We have assumed that $u \neq 0$, so that either there exists $\alpha \in eQ_1f$ such that $\lambda_\alpha \neq 0$ or there exists $\alpha \in fQ_1e$ such that $\lambda_{\alpha^*} \neq 0$, using the first expression of u . We separate the two cases.

Assume that there exists $\alpha \in eQ_1f$ such that $\lambda_\alpha \neq 0$. Then it follows from identity (4.8) that Q_1e is empty. From (4.5), for all $a \in eQ_1$, there exist $\beta \in eQ_1f$ and $b \in Q_1f$ such that $\lambda_\alpha a a^* \alpha = -\mu_\beta \beta b^* b$. Hence $\beta = a = b = \alpha$ and therefore $eQ_1 = \{\alpha\} = eQ_1f$ and $\mu_\beta = -\lambda_\alpha \neq 0$. From (4.9), it follows that fQ_1 is empty. Finally, (4.5) becomes

$$\lambda_\alpha \alpha \alpha^* \alpha = \lambda_\alpha \alpha \alpha^* \alpha + \lambda_\alpha \sum_{\substack{b \in Q_1f \\ b \neq \alpha}} \alpha b^* b$$

so that $\sum_{\substack{b \in Q_1f \\ b \neq \alpha}} \alpha b^* b = 0$ and hence $Q_1f = \{\alpha\}$.

We have proved that $Q_1e = \emptyset = fQ_1$ so that in particular $e \neq f$, and that $Q_1f = \{\alpha\} = eQ_1 = eQ_1f$. Finally, $Q = e \xrightarrow{\alpha} f$ and $\Delta = A_2$.

In the case where there exists $\alpha \in fQ_1e$ such that $\lambda_{\alpha^*} \neq 0$, a similar proof using (4.7), (4.10) and (4.6) shows that $Q = f \xleftarrow{\alpha} e$ and $\Delta = A_2$. \square

As an immediate consequence of Proposition 2.8 and Theorem 4.2, we obtain that in the Koszul calculus of $A(\Delta)$, the Koszul cup product is graded commutative and the Koszul cap product is graded symmetric. The precise statement is the following.

COROLLARY 4.3. *Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. We consider an A -bimodule M . For any $\alpha \in \text{HK}^\bullet(A, M)$, $\beta \in \text{HK}^\bullet(A)$ and $\gamma \in \text{HK}_\bullet(A)$, we have the identities*

$$[\alpha, \beta]_{\overline{K}} = 0, \quad (4.11)$$

$$[\alpha, \gamma]_{\overline{K}} = 0. \quad (4.12)$$

The same conclusion holds if $\Delta = A_1$ (obvious) and if $\Delta = A_2$ (because A is Koszul).

4.3. Duality in Koszul (co)homology of preprojective algebras

There is a remarkable duality between Koszul homology and cohomology for preprojective algebras. This duality is realised as a cap action by a Koszul 2-chain $\omega_0 \in A \otimes_{k^e} R$ defined for any graph Δ by

$$\omega_0 = \sum_{i \in Q_0} e_i \otimes_{k^e} \sigma_i = \sum_{i \in Q_0} e_i \otimes \sigma_i.$$

From $\sigma_i = \sum_{a \in \bar{Q}_1, t(a)=i} \varepsilon(a)aa^*$, we get

$$\omega_0 = \sum_{a \in \bar{Q}_1} 1 \otimes_{k^e} \varepsilon(a)aa^* = - \sum_{a \in \bar{Q}_1} 1 \otimes_{k^e} \varepsilon(a)a^*a. \quad (4.13)$$

Then it is easy to check that ω_0 is a Koszul 2-cycle. Being homogeneous of weight 0, ω_0 is not a 2-boundary whenever $\Delta \neq A_1$.

In the following statement, we need the DG algebra \tilde{A} and the DG \tilde{A} -bimodules $\text{Hom}_{k^e}(W_\bullet, M)$ and $M \otimes_{k^e} W_\bullet$, introduced just before Definition 2.3.

THEOREM 4.4. *Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. Consider the Koszul 2-cycle $\omega_0 = \sum_{i \in Q_0} e_i \otimes \sigma_i \in A \otimes_{k^e} R$. For each Koszul p -cochain f with coefficients in an A -bimodule M , we define the Koszul $(2-p)$ -chain $\theta_M(f)$ with coefficients in M by*

$$\theta_M(f) = \omega_0 \underset{K}{\frown} f. \quad (4.14)$$

Then the equalities

$$\theta_{M \otimes_A N}(f \underset{K}{\smile} g) = \theta_M(f) \underset{K}{\frown} g = f \underset{K}{\frown} \theta_N(g) \quad (4.15)$$

hold for any Koszul cochains f and g with coefficients in A -bimodules M and N respectively.

Moreover the linear map $\theta_M : \text{Hom}_{k^e}(W_\bullet, M) \rightarrow M \otimes_{k^e} W_{2-\bullet}$ is an isomorphism of DG \tilde{A} -bimodules.

It follows that $H(\theta_M) : \text{HK}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}^\bullet(A, M)$ is an isomorphism of graded $\text{HK}^\bullet(A)$ -bimodules and that $H(H(\theta_M)) : \text{HK}_{hi}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}^{hi}(A, M)$ is an isomorphism of graded $\text{HK}_{hi}^\bullet(A)$ -bimodules.

Proof. First we show that $f \underset{K}{\frown} \omega_0 = \omega_0 \underset{K}{\frown} f$ for all $f \in \text{Hom}_{k^e}(W_p, M)$. Using the definition of ω_0 and the equalities (4.13), (2.7) and (2.8), we obtain for $p = 0, 1, 2$,

$$\begin{aligned} f \underset{K}{\frown} \omega_0 &= \sum_{i \in Q_0} (f(1)e_i) \otimes_{k^e} \sigma_i = \sum_{i \in Q_0} (e_i f(1)) \otimes_{k^e} \sigma_i = \omega_0 \underset{K}{\frown} f, \\ f \underset{K}{\frown} \omega_0 &= - \sum_{a \in \bar{Q}_1} \varepsilon(a)(f(a^*)1) \otimes_{k^e} a = \sum_{a \in \bar{Q}_1} \varepsilon(a)(1f(a)) \otimes_{k^e} a^* = \omega_0 \underset{K}{\frown} f, \\ f \underset{K}{\frown} \omega_0 &= \sum_{i \in Q_0} (e_i f(\sigma_i)) \otimes_{k^e} 1 = \sum_{i \in Q_0} (f(\sigma_i)e_i) \otimes_{k^e} 1 = \omega_0 \underset{K}{\frown} f. \end{aligned}$$

Next, for $f \in \text{Hom}_{k^e}(W_p, M)$ and $g \in \text{Hom}_{k^e}(W_q, M)$, we have

$$\theta_{M \otimes_A N}(f \underset{K}{\smile} g) = \omega_0 \underset{K}{\frown} (f \underset{K}{\smile} g) = (\omega_0 \underset{K}{\frown} f) \underset{K}{\frown} g = (f \underset{K}{\frown} \omega_0) \underset{K}{\frown} g = f \underset{K}{\frown} (\omega_0 \underset{K}{\frown} g),$$

providing equalities (4.15). Therefore $\theta_M : \text{Hom}_{k^e}(W_\bullet, M) \rightarrow M \otimes_{k^e} W_{2-\bullet}$ is a morphism of graded \tilde{A} -bimodules, where \tilde{A} is just considered as a graded algebra. It remains to examine what happens for the Koszul differentials.

Assuming $M = A$ in the equalities (4.15), we derive

$$\theta_N([f, g]_{\widehat{K}}) = [\theta_A(f), g]_{\widehat{K}} = [f, \theta_N(g)]_{\widehat{K}}.$$

Combining $\theta_N([\mathbf{e}_A, g]_{\widehat{K}}) = [\mathbf{e}_A, \theta_N(g)]_{\widehat{K}}$ with $b_K = -[\mathbf{e}_A, -]_{\widehat{K}}$ and $b^K = -[\mathbf{e}_A, -]_{\widehat{K}}$, we deduce that θ_M is a morphism of complexes, thus a morphism of DG \tilde{A} -bimodules.

We prove that θ_M is an isomorphism by giving an inverse map $\eta : M \otimes_{k^e} W_{2-\bullet} \rightarrow \text{Hom}_{k^e}(W_{\bullet}, M)$. We define $\eta_p : M \otimes_{k^e} W_{2-p} \rightarrow \text{Hom}_{k^e}(W_p, M)$ for $p = 0, 1, 2$, by

$$\begin{aligned} \eta_0(m \otimes_{k^e} \sigma_i)(e_j) &= \delta_{ij} e_j m e_i \\ \eta_1(m \otimes_{k^e} a)(b) &= \delta_{ba^*} \varepsilon(b) \mathbf{t}(b) m \mathbf{s}(b) \quad \text{for any arrows } a \text{ and } b \text{ of } \overline{Q} \\ \eta_2(m \otimes_{k^e} e_i)(\sigma_j) &= \delta_{ij} e_j m e_i \end{aligned}$$

where δ is the Kronecker symbol. It is routine to verify that these linear maps are well-defined and form an inverse map for θ_M .

Finally the isomorphism $H(\theta_M)$ of graded $\text{HK}^\bullet(A)$ -bimodules satisfies

$$H(\theta_M)(\overline{\mathbf{e}}_A \underset{K}{\frown} \alpha) = \overline{\mathbf{e}}_A \underset{K}{\frown} H(\theta_M)(\alpha)$$

for all $\alpha \in \text{HK}^\bullet(A, M)$. Therefore $H(\theta_M)$ is a morphism of complexes for higher (co)homologies. Taking higher (co)homologies, we get a $\text{HK}_{hi}^\bullet(A)$ -bimodule isomorphism

$$H(H(\theta_M)) : \text{HK}_{hi}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}^{hi}(A, M).$$

□

By analogy with the Poincaré duality in singular (co)homology [31] and with the Van den Bergh duality in Hochschild (co)homology [41, 32], we say that the isomorphism

$$H(\theta_M) = \overline{\omega}_0 \underset{K}{\frown} - : \text{HK}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}(A, M)$$

is a Poincaré Van den Bergh duality for Koszul (co)homology, of fundamental class $\overline{\omega}_0$, where $\overline{\omega}_0 \in \text{HK}_2(A)_0$. In the next subsection, we extract from this duality a generalisation of the 2-Calabi-Yau property.

Unless Δ has no loop and $\text{char } \mathbb{F} = 2$, the class $\overline{\mathbf{e}}_A \in \text{HK}^1(A)_1$ is non-zero, hence

$$H(\theta_A)(\overline{\mathbf{e}}_A) = \overline{\omega}_0 \underset{K}{\frown} \overline{\mathbf{e}}_A = \partial_{\frown}(\overline{\omega}_0)$$

is non-zero in $\text{HK}_1(A)_1$. Consequently, the fundamental class $\overline{\omega}_0$ of the Poincaré Van den Bergh duality is not a cycle for the higher Koszul homology, so that the isomorphism $H(H(\theta_M))$ cannot be naturally expressed as a cap action.

The class $H(\theta_A)(\overline{\mathbf{e}}_A)$ is the class of the Koszul 1-cycle $\omega_0 \underset{K}{\frown} \mathbf{e}_A$ where

$$\omega_0 \underset{K}{\frown} \mathbf{e}_A = \sum_{a \in \overline{Q}_1} \varepsilon(a) a \otimes_{k^e} a^* = \sum_{a \in \overline{Q}_1} (a \otimes_{k^e} a^* - a^* \otimes_{k^e} a).$$

It is interesting to view the last element as the image by the canonical linear map $\text{can} : V \otimes_k V \rightarrow V \otimes_{k^e} V$ of the element

$$w = \sum_{a \in \overline{Q}_1} (a \otimes_k a^* - a^* \otimes_k a) = \sum_{a \in \overline{Q}_1} \varepsilon(a) a a^* \in R \subseteq V \otimes_k V.$$

In the identification $V \otimes_k V \cong \mathbb{F}\overline{Q}_2$, $V \otimes_{k^e} V$ is identified with the subspace of cycles of length 2 and the map can is identified with the projection whose kernel is the space spanned by the non-cyclic paths. Since R is generated by the cycles σ_i , we can make the identification $\omega_0 \underset{K}{\frown} \mathbf{e}_A = w$. The element w was defined in [14, Proposition 8.1.1] as a representative of a bi-symplectic 2-form ω . Bi-symplectic 2-forms were introduced by Crawley-Boevey, Etingof and Ginzburg as

an essential ingredient of the Hamiltonian reduction in noncommutative geometry [14]; they are related to the double Poisson algebras defined by Van den Bergh [42].

REMARK 4.5. Assume that $\Delta = A_2$, so that A is defined by the quiver

$$\overline{Q} \quad 0 \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{a^*} \end{array} 1$$

subject to the relations $\sigma_0 = -a^*a$, $\sigma_1 = aa^*$. Then the statement of Theorem 4.4 is valid in a weaker form, namely the isomorphisms involved are only morphisms. Moreover, θ_M is bijective only in degree q , $0 \leq q \leq 2$, with the same inverse η_q . More generally, for any $p \geq 1$, the Koszul $(2p)$ -cycle

$$\omega_{p-1} = e_1 \otimes (aa^*)^p - e_0 \otimes (a^*a)^p$$

provides a morphism $\omega_{p-1} \underset{K}{\frown} -$ which is bijective only in degree q , $0 \leq q \leq 2p$. From that, we deduce an isomorphism $\overline{\omega}_{p-1} \underset{K}{\frown} -$ from $\text{HK}^q(A, M)$ to $\text{HK}_{2p-q}(A, M)$ for $0 < q < 2p$. Varying p , we obtain the following duality and 2-periodicity

$$\begin{aligned} \text{HK}^q(A, M) &\cong \text{HK}_q(A, M) \cong \text{HK}^1(A, M), \quad q \text{ odd } \geq 1 \\ \text{HK}^q(A, M) &\cong \text{HK}_q(A, M) \cong \text{HK}^2(A, M), \quad q \text{ even } \geq 2. \end{aligned}$$

Using this for $M = A$, it is straightforward to compute explicitly the restricted Koszul calculus of A . We leave the details to the reader. Notice that, since A is Koszul, we have the same duality and 2-periodicity for Hochschild (co)homology, recovering a known result as a consequence of remarkable isomorphisms due to Eu and Schedler [24, Theorem 2.3.27, Theorem 2.3.47], here applied to [24, Example 2.3.10, Corollary 2.1.13].

4.4. Deriving an adapted 2-Calabi-Yau property

Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. Let M be an A -bimodule. Assume that B is a unital associative algebra such that M is a right B -module compatible with the A -bimodule structure (see Subsection 3.1). Denote by $\text{Mod-}B$ the category of right B -modules. Recall that \tilde{A} denotes the DG algebra $(\text{Hom}_{k^e}(W_\bullet, A), b_K, \underset{K}{\smile})$.

According to Subsection 3.2, $M \otimes_{k^e} W_\bullet$ and $\text{Hom}_{k^e}(W_\bullet, M)$ are DG \tilde{A} -bimodules in $\text{Mod-}B$. Moreover, $\text{HK}_\bullet(A, M)$ and $\text{HK}^\bullet(A, M)$ are graded $\text{HK}^\bullet(A)$ -bimodules in $\text{Mod-}B$. Finally, $\text{HK}_\bullet^{hi}(A, M)$ and $\text{HK}_{hi}^\bullet(A, M)$ are graded $\text{HK}_{hi}^\bullet(A)$ -bimodules in $\text{Mod-}B$.

LEMMA 4.6. *The map $\theta_M : \text{Hom}_{k^e}(W_\bullet, M) \rightarrow M \otimes_{k^e} W_{2-\bullet}$ is an isomorphism of DG \tilde{A} -bimodules in $\text{Mod-}B$. Moreover, $H(\theta_M) : \text{HK}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}(A, M)$ is an isomorphism of graded $\text{HK}^\bullet(A)$ -bimodules in $\text{Mod-}B$, and $H(H(\theta_M)) : \text{HK}_{hi}^\bullet(A, M) \rightarrow \text{HK}_{2-\bullet}^{hi}(A, M)$ is an isomorphism of graded $\text{HK}_{hi}^\bullet(A)$ -bimodules in $\text{Mod-}B$.*

Proof. It is enough to prove that $\theta_M : f \mapsto \omega_0 \underset{K}{\frown} f$ is B -linear. For a k -bimodule morphism $f : W_p \rightarrow M$, $z = a \otimes_{k^e} x_1 \dots x_q \in A \otimes_{k^e} W_q$ and $b \in B$, we verify the identities

$$(f.b) \underset{K}{\frown} z = (f \underset{K}{\frown} z).b \text{ and } z \underset{K}{\frown} (f.b) = (z \underset{K}{\frown} f).b.$$

The first one uses the fact that the right actions of A and B on M commute, while the second one uses the fact that M is an A - B -bimodule (see (iii) in Lemma 3.1). Applying the second one to $z = \omega_0$, we obtain that θ_M is B -linear. \square

We specialise this lemma to $M = B = A^e$ and, using the isomorphism $\tilde{\varphi}$ in Subsection 3.3, we identify $A^e \otimes_{k^e} W_\bullet$ with $K(A)$ to get the next proposition.

PROPOSITION 4.7. *Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. The map*

$$\theta_{A^e} : \text{Hom}_{k^e}(W_\bullet, A^e) \rightarrow K(A)_{2-\bullet}$$

is an isomorphism of DG \tilde{A} -bimodules in $A\text{-Bimod}$. Moreover,

$$H(\theta_{A^e}) : \text{HK}^\bullet(A, A^e) \rightarrow H_{2-\bullet}(K(A))$$

is an isomorphism of graded $\text{HK}^\bullet(A)$ -bimodules in $A\text{-Bimod}$.

The homology of $K(A)$ is isomorphic to A in degree 0, and to 0 in degree 1, hence we obtain a generalisation of the 2-Calabi-Yau property, formulated as follows.

THEOREM 4.8. *Let $A = A(\Delta)$ be a preprojective algebra over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$. Then the $\text{HK}^\bullet(A)^e\text{-}A^e$ -bimodules $\text{HK}^\bullet(A, A^e)$ and $H_{2-\bullet}(K(A))$ are isomorphic. In particular, we have the following.*

- (i) *The A -bimodule $\text{HK}^2(A, A^e)$ is isomorphic to the A -bimodule A .*
- (ii) *$\text{HK}^1(A, A^e) = 0$.*
- (iii) *The A -bimodule $\text{HK}^0(A, A^e)$ is isomorphic to the A -bimodule $H_2(K(A))$.*

Since $H_1(K(A)) \cong \text{HK}^1(A, A^e) = 0$, the higher Koszul differentials vanish. Therefore $H_p^{hi}(K(A)) \cong H_p(K(A))$, $\text{HK}_{hi}^p(A, A^e) \cong \text{HK}^p(A, A^e)$ and $H(H(\theta_{A^e})) \cong H(\theta_{A^e})$.

From the generator $1 \otimes_k 1$ of the A -bimodule $H_0(K(A))$, we draw from (i) a generator of the free A -bimodule $\text{HK}^2(A, A^e)$ defined as the class of $f : R \rightarrow A \overset{\circ}{\otimes} A$ with $f(\sigma_i) = e_i \otimes e_i$ for any i .

In (iii), the A -bimodules are never 0 when Δ is Dynkin of types ADE since A is not Koszul in this case. This situation is drastically different from the 2-Calabi-Yau property defined by Ginzburg in terms of the Hochschild cohomology spaces $\text{HH}^p(A, A^e)$ [28, §3.2]. In Ginzburg's definition, $\text{HH}^p(A, A^e) = 0$ for all $p < 2$.

5. Generalisations of Calabi-Yau algebras

5.1. Duality for Koszul complex Calabi-Yau algebras

From Theorem 4.8, we are led to introduce a general definition in the framework of quiver algebras with homogeneous quadratic relations (see Section 2). The notation introduced in Section 2 stands throughout. We are interested in quadratic k -algebras $A = T_k(V)/(R)$ over a finite quiver \mathcal{Q} as defined in Subsection 2.1, and in the Koszul calculus of A as presented in the remainder of Section 2. Note that $\mathcal{Q}_1 = \overline{\mathcal{Q}}_1$ if we want to specialise to preprojective algebras.

For the definition of the bounded derived category $\mathcal{D}^b(\mathcal{C})$ of an abelian category \mathcal{C} , we refer to [44, Chapter 10]. Recall that $A\text{-Bimod}$ denotes the category of A -bimodules.

DEFINITION 5.1. *Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Let $n \geq 0$ be an integer. We say that A is Koszul complex Calabi-Yau (Kc-Calabi-Yau) of dimension n , or n -Kc-Calabi-Yau, if*

- (i) the Koszul bimodule complex $K(A)$ of A has length n , and
- (ii) $\mathrm{RHom}_{A^e}(K(A), A^e) \cong K(A)[-n]$ in $\mathcal{D}^b(A\text{-Bimod})$.

Property (ii) is equivalent to saying that there is an A -bimodule quasi-isomorphism from $\mathrm{Hom}_{k^e}(W_\bullet, A^e)$ to $K(A)[-n]$. According to Theorem 4.2 and Proposition 4.7, a preprojective algebra $A(\Delta)$ over \mathbb{F} with $\Delta \neq A_1$ and $\Delta \neq A_2$ is Kc-Calabi-Yau of dimension 2. In fact, the isomorphism θ_{A^e} induces an isomorphism $\mathrm{RHom}_{A^e}(K(A), A^e) \rightarrow K(A)[-2]$ in $\mathcal{D}^b(A\text{-Bimod})$.

Let us recall Ginzburg’s definition of Calabi-Yau algebras [28, Definition 3.2.3] as reformulated by Van den Bergh [43, Definition 8.2]. We shall apply this definition to quadratic quiver algebras by considering it as \mathbb{F} -algebras.

DEFINITION 5.2. An associative \mathbb{F} -algebra A is said to be Calabi-Yau of dimension n if

- (i) A is homologically smooth, that is, A has a bounded resolution by finitely generated projective A -bimodules,
- (ii) $\mathrm{RHom}_{A^e}(A, A^e) \cong A[-n]$ in $\mathcal{D}^b(A\text{-Bimod})$.

Definition 5.1 is a true generalisation of Definition 5.2 for quadratic quiver algebras. If Δ is Dynkin of type ADE, $A(\Delta)$ is not Calabi-Yau in Ginzburg’s definition since $A(\Delta)$ is not homologically smooth in this case (the minimal projective resolution of $A(\Delta)$ has infinite length). However, the two definitions coincide if A is Koszul.

PROPOSITION 5.3. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Assume that A is Koszul. Then A is n -Kc-Calabi-Yau if and only if A is n -Calabi-Yau.

Proof. Assume that A is n -Kc-Calabi-Yau. Property (i) and the fact that A is Koszul show that A is homologically smooth. Furthermore, $K(A) \cong A$ in $\mathcal{D}^b(A\text{-Bimod})$. Thus $\mathrm{RHom}_{A^e}(A, A^e) \cong A[-n]$ in $\mathcal{D}^b(A\text{-Bimod})$, and we recover Definition 5.2.

Assume that A is n -Calabi-Yau. We know that n is equal to the projective dimension of the A -bimodule A [41] which in turn is equal to the length of a minimal projective resolution of A (see for instance [5]). Hence $K(A)$ has length n and $K(A) \cong A$ in $\mathcal{D}^b(A\text{-Bimod})$, which allows us to conclude that A is n -Kc-Calabi-Yau. \square

If the graph Δ is not Dynkin ADE, we know that $A(\Delta)$ is Koszul (Proposition 4.1), thus we recover the fact that $A(\Delta)$ is 2-Calabi-Yau [14, 9].

In Subsection 2.8, we have seen that $K(A)$ and the minimal projective resolution $P(A)$ coincide up to the homological degree 2. Therefore, if $n \in \{0, 1\}$ and if A is n -Calabi-Yau or n -Kc-Calabi-Yau, then $P(A) = K(A)$ so that A is Koszul, and it follows that the two definitions are equivalent when $n \in \{0, 1\}$.

PROPOSITION 5.4. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Then A is Calabi-Yau of dimension 0 if and only if $\mathcal{Q}_1 = \emptyset$.

We leave the proof as an exercise. If A is Calabi-Yau of dimension 1, then $R = 0$, that is, $A = T_k(V) \cong \mathbb{F}\mathcal{Q}$ with $\mathcal{Q}_1 \neq \emptyset$. It is indeed 1-Calabi-Yau if \mathcal{Q} has only one vertex and one loop, but we have not yet found other examples when \mathcal{Q} is connected.

If A is n -Calabi-Yau, the Van den Bergh duality theorem states that the vector spaces $\mathrm{HH}^p(A, M)$ and $\mathrm{HH}_{n-p}(A, M)$ are isomorphic [41]. From Definition 5.1, we draw an analogous duality theorem for Koszul homology/cohomology.

THEOREM 5.5. *Let A be a Koszul complex Calabi-Yau algebra of dimension n over \mathcal{Q} . Then for any A -bimodule M , the vector spaces $\mathrm{HK}^p(A, M)$ and $\mathrm{HK}_{n-p}(A, M)$ are isomorphic.*

Proof. Denote by $\mathrm{Vect}_{\mathbb{F}}$ the category of \mathbb{F} -vector spaces. For any A -bimodule M , the left derived functor $M \overset{L}{\otimes}_{A^e} -$ and the right derived functor $\mathrm{RHom}_{A^e}(-, M)$ are defined from $\mathcal{D}^b(A\text{-Bimod})$ to $\mathcal{D}^b(\mathrm{Vect}_{\mathbb{F}})$ [44, Chapter 10].

Our proof is based on a natural transformation depending on an A -bimodule M . Let $F : A\text{-Bimod} \rightarrow \mathrm{Vect}_{\mathbb{F}}$ be the functor $F : N \mapsto \mathrm{Hom}_{A^e}(N, M)$ where M and N are seen as right A^e -modules. Specialising to $M = A^e$ in F , we define a functor $G : A\text{-Bimod} \rightarrow A\text{-Bimod}$. Let $H : A\text{-Bimod} \rightarrow \mathrm{Vect}_{\mathbb{F}}$ be the functor $H : N' \mapsto M \otimes_{A^e} N'$ where N' is viewed as a left A^e -module. Then we define a linear map

$$\phi_M : M \otimes_{A^e} \mathrm{Hom}_{A^e}(N, A^e) \rightarrow \mathrm{Hom}_{A^e}(N, M)$$

by $\phi_M(m \otimes_{A^e} g)(x) = m.g(x)$ for $m \in M, g \in \mathrm{Hom}_{A^e}(N, A^e)$ and $x \in N$. This map is functorial in N , defining a natural transformation $\phi_M : H \circ G \Rightarrow F$.

If the A -bimodule P is projective and finitely generated,

$$\phi_M : M \otimes_{A^e} \mathrm{Hom}_{A^e}(P, A^e) \rightarrow \mathrm{Hom}_{A^e}(P, M)$$

is an isomorphism. It is standard, see e.g. [11, Proposition (8.3) (c)]. Then for any bounded chain complex C of finitely generated projective A -bimodules, ϕ_M induces in $\mathcal{D}^b(\mathrm{Vect}_{\mathbb{F}})$ an isomorphism

$$M \overset{L}{\otimes}_{A^e} \mathrm{RHom}_{A^e}(C, A^e) \cong \mathrm{RHom}_{A^e}(C, M). \tag{5.1}$$

Applying it to $C = K(A)$ and using (ii) in Definition 5.1, we get an isomorphism

$$\mathrm{RHom}_{A^e}(K(A), M) \cong M \overset{L}{\otimes}_{A^e} K(A)[-n]$$

in $\mathcal{D}^b(\mathrm{Vect}_{\mathbb{F}})$. Taking homology, we deduce that $\mathrm{HK}^p(A, M) \cong \mathrm{HK}_{n-p}(A, M)$ as vector spaces. \square

5.2. Koszul complex Calabi-Yau algebras versus Calabi-Yau algebras

Recall that if $\Delta \neq A_1$ and $\Delta \neq A_2$, then $A(\Delta)$ is 2-Kc-Calabi-Yau. But observe that if $A(\Delta)$ is moreover 2-Calabi-Yau, then $A(\Delta)$ is Koszul. We are led to the following conjecture.

CONJECTURE 5.6. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . If A is n -Calabi-Yau and n -Kc-Calabi-Yau, then A is Koszul. In other words, if A is not Koszul, the properties n -Calabi-Yau and n -Kc-Calabi-Yau are not simultaneously true.

PROPOSITION 5.7. Let $A = T_k(V)/(R)$ be a quadratic k -algebra over \mathcal{Q} . Conjecture 5.6 holds if $n \leq 3$.

Proof. Assume that A is n -Calabi-Yau and n -Kc-Calabi-Yau. We can assume that $n \geq 2$. Since $\mathrm{HK}^p(A, A^e) \cong \mathrm{HH}^p(A, A^e) = 0$ when $p = 0$ and $p = 1$, we have $H_n(K(A)) \cong$

$H_{n-1}(K(A)) = 0$, hence A is Koszul if $n = 2$. When $n = 3$, we also have $H_1(K(A)) = 0$ because the complex $K(A)$ is always exact in degree 1, therefore A is also Koszul in this case. \square

5.3. Strong Kc-Calabi-Yau algebras

DEFINITION 5.8. Let A be an n -Kc-Calabi-Yau algebra over \mathcal{Q} . The image $c \in \mathrm{HK}_n(A)$ of the unit 1 of the algebra A by the isomorphism $\mathrm{HK}^0(A) \cong \mathrm{HK}_n(A)$ in Theorem 5.5 is called the fundamental class of the n -Kc-Calabi-Yau algebra A .

We shall now define strong Kc-Calabi-Yau algebras. For this, we need the DG algebra $\tilde{A} = \mathrm{Hom}_{A^e}(K(A), A)$ and we work with DG \tilde{A} -bimodules in $A\text{-Bimod}$, as defined in Section 3 and recalled below. The point is that $K(A)$ is such a DG \tilde{A} -bimodule in $A\text{-Bimod}$ (Proposition 3.6). Denote by $\mathcal{C}(\tilde{A}, A\text{-Bimod})$ and $\mathcal{C}(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ the category of DG \tilde{A} -bimodules in $A\text{-Bimod}$ and $\mathrm{Vect}_{\mathbb{F}}$ respectively [45]. Remark that a DG \tilde{A} -bimodule in $\mathrm{Vect}_{\mathbb{F}}$ is just a DG \tilde{A} -bimodule.

DEFINITION 5.9. A DG \tilde{A} -bimodule C in $A\text{-Bimod}$ is a chain complex in $A\text{-Bimod}$ (as usual, C can be viewed as a cochain complex) endowed with a DG \tilde{A} -bimodule structure such that the bimodule actions of A and \tilde{A} are compatible.

For any A -bimodule M , $\mathrm{Hom}_{A^e}(C, M)$ is a (cochain) DG \tilde{A} -bimodule in $\mathrm{Vect}_{\mathbb{F}}$ (in $A\text{-Bimod}$ when $M = A^e$) for the following actions

$$(f.u)(x) = (-1)^p u(x.f), \quad (u.f)(x) = u(f.x)$$

where $f : A \otimes_k W_p \otimes_k A \rightarrow A$, $u : C_q \rightarrow M$ and $x \in C_{p+q}$. Note that $x.f$ and $f.x$ are in C_q by the graded actions of \tilde{A} on C . If $C = K(A)$, we recover the cup actions, that is, $f.u = f \underset{K}{\smile} u$ and $u.f = u \underset{K}{\smile} f$. In particular, $\mathrm{Hom}_{A^e}(K(A), A^e)$ is a DG \tilde{A} -bimodule in $A\text{-Bimod}$.

Similarly, for any cochain DG \tilde{A} -bimodule C' in $A\text{-Bimod}$, $M \otimes_{A^e} C'$ is a cochain DG \tilde{A} -bimodule in $\mathrm{Vect}_{\mathbb{F}}$ for the following actions

$$f.(m \otimes_{A^e} u) = m \otimes_{A^e} (f.u), \quad (m \otimes_{A^e} u).f = m \otimes_{A^e} (u.f)$$

where $f \in \tilde{A}$, $m \in M$ and $u \in C'$.

The bounded derived categories $\mathcal{D}^b(\tilde{A}, A\text{-Bimod})$ and $\mathcal{D}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ are defined in [45, Definition 7.2.7, Definition 7.3.3]. Unfortunately, it is not clear to us if the functors $\mathrm{Hom}_{A^e}(-, M) : \mathcal{C}^b(\tilde{A}, A\text{-Bimod}) \rightarrow \mathcal{C}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ and $M \otimes_{A^e} - : \mathcal{C}^b(\tilde{A}, A\text{-Bimod}) \rightarrow \mathcal{C}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ can be derived. Note that the first one takes values in $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ when $M = A^e$.

DEFINITION 5.10. Let A be a Kc-Calabi-Yau algebra of dimension n . Then A is said to be strong n -Kc-Calabi-Yau if the derived functor of the endofunctor $\mathrm{Hom}_{A^e}(-, A^e)$ of $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ exists and if $\mathrm{RHom}_{A^e}(K(A), A^e) \cong K(A)[-n]$ in $\mathcal{D}^b(\tilde{A}, A\text{-Bimod})$.

The preprojective algebras of connected graphs distinct from A_1 and A_2 are strong 2-Kc-Calabi-Yau algebras if they satisfy the first property in this definition. In fact, using Proposition 4.7, θ_{A^e} provides then an isomorphism $\mathrm{RHom}_{A^e}(K(A), A^e) \rightarrow K(A)[-2]$ in $\mathcal{D}^b(\tilde{A}, A\text{-Bimod})$.

THEOREM 5.11. Let A be a Kc-Calabi-Yau algebra of dimension n over \mathcal{Q} , with fundamental class c . We assume that A is strong Kc-Calabi-Yau and that the derived functors of

the functors $\mathrm{Hom}_{A^e}(-, A)$ and $A \otimes_{A^e} -$ from $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ to $\mathcal{C}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ exist. Then

$$c \underset{K}{\frown} - : \mathrm{HK}^{\bullet}(A) \rightarrow \mathrm{HK}_{n-\bullet}(A)$$

is an isomorphism of $\mathrm{HK}^{\bullet}(A)$ -bimodules, inducing an isomorphism of $\mathrm{HK}_{hi}^{\bullet}(A)$ -bimodules from $\mathrm{HK}_{hi}^{\bullet}(A)$ to $\mathrm{HK}_{n-\bullet}^{hi}(A)$. For all $\alpha \in \mathrm{HK}^p(A)$, we have $c \underset{K}{\frown} \alpha = (-1)^{np} \alpha \underset{K}{\frown} c$.

Proof. Following the proof of Theorem 5.5, we are interested in the morphism of cochain complexes

$$\phi_M : M \otimes_{A^e} \mathrm{Hom}_{A^e}(C, A^e) \rightarrow \mathrm{Hom}_{A^e}(C, M),$$

when the bounded chain complex C of A -bimodules is moreover a DG \tilde{A} -bimodule in $A\text{-Bimod}$. We prove now that ϕ_M is a morphism of DG \tilde{A} -bimodules in $\mathrm{Vect}_{\mathbb{F}}$, that is, a morphism in the category $\mathcal{C}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$ whose objects are viewed as cochain complexes. For this, we need only prove that ϕ_M is a morphism of \tilde{A} -bimodules. Let us prove that ϕ_M is left \tilde{A} -linear, the right linearity being similar. For $f : A \otimes_k W_p \otimes_k A \rightarrow A$, $u : C_q \rightarrow A^e$ and $x \in C_{p+q}$, we have

$$\phi_M(f.(m \otimes_{A^e} u))(x) = \phi_M(m \otimes_{A^e} (f.u))(x) = m.((f.u)(x)) = (-1)^p m.(u(x.f)),$$

while $f.(\phi_M(m \otimes_{A^e} u))(x) = (-1)^p \phi_M(m \otimes_{A^e} u)(x.f) = (-1)^p m.(u(x.f))$, which is what we want.

Continuing as in the proof of Theorem 5.5, the functors F , G and H induce functors on the complexes with enriched structures. Precisely, F and G are now functors from $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$ to $\mathcal{C}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$, and H is now an endofunctor of $\mathcal{C}^b(\tilde{A}, A\text{-Bimod})$. Under these notations, ϕ_M defines a natural transformation $\phi_M : H \circ G \Rightarrow F$.

We specialise to $M = A$. The assumptions in the theorem show that the derived functors of F , G and H exist, so that we can derive the natural transformation ϕ_A [45]. Then for any bounded chain complex DG \tilde{A} -bimodule C in $A\text{-Bimod}$ formed by finitely generated projective A -bimodules, we obtain an isomorphism

$$\phi_A : A \overset{L}{\otimes}_{A^e} \mathrm{RHom}_{A^e}(C, A^e) \cong \mathrm{RHom}_{A^e}(C, A) \quad (5.2)$$

in $\mathcal{D}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$. Applying this to $C = K(A)$ and using Definition 5.10, we get

$$\mathrm{RHom}_{A^e}(K(A), A) \cong A \overset{L}{\otimes}_{A^e} K(A)[-n]$$

in $\mathcal{D}^b(\tilde{A}, \mathrm{Vect}_{\mathbb{F}})$. Taking homology, we deduce an isomorphism $\mathrm{HK}^{\bullet}(A) \cong \mathrm{HK}_{n-\bullet}(A)$ of graded bimodules over the graded algebra $H(\tilde{A}) = \mathrm{HK}^{\bullet}(A)$. Denote this isomorphism by ψ .

The fact that ψ is a morphism of graded $\mathrm{HK}^{\bullet}(A)$ -bimodules translates as

$$\psi(\alpha \underset{K}{\smile} \beta) = \psi(\alpha) \underset{K}{\frown} \beta = (-1)^{np} \alpha \underset{K}{\frown} \psi(\beta) \quad (5.3)$$

for any $\alpha \in \mathrm{HK}^p(A)$ and $\beta \in \mathrm{HK}^{\bullet}(A)$. In accordance with Definition 5.8, define $c \in \mathrm{HK}_n(A)$ by $c = \psi(1)$ where $1 \in \mathrm{HK}^0(A)$ is the unit of A . Applying identities (5.3) to the trivial equalities $\alpha = 1 \underset{K}{\smile} \alpha = \alpha \underset{K}{\smile} 1$, we obtain

$$\psi(\alpha) = c \underset{K}{\frown} \alpha = (-1)^{np} \alpha \underset{K}{\frown} c. \quad (5.4)$$

Finally ψ is a morphism of complexes for higher (co)homologies since we have

$$\psi(\bar{\epsilon}_A \underset{K}{\smile} \alpha) = (-1)^n \bar{\epsilon}_A \underset{K}{\frown} \psi(\alpha).$$

Then $H(\psi) : \mathrm{HK}_{hi}^{\bullet}(A) \rightarrow \mathrm{HK}_{n-\bullet}^{hi}(A)$ is an isomorphism of $\mathrm{HK}_{hi}^{\bullet}(A)$ -bimodules. \square

Except in some particular cases, $\bar{\epsilon}_A \in \text{HK}^1(A)$ is non-zero, so that

$$\psi(\bar{\epsilon}_A) = (-1)^n \bar{\epsilon}_A \underset{K}{\frown} c = (-1)^n \partial_{\frown}(c)$$

is non-zero in $\text{HK}_{n-1}(A)$. Therefore $c \in \text{HK}_n(A)$ is not a cycle for higher Koszul homology and the isomorphism $H(\psi)$ cannot be naturally expressed as a cap action. As suggested by the preprojective algebras (Subsection 4.3), the class $\psi(\bar{\epsilon}_A)$ should be of interest for further investigations.

It is also interesting to remark that the identities (5.3) involving the isomorphism ψ imply that the graded algebra $\text{HK}^\bullet(A)$ is commutative if and only if the graded $\text{HK}^\bullet(A)$ -bimodule $\text{HK}_\bullet(A)$ is symmetric. As seen in Corollary 4.3, we have a stronger result for the preprojective algebras.

6. Koszul calculus of the preprojective algebras of Dynkin ADE type

We shall determine in this section the Koszul calculus and the higher Koszul calculus of any non-Koszul preprojective algebra A , that is, an algebra of type A, D or E with at least 3 vertices.

We first give some general facts and notation.

- (N1) We shall use the dimensions of the Hochschild cohomology and homology spaces of A which can be obtained in all characteristics as a consequence of the work of Etingof, Eu and Schedler in [24, Theorem 3.2.7] and [21]. In particular, by [24, Lemma 3.2.17] the centre of A is independent of the characteristic of \mathbb{F} . Bases of the Hochschild (co)homology spaces in characteristic zero induce free subsets of the Hochschild (co)homology spaces in positive characteristic, but there may be some extra basis elements in some cases.
- (N2) We know from Corollary 4.3 that the cup product on the Koszul cohomology of a preprojective algebra is graded commutative and that the cap product is graded symmetric. Moreover, it follows from Theorem 4.4 that the cap product can be obtained from the cup product. Indeed, if $f \in \text{HK}^p(A)$ and $x \in \text{HK}_q(A)$, we have

$$f \underset{K}{\frown} x = \theta_A(f \underset{K}{\smile} \theta_A^{-1}(x)) = \theta_A((-1)^{pq} \theta_A^{-1}(x) \underset{K}{\smile} f) = (-1)^{pq} x \underset{K}{\frown} f. \quad (6.1)$$

- (N3) Let X and Y be \mathbb{N} -graded spaces and let $f : X \rightarrow Y$ be a homogeneous map of degree 1. Let y_1, \dots, y_p be elements of pairwise different degrees. Then if $\sum_{i=1}^p y_i \in \text{Im } f$, at least one of the y_i is in $\text{Im } f$. We shall use this in the following context. The differentials b_K^1 and b_K^2 are homogeneous of weight 1. If we have a set of cocycles of pairwise different coefficient weights, that are not coboundaries, then they are linearly independent up to coboundaries, that is, they represent linearly independent cohomology classes. This also applies if some of the elements have the same weight but we already know that these elements are linearly independent up to coboundaries.
- (N4) We shall use the map $\kappa : A \rightarrow A$ constructed as follows. Let A be a preprojective algebra over a graph Δ ; let \bar{Q} be its quiver. Consider the map $\bar{Q}_1 \rightarrow \bar{Q}_1$ that sends a to a^* . It induces an anti-automorphism κ of A such that $\kappa(e_i) = e_i$ for all $i \in Q_0$ (since κ sends the relation $\sigma_i = \sum_{\substack{a \in \bar{Q}_1 \\ t(a)=i}} \varepsilon(a) a a^*$ to itself).
- (N5) We shall be using the Nakayama automorphism ν of A defined by Brenner, Butler and King for all preprojective algebras of Dynkin ADE type [10, Section 4]. In order to describe it, we need the Nakayama permutation $\bar{\nu}$ on the set of vertices of Δ . It is known that $\bar{\nu} = \text{id}$ if Δ is D_n with n even or if Δ is E_7 or E_8 , and that otherwise $\bar{\nu}$ is induced by the unique graph automorphism of order 2.

If α is an arrow in \overline{Q}_1 , let β be the unique arrow from $\overline{\nu}(\mathfrak{s}(\alpha))$ to $\overline{\nu}(\mathfrak{t}(\alpha))$. The Nakayama automorphism of [10] is described as follows:

$$\nu(\alpha) = \begin{cases} \beta & \text{if } \alpha \in Q_1^* \text{ or } \beta \in Q_1^* \\ -\beta & \text{if } \alpha \in Q_1 \text{ and } \beta \in Q_1. \end{cases}$$

Given a basis of a selfinjective quiver algebra $A = \mathbb{F}Q/I$ consisting of paths and containing a basis $\{\pi_i; i \in Q_0\}$ of the socle of A , an explicit construction of an associative non-degenerate bilinear form on A was given in [46, Proposition 3.15] (see also Subsection 6.6). We have chosen in each case such a basis so that ν is the Nakayama automorphism corresponding to this bilinear form (characterised on the arrows α in \overline{Q}_1 by $y\alpha = \pi_{\mathfrak{s}(\alpha)} \Leftrightarrow \nu(\alpha)y = \pi_{\mathfrak{t}(\alpha)}$ for all the basis elements y).

- (N6) When we define a cochain $f \in \text{Hom}_{k^e}(X, A)$ with $X \in \{k, V, A\}$, it will be implicit that if $f(x)$ is not defined for some $x \in X$ then $f(x) = 0$.
- (N7) For any cochain $f \in \text{Hom}_{k^e}(W_p, A)$, we shall set $\check{f} = \theta_A(f) \in A \otimes_{k^e} W_{2-p}$.
- (N8) Finally, given a Dynkin graph Δ and a ring L , we shall denote by Λ_L the preprojective algebra of Δ over L , so that $A = \Lambda_{\mathbb{F}}$.

6.1. Koszul calculus for preprojective algebras of type A

The preprojective algebra A of type A_n is defined by the quiver

$$\overline{Q} \quad 0 \begin{array}{c} \xrightarrow{a_0} \\ \xleftarrow{a_0^*} \end{array} 1 \begin{array}{c} \xrightarrow{a_1} \\ \xleftarrow{a_1^*} \end{array} 2 \begin{array}{c} \xrightarrow{a_2} \\ \xleftarrow{a_2^*} \end{array} \cdots \begin{array}{c} \xrightarrow{a_{n-3}} \\ \xleftarrow{a_{n-3}^*} \end{array} n-2 \begin{array}{c} \xrightarrow{a_{n-2}} \\ \xleftarrow{a_{n-2}^*} \end{array} n-1$$

subject to the relations

$$\begin{cases} \sigma_0 = -a_0^*a_0 \\ \sigma_i = a_{i-1}a_{i-1}^* - a_i^*a_i & 1 \leq i \leq n-2 \\ \sigma_{n-1} = a_{n-2}a_{n-2}^* \end{cases}$$

The Nakayama automorphism ν of A described in (N5) is given by $\nu(e_i) = e_{n-1-i}$, $\nu(a_i) = a_{n-2-i}^*$ and $\nu(a_i^*) = a_{n-2-i}$.

Erdmann and Snashall have given in [18] a basis B of A . We shall only need the sets e_iBe_i , e_iBe_{i+1} and $e_{i+1}Be_i$, which can be rewritten as follows: set $m_A = \lfloor \frac{n-1}{2} \rfloor$; then

$$\begin{aligned} e_iBe_i &= \{(a_i^*a_i)^\ell; 0 \leq \ell \leq \min(i, n-1-i)\} \text{ for } 0 \leq i \leq n-1, \text{ with } (a_i^*a_i)^0 = e_i \text{ for all } i, \\ e_{i+1}Be_i &= \{a_i(a_i^*a_i)^\ell; 0 \leq \ell \leq \min(i, n-2-i)\} \quad \text{and} \quad e_iBe_{i+1} = \kappa(e_{i+1}Be_i). \end{aligned}$$

For each i , Ae_i contains precisely one basis element of maximal length $n-1$, which is

$$\pi_i = \begin{cases} a_{n-2} \cdots a_i (a_i^*a_i)^i & \text{if } i < m_A \\ a_{m_A} (a_{m_A}^* a_{m_A})^{m_A} & \text{if } i = m_A \text{ and } n \text{ is even} \\ (a_{m_A}^* a_{m_A})^{m_A} & \text{if } i = m_A \text{ and } n \text{ is odd} \\ a_{n-1}^* \cdots a_{i-1}^* (a_{i-1} a_{i-1}^*)^{n-1-i} & \text{if } i > m_A. \end{cases}$$

They form a basis of the socle of A .

6.1.1. *The Koszul cohomology and homology spaces in type A* The spaces $\text{HK}^0(A) = \text{HH}^0(A) = Z(A)$ and $\text{HK}^1(A) = \text{HH}^1(A)$ are known from [18]. Therefore we only need to compute $\text{HK}^2(A)$. Recall our assumption that $n \geq 3$; then by Theorem 4.2 all the elements in $\text{Hom}_{k^e}(R, A)$ are cocycles. Moreover, using Theorem 4.4 and [24, Theorem 3.2.7], we have $\dim \text{HK}^2(A) = \dim \text{HK}_0(A) = \dim \text{HH}_0(A) = n$. Since every element in $\text{Im } b_K^2$ has coefficient

weight at least 1, the n cocycles h_i defined by $h_i(\sigma_j) = \delta_{ij}e_i$ for all j are linearly independent modulo $\text{Im } b_K^2$. It follows that they form a basis of $\text{HK}^2(A)$.

Combining with the results from [18], we have the following result.

PROPOSITION 6.1. *Let A be a preprojective algebra of type A_n .*

A basis of $\text{HK}^0(A)$ is given by the set $\{z_\ell; 0 \leq \ell \leq m_A\}$ with $z_0 = 1$ and $z_\ell = \sum_{i=1}^{n-2} (a_i^ a_i)^\ell = \sum_{i=\ell}^{n-1-\ell} (a_i^* a_i)^\ell = z_1^\ell$ for $1 \leq \ell \leq m_A$.*

A basis of $\text{HK}^1(A)$ is given by the set $\{\check{\zeta}_\ell; 0 \leq \ell \leq n - 2 - m_A\}$, where $\zeta_\ell \in \text{Hom}_{k^e}(V, A)$ is defined by $\zeta_\ell(a_i) = a_i(a_i^ a_i)^\ell$ for all i (or for $\ell \leq i \leq n - 2 - \ell$).*

A basis of $\text{HK}^2(A)$ is given by the set $\{\check{h}_i; 0 \leq i \leq n - 1\}$ where $h_i \in \text{Hom}_{k^e}(R, A)$ defined by $h_i(\sigma_j) = \delta_{ij}e_i$ for all j .

As a consequence of Theorem 4.4, we obtain bases of the Koszul homology spaces.

COROLLARY 6.2. *A basis of $\text{HK}_0(A)$ is given by the set $\{\check{h}_i; 0 \leq i \leq n - 1\}$ where $\check{h}_i = e_i \otimes e_i$.*

A basis of $\text{HK}_1(A)$ is given by the set $\{\check{\zeta}_\ell; 0 \leq \ell \leq n - m_A - 2\}$ where $\check{\zeta}_\ell = \sum_{i=0}^{n-2} a_i(a_i^ a_i)^\ell \otimes a_i^* = \sum_{i=\ell}^{n-2-\ell} a_i(a_i^* a_i)^\ell \otimes a_i^*$. A basis of $\text{HK}_2(A)$ is given by the set $\{\check{z}_\ell; 0 \leq \ell \leq m_A\}$ where $\check{z}_\ell = \sum_{i=0}^{n-1} (a_i^* a_i)^\ell \otimes \sigma_i$.*

Note that $\check{z}_0 = \omega_0$ is the fundamental class.

6.1.2. *Cup and cap products* We know from Corollary 4.3 that the cup product on $\text{HK}^\bullet(A)$ is graded-commutative. The following result gives all the non zero cup products of elements in $\text{HK}^\bullet(A)$.

PROPOSITION 6.3. *Let A be a preprojective algebra of type A_n . Up to graded commutativity, the non zero cup products in $\text{HK}^\bullet(A)$ are given by*

$$\begin{aligned} z_0 \smile_K \bar{f} &= \bar{f} \quad \text{for all } \bar{f} \in \text{HK}^\bullet(A) \\ z_{\ell_1} \smile_K z_{\ell_2} &= z_{\ell_1 + \ell_2} \quad \text{if } \ell_1 + \ell_2 \leq m_A \\ z_{\ell_1} \smile_K \check{\zeta}_{\ell_2} &= \check{\zeta}_{\ell_1 + \ell_2} \quad \text{if } \ell_1 + \ell_2 \leq n - m_A - 2 \end{aligned}$$

Proof. The first cup product is clear and the other cup products in the statement only involve in $\text{HH}^0(A)$ and $\text{HH}^1(A)$, therefore they are known from [18].

The basis elements of $\text{HK}^2(A)$ have coefficient weight 0, and b_K^2 is homogeneous of weight 1, therefore any element that has positive coefficient weight must be a coboundary. The other cup products (that all vanish) follow from this. \square

We now deduce the cap products from (6.1).

COROLLARY 6.4. *Up to graded symmetry, the non zero cap products are the following.*

$$\begin{aligned} z_0 \frown_K x &= x \quad \text{for all } x \in \text{HK}_\bullet(A) \\ z_{\ell_1} \frown_K \check{z}_{\ell_2} &= \check{z}_{\ell_1 + \ell_2} \quad \text{if } \ell_1 + \ell_2 \leq m_A \end{aligned}$$

$$z_{\ell_1} \underset{K}{\frown} \bar{\zeta}_{\ell_2} = \bar{\zeta}_{\ell_1 + \ell_2} \quad \text{if } \ell_1 + \ell_2 \leq n - m_A - 2$$

6.1.3. *Higher Koszul cohomology and homology* We start with a lemma giving the cohomology class of the fundamental 1-cocycle.

LEMMA 6.5. *The cohomology class of ϵ_A is equal to the cohomology class of $2\zeta_0$.*

Proof. Let $\zeta_0^* \in \text{Hom}_{k^e}(V, A)$ be the cocycle defined by $\zeta_0^*(a_i^*) = a_i^*$ for all $i \in Q_0$. Since $\epsilon_A = \zeta_0 + \zeta_0^*$, we must prove that $\zeta_0^* - \zeta_0$ is a coboundary.

Consider $v = \sum_{i=0}^{n-2} \sum_{j=i}^{n-2} e_i \in \bigoplus_{i \in Q_0} e_i A e_i \cong \text{Hom}_{k^e}(k, A)$. Then $b_K^1(v) = \zeta_0^* - \zeta_0$, as required. \square

As a consequence, the complex defining the higher Koszul cohomology is

$$0 \rightarrow \text{HK}^0(A) \xrightarrow{\partial^1} \text{HK}^1(A) \xrightarrow{\partial^2} \text{HK}^2(A) \rightarrow 0 \cdots$$

with $\partial^1(z_\ell) = 2\bar{\zeta}_\ell$ for $0 \leq \ell \leq n - m_A - 2$ and $\partial^2 = 0$. We then have the following higher Koszul cohomology.

PROPOSITION 6.6. *Let A be a preprojective algebra of type A_n . If $\text{char}(\mathbb{F}) = 2$, then $\text{HK}_{hi}^\bullet(A) = \text{HK}^\bullet(A)$.*

If $\text{char}(\mathbb{F}) \neq 2$ and n is even, then

$$\begin{aligned} \text{HK}_{hi}^2(A) &= \text{HK}^2(A) \\ \text{HK}_{hi}^p(A) &= 0 \text{ if } p \neq 2. \end{aligned}$$

Finally, if $\text{char}(\mathbb{F}) \neq 2$ and n is odd, then

$$\begin{aligned} \text{HK}_{hi}^0(A) &= \text{HK}^0(A)_{2m_A} \text{ has dimension 1 and is spanned by } z_{m_A} \\ \text{HK}_{hi}^2(A) &= \text{HK}^2(A) \\ \text{HK}_{hi}^p(A) &= 0 \text{ if } p \neq 0 \text{ and } p \neq 2. \end{aligned}$$

Higher Koszul homology can then be deduced using duality (Theorem 4.4).

COROLLARY 6.7. *If $\text{char}(\mathbb{F}) = 2$, then $\text{HK}_\bullet^{hi}(A) = \text{HK}_\bullet(A)$.*

If $\text{char}(\mathbb{F}) \neq 2$ and n is even, then


$$\begin{aligned} \text{HK}_0^{hi}(A) &= \text{HK}_0(A) \\ \text{HK}_p^{hi}(A) &= 0 \text{ if } p \neq 0. \end{aligned}$$

Finally, if $\text{char}(\mathbb{F}) \neq 2$ and n is odd, then

$$\begin{aligned} \text{HK}_0^{hi}(A) &= \text{HK}_0(A) \\ \text{HK}_2^{hi}(A) &= \text{HK}_2(A)_{2m_A} \text{ has dimension 1 and is spanned by } \check{z}_{m_A} \\ \text{HK}_p^{hi}(A) &= 0 \text{ if } p \neq 0 \text{ and } p \neq 2. \end{aligned}$$

6.2. Koszul calculus for preprojective algebras of type D

The preprojective algebra A of type D_n is defined by the quiver



subject to the relations

$$\begin{aligned} \sigma_0 &= -a_0^*a_0 & \sigma_i &= a_{i-1}a_{i-1}^* - a_i^*a_i & 3 \leq i \leq n-2 \\ \sigma_1 &= -a_1^*a_1 & \sigma_{n-1} &= a_{n-2}a_{n-2}^* \\ \sigma_2 &= a_0a_0^* + a_1a_1^* - a_2^*a_2 \end{aligned}$$

The Nakayama automorphism ν of A described in (N5) is given by $\nu(e_i) = e_i$, $\nu(a_i) = -a_i^*$ and $\nu(a_i^*) = a_i$ if n is even or if n is odd and $i \geq 2$, and it exchanges e_0 and e_1 , a_0 and $-a_1$, and a_0^* and a_1^* if n is odd.

Eu has given in [22] a basis B of A . Set $m_D = \lfloor \frac{n-2}{2} \rfloor$ and $u = n - m_D - 2$. We shall only need bases of the $e_j A e_i$ when i and j are equal or adjacent vertices, which can be rewritten as follows:

$$\begin{aligned} e_0 B e_0 &= \{(a_0^* a_1 a_1^* a_0)^\ell; 0 \leq \ell \leq m_D\} \\ e_1 B e_1 &= \{(a_1^* a_0 a_0^* a_1)^\ell; 0 \leq \ell \leq m_D\} \\ e_i B e_i &= \{(a_i^* a_i)^\ell; 0 \leq \ell \leq n - i - 1\} \\ &\quad \cup \{(a_i^* a_i)^\ell a_{i-1} \cdots a_2 a_1 a_1^* a_2^* \cdots a_{i-1}^*; 0 \leq \ell \leq n - i - 1\} \quad \text{if } i \geq 2 \\ e_2 B e_i &= \{(a_2^* a_2)^\ell a_i; 0 \leq \ell \leq n - 3\} \quad \text{for } i \in \{0, 1\} \\ e_{i+1} B e_i &= \{(a_{i+1}^* a_{i+1})^\ell a_i; 0 \leq \ell \leq n - i - 2\} \\ &\quad \cup \{(a_{i+1}^* a_{i+1})^\ell a_i \cdots a_2 a_1 a_1^* a_2^* \cdots a_{i-1}^*; 0 \leq \ell \leq n - i - 2\} \quad \text{if } i \geq 2 \\ e_i B e_{i+1} &= \kappa(e_{i+1} B e_i), \quad e_i B e_2 = \kappa(e_2 B e_0) \quad \text{for } i \in \{0, 1\}. \end{aligned}$$

For each i , $A e_i$ contains precisely one basis element of maximal length $2(n-2)$, which is

$$\begin{aligned} \pi_0 &= -a_0^* a_2^* a_3^* \cdots a_{n-2}^* a_{n-2} \cdots a_3 a_2 a_0 = \begin{cases} -(a_0^* a_1 a_1^* a_0)^{m_D} & \text{if } n \text{ is even} \\ -a_1^* a_0 (a_0^* a_1 a_1^* a_0)^{m_D} & \text{if } n \text{ is odd} \end{cases} \\ \pi_1 &= a_1^* a_2^* a_3^* \cdots a_{n-2}^* a_{n-2} \cdots a_3 a_2 a_1 = \begin{cases} (a_1^* a_0 a_0^* a_1)^{m_D} & \text{if } n \text{ is even} \\ a_0^* a_1 (a_1^* a_0 a_0^* a_1)^{m_D} & \text{if } n \text{ is odd} \end{cases} \\ \pi_i &= (a_i^* a_i)^{n-i-1} a_{i-1} \cdots a_2 a_1 a_1^* a_2^* \cdots a_{i-1}^* \quad \text{if } 2 \leq i \leq n-2 \\ \pi_{n-1} &= a_{n-2} a_{n-3} \cdots a_1 a_1^* a_2^* \cdots a_{n-2}^*. \end{aligned}$$

They form a basis of the socle of A .

6.2.1. *The Koszul cohomology and homology spaces in type D* The centre of A does not depend on the characteristic of \mathbb{F} by fact (N1) and was computed in [22] in characteristic 0. A basis of $\text{HH}^1(\Lambda_{\mathbb{C}})$ was determined in [22]. Moreover, $\dim \text{HK}^1(A) = \dim \text{HH}^1(A)$, which is equal to $\dim \text{HH}^1(\Lambda_{\mathbb{C}}) = n$ if $\text{char}(\mathbb{F}) \neq 2$ and to $\dim \text{HH}^1(\Lambda_{\mathbb{C}}) + m$ if $\text{char}(\mathbb{F}) = 2$ by [24].

It also follows from Theorem 4.4 and [24] that $\dim \text{HK}^2(A) = \dim \text{HK}_0(A) = \dim \text{HH}_0(A)$, which is equal to $n - m_D - 2$ if $\text{char}(\mathbb{F}) \neq 2$ and to $n - 2$ if $\text{char}(\mathbb{F}) = 2$.

In order to give bases of the $\text{HK}^p(A)$ for $p = 0, 1, 2$, we define the following cochains:

- the elements $z_0 = 1$ and $z_\ell = (a_0^* a_1 a_1^* a_0)^{\ell} + (a_1^* a_0 a_0^* a_1)^{\ell} + \sum_{i=2}^{n-2} (a_i^* a_i)^{2\ell} = z_1^{\ell}$ for $\ell > 0$ in A . Note that if n is even, then $z_1^u = z_1^{m_D} = -\pi_0 + \pi_1$, but if n is odd then $z_1^u = 0$;
- the elements $\zeta_\ell \in \text{Hom}_{k^e}(V, A)$ with $0 \leq \ell \leq u - 1$ defined by $\zeta_\ell(a_i) = a_i z_\ell$ for all i ;
- the elements $\rho_\ell \in \text{Hom}_{k^e}(V, A)$ with $0 \leq \ell \leq m_D - 1$ where $\rho_\ell(a_i) = (a_2^* a_2)^{2\ell+1} a_i$ for $i = 0, 1$ and $\rho_\ell(a_i^*) = a_i^* (a_2^* a_2)^{2\ell+1}$ for $i = 0, 1$;
- the elements $h_j \in \text{Hom}_{k^e}(R, A)$ for $0 \leq j \leq n - 1$, where $h_j(\sigma_i) = \delta_{ij} e_j$;
- the elements $\gamma_\ell \in \text{Hom}_{k^e}(R, A)$ for $1 \leq \ell \leq m_D$ where $\gamma_\ell(\sigma_0) = (a_0^* a_1 a_1^* a_0)^{\ell}$.

PROPOSITION 6.8. *Let A be a preprojective algebra of type D_n .*

- (i) *The elements in $\{\pi_i; 0 \leq i \leq n - 1 \text{ and } \nu(e_i) = e_i\} \cup \{z_\ell; 0 \leq \ell \leq u - 1\}$ form a basis of $\text{HK}^0(A)$.*
- (ii) *If $\text{char}(\mathbb{F}) \neq 2$, the $\bar{\zeta}_\ell$, for $0 \leq \ell \leq u - 1$, form a basis of $\text{HK}^1(A)$.
If $\text{char}(\mathbb{F}) = 2$, the $\bar{\zeta}_\ell$ for $0 \leq \ell \leq u - 1$ and the $\bar{\rho}_\ell$ for $0 \leq \ell \leq m_D - 1$ form a basis of $\text{HK}^1(A)$.*
- (iii) *If $\text{char}(\mathbb{F}) \neq 2$, the \bar{h}_j for $0 \leq j \leq n - 1$ form a basis of $\text{HK}^2(A)$.
If $\text{char}(\mathbb{F}) = 2$, the \bar{h}_j for $0 \leq j \leq n - 1$ and the $\bar{\gamma}_\ell$ for $1 \leq \ell \leq m_D$ form a basis of $\text{HK}^2(A)$.*

Proof. The results for $\text{HK}^0(A)$ and, when $\text{char}(\mathbb{F}) \neq 2$, for $\text{HK}^1(A)$ follow from the comments before the proposition.

Assume that $\text{char}(\mathbb{F}) = 2$. In order to prove the result for $\text{HK}^1(A)$, we must prove that the elements we have considered in $\text{Hom}_{k^e}(V, A)$ are cocycles that are linearly independent modulo coboundaries. It is in fact enough to prove that the ρ_ℓ are cocycles that are not coboundaries by fact (N3).

First note that, at the level of cochains, $\rho_\ell = \rho_0 \underset{K}{\smile} z_\ell$. Therefore, to prove that ρ_ℓ is a cocycle, it is enough to prove that ρ_0 is a cocycle, and this is easy to check.

Since $\rho_\ell \underset{K}{\smile} z_{m_D-1-\ell} = \rho_{m_D-1}$, in order to prove that ρ_ℓ is not a coboundary for all ℓ , it is enough to prove that ρ_{m_D-1} is not a coboundary. The map ρ_{m_D-1} has coefficient weight $4m_D - 1$. If ρ_{m_D-1} is a coboundary, then it is the image of a morphism in $\text{Hom}_{k^e}(k, A) \cong \bigoplus_{i \in Q_0} e_i A e_i$ whose coefficients are linear combinations of cycles in A of weight $4m_D - 2$, which are known. It is then straightforward to show that the image of any such morphism under b_K^1 is not equal to ρ_{m_D-1} .

For (iii), we first observe that every cochain in $\text{Hom}_{k^e}(R, A)$ is a cocycle. Moreover, the h_j are n cocycles that are clearly linearly independent modulo coboundaries (all coboundaries have coefficient weight at least equal to 1). Therefore if $\text{char}(\mathbb{F}) \neq 2$, the result follows.

If $\text{char}(\mathbb{F}) = 2$, it is enough to prove that the γ_ℓ are not coboundaries by fact (N3). At the level of cochains, $\gamma_\ell \underset{K}{\smile} z_{m_D-\ell} = \gamma_{m_D}$ for $1 \leq \ell \leq m_D$, therefore it is enough to prove that γ_{m_D} is not a coboundary. The map γ_{m_D} has coefficient weight $4m_D$, therefore if γ_{m_D} is a coboundary, then it is the image of a morphism in $\text{Hom}_{k^e}(V, A)$ whose coefficients are linear combinations of elements of weight $4m_D - 2$ in A between two adjacent vertices in Q_0 , which are known. Here again, checking that the image of any such morphism under b_K^2 is not equal to γ_{m_D} is straightforward. \square

Koszul homology follows using duality (Theorem 4.4), as in Corollary 6.2.

6.2.2. *Cup and cap products* We now determine the cup products of the elements in the bases of the Koszul cohomology spaces given above.

LEMMA 6.9. *For $1 \leq i \leq n - 2$, consider the cochains u_i, v_i and w_i in $\text{Hom}_{k^e}(R, A)$ defined by $u_i(\sigma_j) = \delta_{ij} e_i z_1$, $v_i(\sigma_j) = \delta_{ij} \pi_i$ and $w_i(\sigma_j) = \delta_{ij} a_i^* a_i z_1$ for all $j \in Q_0$.*

If $\text{char}(\mathbb{F}) \neq 2$, the u_i , v_i and w_i are all coboundaries.

If $\text{char}(\mathbb{F}) = 2$, then the u_i for $i \geq 2$ are coboundaries, and $\bar{u}_0 = \bar{u}_1 = \bar{\gamma}_1$. Moreover, if n is odd, all the v_i are coboundaries and if n is even, then $\bar{v}_i = \bar{\gamma}_{m_D}$ for all i . Finally, the w_i are all coboundaries.

Proof. Every element in $\text{Hom}_{k^e}(R, A)$ is a cocycle. Moreover, the differential b_K^2 is homogeneous of degree 1 with respect to the coefficient weight, and the coefficient weight of all the basis elements in $\text{HK}^2(A)$ is a multiple of 4, and is 0 if $\text{char}(\mathbb{F}) \neq 2$.

It follows that if $\text{char}(\mathbb{F}) \neq 2$, all the u_i , v_i and w_i must be coboundaries, and if $\text{char}(\mathbb{F}) = 2$, the w_i are coboundaries and so are the v_i if n is odd.

Assume that $\text{char}(\mathbb{F}) = 2$. We must now study the u_i , as well as the v_i when n is even.

Note that $u_{n-2} = 0$ and $u_0 = \gamma_1$. For $0 \leq i \leq n-3$, define $p_i \in \text{Hom}_{k^e}(V, A)$ by $p_i(a_0) = a_1 a_1^* a_0$, $p_i(a_1) = a_0 a_0^* a_1$ and $p_i(a_i) = a_i a_i^* a_i$ if $i \geq 2$. Then, for $2 \leq i \leq n-3$, we have $u_i = b_K^2 \left(\sum_{j=i}^{n-3} p_j \right)$. Moreover, $b_K^2 \left(\sum_{j=0}^{n-3} p_j \right) = u_0 + u_1$. It follows that the cohomology classes of u_0 and u_1 are both equal to that of γ_1 .

We now turn to the v_i . Note that since n is even and $\text{char}(\mathbb{F}) = 2$, the map v_0 is the map γ_{m_D} , which is not a coboundary.

Define $q_i \in \text{Hom}_{k^e}(V, A)$ by $q_i(a_i) = (a_{i+1}^* a_{i+1})^{n-i-2} a_i \cdots a_2 a_1 a_1^* a_2^* \cdots a_{i-1}^*$ for $1 \leq i \leq n-2$ and $q_0(a_0) = (a_2^* a_2)^{n-3}$. Then, for $2 \leq i \leq n-2$, we have $v_i - v_1 = b_K^2 \left(\sum_{j=1}^{i-1} q_j \right)$, and $v_1 - v_0 = b_K^2(q_0 - q_1)$. Therefore $\bar{v}_i = \bar{v}_0 = \bar{\gamma}_{m_D}$ for all i . \square

We now give all the non zero cup products.

PROPOSITION 6.10. *Let A be a preprojective algebra of type D_n . Up to graded commutativity, the non zero cup products of elements in $\text{HK}^\bullet(A)$ are:*

$$\begin{aligned} z_0 \underset{K}{\smile} \bar{f} &= \bar{f} \text{ for all } \bar{f} \in \text{HK}^\bullet(A); & z_{\ell_1} \underset{K}{\smile} z_{\ell_2} &= \begin{cases} z_{\ell_1+\ell_2} & \text{if } \ell_1 + \ell_2 \leq u - 1; \\ -\pi_0 + \pi_1 & \text{if } n \text{ is even and } \ell_1 + \ell_2 = m_D; \end{cases} \\ z_{\ell_1} \underset{K}{\smile} \bar{\zeta}_{\ell_2} &= \bar{\zeta}_{\ell_1+\ell_2} \text{ if } \ell_1 + \ell_2 \leq u - 1; \\ z_{\ell_1} \underset{K}{\smile} \bar{\rho}_{\ell_2} &= \bar{\rho}_{\ell_1+\ell_2} \text{ if } \ell_1 + \ell_2 \leq u - 1; & z_\ell \underset{K}{\smile} \bar{h}_i &= \begin{cases} \bar{\gamma}_\ell & \text{if } \ell \geq 1, \text{ char}(\mathbb{F}) = 2 \text{ and } i \in \{0, 1\} \\ h_i & \text{if } \ell = 0; \end{cases} \\ z_{\ell_1} \underset{K}{\smile} \bar{\gamma}_{\ell_2} &= \bar{\gamma}_{\ell_1+\ell_2} \text{ if } \ell_1 + \ell_2 \leq m_D; & \pi_i \underset{K}{\smile} \bar{h}_j &= \gamma_{m_D} \text{ if } i = j, n \text{ is even and } \text{char}(\mathbb{F}) = 2. \end{aligned}$$

Proof. We use the notation in Lemma 6.9.

For $\ell \geq 1$, we have $z_\ell \underset{K}{\smile} h_i = z_{\ell-1} \underset{K}{\smile} z_1 \underset{K}{\smile} h_i = z_{\ell-1} \underset{K}{\smile} u_i$ and the result follows from Lemma 6.9.

Next, $\pi_i \underset{K}{\smile} h_j = \delta_{ij} v_j$ and again the result is a consequence of Lemma 6.9.

Now assume that $\text{char}(\mathbb{F}) = 2$, so that the $\bar{\rho}_\ell$ occur in the basis of $\text{HK}^1(A)$. At the level of cochains, we have $\rho_{\ell_1} \underset{K}{\smile} \rho_{\ell_2} = w_2 \underset{K}{\smile} z_{\ell_1+\ell_2}$, which is a coboundary.

The map $\rho_{\ell_1} \underset{K}{\smile} \zeta_{\ell_2} = u_2 \underset{K}{\smile} z_{\ell_1+\ell_2+1}$ is also a coboundary, as required.

The remaining cup products are easy to compute. Note that the cup product in $\text{HK}_0(A) \cong Z(A)$ is the ordinary product, and that the elements π_i are in the socle of A , hence are annihilated by the radical of A . \square

The cap products follow using duality, as in Corollary 6.4.

6.2.3. *Higher Koszul (co)homology* As in the case of a preprojective algebra of type A , the cohomology class of the fundamental 1-cocycle is equal to $2\zeta_0$ so that $\partial^1_\zeta(z_\ell) = 2\zeta_\ell$ for $0 \leq \ell \leq u - 1$, $\partial^1_\zeta(\pi_i) = 0$ and $\partial^2_\zeta = 0$. We then have the following higher Koszul cohomology.

PROPOSITION 6.11. *Let A be a preprojective algebra of type D_n .*


- (i) *If $\text{char}(\mathbb{F}) = 2$, then $\text{HK}_{hi}^\bullet(A) = \text{HK}^\bullet(A)$.*
- (ii) *If $\text{char}(\mathbb{F}) \neq 2$, then*

$$\begin{aligned} \text{HK}_{hi}^0(A) &= \text{HK}^0(A)_{>0} \text{ has basis the } \pi_i \text{ that are in } Z(A) \\ \text{HK}_{hi}^2(A) &= \text{HK}^2(A) \\ \text{HK}_{hi}^p(A) &= 0 \text{ if } p \neq 0 \text{ and } p \neq 2. \end{aligned}$$

Higher Koszul homology follows from Theorem 4.4 as in Corollary 6.7.

6.3. Koszul calculus for preprojective algebras of type E_6

The preprojective algebra A of type E_6 is defined by the quiver



subject to the relations

$$\begin{aligned} \sigma_0 &= -a_0^*a_0 & \sigma_3 &= a_0a_0^* + a_2a_2^* - a_3^*a_3 \\ \sigma_1 &= -a_1^*a_1 & \sigma_4 &= a_3a_3^* - a_4^*a_4 \\ \sigma_2 &= a_1a_1^* - a_2^*a_2 & \sigma_5 &= a_4a_4^* \end{aligned}$$

The Nakayama automorphism of (N5) is given by

$$\begin{aligned} \nu(e_i) &= e_i \text{ if } i \in \{0, 3\} & \nu(a_0) &= -a_0 & \nu(a_i) &= a_{5-i}^* \text{ if } i \geq 1 \\ \nu(e_i) &= e_{6-i} \text{ if } i \in \{1, 2, 4, 5\} & \nu(a_0^*) &= a_0^* & \nu(a_i^*) &= a_{5-i} \text{ if } i \geq 1. \end{aligned}$$

To simplify notation, we shall denote by $c_0 = a_0a_0^*$, $c_2 = a_2a_2^*$ and $c_3 = a_3^*a_3$ the three 2-cycles at the vertex 3.

The socle is the part of weight 10 of A , and the set $\{\pi_i; i \in Q_0\}$ where $\pi_0 = a_0^*c_3^2c_0c_3a_0$, $\pi_1 = a_4a_3c_0c_3c_0a_2a_1$, $\pi_2 = a_2^*(c_3c_0)^2a_3^*$, $\pi_3 = c_3(c_0c_3)^2$, $\pi_4 = \kappa(\pi_2)$ and $\pi_5 = \kappa(\pi_1)$.

6.3.1. *The Koszul cohomology and homology spaces in type E_6* We shall follow the same method as in types A and D, using the results from [24] and Theorem 4.4 to determine the dimensions of the spaces, and using results from [22] for the parts that are the same as in characteristic 0.

We define the following elements

- in A : $z_0 = 1$, $z_6 = a_1^*a_2^*a_3^*a_3a_2a_1 + a_2^*c_3^2a_2 - c_0c_3c_0 + a_3c_2^2a_3^* + a_4a_3a_2a_2^*a_3^*a_4^*$ and $z_8 = a_2^*c_0c_3c_0a_2 + c_0c_3^2c_0 + a_3c_0c_3c_0a_3^*$;
- in $\text{Hom}_{k^e}(V, A)$: the maps ζ_ℓ defined by $\zeta_\ell(a_i) = a_iz_\ell$ for $\ell \in \{0, 6, 8\}$, the map ρ_3 defined by $\rho_3(a_2) = c_0a_2$, $\rho_3(a_3) = a_3c_3$ and $\rho_3(a_2^*) = a_2^*c_3$, and the map ρ_5 defined by $\rho_5(a_0) = c_2c_3a_0$, $\rho_5(a_1) = a_2^*c_0a_2a_1$, $\rho_5(a_2) = c_2^2a_2$, $\rho_5(a_0^*) = -a_0^*c_3^2$, $\rho_5(a_1^*) = -a_1^*a_2^*c_0a_2$ and $\rho_5(a_2^*) = a_2^*c_3^2$;
- in $\text{Hom}_{k^e}(R, A)$: the maps h_j defined for $0 \leq j \leq 5$ by $h_j(\sigma_i) = \delta_{ij}e_j$ for all i , the map γ_4 defined by $\gamma_4(\sigma_0) = a_0^*c_3a_0$ and the map γ_6 defined by $\gamma_6(\sigma_0) = a_0^*c_3^2a_0$.

We shall use the following lemma.

LEMMA 6.12. *Assume that $\text{char}(\mathbb{F}) = 3$. Let $\gamma \in \text{Hom}_{k^e}(R, A)$ be an element of coefficient weight 6, so that*

$$\begin{aligned} \gamma(\sigma_0) &= \lambda_0 a_0^* c_3^2 a_0 & \gamma(\sigma_3) &= \lambda_3 c_3^2 c_0 + \lambda'_3 c_3 c_0 c_3 + \lambda''_3 c_0 c_3^2 \\ \gamma(\sigma_1) &= \lambda_1 a_1^* a_2^* c_0 a_2 a_1 & \gamma(\sigma_4) &= \lambda_4 a_3 c_0 c_3 a_3^* + \lambda'_4 a_3 c_3 c_0 a_3^* \\ \gamma(\sigma_2) &= \lambda_2 a_2^* c_0 c_3 a_2 + \lambda'_2 a_2^* c_3 c_0 a_2 & \gamma(\sigma_5) &= \lambda_5 a_4 a_3 c_0 a_3^* a_4^*. \end{aligned}$$

Then γ is a coboundary if, and only if, $\sum_{i=0}^5 \lambda_i + \sum_{i=2}^4 \lambda'_i + \lambda''_3 = 0$.

Proof. The proof is straightforward, once we know that a cochain of weight 5 takes its values in $A_5 = E \oplus \kappa(E)$, where E is the space spanned by $c_3^2 a_0, c_0 c_3 a_0, a_2^* c_0 a_2 a_1, c_3^2 a_2, c_0 c_3 a_2, a_3 c_0 c_3, a_3 c_3 c_0, a_4 a_3 c_0 a_3^*$. \square

PROPOSITION 6.13. *Let A be a preprojective algebra of type E_6 .*

- (i) *The elements in $\{z_0, z_6, z_8, \pi_0, \pi_3\}$ form a basis of $\text{HK}^0(A)$.*
- (ii) *If $\text{char}(\mathbb{F}) \notin \{2, 3\}$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 6, 8\}$ form a basis of $\text{HK}^1(A)$.
If $\text{char}(\mathbb{F}) = 2$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 6, 8\} \cup \{\bar{\rho}_3\}$ form a basis of $\text{HK}^1(A)$.
If $\text{char}(\mathbb{F}) = 3$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 6, 8\} \cup \{\bar{\rho}_5\}$ form a basis of $\text{HK}^1(A)$.*
- (iii) *If $\text{char}(\mathbb{F}) \notin \{2, 3\}$, the elements in $\{\bar{h}_j; j \in Q_0\}$ form a basis of $\text{HK}^2(A)$.
If $\text{char}(\mathbb{F}) = 2$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_4\}$ form a basis of $\text{HK}^2(A)$.
If $\text{char}(\mathbb{F}) = 3$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_6\}$ form a basis of $\text{HK}^2(A)$.*

Proof. The centre was given in [22], so we have (i).

For $\text{HK}^1(A)$ and $\text{HK}^2(A)$, the number of elements in the statement is equal to the dimension of the corresponding cohomology space. Moreover, all the elements in the statement are indeed cocycles.

If $\text{char}(\mathbb{F})$ is not 2 or 3, a basis of $\text{HK}^1(A) = \text{HH}^1(A)$ was given in [22]. It consists of the classes of the ζ'_ℓ with $\ell \in \{0, 6, 8\}$ where $\zeta'_\ell(a_i) = a_i z_\ell$ for $0 \leq i \leq 2$ and $\zeta'_\ell(a_i^*) = a_i^* z_\ell$ for $3 \leq i \leq 4$. Since $\zeta_0 - \zeta'_0$ is equal to $b_K^1(e_3 + 2e_5)$, and $\zeta_\ell - \zeta'_\ell = (\zeta_0 - \zeta'_0) \smile_K z_\ell$ is also a coboundary, ζ_ℓ and ζ'_ℓ represent the same cohomology class for $\ell \in \{0, 6, 8\}$. Moreover, as in types A and D, the elements \bar{h}_j form a basis of $\text{HK}^2(A)$.

If $\text{char}(\mathbb{F}) \in \{2, 3\}$, we need only prove that the extra elements are not coboundaries by fact (N3).

If $\text{char}(\mathbb{F}) = 2$, we have $z_6 \smile_K \rho_3 - \zeta_8 = b_K^1(g)$ where g is defined by $g(e_2) = a_2^* c_3 c_0 c_3 a_2$, and ζ_8 is not a coboundary, therefore ρ_3 cannot be a coboundary. Moreover, assume that $\gamma_4 = b_K^1(g')$ is a coboundary. Then g' would be of coefficient weight 3, and we would necessarily take values in $A_3 = E \oplus \kappa(E)$ where E is spanned by $c_3 a_0, c_0 a_2, c_3 a_2, a_3 c_0, a_3 c_3$. This leads to a contradiction.

If $\text{char}(\mathbb{F}) = 3$, assume that ρ_5 is a coboundary $b_K^1(h)$, then h is of weight 4, and necessarily $h(e_0) = \lambda_0 a_0^* c_3 a_0$, $h(e_2) = \lambda_2 a_2^* c_3 a_2$ and $h(e_3) = \lambda_3 c_3 c_0 + \lambda'_3 c_3^2 + \lambda''_3 c_0 c_3^2$, and by considering $b_K^1(h)(a_0), b_K^1(h)(a_0^*), b_K^1(h)(a_2)$ and $b_K^1(h)(a_2^*)$ we get a contradiction. Finally, the fact that γ_6 is not a coboundary follows from Lemma 6.12. \square

Koszul homology follows using duality (Theorem 4.4), as in Corollary 6.2.

6.3.2. *Cup and cap products* We now determine the cup products of the elements in the bases of the Koszul cohomology spaces given above.

PROPOSITION 6.14. *Let A be a preprojective algebra of type E_6 . Up to graded commutativity, the non zero cup products of elements in $\mathrm{HK}^\bullet(A)$ are:*

$$\begin{aligned} z_0 \underset{K}{\smile} \bar{f} &= \bar{f} \text{ for all } \bar{f} \in \mathrm{HK}^\bullet(A) & z_\ell \underset{K}{\smile} \bar{\zeta}_0 &= \bar{\zeta}_\ell \text{ for } \ell \in \{0, 6, 8\} \\ z_6 \underset{K}{\smile} \bar{h}_i &= \bar{\gamma}_6 \text{ if } i \in \{1, 4\} & z_6 \underset{K}{\smile} \bar{h}_i &= -\bar{\gamma}_6 \text{ if } i \in \{2, 5\} \\ z_6 \underset{K}{\smile} \bar{\rho}_3 &= \bar{\zeta}_8 & \bar{\zeta}_0 \underset{K}{\smile} \bar{\rho}_3 &= \bar{\gamma}_4. \end{aligned}$$

Proof. The first two cup products are clear.

For the cup products of z_6 with the \bar{h}_i and for $\bar{\zeta}_0 \underset{K}{\smile} \bar{\rho}_5$, we use Lemma 6.12.

The last cup product follows from the fact that we have $\zeta_0 \underset{K}{\smile} \rho_3 - \gamma_4 = b_K^1(g')$ where $g'(a_0^*) = a_0^*c_3$ and $g'(a_3) = a_3c_3$. The cup product $z_6 \underset{K}{\smile} \bar{\rho}_3$ was already in the proof of Proposition 6.13.

Consideration of the coefficient weights yields the vanishing of the other cup products. \square

The cap products follow using duality, as in Corollary 6.4.

6.3.3. *Higher Koszul cohomology and homology* As in type A, the differential ∂_\smile sends z_ℓ to $2\bar{\zeta}_\ell$ for $\ell \in \{0, 6, 8\}$ and the differential ∂_\smile^2 is zero.

We then have the following higher Koszul cohomology.

PROPOSITION 6.15. *Let A be a preprojective algebra of type E_6 .*

If $\mathrm{char}(\mathbb{F}) = 2$, then $\mathrm{HK}_{hi}^\bullet(A) = \mathrm{HK}^\bullet(A)$.

If $\mathrm{char}(\mathbb{F}) = 3$, then

$$\begin{aligned} \mathrm{HK}_{hi}^0(A) &= \mathrm{HK}^0(A)_{10} \text{ has dimension 2 and is spanned by } \pi_0 \text{ and } \pi_3 \\ \mathrm{HK}_{hi}^2(A) &= \mathrm{HK}^2(A) \\ \mathrm{HK}_{hi}^1(A) &= \mathrm{span}\{[\bar{\rho}_5]\} \\ \mathrm{HK}_{hi}^p(A) &= 0 \text{ if } p > 2. \end{aligned}$$


If $\mathrm{char}(\mathbb{F}) \notin \{2, 3\}$, then

$$\begin{aligned} \mathrm{HK}_{hi}^0(A) &= \mathrm{HK}^0(A)_{10} \text{ has dimension 2 and is spanned by } \pi_0 \text{ and } \pi_3 \\ \mathrm{HK}_{hi}^2(A) &= \mathrm{HK}^2(A) \\ \mathrm{HK}_{hi}^p(A) &= 0 \text{ if } p \neq 0 \text{ and } p \neq 2. \end{aligned}$$

Higher Koszul homology follows from Theorem 4.4.

6.4. Koszul calculus for preprojective algebras of type E_7

The preprojective algebra A of type E_7 is defined by the quiver



subject to the relations

$$\sigma_0 = -a_0^*a_0$$

$$\sigma_4 = a_3a_3^* - a_4^*a_4$$

$$\begin{aligned}
\sigma_1 &= -a_1^* a_1 & \sigma_5 &= a_4 a_4^* - a_5^* a_5 \\
\sigma_2 &= a_1 a_1^* - a_2^* a_2 & \sigma_6 &= a_5 a_5^* \\
\sigma_3 &= a_0 a_0^* + a_2 a_2^* - a_3^* a_3
\end{aligned}$$

The Nakayama automorphism is given by $\nu(a_i) = -a_i$ and $\nu(a_i^*) = a_i^*$ for $i \in Q_0$.

To simplify notation, we shall denote by $c_0 = a_0 a_0^*$, $c_2 = a_2 a_2^*$ and $c_3 = a_3^* a_3$ the three 2-cycles at the vertex 3.

The socle of A is the part of weight 16 of A . A basis of the socle is given by $\pi_0 = (a_0^* c_3 a_0)^4$, $\pi_1 = -a_1^* a_2^* c_0 c_3 (c_3 c_0)^2 a_2 a_1$, $\pi_2 = -(a_2^* c_0 a_2)^4$, $\pi_3 = (c_3 c_0)^3 c_3^2$, $\pi_4 = -(a_3 c_0 a_3^*)^4$, $\pi_5 = -a_4 a_3 (c_3 c_0)^3 a_3^* a_4^*$ and $\pi_6 = -a_5 a_4 (a_3 c_0 a_3^*)^3 a_4^* a_5^*$.

6.4.1. *The Koszul cohomology and homology spaces in type E_7* We define the following elements

- in A : $z_0 = 1$, $z_8 = a_0^* c_2 c_0 c_2 a_0 - a_2^* c_2 c_0 c_2 a_2 - c_2 c_3^2 c_2 + a_3 c_0 c_2 c_0 a_3^* - a_4 a_3 c_2^2 a_3^* a_4^* + a_5 a_4 a_3 c_0 a_3^* a_4^* a_5^*$ and $z_{12} = a_0^* (c_2 c_0)^2 c_2 a_0^* + a_2^* (c_0 c_2)^2 c_0 a_2 - (c_3 c_0 c_3)^2 + a_3 c_3 (c_0 c_3)^2 a_3^*$;
- in $\text{Hom}_{k^e}(V, A)$: the maps ζ_ℓ defined by $\zeta_\ell(a_i) = a_i z_\ell$ for $\ell \in \{0, 8, 12\}$, the map ρ_3 defined by $\rho_3(a_2) = c_0 a_2$, $\rho_3(a_3) = a_3 c_3$, $\rho_3(a_4) = a_4 a_3 a_3^*$ and $\rho_3(a_2^*) = a_2^* c_3$, the map ρ_7 defined by $\rho_7(a_0) = c_3^3 a_0 + c_3 c_0 c_3 a_0$, $\rho_7(a_3) = a_3 c_3 c_0 c_3$ and $\rho_7(a_3^*) = c_3 c_0 c_3 a_3^*$, the map ρ_{15} defined by $\rho_{15}(a_0) = (c_2 c_0)^3 c_2 a_0$ and $\rho_{15}(a_0^*) = a_0^* c_2 (c_0 c_2)^3$ and the map ρ_5 defined by $\rho_5(a_0) = -c_2 c_3 a_0$, $\rho_5(a_2) = c_3 c_0 a_2$, $\rho_5(a_3) = a_3 c_3^2$, $\rho_5(a_0^*) = a_0^* c_3^2$ and $\rho_5(a_2^*) = -a_2^* c_2 c_0$;
- in $\text{Hom}_{k^e}(R, A)$: the maps h_j defined for $0 \leq j \leq 6$ by $h_j(\sigma_i) = \delta_{ij} e_j$ for all i , the map γ_4 defined by $\gamma_4(\sigma_0) = a_0^* c_3 a_0$, the map γ_8 defined by $\gamma_8(\sigma_0) = a_0^* c_3^3 a_0$, the map γ_{16} defined by $\gamma_{16}(\sigma_0) = \pi_0$ and the map and γ_6 defined by $\gamma_6(\sigma_0) = a_0^* c_3^2 a_0$.

LEMMA 6.16. *First assume that $\text{char}(\mathbb{F}) = 2$.*

- (i) *Let $u_{16} \in \text{Hom}_{k^e}(R, A)$ be an element of weight 16 so that $u_{16}(\sigma_i) = \lambda_i \pi_i$ for $i \in Q_0$. Then u_{16} is a coboundary if, and only if, $\sum_{i=0}^6 \lambda_i = 0$.*
- (ii) *Let $u_8 \in \text{Hom}_{k^e}(R, A)$ be an element of weight 8 so that $u_8(\sigma_0) = \lambda_0 a_0^* c_3^3 a_0$, $u_8(\sigma_2) = \lambda_2 a_2^* c_3 c_0 c_3 a_2 + \lambda_2' a_2^* c_3^2 c_0 a_2$, $u_8(\sigma_3) = \lambda_3 c_3^3 c_0 + \lambda_3' c_3 c_3^2 + \lambda_3'' c_0 c_3^2 c_0 + \lambda_3''' c_3^2 c_0 c_3$, $u_8(\sigma_4) = \lambda_4 a_3 c_3^2 c_0 a_3^* + \lambda_4' a_3 c_3 c_0 c_3 a_3^* + \lambda_4'' a_3 c_0 c_3^2 a_3^*$, $u_8(\sigma_5) = \lambda_5 a_4 a_3 c_3 c_0 a_3^* a_4^* + \lambda_5' a_4 a_3 c_0 c_3 a_3^* a_4^*$ and $u_8(\sigma_6) = a_5 a_4 a_3 a_0 a_0^* a_3^* a_4^* a_5^*$. Then u_8 is a coboundary if, and only if, $\lambda_0 + \lambda_2 + \lambda_2' + \lambda_3 + \lambda_3' + \lambda_3'' + \lambda_3''' + \lambda_4 + \lambda_4' + \lambda_4'' + \lambda_5 + \lambda_5' + \lambda_6 = 0$.*

Now assume that $\text{char}(\mathbb{F}) = 3$.

- (iii) *Let $u_6 \in \text{Hom}_{k^e}(R, A)$ be an element of weight 6 so that $u_6(\sigma_0) = \lambda_0 a_0^* c_3^2 a_0$, $u_6(\sigma_1) = \lambda_1 a_1^* a_2^* c_0 a_2^* a_1^*$, $u_6(\sigma_2) = \lambda_2 a_2^* c_3 c_0 a_2 + \lambda_2' a_2^* c_3^2 a_2$, $u_6(\sigma_3) = \lambda_3 c_3 c_0 c_3 + \lambda_3' c_3^2 c_0 + \lambda_3'' c_0 c_3 c_0 + \lambda_3''' c_3^3$, $u_6(\sigma_4) = \lambda_4 a_3 c_0 c_3 a_3^* + \lambda_4' a_3 c_3 c_0 a_3^*$ and $u_6(\sigma_5) = \lambda_5 a_4 a_3 c_0 a_3^* a_4^*$. Then u_6 is a coboundary if, and only if, $\sum_{i=0}^5 \lambda_i - \lambda_2' + \lambda_3' - \lambda_4' = 0$.*

Proof. For each $\ell \in \{16, 6\}$, if the map u_ℓ were a coboundary, it would be the image of a map $g_\ell \in \text{Hom}_{k^e}(V, A)$ whose coefficients would be in the space generated by the paths between adjacent vertices with weight $\ell - 1$. The proof is then straightforward once we know bases of these spaces. Note that once we have a basis of $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_w e_{s(\alpha)}$, applying κ gives a basis of $e_{s(\alpha)} A_w e_{t(\alpha)}$ for a given weight w .

In weight 15, a basis of $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_{15} e_{s(\alpha)}$ is given by $\bar{a}_0 = (a_0^* c_3 a_0)^3 a_0^* c_3$, $\bar{a}_1 = -a_1^* a_2^* c_0 c_3 (c_3 c_0)^2 a_2$, $\bar{a}_2 = -a_2^* c_0 (c_2 c_0)^3$, $\bar{a}_3 = (c_3 c_0)^3 c_3 a_3^*$, $\bar{a}_4 = -a_3 (c_3 c_0)^3 a_3^* a_4^*$, $\bar{a}_5 = a_4 (a_3 c_0 a_3^*)^3 a_4^* a_5^*$.

In weight 7, a basis of $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_7 e_{s(\alpha)}$ is given by $c_3^3 a_0$, $c_3 c_0 c_3 a_0$, $a_2^* c_3^2 a_2 a_1$, $c_3^2 c_0 a_2$, $c_0 c_3^2 a_2$, $c_3 c_0 c_3 a_2$, $a_3 c_3 c_0 c_3$, $a_3 c_0 c_3^2$, $a_3 c_3^2 c_0$, $a_4 a_3 c_3 c_0 a_3^*$, $a_4 a_3 c_0 c_3 a_3^*$, $a_5 a_4 a_3 c_0 a_3^* a_4^*$.

In weight 5, a basis of $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_5 e_{s(\alpha)}$ is given by $c_3^2 a_0$, $c_0 c_3 a_0$, $a_2^* c_0 a_2 a_1$, $c_3 c_0 a_2$, $c_0 c_3 a_2$, $a_3 c_3 c_0$, $a_3 c_0 c_3$, $a_3 c_3^2$, $a_4 a_3 c_0 a_3^*$. \square

PROPOSITION 6.17. *Let A be a preprojective algebra of type E_7 .*

- (i) *The elements in $\{z_0, z_8, z_{12}\} \cup \{\pi_i; i \in Q_0\}$ form a basis of $\mathrm{HK}^0(A)$.*
- (ii) *If $\mathrm{char}(\mathbb{F}) \notin \{2, 3\}$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 8, 12\}$ form a basis of $\mathrm{HK}^1(A)$.*
If $\mathrm{char}(\mathbb{F}) = 2$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 8, 12\} \cup \{\bar{\rho}_3, \bar{\rho}_7, \bar{\rho}_{15}\}$ form a basis of $\mathrm{HK}^1(A)$.
If $\mathrm{char}(\mathbb{F}) = 3$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 8, 12\} \cup \{\bar{\rho}_5\}$ form a basis of $\mathrm{HK}^1(A)$.
- (iii) *If $\mathrm{char}(\mathbb{F}) \notin \{2, 3\}$, the elements in $\{\bar{h}_j; j \in Q_0\}$ form a basis of $\mathrm{HK}^2(A)$.*
If $\mathrm{char}(\mathbb{F}) = 2$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_4, \bar{\gamma}_8, \bar{\gamma}_{16}\}$ form a basis of $\mathrm{HK}^2(A)$.
If $\mathrm{char}(\mathbb{F}) = 3$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_6\}$ form a basis of $\mathrm{HK}^2(A)$.

Proof. The centre was given in [22], so we have (i).

For $\mathrm{HK}^1(A)$ and $\mathrm{HK}^2(A)$, the number of elements in the statement is equal to the dimension of the corresponding cohomology space. Moreover, all the elements in the statement are indeed cocycles.

If $\mathrm{char}(\mathbb{F})$ is not 2 or 3, a basis of $\mathrm{HK}^1(A) = \mathrm{HH}^1(A)$ was given in [22]. It consists of the classes of the ζ'_ℓ with $\ell \in \{0, 8, 12\}$ where $\zeta'_\ell(a_i) = a_i z_\ell$ for $0 \leq i \leq 2$ and $\zeta'_\ell(a_i^*) = a_i^* z_\ell$ for $3 \leq i \leq 5$. Since $\zeta_0 - \zeta'_0$ is equal to $b_K^1(e_3 + 2e_5 + 3e_6)$, and $\zeta_\ell - \zeta'_\ell = (\zeta_0 - \zeta'_0) \smile_K z_\ell$ is also a coboundary, ζ_ℓ and ζ'_ℓ represent the same cohomology class for $\ell \in \{0, 8, 12\}$. Moreover, as in types A, D and E_6 , the elements \bar{h}_j form a basis of $\mathrm{HK}^2(A)$.

If $\mathrm{char}(\mathbb{F}) \in \{2, 3\}$, we need only prove that the extra elements are not coboundaries by fact (N3).

If $\mathrm{char}(\mathbb{F}) = 2$, it follows from Lemma 6.16 that $\zeta_0 \smile_K \rho_{15} - \gamma_{16}$, $\zeta_8 \smile_K \rho_7 - \gamma_{16}$, $\zeta_{12} \smile_K \rho_3 - \gamma_{16}$, $z_8 \smile_K \gamma_8 - \gamma_{16}$ and $z_{12} \smile_K \gamma_4 - \gamma_{16}$ are coboundaries. Therefore it is enough to check that γ_{16} is not a coboundary, and this also follows from Lemma 6.16.

If $\mathrm{char}(\mathbb{F}) = 3$, again using Lemma 6.16, $\zeta_0 \smile_K \rho_5 - \gamma_6$ is a coboundary and γ_6 is not a coboundary, therefore ρ_5 is not a coboundary either. \square

Koszul homology follows using duality (Theorem 4.4), as in Corollary 6.2.

6.4.2. *Cup and cap products* We now determine the cup products of the elements in the bases of the Koszul cohomology spaces given above.

PROPOSITION 6.18. *Let A be a preprojective algebra of type E_7 . Up to graded commutativity, the non zero cup products of elements in $\mathrm{HK}^\bullet(A)$ are:*

$$\begin{array}{ll}
 z_0 \smile_K \bar{f} = \bar{f} \text{ for all } \bar{f} \in \mathrm{HK}^\bullet(A) & z_8^2 = \pi_0 + \pi_4 - \pi_6 \\
 z_\ell \smile_K \bar{\zeta}_0 = \bar{\zeta}_\ell \text{ for } \ell \in \{0, 8, 12\} & z_8 \smile_K \bar{\rho}_7 = \bar{\rho}_{15} \\
 z_{12} \smile_K \bar{\rho}_3 = \bar{\rho}_{15} & \pi_i \smile_K \bar{h}_i = \bar{\gamma}_{16} \text{ for } i \in Q_0 \\
 z_8 \smile_K \bar{h}_i = \bar{\gamma}_8 \text{ if } i \in \{0, 4, 6\} & z_8 \smile_K \bar{\gamma}_8 = \bar{\gamma}_{16} \\
 z_{12} \smile_K \bar{\gamma}_4 = \bar{\gamma}_{16} & \bar{\zeta}_0 \smile_K \bar{\rho}_\ell = -\bar{\gamma}_{\ell+1} \text{ for } \ell \in \{3, 7, 15, 6\} \\
 \bar{\zeta}_8 \smile_K \bar{\rho}_7 = \bar{\gamma}_{16} & \bar{\zeta}_{12} \smile_K \bar{\rho}_3 = \bar{\gamma}_{16}
 \end{array}$$

Proof. Most of the cup-products are easy to compute, follow from Lemma 6.16 or vanish for weight reasons. The remaining ones are obtained as follows (at the level of cochains):

$$z_8 \smile_K \rho_7 = \rho_{15} + b_K^1([e_3 \mapsto (c_3 c_0)^3 c_3])$$

$$\begin{aligned} z_{12} \underset{K}{\smile} \rho_3 &= \rho_{15} + b_K^1([e_3 \mapsto (c_3 c_0)^3 c_2]) \\ \zeta_0 \underset{K}{\smile} \rho_3 &= \gamma_4 + b_K^1(h) \end{aligned}$$

where $h \in \text{Hom}_{k^e}(V, A_3)$ is defined by $h(a_0) = c_3 a_0$ and $h(a_2) = c_2 a_2$. □

The cap products follow using duality, as in Corollary 6.4.

6.4.3. *Higher Koszul cohomology and homology* As in types A, D and E_6 , the differential ∂_{\smile}^1 sends z_ℓ to $2\bar{\zeta}_\ell$ for $\ell \in \{0, 8, 12\}$ and the differential ∂_{\smile}^2 is zero except when $\text{char}(\mathbb{F}) = 3$ where $\partial_{\smile}^2(\bar{\rho}_5) = \bar{\gamma}_6$.

We then have the following higher Koszul cohomology.

PROPOSITION 6.19. *Let A be a preprojective algebra of type E_7 .*

If $\text{char}(\mathbb{F}) = 2$, then $\text{HK}_{hi}^\bullet(A) = \text{HK}^\bullet(A)$.


If $\text{char}(\mathbb{F}) \neq 2$, then

$$\begin{aligned} \text{HK}_{hi}^0(A) &= \text{HK}^0(A)_{16} \text{ has dimension 7 and is spanned by the } \pi_i \text{ for } i \in Q_0 \\ \text{HK}_{hi}^2(A) &= \text{HK}^2(A)_0 \text{ has dimension 7 and is spanned by the } [\bar{h}_i] \text{ for } i \in Q_0 \\ \text{HK}_{hi}^p(A) &= 0 \text{ if } p \neq 0 \text{ and } p \neq 2. \end{aligned}$$

Higher Koszul homology follows from Theorem 4.4.

6.5. Koszul calculus for preprojective algebras of type E_8

The preprojective algebra A of type E_8 is defined by the quiver



subject to the relations

$$\begin{aligned} \sigma_0 &= -a_0^* a_0 & \sigma_4 &= a_3 a_3^* - a_4^* a_4 \\ \sigma_1 &= -a_1^* a_1 & \sigma_5 &= a_4 a_4^* - a_5^* a_5 \\ \sigma_2 &= a_1 a_1^* - a_2^* a_2 & \sigma_6 &= a_5 a_5^* - a_6^* a_6 \\ \sigma_3 &= a_0 a_0^* + a_2 a_2^* - a_3^* a_3 & \sigma_7 &= a_6 a_6^* \end{aligned}$$

The Nakayama automorphism is given by $\nu(a_i) = -a_i$ and $\nu(a_i^*) = a_i^*$ for $i \in Q_0$.

To simplify notation, we shall denote by $c_0 = a_0 a_0^*$, $c_2 = a_2 a_2^*$ and $c_3 = a_3^* a_3$ the three 2-cycles at the vertex 3.

The socle of A is the part of weight 28 of A . A basis of the socle is given by $\pi_0 = (a_0^* c_2 a_0)^7$, $\pi_1 = a_1^* a_2^* (c_0 c_2)^5 c_2 c_0 a_2 a_1$, $\pi_2 = -(a_2^* c_0 a_2)^7$, $\pi_3 = (c_2 c_0)^7$, $\pi_4 = -a_3 (c_2 c_0)^6 c_2 a_3^*$, $\pi_5 = a_4 a_3 (c_2 c_0)^6 a_3^* a_4^*$, $\pi_6 = a_5 a_4 a_3 (c_0 c_2)^5 c_0 a_3^* a_4^* a_5^*$ and $\pi_7 = -a_6 a_5 a_4 a_3 (c_0 c_2)^3 (c_2 c_0)^2 a_3^* a_4^* a_5^* a_6^*$.

6.5.1. *The Koszul cohomology and homology spaces in type E_8* We define the following elements

- in A :
- $z_0 = 1$,

- $z_{12} = a_5 a_4 a_3 c_0 c_2 c_0 a_3^* a_4^* a_5^* + a_4 a_3 (c_2 c_0)^2 a_3^* a_4^* + a_3 (c_2 c_0)^2 c_2 a_3^* - (c_3 c_2 c_3)^2 + a_2^* (c_0 c_2)^2 c_0 a_2 - a_1^* a_2^* c_0 c_2^2 c_0 a_2 a_1 + a_0^* (c_2 c_0)^2 c_2 a_0$,
- $z_{20} = a_5 a_4 a_3 (c_0 c_2)^3 c_0 a_3^* a_4^* a_5^* + a_4 a_3 (c_0 c_2)^2 (c_2 c_0)^2 a_3^* a_4^* + a_3 c_0 (c_2 c_0)^4 a_3^* + (c_2 c_0)^5 - (c_0 c_2^2)^3 c_0 + (c_0 c_2)^5 + a_2^* (c_2 c_0 c_2)^3 a_2 + a_0^* (c_2 c_0)^4 c_2 a_0$ and
- $z_{24} = z_{12}^2$;
- in $\text{Hom}_{k^e}(V, A)$:
 - the maps ζ_ℓ defined by $\zeta_\ell(a_i) = a_i z_\ell$ for $\ell \in \{0, 12, 20, 24\}$,
 - the map ρ_3 defined by $\rho_3(a_2) = c_0 a_2$, $\rho_3(a_3) = a_3 c_3$, $\rho_3(a_4) = a_4 a_3 a_3^*$, $\rho_3(a_5) = a_5 a_4 a_4^*$ and $\rho_3(a_2^*) = a_2^* c_3$,
 - the map ρ_7 defined by $\rho_7(a_0) = c_0 c_3^2 a_0$, $\rho_7(a_3) = a_3 c_0 c_3^2 + a_3 c_3^2 c_0 + a_3 c_0 c_3 c_0$ and $\rho_7(a_3^*) = c_3 c_0 c_3 a_3^*$,
 - the map ρ_{15} defined by $\rho_{15}(a_3) = a_3 c_2 (c_2 c_0)^2$, $\rho_{15}(a_4) = a_4 a_3 (c_0 c_3)^2 a_3^* + a_4 a_3 (c_0 c_3^2)^2 a_3^* + a_4 a_3 (c_3 c_0)^3 a_3^*$, $\rho_{15}(a_5) = a_5 a_4 a_3 c_0 (c_3 c_0)^2 a_3^* a_4^*$, $\rho_{15}(a_0^*) = a_0^* c_3 (c_3 c_0)^3$ and $\rho_{15}(a_5^*) = a_4 a_3 (c_0 c_3)^2 c_0 a_3^* a_4^* a_5^*$,
 - the map ρ_{27} defined by $\rho_{27}(a_3) = a_3 c_0 (c_3 c_0)^6$ and $\rho_{27}(a_3^*) = (c_3 c_0)^6 c_3 a_3^*$,
 - the map ρ_5 defined by $\rho_5(a_0) = -c_0 c_3 a_0 - c_3^2 a_0$, $\rho_5(a_3) = a_3 c_3 c_0 + a_3 c_3^2$, $\rho_5(a_4) = a_4 a_3 c_3 a_3^*$, $\rho_5(a_0^*) = a_0^* c_3^2$ and $\rho_5(a_3^*) = -c_3 c_0 a_3^*$,
 - the map ρ_{17} defined by defined by $\rho_{17}(a_0) = -c_3^2 (c_0 c_3)^3 a_0 - (c_0 c_3)^4 a_0$, $\rho_{17}(a_3) = a_3 (c_3 c_0)^4 + a_3 (c_0 c_3)^4 + a_3 (c_3 c_0)^3 c_3^2$, $\rho_{17}(a_0^*) = a_0^* (c_2 c_0)^3 c_3^2$ and $\rho_{17}(a_3^*) = -(c_3 c_0)^4 a_3^*$, and
 - the map ρ_9 defined by $\rho_9(a_0) = -2c_3^2 c_0 c_3 a_0 + 2c_3 c_0 c_3^2 a_0 + (c_0 c_3)^2 a_0$, $\rho_9(a_2) = c_2^2 c_0 c_2 a_2 + (c_2 c_0)^2 a_2$, $\rho_9(a_3) = -a_3 (c_0 c_3)^2$, $\rho_9(a_0^*) = 2a_0^* c_3^2 c_0 c_3 - 2a_0^* c_3 c_0 c_3^2 + a_0^* (c_3 c_0)^2$, $\rho_9(a_2^*) = a_2^* c_2 c_0 c_2^2 - a_2^* (c_2 c_0)^2$ and $\rho_9(a_3^*) = (c_0 c_3)^2 a_3^*$;
- in $\text{Hom}_{k^e}(R, A)$: the maps h_j defined for $0 \leq j \leq 7$ by $h_j(\sigma_i) = \delta_{ij} e_j$ for all i , and

$$\begin{array}{ll}
\gamma_4: \sigma_0 \mapsto a_0^* c_3 a_0 & \gamma_8: \sigma_0 \mapsto a_0^* c_3^3 a_0 \\
\gamma_{16}: \sigma_0 \mapsto a_0^* (c_2 c_0)^3 c_2 a_0 & \gamma_{28}: \sigma_0 \mapsto \pi_0 \\
\gamma_6: \sigma_0 \mapsto a_0^* c_3^2 a_0 & \gamma_{18}: \sigma_0 \mapsto a_0^* c_3^2 (c_0 c_3)^3 a_0 \\
\gamma_{10}: \sigma_0 \mapsto a_0^* c_3^2 c_0 c_3 a_0 &
\end{array}$$

LEMMA 6.20. *First assume that $\text{char}(\mathbb{F}) = 2$.*

- (i) *Let $u_{28} \in \text{Hom}_{k^e}(R, A)$ be an element of weight 28, so that $u_{28}(\sigma_i) = \lambda_i \pi_i$ for all $i \in Q_0$. Then u_{28} is a coboundary if, and only if, $\sum_{i \in Q_0} \lambda_i = 0$.*

Now assume that $\text{char}(\mathbb{F}) = 3$.

- (ii) *Let $u_{18} \in \text{Hom}_{k^e}(R, A)$ be an element of weight 18, so that $u_{18}(\sigma_0) = \lambda_0 a_0^* c_3^2 (c_0 c_3)^3 a_0 + \lambda_0' a_0^* c_3 c_0 (c_3 c_0 c_3)^2 a_0$, $u_{18}(\sigma_1) = \lambda_1 a_1^* a_2^* c_0 (c_2 c_0)^3 a_2 a_1$, $u_{18}(\sigma_2) = \lambda_2 a_2^* (c_2 c_0 c_2)^2 c_0 c_2 a_2 + \lambda_2' a_2^* (c_0 c_2)^4 a_2 + \lambda_2'' a_2^* (c_2 c_0)^4 a_2$, $u_{18}(\sigma_3) = \lambda_3 c_0 (c_3 c_0)^4 + \lambda_3' (c_3 c_0)^4 c_3 + \lambda_3'' (c_3 c_0)^2 c_3 (c_3 c_0)^2 + \lambda_3^{(3)} (c_0 c_3)^2 c_3 (c_0 c_3)^2 + \lambda_3^{(4)} (c_3 c_0)^3 c_3^2 c_0 + \lambda_3^{(5)} c_0 c_3^2 (c_0 c_3)^3$, $u_{18}(\sigma_4) = \lambda_4 a_3 (c_3 c_0)^4 a_3^* + \lambda_4' a_3 (c_0 c_3)^4 a_3^* + \lambda_4'' a_3 c_3^2 (c_3 c_0)^3 a_3^* + \lambda_4^{(3)} a_3 (c_0 c_3)^3 c_3^2 a_3^*$, $u_{18}(\sigma_5) = \lambda_5 a_4 a_3 (c_3 c_0)^3 c_3 a_3^* a_4^* + \lambda_5' a_4 a_3 c_3 (c_3 c_0)^3 a_3^* a_4^* + \lambda_5'' a_4 a_3 (c_0 c_3)^3 c_3 a_3^* a_4^*$, $u_{18}(\sigma_6) = \lambda_6 a_5 a_4 a_3 (c_3 c_0)^3 a_3^* a_4^* a_5^* + \lambda_6' a_5 a_4 a_3 (c_0 c_3)^3 a_3^* a_4^* a_5^*$, $u_{18}(\sigma_7) = \lambda_7 a_6 a_5 a_4 a_3 (c_0 c_3)^2 c_0 a_3^* a_4^* a_5^* a_6^*$. Then u_{18} is a coboundary if, and only if, $\lambda_0 + \lambda_0' + \lambda_1 + \lambda_1' + \lambda_2 + \lambda_2' + \lambda_2'' + \lambda_3 + \lambda_3' + \lambda_3'' + \lambda_3^{(3)} + \lambda_3^{(4)} + \lambda_3^{(5)} + \lambda_4 + \lambda_4' - \lambda_4'' - \lambda_4^{(3)} - \lambda_5 - \lambda_5' - \lambda_5'' - \lambda_6 - \lambda_6' - \lambda_7 = 0$.*

- (iii) *The map $\rho_{17} \in \text{Hom}_{k^e}(V, A)$ is not a coboundary.*

Now assume that $\text{char}(\mathbb{F}) = 5$.

- (iv) *Let $u_{10} \in \text{Hom}_{k^e}(R, A)$ be an element of weight 10, so that $u_{10}(\sigma_0) = \lambda_0 a_0^* c_3^2 c_0 c_3 a_0 + \lambda_0' a_0^* c_3 c_0 c_3^2 a_0$, $u_{10}(\sigma_1) = \lambda_1 a_1^* a_2^* c_0 c_2 c_0 a_2 a_1$, $u_{10}(\sigma_2) = \lambda_2 a_2^* (c_0 c_2)^2 a_2 + \lambda_2' a_2^* (c_2 c_0)^2 a_2 + \lambda_2'' a_2^* c_0 c_2^2 c_0 a_2$, $u_{10}(\sigma_3) = \lambda_3 c_0 (c_3 c_0)^2 + \lambda_3' c_3 (c_0 c_3)^2 + \lambda_3'' c_3 (c_3 c_0)^2 + \lambda_3^{(3)} (c_0 c_3)^2 c_3 + \lambda_3^{(4)} c_3 c_0 c_3^2 c_0 + \lambda_3^{(5)} c_0 c_3^2 c_0 c_3$, $u_{10}(\sigma_4) = \lambda_4 a_3 (c_0 c_3)^2 a_3^* + \lambda_4' a_3 (c_3 c_0)^2 a_3^* + \lambda_4'' a_3 c_3 c_0 c_3^2 a_3^* +$*

$\lambda_4^{(3)} a_3 c_3^2 c_0 c_3 a_3^*$, $u_{10}(\sigma_5) = \lambda_5 a_4 a_3 c_0 c_3 c_0 a_3^* a_4^* + \lambda_5' a_4 a_3 c_3^2 c_0 a_3^* a_4^* + \lambda_5'' a_4 a_3 c_0 c_3^2 a_3^* a_4^*$,
 $u_{10}(\sigma_6) = \lambda_6 a_5 a_4 a_3 c_3 c_0 a_3^* a_4^* a_5^* + \lambda_6' a_5 a_4 a_3 c_0 c_3 a_3^* a_4^* a_5^*$, $u_{10}(\sigma_7) = \lambda_7 a_6 a_5 a_4 a_3 c_0 a_3^* a_4^* a_5^* a_6^*$.
 Then u_{10} is a coboundary if, and only if, $\lambda_0 + \lambda_0' + \lambda_1 + \lambda_2 + \lambda_2' + \lambda_2'' + \lambda_3' + \lambda_3'' + \lambda_3^{(3)} + \lambda_3^{(4)} + \lambda_3^{(5)} + \lambda_4 - \lambda_4' + 2\lambda_4'' + 2\lambda_4^{(3)} + \lambda_5 + 2\lambda_5' + 2\lambda_5'' + 2\lambda_6 + 2\lambda_6' + 2\lambda_7 = 0$.

Proof. For each $\ell \in \{28, 18, 10\}$, if the map u_ℓ were a coboundary, it would be the image of a map $g_\ell \in \text{Hom}_{k^e}(V, A)$ whose coefficients would be in the space generated by the paths between adjacent vertices with weight $\ell - 1$. The proof of (i), (ii) and (iv) is then straightforward once we know bases of these spaces.

In weight 27, a basis is given by $\bar{a}_0 = a_0^* c_2 (c_0 c_2)^6$, $\bar{a}_1 = a_1^* a_2^* (c_0 c_2)^5 c_3^2 a_2$, $\bar{a}_2 = -a_2^* c_0 (c_2 c_0)^6$, $\bar{a}_3 = (c_2 c_0)^6 c_2 a_3^*$, $\bar{a}_4 = -a_3 (c_2 c_0)^6 a_3^* a_4^*$, $\bar{a}_5 = -a_4 a_3 c_0 (c_2 c_0)^5 a_3^* a_4^* a_5^*$, $\bar{a}_6 = a_5 a_4 a_3 (c_0 c_2)^3 (c_2 c_0)^2 a_3^* a_4^* a_5^* a_6^*$, and the $\kappa(\bar{a}_i)$ for all $i \in Q_0$.

In weight 17, the space $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_{18} e_{s(\alpha)}$ has basis $(c_0 c_3)^4 a_0$, $c_3^2 (c_0 c_3)^3 a_0$, $c_3 c_0 (c_3 c_0 c_3)^2 a_0$, $a_2^* c_2 c_0 c_2 (c_2 c_0)^2 a_2 a_1$, $a_2^* (c_0 c_2)^3 c_0 a_2 a_1$, $(c_3 c_0)^4 a_2$, $(c_0 c_3)^4 a_2$, $(c_0 c_3)^3 c_3 c_0 a_2$, $c_3 c_0 (c_3 c_0 c_3)^2 a_2$, $a_3 (c_3 c_0)^4$, $a_3 (c_0 c_3)^4$, $a_3 (c_0 c_3)^3 c_3 c_0$, $a_3 c_0 c_3 (c_3 c_0)^3$, $a_3 (c_3 c_0)^3 c_3^2$, $a_4 a_3 (c_0 c_3)^3 c_0 a_3^*$, $a_4 a_3 (c_0 c_3)^3 c_3 a_3^*$, $a_4 a_3 c_3 (c_3 c_0)^3 a_3^*$, $a_5 a_4 a_3 (c_3 c_0)^3 a_3^* a_4^*$, $a_5 a_4 a_3 (c_0 c_3)^3 a_3^* a_4^*$, $a_6 a_5 a_4 a_3 (c_0 c_3)^2 c_0 a_3^* a_4^* a_5^*$ and $\bigoplus_{\alpha \in Q_1} e_{s(\alpha)} A_{18} e_{t(\alpha)} = \kappa \left(\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_{18} e_{s(\alpha)} \right)$.

In weight 9, the space $\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_9 e_{s(\alpha)}$ has basis $c_3^2 c_0 c_3 a_0$, $c_3 c_0 c_3^2 a_0$, $(c_0 c_3)^2 a_0$, $a_2^* c_0 c_2 c_0 a_2 a_1$, $c_2^2 c_0 c_2 a_2$, $(c_2 c_0)^2 a_2$, $(c_0 c_2)^2 a_2$, $c_0 c_2^2 c_0 a_2$, $a_3 c_3 c_0 c_3^2$, $a_3 (c_0 c_3)^2$, $a_3 (c_3 c_0)^2$, $a_3 c_0 c_3^2 c_0$, $a_4 a_3 c_3^2 c_0 a_3^*$, $a_4 a_3 c_0 c_3^2 a_3^*$, $a_4 a_3 c_0 c_3 c_0 a_3^*$, $a_5 a_4 a_3 c_3 c_0 a_3^* a_4^*$, $a_5 a_4 a_3 c_0 c_3 a_3^* a_4^*$, $a_6 a_5 a_4 a_3 c_0 a_3^* a_4^* a_5^*$ and $\bigoplus_{\alpha \in Q_1} e_{s(\alpha)} A_9 e_{t(\alpha)} = \kappa \left(\bigoplus_{\alpha \in Q_1} e_{t(\alpha)} A_9 e_{s(\alpha)} \right)$.

Finally, if ρ_{17} were a coboundary, it would be equal to $b_k^1(g)$ for some $g \in \text{Hom}_{k^e}(k, A_{16})$. Such a map g would necessarily satisfy $g(e_0) \in \text{span} \{a_0^* (c_3 c_0)^3 c_3 a_0\}$, $g(e_3) \in \text{span} \{(c_3 c_0)^4, (c_0 c_3)^4, c_0 c_3 (c_3 c_0)^3, (c_0 c_3)^3 c_3 c_0, c_3^2 (c_0 c_3)^3, c_3 c_0 (c_3 c_0 c_3)^2\}$, and $g(e_4) \in \text{span} \{a_3 c_0 (c_3 c_0)^3 a_3^*, a_3 c_3 (c_0 c_3)^3 a_3^*, a_3 c_3 (c_3 c_0)^3 a_3^*, a_3 (c_0 c_3)^3 c_3 a_3^*\}$. Then, by considering $b_k^1(g)(a_0)$, $b_k^1(g)(a_0^*)$ and $b_k^1(g)(a_3)$ we get a contradiction. \square

PROPOSITION 6.21. *Let A be a preprojective algebra of type E_8 .*

- (i) *The elements in $\{z_0, z_{12}, z_{20}, z_{24}\} \cup \{\pi_i; i \in Q_0\}$ form a basis of $\text{HK}^0(A)$.*
- (ii) *If $\text{char}(\mathbb{F}) \notin \{2, 3, 5\}$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 12, 20, 24\}$ form a basis of $\text{HK}^1(A)$.*
If $\text{char}(\mathbb{F}) = 2$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 12, 20, 24\} \cup \{\bar{\rho}_3, \bar{\rho}_7, \bar{\rho}_{15}, \bar{\rho}_{27}\}$ form a basis of $\text{HK}^1(A)$.
If $\text{char}(\mathbb{F}) = 3$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 12, 20, 24\} \cup \{\bar{\rho}_5, \bar{\rho}_{17}\}$ form a basis of $\text{HK}^1(A)$.
If $\text{char}(\mathbb{F}) = 5$, the elements in $\{\bar{\zeta}_\ell; \ell = 0, 12, 20, 24\} \cup \{\bar{\rho}_9\}$ form a basis of $\text{HK}^1(A)$.
- (iii) *If $\text{char}(\mathbb{F}) \notin \{2, 3, 5\}$, the elements in $\{\bar{h}_j; j \in Q_0\}$ form a basis of $\text{HK}^2(A)$.*
If $\text{char}(\mathbb{F}) = 2$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_4, \bar{\gamma}_8, \bar{\gamma}_{16}, \bar{\gamma}_{28}\}$ form a basis of $\text{HK}^2(A)$.
If $\text{char}(\mathbb{F}) = 3$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_6, \bar{\gamma}_{18}\}$ form a basis of $\text{HK}^2(A)$.
If $\text{char}(\mathbb{F}) = 5$, the elements in $\{\bar{h}_j; j \in Q_0\} \cup \{\bar{\gamma}_{10}\}$ form a basis of $\text{HK}^2(A)$.

Proof. The centre was given in [22], so we have (i).

For $\text{HK}^1(A)$ and $\text{HK}^2(A)$, the number of elements in the statement is equal to the dimension of the corresponding cohomology space. Moreover, all the elements in the statement are indeed cocycles.

If $\text{char}(\mathbb{F})$ is not 2, 3 or 5, a basis of $\text{HK}^1(A) = \text{HH}^1(A)$ was given in [22]. It consists of the classes of the ζ'_ℓ with $\ell \in \{0, 12, 20, 24\}$ where $\zeta'_\ell(a_i) = a_i z_\ell$ for $0 \leq i \leq 2$ and $\zeta'_\ell(a_i^*) = a_i^* z_\ell$ for $3 \leq i \leq 6$. Since $\zeta_0 - \zeta'_0$ is equal to $b_K^1(e_3 + 2e_5 + 3e_6 + 4e_7)$, and $\zeta_\ell - \zeta'_\ell = (\zeta_0 - \zeta'_0) \underset{K}{\smile} z_\ell$

is also a coboundary, ζ_ℓ and ζ'_ℓ represent the same cohomology class for $\ell \in \{0, 12, 20, 24\}$. Moreover, as in types A, D, E₆ and E₇, the elements \bar{h}_j form a basis of $\text{HK}^2(A)$.

The rest of the proof is the same as that of Proposition 6.17, based on Lemma 6.20 and the fact that the following cup products at the level of cochains are all coboundaries: $\zeta_0 \smile_K \rho_{27} - \gamma_{28}$, $\zeta_{12} \smile_K \rho_{15} - \gamma_{28}$, $\zeta_{24} \smile_K \rho_3 - \gamma_{28}$, $\zeta_{20} \smile_K \rho_7 - \gamma_{28}$, $z_{24} \smile_K \gamma_4 - \gamma_{28}$, $z_{12} \smile_K \gamma_{16} - \gamma_{28}$, $z_{20} \smile_K \gamma_8 - \gamma_{28}$, $z_{12} \smile_K \rho_5 - \rho_{17}$, $z_{12} \smile_K \gamma_6 - \gamma_{18}$, $\zeta_0 \smile_K \rho_9 - \gamma_{10}$, whereas γ_{28} , γ_{18} , γ_{10} and ρ_{17} are not. \square

Koszul homology follows using duality (Theorem 4.4), as in Corollary 6.2.

6.5.2. *Cup and cap products* We now determine the cup products of the elements in the bases of the Koszul cohomology spaces given above.

PROPOSITION 6.22. *Let A be a preprojective algebra of type E₈. Up to graded commutativity, the non zero cup products of elements in $\text{HK}^\bullet(A)$ are:*

$$\begin{array}{ll}
z_0 \smile_K \bar{f} = \bar{f} \text{ for all } \bar{f} \in \text{HK}^\bullet(A) & z_{12}^2 = z_{24} \\
z_\ell \smile_K \bar{\zeta}_0 = \bar{\zeta}_\ell \text{ for } \ell \in \{0, 12, 20, 24\} & z_{12} \smile_K \bar{\rho}_3 = \bar{\rho}_{15} \\
z_{24} \smile_K \bar{\rho}_3 = \bar{\rho}_{27} & z_{12} \smile_K \bar{\rho}_{15} = \bar{\rho}_{27} \\
z_{20} \smile_K \bar{\rho}_7 = \bar{\rho}_{27} & z_{12} \smile_K \bar{\rho}_5 = \bar{\rho}_{17} \\
z_{12} \smile_K \bar{\rho}_9 = -\bar{\zeta}_{20} & z_{12} \smile_K \bar{\zeta}_{12} = \bar{\zeta}_{24} \\
\pi_i \smile_K \bar{h}_i = \bar{\gamma}_{28} \text{ for } i \in Q_0 & z_{12} \smile_K \bar{\gamma}_4 = \bar{\gamma}_{16} \\
z_{12} \smile_K \bar{\gamma}_{16} = \bar{\gamma}_{28} & z_{24} \smile_K \bar{\gamma}_4 = \bar{\gamma}_{28} \\
z_{20} \smile_K \bar{\gamma}_8 = \bar{\gamma}_{28} & z_{12} \smile_K \bar{\gamma}_6 = \bar{\gamma}_{18} \\
\bar{\zeta}_0 \smile_K \bar{\rho}_\ell = -\bar{\gamma}_{\ell+1} \text{ for } \ell \in \{3, 7, 15, 27, 9\} & \bar{\zeta}_{12} \smile_K \bar{\rho}_3 = \bar{\gamma}_{16} \\
\bar{\zeta}_{12} \smile_K \bar{\rho}_{15} = \bar{\gamma}_{28} & \bar{\zeta}_{20} \smile_K \bar{\rho}_7 = \bar{\gamma}_{28} \\
\bar{\zeta}_{24} \smile_K \bar{\rho}_3 = \bar{\gamma}_{28}. &
\end{array}$$

Proof. Most of the cup-products are easy to compute, follow from Lemma 6.20 or vanish for weight reasons. The remaining ones are obtained as follows (at the level of cochains): $z_{24} \smile_K \rho_3 = \rho_{27} + b_K^1([e_3 \mapsto c_0(c_3c_0)^6])$, we have $z_{12} \smile_K \rho_3 = \rho_{15} + b_K^1(h)$ where h is defined by $h(a_2) = (c_0c_2)^3c_0a_2$, $h(a_3) = a_3c_0(c_3c_0)^3 + a_3c_3(c_0c_3)^3$, $h(a_4) = a_4a_3(c_0c_3^2)^2a_3^*$ and $h(a_2^*) = a_2^*c_0(c_2c_0)^3 + a_2^*c_2(c_0c_2)^3$, and finally $z_{12} \smile_K \rho_9 + \zeta_{20} = b_K(h')$ where h' is defined by $h'(e_0) = a_0^*(c_3c_0)^4c_3a_0$, $h'(e_2) = 2a_2^*(c_2c_0)^4c_2a_2$, $h'(e_3) = -2(c_0c_3)^5 + 2c_3^2(c_0c_3)^4 + (c_3c_0)^4c_3^2$, $h'(e_4) = -a_3(c_0c_3)^4c_0a_3^* + a_3c_3(c_3c_0)^4a_3^* - a_3(c_0c_3)^4c_3a_3^*$ and $h'(e_5) = -a_4a_3(c_0c_3)^4a_3^*a_4^*$. \square

The cap products follow using duality, as in Corollary 6.4.

6.5.3. *Higher Koszul cohomology and homology* As in types A, D, E₆ and E₇, the differential ∂^1_\smile sends z_ℓ to $2\zeta_\ell$ for $\ell \in \{0, 8, 12\}$ and the differential ∂^2_\smile is zero except when $\text{char}(\mathbb{F}) = 5$ where $\partial^2_\smile(\bar{\rho}_9) = 2\bar{\gamma}_{10}$.

We then have the following higher Koszul cohomology.

PROPOSITION 6.23. *Let A be a preprojective algebra of type E_8 .*

If $\text{char}(\mathbb{F}) = 2$, then $\text{HK}_{hi}^\bullet(A) = \text{HK}^\bullet(A)$.

If $\text{char}(\mathbb{F}) = 3$, then

$$\text{HK}_{hi}^0(A) = \text{HK}^0(A)_{28} \text{ and the } \pi_i \text{ for } i \in Q_0 \text{ form a basis}$$

$$\text{HK}_{hi}^1(A) = \text{span} \{[\bar{\rho}_5], [\bar{\rho}_{17}]\}$$

$$\text{HK}_{hi}^2(A) = \text{HK}^2(A)_0 \text{ and the } [\bar{h}_i] \text{ for } i \in Q_0 \text{ form a basis}$$

$$\text{HK}_{hi}^p(A) = 0 \text{ if } p \neq 0 \text{ and } p \neq 2.$$

If $\text{char}(\mathbb{F}) \notin \{2, 3\}$, then

$$\text{HK}_{hi}^0(A) = \text{HK}^0(A)_{28} \text{ has dimension 8 and is spanned by the } \pi_i \text{ for } i \in Q_0$$

$$\text{HK}_{hi}^2(A) = \text{HK}^2(A)_0 \text{ has dimension 8 and is spanned by the } [\bar{h}_i] \text{ for } i \in Q_0$$

$$\text{HK}_{hi}^p(A) = 0 \text{ if } p \neq 0 \text{ and } p \neq 2.$$

Higher Koszul homology follows from Theorem 4.4.

6.6. Comparison of Koszul and Hochschild (co)homology for preprojective algebras of type ADE

Let A be a preprojective algebra over a Dynkin graph of type ADE. Schofield constructed a minimal projective resolution $(P^\bullet, \partial^\bullet)$ of A as a bimodule over itself, that is periodic (of period at most 6), which was described in [20, 22]. Following Proposition 2.13, the embedding $H(\iota^*)_2$ sends the Hochschild cohomology class of an element in $\text{Ker}(\partial^3 \circ -)$ to its Koszul cohomology class, and the surjection $H(\iota)_2$ induces an isomorphism between $\text{HH}_2(A)$ and $\text{HK}_2(A)/\text{Im}(\text{id}_A \otimes \partial^3)$.

We first transport the maps $\partial^3 \circ -$ and $\text{id}_A \otimes \partial^3$ via the natural isomorphisms $A \otimes_{A^e} (Ae_j \otimes e_i A) \rightarrow e_i Ae_j$ that sends $\lambda \otimes (a \otimes b)$ to $b\lambda a$ and $\text{Hom}_{A^e}(Ae_i \otimes e_j A, A) \cong e_i Ae_j$ that sends f to $f(e_i \otimes e_j)$.

The associative non degenerate bilinear form on the selfinjective preprojective A can be defined as follows, see [20] and [46, Proposition 3.15]: let B be a basis of A consisting of homogeneous elements, that contains the idempotents e_i , $i \in Q_0$ and a basis $\{\pi_i; i \in Q_0\}$ of the socle of A , and such that each $v \in B$ belongs to $e_j Ae_i$ for some i, j in Q_0 . Then if $x \in Ae_i$, (y, x) is the coefficient of π_i in the expression of yx as a linear combination of elements in B . The Nakayama automorphism ν of A satisfies $(y, x) = (\nu(x), y)$ for all x, y in A , and induces a permutation of the indices, the Nakayama permutation $\bar{\nu}$, that is, a permutation of Q_0 such that $\text{top}(Ae_i) \cong \text{soc}(Ae_{\bar{\nu}(i)})$, characterised by $\nu(e_i) = e_{\bar{\nu}(i)}$.

Let \hat{B} be the dual basis of B with respect to the non degenerate form $(-, -)$, so that $(\hat{w}, v) = \delta_{vw}$ for all v, w in B . In particular, if $v \in Be_i$, the coefficient of π_i in $\hat{w}v$ is 1. Note that $v \in e_j Be_i$ if and only if $\hat{v} \in e_{\bar{\nu}(i)} \hat{B} e_j$.

Then the maps $\partial^3 \circ -$ and $\text{id}_A \otimes_{A^e} \partial^3$ become respectively

$$\delta^3: \bigoplus_{i \in Q_0} e_i Ae_i \rightarrow \bigoplus_{i \in Q_0} e_i Ae_{\bar{\nu}(i)} \text{ defined by } y \mapsto \sum_{x \in B} \hat{x} y x$$

$$\text{and } \delta_3: \bigoplus_{i \in Q_0} e_i Ae_{\bar{\nu}(i)} \rightarrow \bigoplus_{i \in Q_0} e_i Ae_i \text{ defined by } y \mapsto \sum_{x \in B} x y \hat{x}.$$

It then follows as in [24, Proposition 3.2.25] that $\text{Im } \delta_3$ is the span of the $\delta_3(e_i)$ such that $\bar{\nu}(i) = i$, and that for such an i we have $\delta_3(e_i) = \sum_{j \in Q_0, \bar{\nu}(j)=i} \text{tr}(\nu|_{e_j Ae_i}) \pi_j$. Moreover, the matrix whose coefficients are the $\text{tr}(\nu|_{e_j Ae_i})$ is either easy to compute or given in [22] for types D and E_7 . It is also known from [24] that the set of elements of weight 0 in $\text{Ker } \delta^3$ identifies with the kernel of the Cartan matrix of A . Moreover, for any element of positive weight $a \in A$,

we have $\delta^3(a) = 0$. Therefore $\mathrm{HH}^2(A)$ is obtained by taking all the elements of positive weight in $\mathrm{HK}^2(A)$ and adding the kernel of the Cartan matrix.

We shall use this as well as the dimensions of the Hochschild and Koszul (co)homology spaces to compare $\mathrm{HH}^2(A)$ with $\mathrm{HK}^2(A)$ and $\mathrm{HH}_2(A)$ with $\mathrm{HK}_2(A)$ in each case. In particular, the injection $\mathrm{HH}^2(A) \rightarrow \mathrm{HK}^2(A)$ is not surjective except in type E_8 with $\mathrm{char}(\mathbb{F}) = 2$.

6.6.1. Comparison of the second Koszul and Hochschild cohomology groups In type A, the space $\mathrm{HH}^2(A)$ was completely described by Erdmann and Snashall in [18], and they proved that $\dim \mathrm{HH}^2(A) = n - m_A - 1$ and gave a basis $\{\tilde{h}_i; 0 \leq i \leq n - m_A - 2\}$ with $\tilde{h}_i = \bar{h}_i + \bar{h}_{n-1-i}$. The morphism of complexes ι_2^* sends \tilde{h}_i to $h_i + h_{n-1-i}$, and this describes the injection $\mathrm{HH}^2(A) \rightarrow \mathrm{HK}^2(A)$.

In type D, if $\mathrm{char}(\mathbb{F}) \neq 2$ and n is even, there is nothing to do since $\mathrm{HH}^2(A) = 0$. If $\mathrm{char}(\mathbb{F}) \neq 2$ and n is odd, then $\dim \mathrm{HH}^2(A) = 1$, the basis given in [22] for $\mathrm{HH}^2(\Lambda_{\mathbb{C}})$ also gives a basis of $\mathrm{HH}^2(A)$, and it is the cohomology class of the map ψ_0 defined by $\psi_0(\sigma_0) = e_0$ and $\psi_0(\sigma_1) = -e_1$. The embedding $\mathrm{HH}^2(A) \rightarrow \mathrm{HK}^2(A)$ is therefore given by $\psi_0 \mapsto h_0 - h_1$.

Now assume that $\mathrm{char}(\mathbb{F}) = 2$. Then $\dim \mathrm{HH}^2(A) = n + m_D - 2$. As we explained above, a basis of $\mathrm{HH}^2(A)$ may be obtained from a basis of the set of elements of positive coefficient weight in $\mathrm{HK}^2(A)$ to which we add elements obtained by determining a basis of the kernel of the Cartan matrix of A . It follows that a basis of $\mathrm{HH}^2(A)$ is given by

$$\begin{aligned} &\{\bar{\gamma}_\ell; 1 \leq \ell \leq m\} \cup \{\bar{\varphi}_i; 2 \leq i \leq n - 1\} \quad \text{if } n \text{ is even} \\ &\{\bar{\gamma}_\ell; 1 \leq \ell \leq m\} \cup \{\bar{\psi}_0\} \cup \{\psi_i; 3 \leq i \leq n - 1\} \quad \text{if } n \text{ is odd} \end{aligned}$$

where $\psi_{2p+1}(\sigma_{2p+1}) = e_{2p+1}$, $\psi_{2p+2}(\sigma_{2p+2}) = e_{2p+2}$ and $\psi_{2p+2}(\sigma_2) = e_2$, $\varphi_{2p}(\sigma_{2p}) = e_{2p}$, $\varphi_{2p+1}(\sigma_{2p+3}) = e_{2p+3}$ and $\varphi_{2p+1}(\sigma_3) = e_3$, for $p \geq 1$. Therefore, the embedding $\mathrm{HH}^2(A) \rightarrow \mathrm{HK}^2(A)$ fixes the $\bar{\gamma}_\ell$ and sends $\bar{\varphi}_{2p}$ to \bar{h}_{2p} and $\bar{\varphi}_{2p+3}$ to $\bar{h}_3 + \bar{h}_{2p+3}$ when n is even, and $\bar{\psi}_0$ to $\bar{h}_0 + \bar{h}_1$, $\bar{\psi}_{2p+1}$ to \bar{h}_{2p+1} and $\bar{\psi}_{2p+2}$ to $\bar{h}_2 + \bar{h}_{2p+2}$ when n is odd.

In type E_6 , the Cartan matrix is equivalent, through row operations, to

$$\begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \end{pmatrix}$$

Let $\varphi_0, \varphi_1, \varphi_2$ and φ_3 be the maps in $\mathrm{Hom}_{k^e}(R, A)$ defined by $\varphi_0(\sigma_0) = e_0$, $\varphi_1(\sigma_1) = e_1$, $\varphi_1(\sigma_5) = -e_5$, $\varphi_2(\sigma_2) = e_2$, $\varphi_2(\sigma_4) = -e_4$ and $\varphi_3(\sigma_3) = e_3$. Then $\mathrm{HH}^2(A)$ is the subspace of $\mathrm{HK}^2(A)$ spanned by

$$\begin{aligned} &\bar{\varphi}_1, \bar{\varphi}_2 \text{ if } \mathrm{char}(\mathbb{F}) \notin \{2, 3\} \\ &\bar{\varphi}_0, \bar{\varphi}_1, \bar{\varphi}_2, \bar{\varphi}_3, \bar{\gamma}_4 \text{ if } \mathrm{char}(\mathbb{F}) = 2 \\ &\bar{\varphi}_1, \bar{\varphi}_2, \bar{\gamma}_6 \text{ if } \mathrm{char}(\mathbb{F}) = 3. \end{aligned}$$

In type E_7 , if $\mathrm{char}(\mathbb{F}) \neq 2$, we have $\dim \mathrm{HH}^2(A) = \dim \mathrm{HK}^2(A)_{>0}$, so that $\mathrm{HH}^2(A)$ is precisely the subspace of $\mathrm{HK}^2(A)$ of elements of positive weight (which is zero unless $\mathrm{char}(\mathbb{F}) \in \{2, 3\}$). If $\mathrm{char}(\mathbb{F}) = 2$, then the Cartan matrix of A is equivalent, through row operations, to the matrix $\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}$ so that $\mathrm{HH}^2(A)$ is the subspace of $\mathrm{HK}^2(A)$ spanned by $\bar{\gamma}_4, \bar{\gamma}_8, \bar{\gamma}_{16}, \bar{h}_1, \bar{h}_2, \bar{h}_3, \bar{h}_5, \bar{h}_0 + \bar{h}_4$ and $\bar{h}_0 + \bar{h}_6$.

In type E_8 , we only need to look at dimensions. Indeed, if $\mathrm{char}(\mathbb{F}) = 2$, then $\dim \mathrm{HH}^2(A) = \dim \mathrm{HK}^2(A)$ so that $\mathrm{HH}^2(A) \cong \mathrm{HK}^2(A)$, and if $\mathrm{char}(\mathbb{F}) \neq 2$ then $\dim \mathrm{HH}^2(A) = \dim \mathrm{HK}^2(A)_{>0}$ so that $\mathrm{HH}^2(A)$ is precisely the subspace of $\mathrm{HK}^2(A)$ of elements of positive weight (which is zero unless $\mathrm{char}(\mathbb{F}) \in \{2, 3, 5\}$).

6.6.2. Comparison of the second Koszul and Hochschild homology groups In type A, if n is even or if n is odd and $\mathrm{char}(\mathbb{F})$ divides $(m_A + 1)$, we have seen that $\dim \mathrm{HK}_2(A) = \dim \mathrm{HH}_2(A)$ and therefore $\mathrm{HK}_2(A) \cong \mathrm{HH}_2(A)$. Now assume that n is odd and $\mathrm{char}(\mathbb{F}) \nmid (m_A +$

1). Then $\dim \text{HK}_2(A) = \dim \text{HH}_2(A) + 1$. We must determine the image of the map δ_3 given in Subsection 6.6.

The Nakayama permutation $\bar{\nu}$ has precisely one fixed point, which is m_A . The matrix of ν restricted to $e_{m_A} A e_{m_A}$ is the identity matrix I_{m_A+1} . Consequently, $\delta_3(e_{m_A}) = (m_A + 1)\pi_{m_A}$ spans the image of δ_3 . Via the isomorphism $\bigoplus_{i \in Q_0} e_i A e_i \cong A \otimes_{k^e} R$, π_{m_A} corresponds to \check{z}_{m_A} so that we have $\text{HH}_2(A) \cong \text{HK}_2(A) / \text{span} \{ \check{z}_{m_A} \}$.

In type D, if n is odd and $\text{char}(\mathbb{F}) = 2$, then we know that $\dim \text{HK}_2(A) = \dim \text{HH}_2(A)$ and therefore $\text{HK}_2(A) \cong \text{HH}_2(A)$. In the other cases, we need to determine the image of the map δ_3 . We should note that the Nakayama automorphism in [22] differs from ν by composition with the inner automorphism given by the invertible element $u = -e_0 + \sum_{i=1}^{n-1} (-1)^i e_i$ (after changing the labelling of vertices and arrows so that they are the same as ours).

If n is odd and $\text{char}(\mathbb{F}) \neq 2$, the fixed points of the Nakayama permutation are the integers i with $2 \leq i \leq n - 1$. The matrix $(n - 2) \times (n - 2)$ matrix H^ν whose (i, j) -coefficient is the trace of ν restricted to $e_j A e_i$ was given in [22, paragraph 11.2.3]; we need only change the signs of coefficients (i, j) with i or j (but not both) equal to 0 or odd, so that $\text{tr}(\nu|_{e_j A e_i}) = 2$ if i and j are even and is equal to 0 otherwise. Therefore for all i fixed by $\bar{\nu}$, we have $\delta_3(e_i) = \sum_{p=1}^{m_D+1} 2\pi_{2p}$. Using the isomorphism $\bigoplus_{i \in Q_0} e_i A e_i \cong A \otimes_{k^e} R$, we obtain $\text{HH}_2(A) \cong \text{HK}_2(A) / \text{span} \left\{ \sum_{p=1}^{m_D+1} \check{\pi}_{2p} \right\}$.

If n is even, all the integers i with $0 \leq i \leq n - 1$ are fixed points of $\bar{\nu}$. The matrix H^ν was given in [22, paragraph 11.2.2] and the same adaptations as in the case n odd gives the $n \times n$ symmetric matrix

$$H^\nu = \begin{pmatrix} m+1 & -m & 0 & 1 & 0 & 1 & \dots & 0 & 1 \\ -m & m+1 & 0 & 1 & 0 & 1 & \dots & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & 2 & 0 & 2 & \dots & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & 2 & 0 & 2 & \dots & 0 & 2 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & 2 & 0 & 2 & \dots & 0 & 2 \end{pmatrix} \text{ which is column equivalent to } \begin{pmatrix} m_D+1 & 1-n \\ -m_D & n-1 \\ 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

It follows that, if n is even,

$$\text{HH}_2(A) \cong \begin{cases} \text{HK}_2(A) / \text{span} \left\{ (m_D + 1)\check{\pi}_0 - m_D\check{\pi}_1 + \sum_{p=1}^{m_D} \check{\pi}_{2p+1} \right\} & \text{if } \text{char}(\mathbb{F}) | (n - 1) \\ \text{HK}_2(A) / \text{span} \left\{ \check{\pi}_0 - \check{\pi}_1; \check{\pi}_0 + \sum_{p=1}^{m_D} \check{\pi}_{2p+1} \right\} & \text{if } \text{char}(\mathbb{F}) \nmid (n - 1) \end{cases}$$

In types E_6 and E_8 , since they have the same dimensions, the homology spaces $\text{HH}_2(A)$ and $\text{HK}_2(A)$ are isomorphic.

Finally, in type E_7 , if $\text{char}(\mathbb{F}) = 3$, then $\text{HH}_2(A) \cong \text{HK}_2(A)$. Now assume that $\text{char}(\mathbb{F}) \neq 3$. The matrix H^ν was given in [22]; here again, the Nakayama automorphism in [22] differs from ν by composing with the inner automorphism associated with the element $-e_0 + e_1 - e_2 + e_3 + e_4 + e_5 + e_6$ (after change of labelling) and therefore the non zero rows of the matrix H^ν are those corresponding to vertices 0, 4 and 6 and are all equal to $(3 \ 0 \ 0 \ 0 \ 3 \ 0 \ 3)$. It follows that $\text{HH}_2(A) \cong \text{HK}_2(A) / \text{span} \{ \check{\pi}_0 + \check{\pi}_4 + \check{\pi}_6 \}$.

6.7. A minimal complete list of cohomological invariants

THEOREM 6.24. *Let A be the preprojective algebra of a Dynkin graph Δ over \mathbb{F} . Assume that A has type either A_n with $n \geq 3$, or D_n with $n \geq 4$, or E_n with $n = 6, 7, 8$. Let A' be a preprojective algebra of type ADE, where the integer n' concerning A' is subjected to the same assumptions. Denote by (d_p) the equality $\dim \text{HK}_{hi}^p(A) = \dim \text{HK}_{hi}^p(A')$. If (d_p) holds for $p = 0, 1$ and 2, then $n = n'$, and A and A' have the same type. The conclusion of this implication does not hold if (d_2) is removed from the assumption.*

Proof. We apply the results contained in Propositions 6.6, 6.11, 6.15, 6.19 and 6.23. The implication is a consequence of the following items.

- (1) Assume that A and A' have types A or D. If $\text{char } \mathbb{F} \neq 2$, then $n = n'$ by (d_2) , and A and A' have the same type by (d_0) . If $\text{char } \mathbb{F} = 2$ and A and A' both have type A, then $n = n'$ by (d_2) . If $\text{char } \mathbb{F} = 2$, A is of type A_n and A' is of type $D_{n'}$, then the sum of (d_0) and (d_1) shows that $n = 2n' + m'_D - 2$ or -3 which contradicts (d_2) : $n = n' + m'_D$. If $\text{char } \mathbb{F} = 2$ and if A and A' both have type D, then $n = n'$ by (d_1) .
- (2) If A and A' have type E, then $n = n'$ by (d_0) .
- (3) If A is of type A_n and A' of type E_6 , then (d_2) implies either that $n = 6$ if $\text{char } \mathbb{F} \neq 2, 3$ or that $n = 7$ if $\text{char } \mathbb{F} = 2, 3$, but each case is excluded by (d_0) . Similarly when A' is of type E_7 and E_8 .
- (4) If A is of type D_n and A' is of type E_6 , then (d_2) implies one of the three following cases: $n = 6$ if $\text{char } \mathbb{F} \neq 2, 3$, $n = 7$ if $\text{char } \mathbb{F} = 3$ or $n + m_D = 7$ if $\text{char } \mathbb{F} = 2$, but each case is excluded by (d_0) . Similarly when A' is of type E_7 and E_8 (we also use (d_1) for E_8).

Let us show that we cannot remove assumption (d_2) . It is clear if $\text{char } \mathbb{F} \neq 2$ because when A is of type A_3 and A' is of type A_5 , both (d_0) and (d_1) hold. If $\text{char } \mathbb{F} = 2$, we check that when A is of type A_9 and A' is of type E_6 then (d_0) and (d_1) hold. \square

It is obvious that $\text{HK}_{hi}^\bullet(A) \cong \mathbb{F}$ if A is of type A_1 and it is easy to check that $\text{HK}_{hi}^\bullet(A) = 0$ when A is of type A_2 , therefore we have obtained a minimal complete list of cohomological invariants for all the ADE preprojective algebras.

Another direct application of our computations is the following. If A is as in Theorem 6.24 and if $\text{char } \mathbb{F} \neq 2$, the product of the algebra $\text{HK}_{hi}^\bullet(A)$ is identically zero. If $\text{char } \mathbb{F} = 2$, $\text{HK}_{hi}^\bullet(A) = \text{HK}^\bullet(A)$ is a unital algebra whose product is fully described in our results.

REMARK 6.25. In the one vertex case, the higher Koszul homology and cohomology play an essential role in a specific formulation of a Koszul Poincaré Lemma (a Koszul Poincaré duality), see Conjectures 6.5 and 7.2 in [7]. For this reason, we have formulated Theorem 6.24 in terms of the higher Koszul cohomology. An analogous statement with Koszul cohomology is also true and follows in the same way from our results, but in this case the minimality depends on the characteristic.

References

1. D. BAER, W. GEIGLE and H. LENZING, The preprojective algebra of a tame hereditary Artin algebra, *Comm. Algebra* 15 (1987), 425-457.
2. A. BEILINSON, V. GINZBURG and W. SOERGEL, Koszul duality patterns in representation theory, *J. Amer. Math. Soc.* 9 (1996), 473-527.
3. J. BELL and J. J. ZHANG, An isomorphism lemma for graded rings, *Proc. Amer. Math. Soc.* 145 (2017), 989-994.
4. D. J. BENSON, *Representations and cohomology I*, Cambridge Univ. Press, 1991.
5. R. BERGER, Dimension de Hochschild des algèbres graduées (French) [Hochschild dimension for graded algebras], *C. R. Math. Acad. Sci. Paris* 341 (2005), 597-600.
6. R. BERGER, Koszul calculus for N-homogeneous algebras, *J. Algebra* 519 (2019), 149-189.
7. R. BERGER, T. LAMBRE and A. SOLOTAR, Koszul calculus, *Glasg. Math. J.* 60 (2018), 361-399.
8. J. BIALKOWSKI, K. ERDMANN and A. SKOWROŃSKI, Deformed preprojective algebras of generalized Dynkin type Ln: classification and symmetry, *J. Algebra* 345 (2011), 150-170.
9. R. BOCKLANDT, Graded Calabi Yau algebras of dimension 3. *J. Pure Appl. Algebra* 212 (2008), 14-32.
10. S. BRENNER, M.C.R. BUTLER and A. KING, Periodic algebras which are almost Koszul. *Algebr. Represent. Theory* 5 (2002), 331-367.
11. K. S. BROWN, *Cohomology of Groups*, Graduate Texts in Mathematics 87, Springer, 1982.
12. C. CIBILIS, Rigidity of truncated quiver algebras. *Adv. Math.* 79 (1990), 18-42.
13. W. CRAWLEY-BOEVEY, DMV lectures on Representations of quivers, preprojective algebras and deformations of quotient singularities. Available at <https://www.math.uni-bielefeld.de/~wcrawley/dmvlecs.pdf> (1999).

14. W. CRAWLEY-BOEVEY, P. ETINGOF and V. GINZBURG, Noncommutative geometry and quiver algebras, *Adv. Math.* 209 (2007), 274-336.
15. W. CRAWLEY-BOEVEY and M.P. HOLLAND, Noncommutative deformations of Kleinian singularities, *Duke Math. Journal* 92 (1998), 605-635.
16. W. CRAWLEY-BOEVEY and M. VAN DEN BERGH, Absolutely indecomposable representations and Kac-Moody Lie algebras. With an appendix by Hiraku Nakajima. *Invent. Math.* 155 (2004), 537-559.
17. V. DLAB and C. M. RINGEL, The preprojective algebra of a modulated graph, in Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), 216-231, *Lecture Notes in Math.*, 832, Springer, Berlin-New York, 1980.
18. K. ERDMANN and N. SNASHALL, Hochschild cohomology of preprojective algebras, I, *J. Algebra* 205 (1998), 391-412.
19. K. ERDMANN and N. SNASHALL, Hochschild cohomology of preprojective algebras, II, *J. Algebra* 205 (1998), 413-434.
20. K. ERDMANN and N. SNASHALL, Preprojective algebras of Dynkin type, periodicity and the second Hochschild cohomology, *CMS Conf. Proc.* 24 (1998).
21. P. ETINGOF and C-H. EU, Hochschild and cyclic homology of preprojective algebras of ADE quivers, *Mosc. Math. J.* 7 (2007), 601-612.
22. C-H. EU, The product in the Hochschild cohomology ring of preprojective algebras of Dynkin quivers, *J. Algebra* 320 (2008), 1477-1530.
23. C-H. EU, The calculus structure of the Hochschild homology/cohomology of preprojective algebras of Dynkin quivers, *J. Pure Appl. Algebra* 214 (2010), 28-46.
24. C-H. EU and T. SCHEDLER, Calabi-Yau Frobenius algebras, *J. Algebra* 321 (2009), 774-815.
25. J. GADDIS, Isomorphisms of graded path algebras, [arXiv:1712.01650](https://arxiv.org/abs/1712.01650).
26. C. GEISS, B. LECLERC and J. SCHRÖER, Preprojective algebras and cluster algebras, in Trends in representation theory of algebras and related topics, 253-283, *EMS Ser. Congr. Rep.*, Eur. Math. Soc., Zürich, 2008.
27. I. M. GELFAND and V. A. PONOMAREV, Model algebras and representations of graphs (Russian), *Funktsional. Anal. i Prilozhen* 13 (1979), 1-12.
28. V. GINZBURG, Calabi-Yau algebras, [arXiv:math.AG/0612139v3](https://arxiv.org/abs/math/0612139v3).
29. J. GRANT and O. IYAMA, Higher preprojective algebras, Koszul algebras, and superpotentials [arXiv:1902.07878v2](https://arxiv.org/abs/1902.07878v2).
30. E. L. GREEN, Introduction to Koszul algebras, in Representation theory and algebraic geometry (Waltham MA, 1995), 45-62, *LMS Lect. Notes Ser.* 238, Cambridge Univ. Press, 1997.
31. A. HATCHER, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
32. T. LAMBRE, Dualité de Van den Bergh et structure de Batalin-Vilkoviskii sur les algèbres de Calabi-Yau, *J. Noncommut. Geom.* 4 (2010), 441-457.
33. R. MARTÍNEZ-VILLA, Applications to Koszul algebras: the preprojective algebra, in Representation theory of algebras (Cocoyoc, 1994), 487-504, *CMS Conf. Proc.* 18, Amer. Math. Soc., Providence RI, 1996.
34. R. MARTÍNEZ-VILLA, Introduction to Koszul Algebras, *Revista Union Matematica Argentina* 48 (2007), 67-95
35. A. POLISHCHUK and L. POSITSIELSKI, *Quadratic algebras*, University Lecture Series 37, AMS, 2005.
36. S. PRIDDY Koszul resolutions, *Trans. Amer. Math. Soc.* 152 (1970), 39-60.
37. M.L. REYES and D. ROGALSKI A twisted Calabi-Yau toolkit, [arXiv:1807.10249](https://arxiv.org/abs/1807.10249).
38. P.H.C. SHAW, Generalisations of preprojective algebras. *PhD thesis of the University of Leeds, 2005*, available at <https://www.math.uni-bielefeld.de/~wrcrawley/shaw-thesis.pdf>.
39. D. TAMARKIN and B. TSYGAN, The ring of differential operators on forms in noncommutative calculus, *Proc. Sympos. Pure Math.* 73, Amer. Math. Soc. (2005), 105-131.
40. M. VAN DEN BERGH, Noncommutative homology of some three-dimensional quantum spaces, *K-Theory* 8 (1994), 213-230.
41. M. VAN DEN BERGH, A relation between Hochschild homology and cohomology for Gorenstein rings, *Proc. Amer. Math. Soc.* 126 (1998), 1345-1348; erratum *ibid.* 130 (2002), 2809-2810.
42. M. VAN DEN BERGH, Double Poisson algebras, *Trans. Amer. Math. Soc.* 360 (2008), 5711-5769.
43. M. VAN DEN BERGH, Calabi-Yau algebras and superpotentials, *Selecta Math. (N.S.)* 21 (2015), 555-603.
44. C. A. WEIBEL, *An introduction to homological algebra*, Cambridge Univ. Press, 1994.
45. A. YEKUTIELI, *Derived Categories*, to appear in Cambridge Univ. Press, [arXiv:1610.09640v4](https://arxiv.org/abs/1610.09640v4).
46. A. ZIMMERMANN, On the use of Külshammer type invariants in representation theory, *Bull. Iran. Math. Soc.* 37 (2011), 291-341.

Roland Berger
 Univ Lyon, UJM-Saint-Étienne, CNRS
 UMR 5208, Institut Camille Jordan,
 F-42023, Saint-Étienne
 France

Roland.Berger@univ-st-etienne.fr

Rachel Taillefer
 Université Clermont Auvergne, CNRS,
 LMBP, F-63000 Clermont-Ferrand
 France

Rachel.Taillefer@uca.fr