



HAL
open science

On subvarieties of singular quotients of bounded domains

Benoit Cadorel, Simone Diverio, Henri Guenancia

► **To cite this version:**

Benoit Cadorel, Simone Diverio, Henri Guenancia. On subvarieties of singular quotients of bounded domains. 2020. hal-02134317v1

HAL Id: hal-02134317

<https://hal.science/hal-02134317v1>

Preprint submitted on 18 Nov 2020 (v1), last revised 21 Jul 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON SUBVARIETIES OF SINGULAR QUOTIENTS OF BOUNDED DOMAINS

BENOÎT CADOREL, SIMONE DIVERIO, AND HENRI GUENANCIA

ABSTRACT. Let X be a quotient of a bounded domain in \mathbb{C}^n . Under suitable assumptions, we prove that every subvariety of X not included in the branch locus of the quotient map is of log general type in some orbifold sense. This generalizes a recent result by Boucksom and Diverio, which treated the case of compact, étale quotients.

Finally, in the case where X is compact, we give a sufficient condition under which there exists a proper analytic subset of X containing all entire curves and all subvarieties not of general type (meant this time in the usual sense as opposed to the orbifold sense).

1. INTRODUCTION

Let X be a quotient of a bounded domain $\Omega \subset \mathbb{C}^n$ by some discrete automorphism group $\Gamma \subset \text{Aut}(\Omega)$. A lot of recent work has been devoted to the research of complex hyperbolicity properties of these quotients X , *i.e.* of restrictions on the geometry of entire curves in X , or on the type of its subvarieties. These quotients provide indeed basic examples to test the general conjectures in complex hyperbolicity, in particular the *Green-Griffiths-Lang conjecture*:

Conjecture 1.1 (Green, Griffiths [GG80], Lang [Lan86]). *Let M be a complex projective manifold of general type. Then there exists a proper algebraic subset $\text{Exc}(M) \subsetneq M$ containing all the images of non constant holomorphic maps $\mathbb{C} \rightarrow M$, and all the subvarieties of M which are not of general type.*

Notably, in the case where $X = \Gamma \backslash \Omega$ is a quotient by a cocompact group acting *freely* and properly discontinuously on Ω , X will be a complex projective manifold; it is an easy application of Liouville theorem that X cannot contain any entire curve. The first statement of Conjecture 1.1 is trivial in this case while the second statement was recently obtained by Boucksom and the second author [BD18] where they show that all subvarieties of X are of general type.

When the action of Γ is not more free or cocompact, X itself may already not be of general type (cf. *e.g.* Section 3.3) and it may not enjoy such nice hyperbolicity properties. However, there is a general philosophy that statements true in the smooth compact case should continue to hold when dealing with the correct *orbifold* or *logarithmic* structures in the singular or non-compact case. For example, our methods yield the following archetypical result.

Model Theorem. *Assume that \overline{X} is a smooth projective manifold, and that $D \subset \overline{X}$ is a reduced divisor such that $X = \overline{X} \setminus D$ is uniformized by a bounded domain $\Omega \subset \mathbb{C}^n$. Then the pair (\overline{X}, D) is of log general type, *i.e.* $K_{\overline{X}} + D$ is big.*

This theorem will appear as a particular case of several results (for instance, Theorem A or Theorem B below) whose main point of focus will be the analysis of the defect of hyperbolicity of such quotients in the spirit of the philosophy above.

Before continuing, let us point out two remarkable enough instances where our Model Theorem applies.

Date: January 10, 2020.

2010 Mathematics Subject Classification. Primary: 32J25; Secondary: 32Q45, 32Q30, 14E20, 14E22.

Key words and phrases. Logarithmic variants of Lang's conjecture, quotients of bounded domains, varieties and orbifolds of log-general type, Bergman metric, L^2 -estimates for $\bar{\partial}$ -equation.

Example 1.2. By a classical result of Griffiths [Gri71], in any projective n -dimensional manifold \overline{X} , there exist nonempty Zariski open subsets $X = \overline{X} \setminus D$ which are uniformized by a bounded domain (moreover, of holomorphy) in \mathbb{C}^n . Even more, any point of any smooth, irreducible, quasi-projective variety over the complex numbers has a Zariski open neighborhood with this property. Thus, in principle, *every* smooth projective manifold falls in the scope of the theorem above. Of course, the main interest of our result resides in situations where we have a precise description of the boundary divisor $D \subset \overline{X}$.

Let us now give a perhaps more concrete example.

Example 1.3. Fix integers $g \geq 2$, and $n \geq 0$, and consider the Teichmüller space $\mathcal{T}_{g,n}$ of compact Riemann surfaces of genus g with n marked points. It is well-known that the Bers embedding realizes $\mathcal{T}_{g,n}$ as a (contractible) bounded domain $\mathcal{B}_{g,n} \subset \mathbb{C}^{3g-3+n}$, which we call the *Bers domain*, and actually provides an isometry between the Kobayashi distance of $\mathcal{B}_{g,n}$ and the Teichmüller metric on $\mathcal{T}_{g,n}$. This domain is very far from being symmetric since it is indeed even not homogeneous: its isometry group is the Mapping Class Group $\text{MCG}_{g,n}$, whose action is properly discontinuously and whose orbits are precisely the equivalent complex structures, so that the quotient $\mathcal{T}_{g,n}/\text{MCG}_{g,n}$ is exactly the moduli space $\mathcal{M}_{g,n}$.

Now, the action of $\text{MCG}_{g,n}$ is not free in general (still, with finite stabilizers), but it is easy to see that it is if $n > 2g + 2$, so that in this case $\mathcal{T}_{g,n}$ is the universal cover of the smooth quasi-projective manifold $\mathcal{M}_{g,n}$. So therefore, in this situation our Model Theorem applies to the Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$ and gives a relatively elementary and direct proof that it is of log-general type with respect to the boundary divisor.

This (and more indeed: the canonical bundle plus the boundary divisor is not only big but even ample) was of course previously known, but to the best of our knowledge the proof relies at least on a precise description of the Picard group of $\overline{\mathcal{M}}_{g,n}$, together with the Cornalba–Harris ampleness criterion, which in turn employs hard GIT. Unfortunately, for the time being, we don't know if it is actually possible with our method to prove the ampleness of the logarithmic canonical bundle.

All this can be obtained with our Model Theorem, which is actually some sort of oversimplified statement with respect to our Theorems A, B, and C. See Examples 1.5, 1.7 and Remark 1.6 below for further comments on this.

Remark 1.4. We would like to further observe that the Weil–Petersson metric is mapping class group invariant and thus descends to $\mathcal{M}_{g,n}$. It has indeed negative sectional curvature. It is known that its behaviour near the boundary gives that its Riemannian sectional curvature has as infimum negative infinity and as supremum zero. But, on the other hand, its holomorphic sectional, Ricci and scalar curvatures are all bounded above by genus-dependent negative constants. Therefore, we can also obtain the log-general type property of $\mathcal{M}_{g,n}$ by a nice application of Guenancia's theorem [Gue18, Theorem B].

1.1. Main results. The general set-up in which our results are stated involves a variety \overline{X} containing a Zariski open subset X , uniformized by a variety Ω on which the curvature of the *Bergman metric* on the canonical bundle K_Ω is positive definite at a generic point. We will call such manifolds Ω *weakly Bergman manifolds*: this class of manifolds contain bounded domains, and more generally complex manifolds of *bounded type* in the sense of [BD18], i.e. the manifolds admitting a bounded, strictly psh function.

- A CRITERION FOR PAIRS TO BE OF LOG GENERAL TYPE

Consider a compact Kähler manifold \overline{X} , and let D be a reduced divisor on \overline{X} such that $X = \overline{X} \setminus D$ is uniformized by a *weakly Bergman manifold* via a covering map $p : \Omega \rightarrow X$. In this situation, we can then endow each component D_i of D with a multiplicity $m_i \in \mathbb{N}^* \cup \{\infty\}$, representing in some sense the order of ramification of p around a general point of D_i . We will

call $\Delta_{\overline{X}} = \sum_i (1 - \frac{1}{m_i}) D_i$ the *covering divisor* associated with our data: the pair $(\overline{X}, \Delta_{\overline{X}})$ is then an orbifold pair in the sense of Campana [Cam04].

Then, our first main result can be stated as follows, cf. Theorem 2.1.

Theorem A. *Let \overline{X} be a compact Kähler manifold endowed with a reduced divisor D such that $\overline{X} \setminus D$ is uniformized by a weakly Bergman manifold. Let $\Delta_{\overline{X}}$ be the associated covering divisor on \overline{X} . Then, the \mathbb{Q} -line bundle $K_{\overline{X}} + \Delta_{\overline{X}}$ is big.*

Example 1.5. If we wanted to try to refine the situation of Example 1.3 using the more precise Theorem A, we would obtain in this particular setting the same result, since the monodromy around the boundary divisor is infinite cyclic, with generator a Dehn twist, so that all the m_i 's are infinite here.

• QUOTIENTS OF MANIFOLDS OF BOUNDED TYPE

Let Ω be a manifold of bounded type in the sense of [BD18]. We want to study the situation where X is a quotient of Ω by a discrete subgroup $\Gamma \subset \text{Aut}(\Omega)$ acting properly discontinuously.

In general, X is neither smooth nor compact. Given $V \subset X$ a subvariety, we explained above that one cannot expect V to be of (log) general type in full generality. However, since immersed subvarieties of Ω are still *weakly Bergman* by [BD18], it is then possible to apply Theorem A to the situation where \overline{X} is replaced by a compactification \overline{V} of V , and Ω by the fiber product $V \times_X \Omega$, which is still a manifold of bounded type.

The theorem below is an application of the idea above: it provides a particular setting where we can obtain that the *orbifold pair* naturally associated to a modification of V is of general type, cf. Theorem 3.1.

Theorem B. *Let \overline{X} be a normal, compact complex space admitting a Kähler resolution. Assume that it admits a Zariski open subset $X = \overline{X} \setminus D$ which is a quotient of a manifold Ω of bounded type and let $p: \Omega \rightarrow X$ be the quotient map.*

Let $V \subset \overline{X}$ be a closed subvariety such that $V \not\subset D \cup \text{Sing}(p)$, where $\text{Sing}(p)$ is the locus of singular values of p . Let $\pi: \widehat{V} \rightarrow V$ be any resolution of singularities of V .

Then \widehat{V} supports a natural covering divisor $\Delta_{\widehat{V}} = \sum_i (1 - \frac{1}{n_i}) F_i$ such that $K_{\widehat{V}} + \Delta_{\widehat{V}}$ is big. Moreover, $\Delta_{\widehat{V}}$ is supported over $D \cup \text{Sing}(p)$ via π .

We will actually prove a more general variant of this result, where we do not need \widehat{V} to be *smooth*, but merely \mathbb{Q} -factorial (see Theorem 3.1). This variant will be applicable to several different contexts: for example, if $X = \Gamma \backslash \Omega$ is a compact quotient and Γ has no fixed point in codimension 1, it will imply that K_X is *big* (which is not equivalent to X being of general type, see Section 3.2 (4) and Section 3.3 (1)).

Remark 1.6. Let us revisit again the situation of Example 1.3 in light of Theorem B: in principle this theorem should allow to treat the more general case with no conditions on n , since here nothing is required about the freeness of the action.

Given a resolution $\widehat{\mathcal{M}}_{g,n}$ of $\overline{\mathcal{M}}_{g,n}$, this theorem yields the existence of a natural orbifold structure $(\widehat{\mathcal{M}}_{g,n}, \Delta_{\widehat{\mathcal{M}}_{g,n}})$, which is of log-general type. Moreover, observe that the \mathbb{Q} -divisor $\Delta_{\widehat{\mathcal{M}}_{g,n}}$ is intrinsically completely described by the action of $\text{MCG}_{g,n}$ on $\mathcal{T}_{g,n}$.

• THE SINGULAR, COMPACT CASE.

In the case where Ω is a bounded domain admitting a cocompact lattice (possibly distinct from Γ), it is possible to give a refinement of Theorem B, in the setting where V is not necessarily compact, but *weakly pseudoconvex*. The next theorem formulates a positivity result for $K_{\widehat{V}} + \Delta_{\widehat{V}}$ in terms of the existence of a singular metric with positive curvature on this line bundle. Applying this result to $V = \mathbb{C}$ will later on allow us to obtain a hyperbolicity criterion for the manifold \overline{X} , cf. Theorem 5.1.

Theorem C. *Assume that Ω is a bounded domain admitting a cocompact lattice. Let $q : V \rightarrow \overline{X}$ be a generically immersive holomorphic map from a weakly pseudoconvex Kähler manifold V , such that $q(V) \not\subset D \cup \text{Sing}(p)$.*

Then, there exists a modification $\sigma : \widehat{V} \rightarrow V$ such that the \mathbb{Q} -divisor $K_{\widehat{V}} + \Delta_{\widehat{V}}$ admits a singular metric $g_{\widehat{V}}$ with non-negative curvature, positive definite at a general point of \widehat{V} , and with an explicit lower bound on this curvature in terms of the Bergman metric of Ω .

Applied to $V = \mathbb{C}$, this refinement can be used to provide a hyperbolicity result in the case where $X = \overline{X}$ is compact. We obtain the following partial generalization to the non-symmetric case of a previous work of the first author with Rousseau and Taji [CRT17], cf. Theorem 5.2.

Example 1.7. Getting back to the case of Example 1.3, but still in the general situation of arbitrary n , we see that Theorem C yields the following.

Let V be a weakly pseudoconvex Kähler manifold supporting a family of curves of genus g with n marked points, with no non-trivial automorphisms. Assume that this family has maximal variation. Then, we have a canonical generically immersive holomorphic map $f : V \rightarrow \mathcal{M}_{g,n}^{\text{reg}}$, and this ensures, by Theorem C, that V is of log-general type. This can be seen as a very special case of Viehweg's hyperbolicity conjecture.

Theorem D. *Assume that Ω is a bounded domain admitting a cocompact lattice. Then there exists a constant α_0 , depending only on Ω , such that the following holds.*

Let $X = \Gamma \backslash \Omega$ be a compact quotient and let $\widehat{X} \xrightarrow{\pi} X$ be a projective resolution of singularities. If for some $\alpha > \alpha_0$, the \mathbb{Q} -divisor

$$L_\alpha := \pi^*(K_X + \Delta_X) - \alpha \Delta_{\widehat{X}}$$

is effective, where Δ_X is the covering divisor associated to $p : \Omega \rightarrow X$, then

- (1) *any subvariety $W \subseteq X$ such that $W \not\subset \pi(\mathbb{B}(L_\alpha)) \cup \text{Sing}(p)$ is of general type.*
- (2) *any entire curve $f : \mathbb{C} \rightarrow X$ has its image included in $\pi(\mathbb{B}(L_\alpha)) \cup \text{Sing}(p)$.*

Here, $\mathbb{B}(L_\alpha) = \bigcap_p \text{Bs}(L_\alpha^{\otimes p})$ denotes the stable base locus of L_α and the intersection is taken over all positive integers p divisible enough so that $L^{\otimes p}$ is a genuine line bundle. Note that Lemma 2.10 guarantees the existence of projective resolutions for X provided that it fulfils the assumptions of Theorem D.

1.2. Further comparison to previous results.

- As already explained, the techniques of the papers are inspired by [BD18] where it is proved that any subvariety $V \subset X$ of a compact étale quotient $X = \Gamma \backslash \Omega$ of a manifold of bounded type is of general type. One of their key observations is that if $\widehat{V} \rightarrow V$ is a resolution of singularities, then one can construct a natural étale, Galois cover $Z \rightarrow \widehat{V}$ where Z is a Bergman manifold i.e. the Bergman kernel on K_Z is well-defined and has strictly positive curvature on a non-empty open set of Z ; this kernel descends to define a metric with the same properties on $K_{\widehat{V}}$, from which the bigness of $K_{\widehat{V}}$ follows.

When the action of Γ is not assumed to be free anymore, one can still get a Galois cover $Z \rightarrow \widehat{V}$ where the Bergman metric on K_Z has similar positivity properties as before, but that metric will *not* descend to a metric on $K_{\widehat{V}}$ anymore but rather on an adjoint bundle $K_{\widehat{V}} + \Delta_{\widehat{V}}$ for some suitable boundary divisor $\Delta_{\widehat{V}}$.

- In the case where Ω is a bounded symmetric domain, a great variety of viewpoints have been recently used to investigate the hyperbolicity properties of these quotients $X = \Gamma \backslash \Omega$. They can be studied by means of Hodge theory [Bru16b, Bru16a], Monge-Ampère equations and negative holomorphic sectional curvature [WY16, Gue18, DT19], or other metric methods [Rou16, Cad16, CRT17, Cad18]. Unfortunately, all these techniques rely to some extent on the precise curvature properties of the Bergman metric on a bounded symmetric domain, which totally break down if the domain is not symmetric. To our knowledge, the best thing that can be

said for a general bounded domain is that the holomorphic sectional curvature of its Bergman metric is bounded above by 2 [Kob59] (but it has no sign in general). It would be anyway interesting to understand if the greater symmetry of bounded domains admitting a cocompact lattice might allow one to infer something more precise about the holomorphic sectional curvature of the Bergman metric (see Section 5 where such a symmetry is exploited to obtain information on its Ricci curvature).

1.3. Outline of the proof.

Let us briefly describe the idea of the proof of Theorem A. Suppose that \bar{X} is a compact Kähler manifold, and that $X = \bar{X} \setminus D$ is a Zariski open subset uniformized by a manifold Ω which is *weakly Bergman*, i.e. on which the Bergman metric is defined at a generic point. We wish to find a \mathbb{Q} -divisor $\Delta_{\bar{X}}$ supported on D , such that $K_{\bar{X}} + \Delta_{\bar{X}}$ is big. The main idea is similar to the metric techniques employed in [CRT17]: we first construct a smooth metric g on K_X with positive definite curvature. Then, we control the divergence of the metric g on the boundary D , to show it extends as a singular metric with positive curvature on $K_{\bar{X}} + \Delta_{\bar{X}}$, for some suitable \mathbb{Q} -divisor $\Delta_{\bar{X}}$ supported on D . The conclusion then comes from a criterion of bigness due to Boucksom [Bou02].

In this situation, the metric g will come directly from the Bergman kernel on Ω , which descends to X to define a positively curved, singular metric g on K_X , with positive definite curvature at a generic point.

Finally, we have to control the divergence of g near the boundary D . To understand this divergence, we will use a geometric construction which is very convenient to determine the adequate orbifold multiplicities to put on the components of D , and which was applied by several authors to extend algebraic orbifold objects on resolutions of quotient singularities (see Tai [Tai82], Weissauer [Wei86]).

In the case where $V \subset \bar{X}$ is a subvariety of a compactification $\bar{X} = X \cup D$ of a quotient of a bounded domain $X = \Gamma \backslash \Omega$, we proceed with the same arguments, replacing essentially X by a resolution of singularities $\widehat{V} \rightarrow V$ of V , and Ω by $\widehat{V} \times_X \Omega$. One technical part of the proof of Theorem C is to bound from below the curvature of the Bergman metric on the open part $\widehat{V} \setminus \text{Supp}(\Delta_{\widehat{V}})$: if we assume that Ω is a bounded domain acted upon by a cocompact lattice, it is possible to use general comparison results between the Carathéodory and Bergman metrics, due to Hahn [Hah78]. Our general method will follow closely the L^2 technique employed in [BD18], which was in turn inspired by [CZ02]; however, it will be slightly more elaborated since we want to be able to deal with the case where V is no more compact, and thus non-necessarily complete Kähler (see Theorem 5.1).

In this situation, the orbifold multiplicities to put on $\Delta_{\widehat{V}}$ give a slightly refined version of what was done in [CRT17], where this technique of construction of an orbifold pair $(\widehat{V}, \Delta_{\widehat{V}})$ was also used to extend singular metrics as well as orbifold symmetric differentials. Our explicit description will allow us to compare the divisors $\Delta_{\widehat{V}}$ and $\Delta_{\widehat{X}}$ appearing in this setting, cf. Proposition 4.9. These comparison result will be of particular importance to prove the hyperbolicity criterion of Theorem D.

1.4. Organization of the paper.

- § 2. We give a definition of the *covering divisors* which will be used throughout the text. After recalling some useful information concerning the Bergman metric, and the existence of projective resolutions, we prove Theorem A.

- § 3. We apply Theorem A to the case of subvarieties of quotients of bounded domains. Theorem B appears as a particular case of Theorem 3.1, which is the main result of this section.

- § 4. Let $V \subset X = \Gamma \backslash \Omega$, and let $\widehat{X} \rightarrow X$ and $\widehat{V} \rightarrow V$ be adequate log resolutions. The main result of this section is the comparison result between $\Delta_{\widehat{V}}$ and $\Delta_{\widehat{X}}$ given by Proposition 4.9.

- §5. In the case where Ω is a bounded domain, we give a uniform bound from below for the curvature of the Bergman metric of subvarieties of Ω , cf. Lemma 5.7. We apply this estimate to derive a lower bound of a natural singular metric with positive curvature on $K_{\mathbb{P}^1} + \Delta_{\mathbb{P}^1}$ in a very general setting, cf. Theorem 5.1. Finally, we go back to the compact case and spell out a criterion for X to satisfy the Green-Griffiths-Lang conjecture, cf. Theorem D.

Acknowledgements. The authors would like to thank Sébastien Boucksom, Junyan Cao, Gabriele Mondello, Erwan Rousseau, Behrouz Taji, Stefano Trapani and Shengyuan Zhao for enlightening discussions about this paper.

B.C. is partially supported by the ANR Programme: Défi de tous les savoirs (DS10) 2015, “GRACK”, Project ID: ANR-15-CE40-0003.

S.D. is partially supported by the ANR Programme: Défi de tous les savoirs (DS10) 2015, “GRACK”, Project ID: ANR-15-CE40-0003ANR and by the ANR Programme: Défi de tous les savoirs (DS10) 2016, “FOLIAGE”, Project ID: ANR-16-CE40-0008.

H.G. is partially supported by the National Science Foundation through the NSF Grant DMS-1510214.

2. A CRITERION FOR A PAIR TO BE OF LOG GENERAL TYPE

2.1. Covering multiplicities. Let us begin the present section with a definition which, although perhaps not completely standard, is well adapted to our purposes.

Definition 2.1 (Covers). Let X, Y be two irreducible and reduced complex spaces of the same dimension and let $p: X \rightarrow Y$ be a surjective holomorphic map.

- One says that p is a *cover* if there exists a discrete subgroup $\Gamma \subset \text{Aut}(X)$ acting properly and discontinuously on X such that p is isomorphic to the quotient map $X \rightarrow \Gamma \backslash X$.
- The *singular locus* of p , $\text{Sing}(p)$, is defined to be the locus of singular values of p , i.e. the locus of points $y \in Y$ such that there exists $x \in p^{-1}(y)$ such that p is not a local biholomorphism around x .
- If p is étale, or equivalently if $\text{Sing}(p) = \emptyset$, one says that Y is *uniformized* by X .

Let \bar{X} be a n -dimensional connected normal complex space and let $X \subseteq \bar{X}$ be some non-empty analytic Zariski open subset of \bar{X} endowed with an étale cover $p: \tilde{X} \rightarrow X$. We denote by $D = \sum_{i \in I} D_i$ the union of the codimension one irreducible components of $\bar{X} \setminus X$.

Given a general point $x_i \in D_i$, one can choose an euclidean neighborhood \bar{U}_i of x_i in \bar{X} such that $U_i := \bar{U}_i \setminus D_i \simeq \mathbb{D}^* \times \mathbb{D}^{n-1}$. One denotes by \tilde{U}_i a connected component of $p^{-1}(U_i)$.

Definition 2.2 (Covering divisor). Let (\bar{X}, X, p) as above. The *covering multiplicity* $m_i = m_i(p) \in \mathbb{N}^* \cup \{\infty\}$ of the divisor D_i is defined to be the degree of the cover $p|_{\tilde{U}_i}: \tilde{U}_i \rightarrow U_i$ above. One then defines the *covering divisor* $\Delta = \Delta(p) := \sum_{i \in I} (1 - \frac{1}{m_i(p)}) D_i$.

If no ambiguity is possible, we will just write m_i and Δ .

Remark 2.3. The number m_i above is independent of the choice of the general point $x_i \in D_i$ and the neighborhood \bar{U}_i . Moreover, it is also clearly independent of the choice of the connected component of $p^{-1}(U_i)$, since the deck transformation group acts transitively on the various components.

2.2. The Bergman metric. Let X be a n -dimensional connected complex manifold and let \mathcal{H}_X be the Hilbert space of holomorphic sections $\sigma \in H^0(X, K_X)$ with finite L^2 norm

$$\|\sigma\|^2 := \int_X i^{n^2} \sigma \wedge \bar{\sigma}.$$

Assuming that $\mathcal{H}_X \neq \{0\}$, one can define a singular metric h_X on K_X as follows. Choose a Hilbertian basis $(e_i)_i$ for \mathcal{H} and let e be a local holomorphic frame for K_X . For each i , we have

$e_i = s_i e$, for some local holomorphic function s_i . Then, we can define

$$\|e\|_{h_X}^2 = \frac{1}{\sum_i |s_i|^2}.$$

It is easy to check that the Chern curvature current $i\Theta(h_X)$ is a closed, positive $(1, 1)$ -current. Moreover, the metric h_X is clearly invariant under the action of $\text{Aut}(X)$ on K_X .

Definition 2.4 (Bergman metric). Let X be a complex manifold such that $\mathcal{H}_X \neq \{0\}$.

The singular hermitian metric h_X on K_X defined above is called the *Bergman metric* on X . Its Chern curvature form $i\Theta(h_X)$ is a positive current.

Remark 2.5. Please notice that the terminology introduced above is not completely standard: usually the term Bergman metric is used for the Chern curvature form $i\Theta(h_X)$ (whenever defined), and what we call here Bergman metric is instead (a manifestation of, using the correspondence between volume forms and metrics on the canonical bundle) the Bergman kernel.

Nevertheless, in the highly non-smooth situations we consider here, what is usually referred to as Bergman metric will be in general merely a current. Such current is very far from being a genuine smooth metric on the tangent bundle, so that we prefer to reserve the term *metric* for its incarnation as a singular metric on the canonical bundle.

The Bergman metric h_X , along with its curvature current $i\Theta(h_X)$ are smooth on the analytic Zariski open subset of X corresponding to the complement of the base locus of the linear system of L^2 integrable sections of K_X .

Definition 2.6 ((weakly) Bergman manifolds). We introduce the following notions.

- (1) A complex manifold X is called *weakly Bergman* if its Bergman metric h_X is well-defined and if the curvature current $i\Theta(h_X)$ is smooth and positive definite on some euclidean open subset $\emptyset \neq U \subset X$.
- (2) A *Bergman manifold* is a weakly Bergman manifold X where the open set U above can be taken to be the whole X .
- (3) A reduced, irreducible complex space is called weakly Bergman if X_{reg} is a weakly Bergman manifold. Equivalently, one (or any) resolution of X is weakly Bergman.

Example 2.7. The following examples are Bergman manifolds: bounded domains in \mathbb{C}^n and their submanifolds, bounded domains in Stein manifolds and their submanifolds (or, more generally, manifolds of bounded type, cf. [BD18]), projective manifolds with very ample canonical bundle.

Remark 2.8. Let X, Y be two irreducible complex spaces and let $f : X \rightarrow Y$ be a bimeromorphic map. Then X is a weakly Bergman space if and only if Y is a weakly Bergman space. Indeed, f induces an isometry $f^* : \mathcal{H}_{Y_{\text{reg}}} \rightarrow \mathcal{H}_{X_{\text{reg}}}$ because any L^2 holomorphic n -form defined on an analytic Zariski open subset of a complex manifold extends automatically to the whole manifold.

Remark 2.9. One can show easily that the condition " $i\Theta(h_X)$ is smooth and positive definite on some open set" is equivalent to the fact that \mathcal{H}_X generates the 1-jets at a generic point of X . This implies that any finite (possibly ramified) cover of a Bergman manifold is again a Bergman manifold.

2.3. Existence of projective resolutions. The aim of this section is to prove the following technical yet useful result.

Lemma 2.10. *Let M be a Bergman manifold. Assume that there exists a discrete subgroup $\Gamma \subset \text{Aut}(M)$, acting properly discontinuously and let $X := \Gamma \backslash M$.*

If X is compact, then it admits a projective resolution. More generally, any compact complex space V admitting a generically immersive map to X , whose image is not entirely contained in the singular locus of X , admits a projective resolution.

Proof. We proceed in two steps.

Step 1. Case of X .

We denote by $p : M \rightarrow X$ the quotient map. The complex space X is normal with quotient singularities; in particular, it is \mathbb{Q} -factorial. Moreover, there exists an effective \mathbb{Q} -divisor Δ supported on the branch locus of p such that $K_M = p^*(K_X + \Delta)$. The Bergman metric h_M descends to X and induces a (singular) hermitian metric g_X on $K_X + \Delta$ with positive curvature current.

Let θ_M (resp. θ_X) be the curvature form $i\Theta(h_M)$ (resp. $i\Theta(g_X)$) of (K_M, h_M) (resp. $(K_X + \Delta, g_X)$). The form θ_M is a smooth Kähler form on M and one has $\theta_M = p^*\theta_X$.

Let ω_X be a hermitian metric on X and let $U \Subset M$ be a relatively compact open subset of M containing a fundamental domain for the action of Γ . Up to rescaling the metric ω_X , one can assume that

$$\theta_M \geq p^*\omega_X$$

holds on U . As both quantities are Γ -invariant, the inequality above is actually valid on the whole M . This implies that $\theta_X = p_*\theta_M$ is a Kähler current; more precisely, one has

$$\theta_X \geq \omega_X.$$

Next, we claim that the positive $(1,1)$ -current θ_X has bounded local potentials. Indeed, let $V \subset X$ a small open set where $\theta_X = dd^c\phi$. On $U \cap p^{-1}(V)$, the Kähler form θ_M can be written $\theta_M = dd^c(p^*\phi)$, hence $p^*\phi$ is smooth and thus locally bounded on $U \cap p^{-1}(V)$. As p maps that open set surjectively to V , our claim is proved.

Now, let $\pi : \widehat{X} \rightarrow X$ be a log resolution obtained by blowing up only smooth centers. It is well-known (cf. e.g. [DP04, Lemma 3.5]) that there exists a smooth $(1,1)$ -form $\beta \in c_1(E)$ where E is a (positive) rational combination of exceptional divisors of π such that

$$\pi^*\theta_X - \beta$$

is a Kähler current on \widehat{X} . By the observation above, this Kähler current has vanishing Lelong numbers, hence Demailly's regularization theorem [Dem92] enables us to find a Kähler form in the same cohomology class as $\pi^*\theta_X - \beta$. In particular, \widehat{X} is Kähler. Moreover, as the cohomology class of $\pi^*\theta_X - \beta$ is rational, \widehat{X} is projective thanks to Kodaira's embedding theorem.

Step 2. Case of V .

Let us call $j : V \rightarrow X$ the generically immersive map from the assumptions and let us denote by $\pi : \widehat{X} \rightarrow X$ the projective resolution obtained in Step 1. The strict transform of $\widehat{V} \subset \widehat{X}$ of V by π is a projective variety that maps bimeromorphically to $j(V)$ by π . As a result, $j(V)$ admits a projective resolution $\widetilde{V} \rightarrow j(V)$. Now, the bimeromorphic map

$$\widetilde{V} \rightarrow V$$

can be resolved by a finite sequence of blow ups along smooth centers; in particular, there exists a projective manifold endowed with a surjective, proper bimeromorphic map to V . \square

Remark 2.11. When the group Γ is linear, one can say more. Indeed, Γ is finitely generated as being a quotient of the fundamental group of the Zariski open set $X^\circ \subset X$ of regular values of p . By Selberg's lemma, there exists a finite index subgroup $\Gamma' \subset \Gamma$ with no torsion element. As Γ' acts properly discontinuously on M , the action must be free. In particular, $X' := \Gamma' \backslash M$ is smooth and $K_{X'}$ is positive by the argument above, hence X' is projective. As a result, X admits a finite cover by a smooth projective manifold; in particular, it is projective too.

2.4. The criterion.

Theorem 2.1. *Let \overline{X} be a compact Kähler manifold endowed with a reduced divisor D such that $\overline{X} \setminus D$ is uniformized by a weakly Bergman manifold. Let Δ_X be the associated covering divisor on \overline{X} .*

Then, the \mathbb{Q} -line bundle $K_{\overline{X}} + \Delta_{\overline{X}}$ is big.

Remark 2.12. An immediate corollary of the theorem is that under those assumptions, the logarithmic canonical bundle $K_{\overline{X}} + D$ is big. In particular, X is of log general type provided that D has simple normal crossings.

Proof. We proceed in two steps.

Step 1. Finding a metric on K_X

Let $X := \overline{X} \setminus D$ and let $p: \tilde{X} \rightarrow X$ the (Galois) cover from the assumptions. The Bergman metric $h_{\tilde{X}}$ is invariant under $\text{Aut}(p) \subset \text{Aut}(X)$ and $K_{\tilde{X}} = p^* K_X$, hence it descends to a singular metric g_X on K_X whose curvature $i\Theta(g_X)$ is a Kähler form on some euclidean open set of X . If one can show that g_X extends across D as a positively curved, singular metric on $K_{\overline{X}} + \Delta_{\overline{X}}$, then Boucksom's theorem [Bou02, Thm. 1.2] will show that $K_{\overline{X}} + \Delta_{\overline{X}}$ is indeed big, as expected.

Step 2. Extending the metric to $K_{\overline{X}} + \Delta_{\overline{X}}$

Next, we want to analyze the behavior of g_X near a general point of each irreducible component of D . Let $x \in D$ be a such a point; we denote by m the covering multiplicity attached to that point (or equivalently the chosen component). There exist a small neighborhood $x \in \overline{U} \simeq \mathbb{D}^n$ in \overline{X} and a system of coordinates (z_1, \dots, z_n) on \overline{U} centered at x such that $\overline{U} \cap D = (z_1 = 0)$. In particular, $U := \overline{U} \setminus D \simeq \mathbb{D}^* \times \mathbb{D}^{n-1}$. Let \tilde{U} a connected component of $p^{-1}(U)$.

$$\begin{array}{ccc} \tilde{U} & \hookrightarrow & \tilde{X} \\ \downarrow & & \downarrow p \\ U & \hookrightarrow & X \hookrightarrow \overline{X} \end{array}$$

By the very definition of Bergman metrics (see [Kob98, (4.10.4) Corollary]), it is easy to see that one has

$$h_{\tilde{U}} \leq h_{\tilde{X}}$$

where $h_{\tilde{U}}$ is the Bergman metric on (the canonical bundle of) \tilde{U} . Moreover, that same metric is invariant under $\text{Aut}(\tilde{U})$ and it descends to a metric g_U on $U \simeq \Delta^* \times \Delta^{n-1}$. One has

$$(2.1) \quad g_U \leq g_X \quad \text{on } U$$

If $m = \deg(p|_{\tilde{U}})$ is finite, then $p|_{\tilde{U}}$ is isomorphic to

$$\begin{aligned} \Delta^* \times \Delta^{n-1} &\rightarrow \Delta^* \times \Delta^{n-1} \\ (w_1, \dots, w_n) &\mapsto (w_1^m, w_2, \dots, w_n) \end{aligned}$$

and otherwise it is isomorphic to the universal cover

$$\begin{aligned} \Delta^n &\rightarrow \Delta^* \times \Delta^{n-1} \\ (w_1, \dots, w_n) &\mapsto (e^{\frac{w_1+1}{w_1-1}}, w_2, \dots, w_n) \end{aligned}$$

Accordingly, one finds

$$(2.2) \quad |dz_1 \wedge \dots \wedge dz_n|_{g_U}^2 = \begin{cases} m^2 \cdot |z_1|^{2(1-\frac{1}{m})} (1 - |z_1|^{2/m})^2 \cdot \prod_{k=2}^n (1 - |z_k|^2)^2 \\ |z_1|^2 \log^2 |z_1|^2 \cdot \prod_{k=2}^n (1 - |z_k|^2)^2 \end{cases}$$

By the formula above, the quantity

$$-\log |dz_1 \wedge \dots \wedge dz_n|_{g_U}^2 + \left(1 - \frac{1}{m}\right) \cdot \log |z_1|^2$$

is locally bounded above near $\{0\} \times \mathbb{D}^{n-1}$ in both cases. Now, let us view $\sigma := dz_1 \wedge \dots \wedge dz_n$ as an element in $H^0(\overline{U}, K_{\overline{X}})$. By what was just said, the psh weight $-\log |\sigma|_{g_U}^2 + (1 - \frac{1}{m}) \log |z_1|^2$ on $(K_{\overline{X}} + \Delta_{\overline{X}})|_U$ is bounded above locally near $\overline{U} \setminus U$, hence extends across that hypersurface

to a psh weight on $(K_{\overline{X}} + \Delta_{\overline{X}})|_{\overline{U}}$. Letting $h_{\Delta_{\overline{X}}}$ be a singular metric on $\mathcal{O}_{\overline{X}}(\Delta_{\overline{X}})$ with curvature current $[\Delta_{\overline{X}}]$, the process above shows that $g_X \otimes h_{\Delta_{\overline{X}}}$ extends in codimension one with positive curvature, hence everywhere. This ends the proof. \square

Corollary 2.13. *Let X be a normal, compact complex space and let X° be some non-empty analytic Zariski open subset. Assume that one of the following conditions holds.*

- (i) X is a quotient of a Bergman manifold, étale over X° .
- (ii) X is \mathbb{Q} -factorial, admits a Kähler resolution and X° is uniformized by a weakly Bergman complex space.

Let $\Delta_X \subset X \setminus X^\circ$ be the covering divisor associated to the above étale cover of X° . Then, the \mathbb{Q} -line bundle $K_X + \Delta_X$ is big.

Proof. In both cases, X admits a Kähler resolution $\pi : \widehat{X} \rightarrow X$. This is a consequence of Lemma 2.10 in case (i) and it is an assumption in case (ii).

By Theorem 2.1, the covering divisor $\Delta_{\widehat{X}} \subset \widehat{X}$ associated to the étale cover of $X^\circ \cap X_{\text{reg}} \hookrightarrow \widehat{X}$ by a weakly Bergman manifold satisfies that $K_{\widehat{X}} + \Delta_{\widehat{X}}$ is big. As π is an isomorphism over the generic point of each components of Δ_X , there is a π -exceptional effective \mathbb{Q} -divisor E such that $\Delta_{\widehat{X}} = \pi^* \Delta_X + E$. In particular, one has a \mathbb{Q} -linear equivalence $K_{\widehat{X}} + \Delta_{\widehat{X}} \sim_{\mathbb{Q}} \pi^*(K_X + \Delta_X)$ valid over $\widehat{X} \setminus \text{Exc}(\pi)$. Now, the following restriction map induces an injection for any divisible enough integer m :

$$\begin{aligned} H^0(\widehat{X}, m(K_{\widehat{X}} + \Delta_{\widehat{X}})) &\longrightarrow H^0(\widehat{X} \setminus \text{Exc}(\pi), m(K_{\widehat{X}} + \Delta_{\widehat{X}})) \\ &\simeq H^0(\pi(\widehat{X} \setminus \text{Exc}(\pi)), m(K_X + \Delta_X)) \\ &\simeq H^0(X, m(K_X + \Delta_X)), \end{aligned}$$

and it follows that $K_X + \Delta_X$ is big. \square

3. APPLICATIONS TO QUOTIENTS OF MANIFOLDS OF BOUNDED TYPE

3.1. Main result. Let Ω be a complex manifold of *bounded type*, i.e. a complex manifold admitting a bounded strictly psh function, as defined in [BD18]. This category includes bounded domains, and is stable by taking étale covers or open/closed subvarieties; the reader may wish to think of Ω as a bounded domain.

Our main object of study will be a suitable compactification of a quotient of Ω . Throughout the text, we will make various assumptions on this compactification; our general hypotheses will be as follows.

Assumption 3.1. We fix a reduced, irreducible, compact complex space \overline{X} , and an open (dense) Zariski subset $X = \overline{X} \setminus D$. We assume that X is a quotient of Ω , i.e. there exists a discrete subgroup $\Gamma \subset \text{Aut}(\Omega)$, acting properly discontinuously, and a fixed identification of complex spaces

$$X = \Gamma \backslash \Omega.$$

We denote by $p : \Omega \rightarrow X$ the projection map.

We let $X^\circ := X \setminus \text{Sing}(p) \subset X$ be the locus of regular values of p and we set $\Omega^\circ := p^{-1}(X^\circ)$. The set X° is a Zariski-analytic open subset of X_{reg} . Note that the inclusion $X^\circ \subset X_{\text{reg}}$ is strict if e.g. p ramifies in codimension one.

$$\begin{array}{ccc} \Omega^\circ & \hookrightarrow & \Omega \\ \downarrow & & \downarrow p \\ X^\circ & \hookrightarrow & X \hookrightarrow \overline{X} \end{array}$$

Theorem 3.1. *With the notation above, let V be a normal, \mathbb{Q} -factorial compact complex space admitting a Kähler resolution and let $j : V \rightarrow \overline{X}$ be a generically immersive map such that $j(V) \not\subset \overline{X} \setminus X^\circ$. Set $V^\circ := j^{-1}(X^\circ)$.*

Then, V admits a natural covering divisor Δ_V supported on $V \setminus V^\circ$ and $K_V + \Delta_V$ is big.

Proof. Let $\pi : \widehat{V} \rightarrow V$ be a Kähler resolution, and let $\widehat{V}^\circ := \pi^{-1}(V^\circ)$. Let Z° be a connected component of $p^{-1}(j(V^\circ))$, let $W^\circ := V^\circ \times_{j(V^\circ)} Z^\circ$ and let $\widehat{W}^\circ := \widehat{V}^\circ \times_{V^\circ} W^\circ$. In the following, we replace W° and \widehat{W}° with their irreducible component dominating V° so that the map $f : \widehat{W}^\circ \rightarrow Z^\circ$ below is a bimeromorphic map of irreducible complex spaces:

$$\begin{array}{ccccccc}
 & & & f & & & \\
 & & & \curvearrowright & & & \\
 \widehat{W}^\circ & \longrightarrow & W^\circ & \longrightarrow & Z^\circ & \hookrightarrow & \Omega^\circ \\
 \downarrow q & & \downarrow & & \downarrow p|_{Z^\circ} & & \downarrow p \\
 \widehat{V}^\circ & \xrightarrow{\pi|_{\widehat{V}^\circ}} & V^\circ & \xrightarrow{j|_{V^\circ}} & j(V^\circ) & \hookrightarrow & X^\circ
 \end{array}$$

As an analytic Zariski open subset of a compact Kähler manifold, \widehat{V}° can be endowed with a complete Kähler metric, cf. Lemma 3.2 below. As q is étale, the same property holds for the smooth manifold \widehat{W}° . The pull back by the bimeromorphic map f of the globally bounded, strictly psh function on Z° induced by the restriction of the one living on Ω is still globally bounded, psh and strictly psh on a non-empty Zariski open subset of \widehat{W}° . By [BD18, Lemma 1.2], \widehat{W}° is a weak Bergman manifold, hence W° is a weakly Bergman complex space, cf. Remark 2.8. The theorem now follows from Corollary 2.13 (ii). \square

We used the following standard result, which we recall for the reader's convenience.

Lemma 3.2. *Let Y be a compact Kähler manifold and let $Y^\circ \subset Y$ be a Zariski open subset. There exists a complete Kähler metric ω on Y° .*

Proof. Up to taking a log resolution of $(Y, Y \setminus Y^\circ)$ leaving Y° untouched, one can assume that the complement of Y° in Y is a simple normal crossings divisor. Then, it is standard to construct Poincaré type metrics on Y° , cf. e.g. [CG72], or [Dem82, Théorème 1.5]. \square

Remark 3.3. As already observed in the introduction, by the main theorem of [Gri71], any projective manifold is covered by Zariski open subsets which are uniformized by pseudoconvex bounded domains. Thus, all projective varieties fall in the scope of Assumption 3.1. Theorem 3.1 implies in particular that for any projective manifold \overline{X} , there exists a Zariski open subset $X \subset \overline{X}$ such that for any subvariety $V \subset \overline{X}$, the quasi-projective variety $V \cap X$ is of log general type.

Of course, the most interesting results will be obtained in settings where we are able to obtain a good description of this open subset X .

3.2. Examples of applications of Theorem 3.1. The following are the main examples of applications.

(1) [COMPACT ÉTALE]

In the setting of Assumption 3.1, assume furthermore that $\overline{X} = X$ is a smooth manifold and that p is étale. Let $W \subset X$ be an irreducible variety of X and let $j : V \rightarrow W$ be a resolution of singularities. Then K_V is big; i.e. W is of general type. This is the content of the main result of [BD18].

(2) [NON-COMPACT ÉTALE]

More generally, assume only that \overline{X} is a smooth compact Kähler manifold and that p is étale. Let $W \subset \overline{X}$ be a compact subvariety not included in $\overline{X} \setminus X$ and let $j : V \rightarrow W$ be a log resolution of $(W, W \cap (\overline{X} \setminus X))$. Then there exists a divisor Δ_V on V , supported

over $\overline{X} \setminus X$ via j such that $K_V + \Delta_V$ is big. In particular, $W \cap X$ is of log general type.

(3) [COMPACT NON-ÉTALE]

Assume that $\overline{X} = X$, let $W \subset X$ be an irreducible subvariety not included in the branch locus of p and let $j: V \rightarrow W$ be a resolution of W . Then, there exists a natural divisor Δ_V on V with coefficients in $(0, 1)$, supported over the branch locus of p via j such that $K_V + \Delta_V$ is big.

(4) [Q-FACTOREAL SUBVARIETIES]

In the setting 3.1, assume that \overline{X} admits a Kähler resolution. Let $V \subset \overline{X}$ be a normal, \mathbb{Q} -factorial subvariety not included in the branch locus of p or $\overline{X} \setminus X$. Then, there exists a reduced divisor Δ_V on V , supported on $\overline{X} \setminus X^\circ$, such that $K_V + \Delta_V$ is big.

(5) [Q-FACTORIZATION OF SUBVARIETIES]

Assume that $\overline{X} = X$, let $W \subset X$ be an irreducible closed subvariety not included in the branch locus of p , let \widetilde{W} be a connected component of $p^{-1}(W)$ and let $\Gamma_{\widetilde{W}} \subset \Gamma$ be the subgroup of Γ preserving \widetilde{W} . Let $\widetilde{V} \rightarrow \widetilde{W}$ be a $\Gamma_{\widetilde{W}}$ -equivariant resolution of singularities of \widetilde{W} and let $V := \Gamma_{\widetilde{W}} \backslash \widetilde{V}$. This normal variety has quotient singularities, in particular it is \mathbb{Q} -factorial, and it comes equipped with a birational map $j: V \rightarrow W$.

$$\begin{array}{ccccc} \widetilde{V} & \longrightarrow & \widetilde{W} & \hookrightarrow & \Omega \\ \downarrow & & \downarrow & & \downarrow p \\ V & \xrightarrow{j} & W & \hookrightarrow & X \end{array}$$

Moreover, there is a natural branching divisor Δ_V on V attached to the cover $\widetilde{V} \rightarrow V$, supported over $\text{Sing}(p)$ via j and satisfying that $K_V + \Delta_V$ is big.

3.3. Two remarks about the singular vs smooth case. The following two examples show that the situation in the singular case is rather subtle.

(1) [K_X BIG v. X OF GENERAL TYPE]

Assume for simplicity that X is compact and p is quasi-étale, that is, $\text{codim}_X(X \setminus X^\circ) \geq 2$. Then, Example (4) in Section 3.2 shows that K_X is big. Unless X has only canonical singularities, this property is weaker than saying that X is of general type, i.e. that the canonical bundle $K_{\widehat{X}}$ of a (or any) resolution $\widehat{X} \rightarrow X$ is big.

For instance, there exist surfaces S which are a quotient of the bi-disk $\Delta^2 \subset \mathbb{C}^2$ such that K_S is ample and yet S is not of general type. One can realize such surfaces as $S = (C_1 \times C_2)/G$ where C_1, C_2 are curves of genus at least two and G is a finite group acting diagonally, cf. [BP16, Table 1].

(2) [X OF GENERAL TYPE BUT NOT ALL ITS SUBVARIETIES]

Assume again that X is compact and p is quasi-étale. The example above shows that X needs not be of general type. Even if we assume that X is of general type, it may still happen that X contains subvarieties $V \not\subset X_{\text{sing}}$ such that V is not of general type.

Indeed, let C be a hyperelliptic curve and let $f: C \rightarrow \mathbb{P}^1$ be the double cover; it induces an involution $\iota \in \text{Aut}(C)$. The transformation $C \times C \ni (z, w) \mapsto (\iota(w), \iota(z))$ induces an action of $\mathbb{Z}/4\mathbb{Z}$ on $C \times C$. Let $X := \mathbb{Z}/4\mathbb{Z} \backslash C \times C$; it is a projective variety with canonical singularities and ample canonical bundle admitting a cover by the bidisk in \mathbb{C}^2 . Yet, the diagonal map $C \rightarrow X$ factors through \mathbb{P}^1 as showed below.

$$\begin{array}{ccccc}
C & \xleftarrow{\Delta} & C \times C & \longrightarrow & X \\
& \searrow f & & \nearrow j & \\
& & \mathbb{P}^1 & &
\end{array}$$

4. COMPARISON OF COVERING DIVISORS

In this section, we work under the general Assumption 3.1. Given any generically immersive map $V \rightarrow \overline{X}$, Definition 2.2 allows us to attach a natural covering divisor $\Delta_{\widehat{V}}$ to any resolution of singularities of $\widehat{V} \rightarrow V$. In this section, we will gather a few facts allowing us to compare the natural orbifold structures on adequate resolution of singularities of V and \overline{X} .

Let us recall how Definition 2.2 permits to construct the orbifold structures in this context.

4.1. Natural orbifold structure on a resolution of singularities of a singular quotient.

Let us fix a log-resolution $\widehat{X} \xrightarrow{\pi} \overline{X}$, such that the preimage of the union of D and the closure of $\text{Sing}(p)$ is a divisor E with simple normal crossings. Let $E = \sum_i E_i$ be the decomposition of E into its irreducible components. Also, let g_X be the natural smooth metric induced on K_X by the Bergman kernel on K_Ω .

Let Y be the normalization of the component $T_{\widehat{X}}$ of the fiber product $\widehat{X} \times_{\overline{X}} \Omega$ dominating \widehat{X} (or, equivalently, \overline{X}). It sits in the following commutative diagram.

$$\begin{array}{ccc}
Y & \xrightarrow{f_{\widehat{X}}} & \widehat{X} \\
\downarrow \sigma & & \downarrow \pi \\
\Omega & \xrightarrow{p} & \overline{X}
\end{array}$$

FIGURE 1.

Remark that Γ acts naturally on the product $\Omega \times_X \widehat{X}$, by having its natural action on the first factor, and leaving the second one invariant. Hence it also acts on Y .

Let $U \subset \widehat{X}$ be a sufficiently small neighborhood of the generic point of E_i . The map $f_{\widehat{X}} : f_{\widehat{X}}^{-1}(U \setminus E_i) \rightarrow U \setminus E_i$ is an étale cover. This map induces a cyclic cover when restricted to any of the connected components of its source: the Galois group of this cover is isomorphic to $\mathbb{Z}/m_i\mathbb{Z}$ for some $m_i \in \mathbb{N}^* \cup \{\infty\}$ (with $\infty \cdot \mathbb{Z} = 0$).

Then, Definition 2.2 gives us the following natural orbifold structure on \widehat{X} .

Definition 4.1. We let $\Delta_{\widehat{X}}$ be the covering divisor associated to the data $(\widehat{X}, \widehat{X} \setminus E, f_{\widehat{X}}|_{\widehat{X} \setminus E})$ by means of Definition 2.2. By the previous discussion, it is equal to the \mathbb{Q} -divisor with simple normal crossing support $\sum_i (1 - \frac{1}{m_i})E_i$, where the m_i are defined as above.

One way to think about this orbifold structure is provided by the following formula which is direct consequence of the definition above.

Lemma 4.2. *With the notation above, one has*

$$K_Y = f_{\widehat{X}}^*(K_{\widehat{X}} + \Delta_{\widehat{X}}).$$

Remark 4.3. A similar, but coarser way of forming an orbifold pair (\widehat{X}, Δ) is used in [CRT17]. In that article, each component E_i is endowed with the multiplicity ∞ if $\pi(E_i) \subset \overline{X} \setminus X$, and with the multiplicity $\mu_i = |S_{\pi(E_i)}|$ otherwise (where $S_{\pi(E_i)}$ is the isotropy group of the generic point of $\pi(E_i)$). With our convention, if $\pi(E_i) \cap X \neq \emptyset$, the Galois group $\mathbb{Z}/m_i\mathbb{Z}$ identifies with a subgroup of the stabilizer of any inverse image of $\pi(E_i)$ in Ω , i.e. it is a subgroup of the isotropy group $S_{\pi(E_i)}$. Consequently, we have $m_i \leq \mu_i$.

Clearly, a component E_i such that $\pi(E_i) \cap X \neq \emptyset$ satisfies $m_i < \infty$. Conversely, one can prove the following:

Lemma 4.4. *Assume that Ω is a bounded domain satisfying the following property: for all $z \in \partial\Omega$, there exist a neighborhood $U_z \subset \mathbb{C}^n$ of z and a psh function φ_z on U_z such that $\varphi_z^{-1}((-\infty, 0)) = \Omega \cap U_z$.*

Then, any component E_i such that $\pi(E_i) \cap X = \emptyset$ will satisfy $m_i = \infty$.

Proof. Note that by upper semicontinuity of φ_z , one has $\varphi_z|_{\partial\Omega \cap U} \equiv 0$. Let $E_i \subset \widehat{X}$ be a divisor with finite multiplicity and let us consider the étale cover $q: q^{-1}(U \setminus E_i) \rightarrow U \setminus E_i \simeq \Delta^* \times \Delta^{n-1}$ as above. Let $V^\circ \subset Y$ be a connected component of $q^{-1}(U \setminus E_i)$, so that $q|_{V^\circ}$ can be compactified as a (surjective) ramified finite cover $\bar{q}: V \rightarrow U \simeq \Delta^n$ of order m_i where V is some smooth manifold containing V° as a Zariski open subset. In particular, one has

$$(4.1) \quad (\bar{q} \circ \pi)(V \setminus V^\circ) = \pi(E_i \cap U).$$

As $\Omega \subset \mathbb{C}^n$ is bounded, the map $f^\circ := \sigma|_{V^\circ}: V^\circ \rightarrow \Omega$ extends to a holomorphic map $f: V \rightarrow \overline{\Omega}$. We claim that

$$(4.2) \quad \text{Im}(f) \subset \Omega$$

from which the Lemma follows. Indeed one would then have $\bar{q} \circ \pi = p \circ f$ on V by density of V° in V and therefore one would get $\pi(E_i \cap U) \subset \text{Im}(p) \subset X$ given (4.1).

We now prove (4.2) arguing by contradiction. Suppose that there exists $v \in V$ such that $f(v) \in \partial\Omega$. Let $z := f(v)$ and let (U_z, φ_z) be provided by our assumption on Ω . There exists a small neighborhood W of $v \in V$ such that $f(W) \subset U_z$. Then, the psh function $\varphi_z \circ f|_W$ is non-negative and attains its maximum 0 at the interior point $v \in W$. By the maximum principle, $\varphi_z \circ f|_W$ is constant, identically equal to 0. This is in contradiction with the fact that $(\varphi_z \circ f|_W)(W \cap V^\circ) \subset \varphi_z(\Omega \cap U_z) \subset (-\infty, 0)$. \square

Remark 4.5. Lemma 4.4 fails for a general bounded domain. Indeed, let $\pi: \widehat{X} \rightarrow X' \simeq \Omega'/\Gamma$ be a resolution of a singular compact quotient X' of some bounded domain Ω' and let $E \subset \widehat{X}$ be an irreducible, π -exceptional divisor. Then, define $\Omega := \Omega' \setminus p^{-1}(\pi(E))$, $X := X' \setminus \pi(E)$ so that $X \simeq \Omega/\Gamma$ is naturally compactified by X' . Then, the multiplicity of E associated to $\pi: \widehat{X} \rightarrow X'$ is finite and yet $\pi(E) \cap X = \emptyset$.

4.2. Relative orbifold construction. The previous construction has a relative variant, which uses Definition 2.2 to construct a particular model $(\widehat{V}, \Delta_{\widehat{V}})$, once we are given a generically immersive map $q: V \rightarrow \overline{X}$ and a resolution $\widehat{X} \rightarrow \overline{X}$.

Suppose here that V is an m -dimensional complex manifold, and that the generically immersive map q is such that $q(V) \not\subset D \cup \text{Sing}(p)$. Then, we will construct \widehat{V} as follows.

Let $V' \xrightarrow{j} \widehat{X}$ be the component of the fiber product $V \times_X \widehat{X}$ that dominates V . Let $\widehat{V} \rightarrow V'$ be a resolution of singularities; it induces a birational map $\sigma: \widehat{V} \rightarrow V'$. Let us denote by $F \subset \widehat{V}$ the non-étale locus of σ , and define $F^{(1)} \subset F$ to be the union of all irreducible components of F with codimension one. We also introduce the fiber product $\widetilde{V} = V \times_X \Omega$. Note that this complex space may have infinitely many connected components, all isomorphic under the action of Γ .

Finally, we let $T_{\widehat{V}}$ be the union of all irreducible components of the product $\widetilde{V} \times_V \widehat{V}$ dominating \widehat{V} , and we denote by Z be the normalization of $T_{\widehat{V}}$. All these operations lead to the diagram showed in Figure 2.

Let $\Gamma_V \subset \Gamma$ be the stabilizer of $\iota(\widetilde{V}) \subset \Omega$. Then Γ_V acts on $V \times_X \Omega$, by its natural action on the second factor, and by the trivial action on the first. Thus, it induces a *natural action on \widetilde{V} , making ι a Γ_V -equivariant map.*

Under these conditions, the group Γ_V has a natural action on the fiber product $\widetilde{V} \times_V \widehat{V}$, by operating on the first factor, and leaving the second one invariant. This action leaves $T_{\widehat{V}}$

$$\begin{array}{ccccc}
& & \widehat{V} & \xrightarrow{r} & \widehat{X} \\
& \nearrow f_{\widehat{V}} & \downarrow \sigma & & \downarrow \pi \\
Z & & V & \xrightarrow{q} & \overline{X} \\
\downarrow \rho & & & & \\
\widetilde{V} & \nearrow \iota & & \xrightarrow{\nu} & \Omega & \nearrow p
\end{array}$$

FIGURE 2.

invariant and, therefore, it induces a natural action on Z . Again, Z may have more than one connected component in general, all equivalent under the action of Γ .

By construction, the map $f_{\widehat{V}}$ is étale over $\widehat{V} \setminus F$. By purity of the branch locus, it is actually étale over $\widehat{V} \setminus F^{(1)}$. Therefore, one can apply Definition 2.2, and endow each component $F_i \subset F^{(1)}$ with a natural multiplicity $n_i \in \mathbb{N}^* \cup \{\infty\}$.

Definition 4.6. We let $\Delta_{\widehat{V}}$ be the covering divisor that Definition 2.2 associates to the data $(\widehat{V}, \widehat{V} \setminus F, f_{\widehat{V}}|_{\widehat{V} \setminus F})$. We have $\Delta_{\widehat{V}} = \sum_i (1 - \frac{1}{n_i}) F_i$, for some $n_i \in \mathbb{N}^* \cup \{\infty\}$.

Similarly to Lemma 4.2, one has

Lemma 4.7. *With the notation above, one has*

$$K_Z = f_{\widehat{V}}^*(K_{\widehat{V}} + \Delta_{\widehat{V}}).$$

4.3. The comparison result. Our next goal is to relate the orbifold multiplicities given by the divisor $\Delta_{\widehat{V}}$ with the ones inherited from the pair $(\widehat{X}, \Delta_{\widehat{X}})$: this will be the content of Proposition 4.9. Before this, we need a lemma.

Lemma 4.8. *The variety Z is naturally isomorphic to the normalization of the union of the components of $\widehat{V} \times_{\widehat{X}} Y$ dominating \widehat{V} .*

Proof. Note that the associativity of fiber products yields

$$\widehat{V} \times_V \widetilde{V} = \widehat{V} \times_V (V \times_{\overline{X}} \Omega) \simeq \widehat{V} \times_{\overline{X}} \Omega \simeq \widehat{V} \times_{\widehat{X}} (\widehat{X} \times_{\overline{X}} \Omega).$$

From this, we get that $T_{\widehat{V}}$, the disjoint union of the components of $\widehat{V} \times_V \widetilde{V}$ dominating \widehat{V} , identifies with the disjoint union of the components of $\widehat{V} \times_{\widehat{X}} T_{\widehat{X}}$ dominating \widehat{V} . Now, the universal property of the normalization functor \bullet^ν allows us to complete the square as follows

$$\begin{array}{ccc}
(\widehat{V} \times_{\widehat{X}} T_{\widehat{X}}^\nu)^\nu & \dashrightarrow & (\widehat{V} \times_{\widehat{X}} T_{\widehat{X}})^\nu \\
\downarrow & & \downarrow \\
\widehat{V} \times_{\widehat{X}} T_{\widehat{X}}^\nu & \longrightarrow & \widehat{V} \times_{\widehat{X}} T_{\widehat{X}}
\end{array}$$

Now, the dotted arrow represents a finite bimeromorphic map between two normal reduced complex analytic spaces hence it is an isomorphism. As $Y = T_{\widehat{X}}^\nu$, the normalization of the disjoint union of components of $\widehat{V} \times_{\widehat{X}} Y$ dominating \widehat{V} is the same thing as the disjoint union of components of $(\widehat{V} \times_{\widehat{X}} T_{\widehat{X}})^\nu$ dominating \widehat{V} . By what was said previously, this is nothing but saying that $T_{\widehat{V}}^\nu = Z$. \square

The natural orbifold structures on \widehat{V} and \widehat{X} are now comparable in the following manner.

Proposition 4.9. *With the notation above, one has*

$$\Delta_{\widehat{V}} \leq r^* \Delta_{\widehat{X}},$$

i.e. the difference $r^ \Delta_{\widehat{X}} - \Delta_{\widehat{V}}$ between these two \mathbb{Q} -divisors is effective.*

Proof. We have the following commutative diagram:

$$\begin{array}{ccccc}
 & & \widehat{V} & \xrightarrow{r} & \widehat{X} \\
 & f_{\widehat{V}} \nearrow & \downarrow & \xrightarrow{f_{\widehat{X}}} & \downarrow \\
 Z & \xrightarrow{s} & Y & & \\
 \downarrow & & \downarrow & \xrightarrow{\quad} & \downarrow \\
 \widetilde{V} & \xrightarrow{\quad} & V & \xrightarrow{\quad} & \overline{X} \\
 & & \downarrow & \nearrow & \\
 & & \Omega & &
 \end{array}$$

We claim that over $\widehat{V} \setminus F$, we have $Z \simeq \widehat{V} \times_{\widehat{X}} Y$. Given Lemma 4.8, it is sufficient to prove that $Z|_{f_{\widehat{V}}^{-1}(\widehat{V} \setminus F)}$ is smooth and that each of its connected components dominates \widehat{V} . As σ is an isomorphism over $\widehat{V} \setminus F$, it suffices to check those properties for $\widetilde{V} \rightarrow V$ over that same set but this is then straightforward.

Let η be a general point of a component F_i of F such that $2 \leq n_i \leq +\infty$ and let $U_{\widehat{V}} \subset \widehat{V}$ be a small neighborhood of η on which F_i admits the equation $v_i = 0$. Let $W_{\widehat{V}}$ be a connected component of $f_{\widehat{V}}^{-1}(U_{\widehat{V}} \setminus F_i)$. By Definition 2.2, the map $W_{\widehat{V}} \xrightarrow{f_{\widehat{V}}} U_{\widehat{V}} \setminus F_i$ is an étale cover, with Galois group $G_i = \mathbb{Z}/n_i\mathbb{Z}$. As $n_i \geq 2$, F_i sit above $\text{Branch}(p)$ as otherwise, $f_{\widehat{V}} : W_{\widehat{V}} \rightarrow U_{\widehat{V}} \setminus F_i \simeq \Delta^m$ is étale hence an isomorphism.

Set $\zeta = r(\eta)$, and let $U_{\widehat{X}}$ be a small neighborhood of ζ containing $r(U_{\widehat{V}})$. Denote also by $W_{\widehat{X}}$ the connected component of $f_{\widehat{X}}^{-1}(U_{\widehat{X}} \setminus E)$ containing $s(W_{\widehat{V}})$. We get a map of étale covers

$$(4.3) \quad \begin{array}{ccc}
 W_{\widehat{V}} & \xrightarrow{s} & W_{\widehat{X}} \\
 f_{\widehat{V}} \downarrow & & \downarrow f_{\widehat{X}} \\
 U_{\widehat{V}} \setminus F_i & \xrightarrow{r} & U_{\widehat{X}} \setminus E
 \end{array}$$

and we know from our observation at the beginning of the proof that the diagram

$$(4.4) \quad \begin{array}{ccc}
 s^{-1}(W_{\widehat{X}}) & \xrightarrow{s} & W_{\widehat{X}} \\
 f_{\widehat{V}} \downarrow & & \downarrow f_{\widehat{X}} \\
 U_{\widehat{V}} \setminus F_i & \xrightarrow{r} & U_{\widehat{X}} \setminus E
 \end{array}$$

is a fiber product. Therefore, we have

$$\begin{aligned}
 n_i &= \deg \left(W_{\widehat{V}} \xrightarrow{f_{\widehat{V}}} U_{\widehat{V}} \setminus F_i \right) \\
 &\leq \#(\text{fiber of } s^{-1}(W_{\widehat{X}}) \xrightarrow{f_{\widehat{V}}} U_{\widehat{V}} \setminus F_i) \\
 &= \deg \left(W_{\widehat{X}} \xrightarrow{f_{\widehat{X}}} U_{\widehat{X}} \setminus E \right)
 \end{aligned}$$

If H is the Galois group of $f_{\widehat{X}} : W_{\widehat{X}} \rightarrow U_{\widehat{X}} \setminus E$, we have $n_i \leq |H|$. Let $(E_j)_{j \in J}$ be the components of E passing through ζ . Since m_j is the order of the element of H associated to the meridian loop around E_j , the proof of [Kol07, Theorem 2.23] shows that H is an abelian group satisfying

$$|H| \leq \prod_{j \in J} m_j.$$

Given $j \in J$, let us introduce z_j a local equation for E_j . Since v_i divides each $r^* z_j$ in \mathcal{O}_η , to finish the proof, it suffices to show that $1 - \frac{1}{n_i} \leq \sum_{j \in J} \left(1 - \frac{1}{m_j}\right)$. But this is now an easy consequence of the inequality $n_i \leq \prod_{j \in J} m_j$ obtained previously. \square

5. A CRITERION FOR HYPERBOLICITY

The main goal of this section is to present a hyperbolicity result for the complex space \overline{X} , provided that the manifold Ω in the general Assumption 3.1 is actually a bounded domain. The section is organized as follows:

- In Section 5.1, we give some complements on the Bergman metric and how to compute its curvature, cf. (5.1).
- In Section 5.2, we gather a few results allowing us to estimate the curvature of the Bergman kernel on Ω , close to the classical comparison theorems between the Bergman, Carathéodory and Kobayashi metrics (see [Hah78, Kob98]). The main result of that section is Proposition 5.7.
- In Section 5.3, we build on the previous section to construct a singular hermitian metric on a modification of a weakly pseudoconvex Kähler manifold V admitting a generic immersive map to \overline{X} . The main result of the section is a curvature inequality for that metric, cf. Theorem 5.9.
- In Section 5.4, we exploit the previous results to state and prove a hyperbolicity criterion for \overline{X} , cf. Theorem 5.2.

Throughout the rest of this section, we assume that $\Omega \subset \mathbb{C}^n$ is a bounded domain.

5.1. Computation of the curvature of the Bergman metric. Let Y be a complex manifold of dimension n . Let us give some complements on the discussion of Section 2.2, and briefly recall how to compute the curvature $i\Theta(h_Y)$ of the Bergman metric h_Y on K_Y (when it is defined).

Let $\mathcal{H}_Y = \left\{ \sigma \in H^0(Y, K_Y) \mid \int_Y i^{n^2} \sigma \wedge \bar{\sigma} < +\infty \right\}$ be the Hilbert space of holomorphic square integrable n -forms on Y . If e is some local trivialization of K_Y , the norm of e for the metric h_Y has the following value at a point $x \in Y$:

$$\|e\|_{h_Y, x} = \frac{1}{\|\text{ev}_x\|_{\mathcal{H}_Y^*}},$$

where $\text{ev}_x : \mathcal{H}_Y \rightarrow \mathbb{C}$ is the evaluation form which to $\sigma \in \mathcal{H}_Y$ associates λ such that $\sigma_x = \lambda e_x$, and $\|\cdot\|_{\mathcal{H}_Y^*}$ is the natural dual norm on \mathcal{H}_Y^* . Thus, the Bergman metric at x is well defined provided there exists $\sigma \in \mathcal{H}_Y$ such that $\sigma_x \neq 0$.

Consider now a point $x \in Y$ such that $\|\cdot\|_{h_Y, x}$ is defined. By definition of h_Y , there exists a section $e \in \mathcal{H}_Y$ such that $\|e\|_{\mathcal{H}_Y} = 1$ and $\|e_x\|_{h_Y, x} = 1$. Now, if $v \in T_{Y, x}$, the curvature of h_Y in the direction v can be computed by the following formula (see [Kob98, Proposition 4.10.10]).

$$(5.1) \quad i\Theta(h_Y)(v, \bar{v}) = \max_f |df(v)|^2,$$

where $f \in \mathfrak{m}_{Y, x} \subset \mathcal{O}_{Y, x}$ runs among all germs of holomorphic functions at x such that there exists $\sigma \in \mathcal{H}_Y$ with $\|\sigma\|_{\mathcal{H}_Y} = 1$, $\sigma(x) = 0$ and $\sigma \stackrel{\text{loc}}{=} f e$.

5.2. Curvature inequalities on subvarieties. We will now use the previous description of the curvature of h_Y to state a comparison result between the curvature of the Bergman metric of a bounded domain and that of a bounded *symmetric* domain included in it. We will then use this result to obtain a curvature estimate for the subvarieties of Ω .

Let \mathcal{D} be a bounded *symmetric* domain of dimension n , centered at $0 \in \mathbb{C}^n$, with coordinates (t_1, \dots, t_n) . Since \mathcal{D} is S^1 -invariant, we see immediately that two polynomials $t^\alpha = t_1^{\alpha_1} \dots t_n^{\alpha_n}$ ($\alpha = (\alpha_1, \dots, \alpha_n)$) and $t^\beta = t_1^{\beta_1} \dots t_n^{\beta_n}$ ($\beta = (\beta_1, \dots, \beta_n)$) are orthogonal for the standard scalar product, whenever $\alpha \neq \beta$. After renormalizing the family $(t^\alpha dt_1 \wedge \dots \wedge dt_n)_{\alpha \in \mathbb{N}^n}$, we get a unitary basis $(e_i)_{i \in \mathbb{N}}$ of $\mathcal{H}_{\mathcal{D}}$, of the form $e_i = f_i dt_1 \wedge \dots \wedge dt_n$, with

$$f_0 = \frac{1}{\text{Vol}(\mathcal{D})^{\frac{1}{2}}}, \quad f_1 = \frac{1}{a_1} t_1, \quad \dots, \quad f_n = \frac{1}{a_n} t_n,$$

where $a_i^2 = \int_{\mathcal{D}} |t_i|^2 d\text{Vol}$, all other f_i being polynomials in t with vanishing 1-jet at 0.

This implies that

$$\|dt_1 \wedge \dots \wedge dt_n\|_{h_{\mathcal{D},0}}^2 = \frac{1}{\sum_{i \in \mathbb{N}} |f_i(0)|^2} = \text{Vol}(\mathcal{D}),$$

Let $v \in T_{\mathcal{D},0}$. Taking $e = e_0$, the equality case in Cauchy-Schwarz inequality shows that the maximum in (5.1) is attained for $\sigma = fe$ with

$$(5.2) \quad f = \frac{\sum_i \bar{v}_i t_i / a_i^2}{(\sum_i |v_i|^2 / a_i^2)^{1/2}} \cdot \text{Vol}(\mathcal{D})^{1/2}.$$

This yields, by (5.1):

$$(5.3) \quad i\Theta(h_{\mathcal{D}})(v, \bar{v}) = \left(\sum_i \frac{|v_i|^2}{a_i^2} \right) \cdot \text{Vol}(\mathcal{D})$$

We are now ready to state our first comparison result.

Lemma 5.1. *Let $x \in \Omega$. Let $j : \mathcal{D} \hookrightarrow \Omega$ be an open embedding, such that $j(0) = x$. Then, we have*

$$j^*(i\Theta(h_{\Omega})_x) \leq \frac{\text{Vol}(\Omega)}{\text{Vol}(\mathcal{D})} \frac{1}{|\text{Jac}(j)(0)|^2} i\Theta(h_{\mathcal{D}})_0.$$

Proof. Let $w \in T_{\mathcal{D},0}$, and let $v = j_*(w)$. We are going to show that the inequality holds when applied to w .

We first gather a few objects allowing us to compute the left hand side. Accordingly to (5.1), we let $e, \sigma \in \mathcal{H}_{\Omega}$ be such that $\|e\|_{\mathcal{H}_{\Omega}} = \|\sigma\|_{\mathcal{H}_{\Omega}} = 1$ and $\|e_x\|_{h_{\Omega,x}} = 1$, $\sigma(x) = 0$, and we finally require that $i\Theta(h_{\Omega})(v, \bar{v}) = |df(v)|^2$, where $\sigma \stackrel{\text{loc}}{=} fe$ near x . Writing $\sigma = g dz_1 \wedge \dots \wedge dz_n$, and $e = g_0 dz_1 \wedge \dots \wedge dz_n$, we get the alternate expression

$$(5.4) \quad i\Theta(h_{\Omega})_x(v, \bar{v}) = \frac{|dg(v)|^2}{|g_0(x)|^2}.$$

Remark that since $\int_{\Omega} \frac{d\text{vol}_{\mathbb{C}^n}}{\text{Vol}(\Omega)} = 1$, we must have

$$(5.5) \quad |g_0(x)|^2 \geq \frac{1}{\text{Vol}(\Omega)}$$

since g_0 realizes the supremum of the evaluation function at x on $B(0,1) \subset \mathcal{H}_Y$.

To compute the right hand side, remark first that

$$j^*\sigma = \text{Vol}(\mathcal{D})^{1/2} (g \circ j) \text{Jac}(j) \left[\frac{dt_1 \wedge \dots \wedge dt_n}{\text{Vol}(\mathcal{D})^{1/2}} \right].$$

Denote by $e_{\mathcal{D}}$ the term between brackets. We have seen previously that $\|e_{\mathcal{D}}\|_{\mathcal{H}_{\mathcal{D}}} = \|e_{\mathcal{D},0}\|_{h_{\mathcal{D}}} = 1$. Moreover, since j is an open immersion, we have $\|j^*\sigma\|_{\mathcal{H}_{\mathcal{D}}} \leq 1$.

These two facts allow us to use (5.1) to bound the curvature of $h_{\mathcal{D}}$ from below, writing $j^*\sigma = f_{\mathcal{D}} e_{\mathcal{D}}$, with $f_{\mathcal{D}} = \text{Vol}(\mathcal{D})^{1/2} (g \circ j) \text{Jac}(j)$. We get

$$\begin{aligned} i\Theta(h_{\mathcal{D}})(w, \bar{w}) &\geq |df_{\mathcal{D}}(w)|^2 \\ &\geq |d(\text{Vol}(\mathcal{D})^{1/2} (g \circ j) \text{Jac}(j)) \cdot w|^2 \\ &= \text{Vol}(\mathcal{D}) |\text{Jac}(j)(0)| |d(g \circ j)(w)|^2 \\ &\geq \frac{\text{Vol}(\mathcal{D})}{\text{Vol}(\Omega)} |\text{Jac}(j)(0)|^2 \frac{|dg(v)|^2}{|g_0(x)|^2} \end{aligned}$$

where at the second line, we used the fact that $g(x) = 0$, and at the last line, we used (5.5). The last equation, combined with (5.4), allows us to end the proof. \square

Remark 5.2. In particular, if $r = d(x, \partial\Omega)$, we can apply the previous lemma to the open embedding of the ball $B(x, r) \hookrightarrow \Omega$, with $j = \text{Id}$. This gives, for any $v \in T_{\Omega, x}$:

$$\begin{aligned} i\Theta(h_\Omega)_x(v, \bar{v}) &\leq \frac{\text{Vol}(\Omega)}{\text{Vol}(B(x, r))} i\Theta(h_{B(x, r)})(v, \bar{v}) \\ &= \frac{\text{Vol}(\Omega)}{\text{Vol}(B(x, r))^2} \cdot \frac{n+1}{r^2} \|v\|_{\mathbb{C}^n}^2 \end{aligned}$$

using (5.3) and the fact that for $\mathcal{D} = B(x, r)$, we have $a_i^2 = \text{Vol}(B(x, r)) \cdot \frac{r^2}{n+1}$ for any i .

The next lemma will be used later on to estimate the curvature of the Bergman metric on subvarieties of Ω .

Lemma 5.3. *Assume that $0 \in \mathcal{D} \subset \Omega$, and that \mathcal{D} is centered at 0. Let Y be a complex manifold, and let $Y \xrightarrow{q} \Omega$ be a generically immersive holomorphic map passing through 0. Choose $y \in q^{-1}(0)$, and let $d_{\Omega, 0} = \max_{z \in \partial\Omega} d(0, z)$.*

Suppose that h_Y is defined at y . Then, we have:

$$i\Theta(h_Y)_y \geq \frac{\min_i a_i^2}{\text{Vol}(\Omega) d_{\Omega, 0}^2} q^*(i\Theta(h_\Omega)_0).$$

Proof. Fix a vector $v \in T_{Y, y}$, and let $w = q_*(v)$. We want to show that the inequality holds when applied to v . We may suppose that $w \neq 0$, the inequality being trivial otherwise.

Since h_Y is defined at y , there exists $\eta \in \mathcal{H}_Y$ such that $\|\eta\|_{\mathcal{H}_Y} = 1$, and $\|\eta_y\|_{h_{Y, y}} = 1$. Besides, by (5.2) and (5.3), we have

$$i\Theta(h_{\mathcal{D}})(w, \bar{w}) = |df(w)|^2,$$

with $f(z) = \frac{\sum_i \bar{w}_i z_i / a_i^2}{(\sum_i |w_i|^2 / a_i^2)^{1/2}} \cdot \text{Vol}(\mathcal{D})^{1/2}$. Note that by Cauchy-Schwarz inequality, we get the following upper bound:

$$|f(z)|^2 \leq \left(\sum_i \frac{|z_i|^2}{a_i^2} \right) \text{Vol}(\mathcal{D}) \leq \frac{d_{\Omega, 0}^2}{\min_i a_i^2} \text{Vol}(\mathcal{D}).$$

Define

$$\begin{aligned} g: \mathbb{C}^n &\longrightarrow \mathbb{C} \\ z &\longmapsto \frac{\min_i a_i}{\text{Vol}(\mathcal{D})^{1/2} d_{\Omega, 0}} f(z). \end{aligned}$$

Then, we have $\sup_Y |g \circ q| \leq 1$, so $\sigma = (g \circ q)\eta \in \mathcal{H}_Y$, and by (5.1), we get

$$\begin{aligned} i\Theta(h_Y)(v, \bar{v}) &\geq |d(g \circ q)(v)|^2 \\ &\geq \frac{\min_i a_i^2}{\text{Vol}(\mathcal{D}) d_{\Omega, 0}^2} |df(w)|^2 \end{aligned}$$

This shows that $i\Theta(h_Y)_y \geq \frac{\min_i a_i^2}{\text{Vol}(\mathcal{D}) d_{\Omega, 0}^2} i q^* \Theta(h_{\mathcal{D}})_0$. Using Lemma 5.1 with $j = \text{Id}_{\mathbb{C}^n}$, we see that $i\Theta(h_{\mathcal{D}})_0 \geq \frac{\text{Vol}(\mathcal{D})}{\text{Vol}(\Omega)} i\Theta(h_\Omega)_0$. This ends the proof. \square

Remark 5.4. When Ω is itself a bounded symmetric domain, we can take $\mathcal{D} = \Omega$. Up to scaling \mathbb{C}^n linearly, we can moreover assume that $a_i = 1$ for each i . Then, using the polydisk theorem (see [Mok89]), we easily see that $-\gamma = -\frac{1}{\text{Vol}(\Omega) d_{\Omega, 0}^2}$ is an upper bound for the holomorphic sectional curvature of the curvature form of the Bergman metric on T_Ω . In this situation, Ω is homogeneous, so the previous lemma gives

$$i\Theta(h_Y) \geq \gamma q^*(i\Theta(h_\Omega))$$

at any point where h_Y is defined.

We now make the following regularity assumption on the bounded domain Ω .

Assumption 5.5. The manifold Ω is a bounded domain admitting a *cocompact* discrete subgroup $\Gamma_0 \subset \text{Aut}(\Omega)$. Let $\mathfrak{U}_0 \subset \Omega$ be a compact fundamental domain for Γ_0 , and let $r_0 = d(\mathfrak{U}_0, \partial\Omega)$, $d_0 = \max_{x \in \mathfrak{U}_0, z \in \partial\Omega} d(x, z)$.

Under this assumption, we can obtain a uniform bound in Lemma 5.3, in terms of some constant depending on Γ_0 .

Definition 5.6. Under the hypothesis of Assumption 5.5, we introduce the following constant

$$C_0 = \frac{1}{d_0^2 \text{Vol}(\Omega)} \sup_{x \in \mathcal{D} \subset \Omega} (\min_i a_i^2),$$

where x runs among the points of \mathfrak{U}_0 , \mathcal{D} runs among the bounded symmetric domains centered at x and included in Ω , and the a_i are the constants associated to \mathcal{D} .

Remark that since $d(\mathfrak{U}_0, \partial\Omega) = r_0 > 0$, we can always take $\mathcal{D} = B(x, r_0)$ in the previous definition. Then, an easy computation shows that

$$C_0 \geq \frac{1}{n+1} \frac{\text{Vol}(B(0, r_0))}{\text{Vol}(\Omega)} \frac{r_0^2}{d_0^2}.$$

Note that we also have the trivial upper bound $d_0 \leq \text{diam}(\Omega)$.

Proposition 5.7. *Let Y be a complex manifold, and let $Y \xrightarrow{q} \Omega$ be a generically immersive holomorphic map. Suppose that h_Y is well defined at a generic point of Y . Then, we have*

$$i\Theta(h_Y) \geq C_0 q^*(i\Theta(h_\Omega))$$

in the sense of currents.

Proof. Since the right hand side is continuous on Y , it suffices to prove the inequality at any point y where h_Y is non-degenerate. The right hand side being invariant under the action of $\Gamma_0 \subset \text{Aut}(\Omega)$, we can let this lattice act on Ω and assume that $x = q(y) \in \mathfrak{U}_0$. One can now apply Lemma 5.3 to any bounded symmetric domain included in Ω and centered at x ; this concludes the proof of the Proposition. \square

5.3. A uniform curvature inequality. We keep working under the Assumption 5.5 on Ω , and we keep using the symbols $\Gamma_{\widehat{V}}$, \mathfrak{U}_0 and C_0 , with the same meaning as before, cf. Section 4 and Figure 2 that we reproduce below as Figure 3 for the reader's convenience. In particular, the

$$\begin{array}{ccccc}
 & & \widehat{V} & \xrightarrow{r} & \widehat{X} \\
 & & \downarrow \sigma & & \downarrow \pi \\
 \widehat{Z} & \xrightarrow{\sigma_Z} & Z & & \overline{X} \\
 & & \downarrow \rho & & \downarrow q \\
 & & \widetilde{V} & \xrightarrow{\iota} & \Omega \\
 & & \uparrow \ell & & \uparrow p
 \end{array}$$

FIGURE 3.

map $q : V \rightarrow \overline{X}$ is a generically immersive map and $\sigma : \widehat{V} \rightarrow V$ is a suitable modification making the diagram commutative. In the following, we will assume that V is a weakly pseudoconvex m -dimensional Kähler manifold. This means that there exists a smooth plurisubharmonic exhaustive function $\psi : V \rightarrow \mathbb{R}$. We want to show that the \mathbb{Q} -line bundle $K_{\widehat{V}} + \Delta_{\widehat{V}}$ admits a natural singular metric with positive curvature. We first make the following remark, which follows directly from Proposition 5.7.

Lemma 5.8. *Suppose that the Bergman metric $h_{Z_{\text{reg}}}$ is well defined at a generic point of Z_{reg} . Then the Γ_V -invariant metric $h_{Z_{\text{reg}}}$ on $K_{Z_{\text{reg}}}$ has positive curvature, satisfying*

$$i\Theta(h_{Z_{\text{reg}}}) \geq C_0 j^*(i\Theta(h_\Omega)),$$

where $j := (i \circ \rho)|_{Z_{\text{reg}}} : Z_{\text{reg}} \rightarrow \Omega$ is the natural map.

The next lemma relies on an adaptation to the non-compact case of some classical arguments in Kähler geometry (see e.g. [DP04]). Let $\widehat{Z} \xrightarrow{\sigma_Z} Z$ be some resolution of singularities to be fixed later and let $f_V : \widehat{Z} \rightarrow V$ be defined as the composition $f_V := \sigma \circ f_{\widehat{V}} \circ \sigma_Z$.

Lemma 5.9. *Assume that V admits a Kähler metric ω , let $(V_i)_{i \in \mathbb{N}}$ be an exhaustive sequence of relatively compact open subsets of V and set $\widehat{Z}_i := f_V^{-1}(V_i)$. Then, for an adequate choice of desingularizations \widehat{V} and \widehat{Z} , each manifold \widehat{Z}_i admits a Kähler metric ω_i .*

Moreover, we can choose the metrics ω_i so that

$$(5.6) \quad \omega_i \xrightarrow{i \rightarrow +\infty} f_V^* \omega$$

where the convergence holds uniformly on compact subsets of \widehat{Z} .

Proof. We may replace \widehat{V} by a resolution of indeterminacies of the bimeromorphic map $V \dashrightarrow \widehat{V}$, to suppose that $\sigma : \widehat{V} \rightarrow V$ is obtained by a sequence of blow-ups along smooth centers. Remark that this sequence may be infinite; however, the centers project onto a locally finite family of subsets of V .

Let E be the exceptional divisor of σ , with irreducible components $E = \sum_{k \in \mathbb{N}} E_k$. A classical argument allows one to find smooth $(1,1)$ -forms $\theta_{E_k} \in c_1(E_k)$ with support in an arbitrarily small neighborhood of E_k and a sequence of positive numbers (a_k) such that the (locally finite) sum $\theta_E = \sum_k a_k \theta_k$ defines a $(1,1)$ -form on \widehat{V} which is negative definite along the fibers of σ . Fix now some $i \in \mathbb{N}$. Since V_i is relatively compact in V , for $\epsilon_i > 0$ small enough, the closed $(1,1)$ -form

$$\omega_{\widehat{V}_i} = \sigma^* \omega - \epsilon_i \theta_E$$

defines a Kähler metric on $\sigma^{-1}(V_i)$.

Now, let $\widehat{Z} \xrightarrow{\sigma_Z} Z$ be a resolution of singularities obtained by blowing-up smooth centers, and let $\widehat{p} : \widehat{Z} \rightarrow \widehat{V}$ be the induced map. We ask that the strict transform $F = \widehat{p}_*^{-1}(\Delta_{\widehat{V}})$ is a *disjoint* union of smooth hypersurfaces, and that F has simple normal crossings with the exceptional divisor E' of the map σ_Z . Using partitions of unity, we can easily construct a smooth function ϕ on \widehat{Z} so that $i\partial\bar{\partial}\phi$ is positive in the directions transverse to the ramification divisor F . As before, we also let $\theta_{E'} \in c_1(E')$ be negative definite along the fibers of σ_Z .

With these definitions, for $\epsilon'_i > 0$ small enough, the closed $(1,1)$ -form

$$\omega_i = \widehat{p}^* \omega_{\widehat{V}_i} + \epsilon'_i (i\partial\bar{\partial}\phi - \theta_{E'}),$$

defines a Kähler metric on \widehat{Z}_i .

For the second requirement to be satisfied, we just need to take ϵ_i and ϵ'_i decreasing to 0 as $i \rightarrow +\infty$. \square

The next proposition is an adaptation to the non-compact case of the main argument of [BD18]. It is the last step towards Theorem 5.1, which is the main result of this section.

Proposition 5.10. *Assume that V is a weakly pseudoconvex Kähler manifold. Then, we can choose \widehat{V} and \widehat{Z} so that \widehat{Z} is a weakly Bergman manifold.*

Remark 5.11. If V is compact Kähler, then \widehat{V} is complete Kähler and admits a generically immersive map towards the manifold of bounded type Ω , and hence the conclusion follows directly from [BD18].

Proof. Let $\psi : V \rightarrow \mathbb{R}$ be a smooth exhaustive plurisubharmonic function. For each $i \in \mathbb{N}$, we let $V_i = \psi^{-1}([0, i[)$, and we fix \widehat{Z} , \widehat{V} and $(\omega_i)_{i \in \mathbb{N}}$ as provided by Lemma 5.9. We will show that $i\Theta(h_{\widehat{Z}})$ is positive definite at a generic point of Z_{reg} . By definition of the Bergman metric, it suffices to show that the L^2 holomorphic m -forms on \widehat{Z} generate the 1-jets at any generic point of Z_{reg} .

Let $z \in Z_{\text{reg}}$ be a point belonging to the regular loci of the maps $j = \iota \circ \rho$ and $\sigma \circ f_{\widehat{V}}$. One picks a germ τ of holomorphic m -form at z (recall that $m = \dim Z$). Remark now that each \widehat{Z}_i is weakly pseudoconvex because V_i is weakly pseudoconvex and the natural maps $\widehat{Z}_i \rightarrow V_i$ are proper. Since each \widehat{Z}_i admits the Kähler metric ω_i , this allows us to use the L^2 -method on \widehat{Z}_i with ω_i (see [Dem12a, Theorem 6.1]).

To do this, we choose a cutoff function χ on \widehat{Z} , equal to 1 in a neighborhood of z , and with compact support $L \subset \widehat{Z}_{i_0} \cap \sigma_Z^{-1}(Z_{\text{reg}})$ for some $i_0 \geq 0$. Without loss of generality, one can assume that L is contained in the regular locus of $f_V : \widehat{Z} \rightarrow V$. Let us define

$$\begin{aligned} \varphi : \Omega &\longrightarrow \mathbb{R} \\ x &\longmapsto (n+1) \log |x - s(z)| + |x|^2. \end{aligned}$$

The function is psh on Ω , and $\widehat{\varphi} := \varphi \circ \iota \circ \rho \circ \sigma_Z$ is strictly psh at $\sigma_Z^{-1}(z)$. Note that both χ and φ are independent of i .

As explained above, we can apply the L^2 method on \widehat{Z}_i to deduce that there exists a smooth $(m, 0)$ -form f_i on each \widehat{Z}_i , satisfying

$$(5.7) \quad i^{m^2} \int_{\widehat{Z}_i} f_i \wedge \bar{f}_i e^{-\widehat{\varphi}} \leq \int_L |\bar{\partial}(\chi\tau)|_{\omega_i}^2 e^{-\widehat{\varphi}} dV_{\omega_i}.$$

Thanks to (5.6), and since the metric $f_V^* \omega$ is non-degenerate on the compact set $L \subset Z_{\text{reg}}$, the right hand side of the above equation is uniformly bounded by some constant C for any $i \geq i_0$. Since φ is bounded from above, this implies a uniform bound

$$(5.8) \quad \|f_i\|_{L^2(\widehat{Z}_i)}^2 = \int_{\widehat{Z}_i} i^{m^2} f_i \wedge \bar{f}_i \leq C e^{\sup_{\Omega} \varphi}.$$

The expression of φ is chosen so that the bound (5.7) implies that f_i has a vanishing 1-jet at z . Thus, for any i , $\eta_i = \chi\tau - f_i$ is a holomorphic m -form on \widehat{Z}_i with $\text{jet}(d\eta_i)_z = d\tau_z$ at z . Also, (5.8) provides a uniform bound

$$\sup_i \|\eta_i\|_{L^2(\widehat{Z}_i)}^2 \leq C'.$$

Thus, we can extract a sequence converging uniformly on compact subsets towards a holomorphic form η . This form η satisfies $d\eta_z = d\tau_z$. Also, by Fatou lemma, we have $\|\eta\|_{L^2(\widehat{Z})}^2 \leq C'$. Thus, η satisfies our requirements. \square

We are now ready to prove the main result of this section.

Theorem 5.1. *Let $q : V \rightarrow \overline{X}$ be a generically immersive map from a weakly pseudoconvex Kähler manifold V such that $q(V) \not\subset D \cup \text{Sing}(p)$.*

Provided that Assumption 5.5 is satisfied, we can choose \widehat{V} so that the \mathbb{Q} -line bundle $\mathcal{O}_{\widehat{V}}(K_{\widehat{V}} + \Delta_{\widehat{V}})$ admits a natural singular metric $g_{\widehat{V}}$ with positive curvature, satisfying

$$(5.9) \quad i\Theta(g_{\widehat{V}}) \geq C_0 (q \circ \sigma)^* i\Theta(g_X)$$

over $V \setminus (q^{-1}(D \cup \text{Sing}(p)))$.

Remark 5.12. In the particular case where V is a compact Kähler manifold, we recover to the case obtained in Theorem 3.1 : $K_{\widehat{V}} + \Delta_{\widehat{V}}$ is big by [Bou02].

Proof. Let $\widehat{V}^\circ \subset \widehat{V}$ be the étale locus of the cover $f_{\widehat{V}}$ and let $F := \widehat{V} \setminus \widehat{V}^\circ$. By purity of the branch locus, F has pure codimension one. Let $n_i \in \mathbb{N}^* \cup \{\infty\}$ be the covering multiplicity attached to an irreducible component F_i of F and let $\Delta_{\widehat{V}} := \sum_i (1 - \frac{1}{n_i}) F_i$.

Since the Bergman metric $h_{Z_{\text{reg}}}$ is invariant under the group Γ_V , it descends to define a singular metric $g_{\widehat{V}}$ on the \mathbb{Q} -line bundle $K_{\widehat{V}} + \Delta_{\widehat{V}}$ with positive curvature by the same arguments as those provided in the proof of Theorem 2.1 where compactness plays no role. A priori, $g_{\widehat{V}}$ is only defined on $f_{\widehat{V}}(Z_{\text{reg}})$ but since that open set has complement whose codimension is at least two in \widehat{V} , the metric extends canonically across $f_{\widehat{V}}(Z_{\text{sing}})$.

It remains to see that the curvature of $g_{\widehat{V}}$ satisfies the required lower bound (5.9). Outside $\text{Sing}(p) \cup D$, the maps $p : \Omega \rightarrow X$ and $f_{\widehat{V}} : Z \rightarrow \widehat{V}$ are étale covers. In particular, one has on that locus an equality of smooth forms

$$\Theta(h_\Omega) = p^* \Theta(g_X) \quad \text{and} \quad \Theta(h_{Z_{\text{reg}}}) = f_{\widehat{V}}^* \Theta(g_{\widehat{V}})$$

and the differential of $f_{\widehat{V}}$ induces an isomorphism $f_{\widehat{V}}^* : \Omega_{\widehat{V},y}^{1,1} \xrightarrow{\sim} \Omega_{Z,x}^{1,1}$ whenever $f_{\widehat{V}}(x) = y$. By commutativity of the diagram in Figure 3, one has that $j^* \Theta(h_\Omega) = f_{\widehat{V}}^* (q \circ \sigma)^* \Theta(h_X)$ and therefore, (5.9) follows from Lemma 5.8. \square

Remark 5.13. As the reader will easily see, if we drop Assumption 5.5 (in particular if we only assume that Ω is a manifold of bounded type), the same proof shows that $K_{\widehat{V}} + \Delta_{\widehat{V}}$ admits a singular metric with positive curvature, but we cannot obtain the bound (5.9) anymore.

5.4. Statement of the criterion. In this section, we will state a hyperbolicity criterion for X , assuming it is now a *compact, non-necessarily smooth* quotient. The precise assumption is as follows.

Assumption 5.14. Under the hypotheses of Assumption 3.1, we assume moreover that X itself is a *compact* complex space, i.e. $\overline{X} = X$.

We resume the notations of the previous section.

Theorem 5.2. *Assume that $X = \overline{X}$ is as in Assumption 5.14, and let $\pi : \widehat{X} \rightarrow X$ be a projective resolution of singularities of X , which exists accordingly to Lemma 2.10. Let $\alpha > \frac{1}{C_0}$, and assume that the \mathbb{Q} -divisor*

$$L_\alpha = \pi^*(K_X + \Delta_X) - \alpha \Delta_{\widehat{X}}$$

is effective, where Δ_X is the covering divisor associated to $p : \Omega \rightarrow X$. Then

- (1) *any subvariety $W \subseteq X$ such that $W \not\subseteq \pi(\mathbb{B}(L_\alpha)) \cup \text{Sing}(p)$ is of general type.*
- (2) *any entire curve $f : \mathbb{C} \rightarrow X$ has its image included in $\pi(\mathbb{B}(L_\alpha)) \cup \text{Sing}(p)$.*

Proof. Let us prove first the statement concerning subvarieties. Suppose that $W \subset X$ is a subvariety as in the theorem, and let V be a resolution of singularities of W . Since the natural map $V \xrightarrow{q} \overline{X}$ is generically immersive, the relative construction of Section 4.2 can be applied, which yields a smooth bimeromorphic model \widehat{V} of V , which we can assume to be projective since \widehat{X} is, and a map $r : \widehat{V} \rightarrow \widehat{X}$ as in Figure 2.

By Theorem 5.1, the \mathbb{Q} -divisor $\mathcal{O}(K_{\widehat{V}} + \Delta_{\widehat{V}})$ admits a metric with positive curvature, and it is controlled as in (5.9) on $(\pi \circ r)^{-1}(X \setminus \text{Sing}(p))$. By assumption, $W \not\subseteq \mathbb{B}(L_\alpha)$, so for m large enough, there exists a section

$$\sigma \in H^0(\widehat{X}, m(\pi^*(K_X + \Delta_X) - \alpha \Delta_{\widehat{X}}))$$

such that $\sigma|_W \neq 0$.

We denote by ϕ_X (resp. $\phi_{\widehat{V}}$) the psh weight on $K_X + \Delta_X$ (resp. $K_{\widehat{V}} + \Delta_{\widehat{V}}$) associated to the metric g_X (resp. $g_{\widehat{V}}$). Next, we introduce the canonical singular weights $\phi_{\Delta_{\widehat{X}}}$ (resp. $\phi_{\Delta_{\widehat{V}}}$) on $\mathcal{O}_{\widehat{X}}(\Delta_{\widehat{X}})$ (resp. $\mathcal{O}_{\widehat{V}}(\Delta_{\widehat{V}})$) whose curvature current is $[\Delta_{\widehat{X}}]$ (resp. $[\Delta_{\widehat{V}}]$).

Because $p^*\phi_X$ is the weight associated to the Bergman metric on Ω , the weight ϕ_X is locally bounded. One introduces the quantity

$$F := |\sigma|^2 e^{-m\pi^*\phi_X} e^{m\alpha\phi_{\Delta_{\widehat{X}}}},$$

it is a *function* on \widehat{X} . Therefore, for any positive number $\beta > 0$, the quantity

$$(5.10) \quad e^{-\phi} := (r^*F)^{-\beta/m} e^{-\phi_{\widehat{V}} + \phi_{\Delta_{\widehat{V}}}}$$

defines a singular hermitian metric on $K_{\widehat{V}}$. The first item in the Theorem is a consequence of the following claim thanks to standard results in pluripotential theory together with [Bou02].

Claim 5.15. *For $\beta \in (\frac{1}{\alpha}, C_0)$, the following two properties hold:*

(i) *The weight ϕ is locally bounded above*

(ii) *The weight ϕ is smooth and strictly psh on $(\pi \circ r)^{-1}(X \setminus \text{Sing}(p))$.*

Proof of Claim 5.15. To prove (i), one observes that

$$\phi = \left(\phi_{\widehat{V}} + \frac{\beta}{m} \log |r^*\sigma|^2 \right) - \beta \cdot (\pi \circ r)^*\phi_X + (\alpha\beta \cdot \phi_{r^*\Delta_{\widehat{X}}} - \phi_{\Delta_{\widehat{V}}})$$

can be decomposed as the sum of two psh weights, one locally bounded weight and another weight which is psh thanks to Proposition 4.9 together with the fact that $\alpha\beta > 1$. In order to prove (ii), one computes the curvature of the weight ϕ on $(\pi \circ r)^{-1}(X \setminus \text{Sing}(p))$ as follows

$$\begin{aligned} dd^c\phi &= i\Theta(g_{\widehat{V}}) - \beta \cdot (\pi \circ r)^*i\Theta(g_X) + \frac{\beta}{m} dd^c \log |r^*\sigma|^2 \\ &\geq i\Theta(g_{\widehat{V}}) - \beta \cdot (\pi \circ r)^*i\Theta(g_X) \\ &\geq (C_0 - \beta) \cdot (\pi \circ r)^*i\Theta(g_X) \end{aligned}$$

where the last inequality follows from the inequality (5.9) in Theorem 5.1. \square

The proof of the second point is very similar: we just have to perform the previous steps with $V = \mathbb{C}$ in a slightly more explicit manner, and then use the Ahlfors-Schwarz lemma (see e.g. [Dem12b]).

Suppose then that there exists a non-constant holomorphic map $f : \mathbb{C} \rightarrow X$ such that $f(\mathbb{C}) \not\subset \pi(\mathbb{B}(L_\alpha)) \cup \text{Sing}(p)$. Now, if we perform the relative orbifold construction with $V = \mathbb{C}$, we see that $\widehat{V} \cong \mathbb{C}$, since this manifold admits a bimeromorphic map onto \mathbb{C} .

Moreover, since $\iota : \widehat{V} \rightarrow \Omega$ is non constant, we see that the universal cover of \widehat{V} must be isomorphic to the disk, and thus $i\Theta(h_{\widehat{V}}) = h_{\widehat{V}}^{-1}$. Pushing forward to \widehat{V} , we get that

$$i\Theta(g_{\widehat{V}}) = g_{\widehat{V}}^{-1}$$

in restriction to the regular locus $V^\circ = \widehat{V} \setminus (f \circ \sigma)^{-1}(\text{Sing}(p))$.

Construct now the metric $h = e^{-\phi}$ on $\mathbb{C} = \widehat{V}$ using (5.10). By Claim 5.15 (ii), valid for V pseudoconvex Kähler, we see that there exists a constant $\delta > 0$ such that $i\Theta(h) \geq \delta i\Theta(g_{\widehat{V}})$ in restriction to V° . Now, we have, again in restriction to V° :

$$\Theta(g_{\widehat{V}}) = g_{\widehat{V}}^{-1} \geq \frac{1}{\sup_{\widehat{X}}(r^*F)^{\frac{\beta}{m}}} h^{-1}$$

Thus, in restriction to V° , one gets

$$(5.11) \quad i\Theta(h) \geq Ch^{-1}$$

where $C = \frac{\delta}{\sup_{\widehat{X}}(r^*F)^{\frac{\beta}{m}}}$.

Finally, we see as in Claim 5.15 (i) above that ϕ is locally bounded above near $\widehat{V} \setminus V^\circ$ and therefore $h = e^{-\phi}$ induces a positively curved metric on $K_{\widehat{V}} \simeq \mathcal{O}_{\mathbb{C}}$. This implies that (5.11) holds *everywhere* on $\widehat{V} = \mathbb{C}$ in the sense of currents. This is however absurd because of the Ahlfors-Schwarz lemma, cf. [Dem95, Lem. 3.2]. \square

REFERENCES

- [BD18] Sébastien Boucksom and Simone Diverio, *A note on Lang’s conjecture for quotients of bounded domains*, arXiv:1809.02398, 2018. ↑ [1](#), [2](#), [3](#), [4](#), [5](#), [7](#), [10](#), [11](#), [21](#)
- [Bou02] Sébastien Boucksom, *On the volume of a line bundle*, Internat. J. Math. **13** (2002), no. 10, 1043–1063. ↑ [5](#), [9](#), [22](#), [24](#)
- [BP16] Ingrid Bauer and Roberto Pignatelli, *Product-quotient surfaces: new invariants and algorithms*, Groups Geom. Dyn. **10** (2016), no. 1, 319–363. MR 3460339 ↑ [12](#)
- [Bru16a] Yohan Brunebarbe, *A strong hyperbolicity property of locally symmetric varieties*, arXiv:1606.03972 (2016). ↑ [4](#)
- [Bru16b] ———, *Symmetric differentials and variations of Hodge structures*, J. Reine Angew. Math. (2016). ↑ [4](#)
- [Cad16] Benoît Cadorel, *Symmetric differentials on complex hyperbolic manifolds with cusps*, arXiv:1606.05470 (2016). ↑ [4](#)
- [Cad18] ———, *Subvarieties of quotients of bounded symmetric domains*, arxiv.org/abs/1809.10978 (2018). ↑ [4](#)
- [Cam04] Frédéric Campana, *Orbifolds, special varieties and classification theory*, Ann. Inst. Fourier (Grenoble) **54** (2004), no. 3, 499–630. ↑ [3](#)
- [CG72] James Carlson and Phillip Griffiths, *A defect relation for equidimensional holomorphic mappings between algebraic varieties.*, Ann. Math. **95** (1972), 557–584. ↑ [11](#)
- [CRT17] Benoît Cadorel, Erwan Rousseau, and Behrouz Taji, *Hyperbolicity of singular spaces*, to appear in J. Éc. polytech. Math., 2017. ↑ [4](#), [5](#), [13](#)
- [CZ02] Bo-Yong Chen and Jin-Hao Zhang, *The Bergman metric on a Stein manifold with a bounded plurisubharmonic function*, Trans. Amer. Math. Soc. **354** (2002), no. 8, 2997–3009. MR 1897387 ↑ [5](#)
- [Dem82] Jean-Pierre Demailly, *Estimations L^2 pour l’opérateur $\bar{\partial}$ d’un fibré vectoriel holomorphe semi-positif au-dessus d’une variété Kählerienne complète*, Ann. Sci. Éc. Norm. Supér.(4) **15** (1982), 457–511. ↑ [11](#)
- [Dem92] Jean-Pierre Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. **1** (1992), no. 3, 361–409. MR 1158622 (93e:32015) ↑ [8](#)
- [Dem95] ———, *Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials*, Proceedings of the Symposia in Pure Maths., (1995), no. 62.2. ↑ [24](#)
- [Dem12a] Jean-Pierre Demailly, *Complex Analytic and Differential Geometry*, September 2012, OpenContent Book, freely available from the author’s web site. ↑ [22](#)
- [Dem12b] ———, *Hyperbolic algebraic varieties and holomorphic differential equations*, Acta Math. Vietnam. **37** (2012), no. 4, 441–512. ↑ [24](#)
- [DP04] Jean-Pierre Demailly and Mihai Păun, *Numerical characterization of the Kähler cone of a compact Kähler manifold*, Ann. of Math. (2) **159** (2004), no. 3, 1247–1274. MR 2113021 ↑ [8](#), [21](#)
- [DT19] Simone Diverio and Stefano Trapani, *Quasi-negative holomorphic sectional curvature and positivity of the canonical bundle*, J. Differential Geom. **111** (2019), no. 2, 303–314. ↑ [4](#)
- [GG80] Mark Green and Phillip Griffiths, *Two applications of algebraic geometry to entire holomorphic mappings*, The Chern Symposium 1979, Proc. Internal. Sympos. Berkeley, CA, 1979 (New York), Springer-Verlag, 1980, pp. 41–74. ↑ [1](#)
- [Gri71] Phillip A. Griffiths, *Complex-analytic properties of certain zariski open sets on algebraic varieties*, Annals of Mathematics **94** (1971), no. 1, 21–51. ↑ [2](#), [11](#)
- [Gue18] Henri Guenancia, *Quasi-projective manifolds with negative holomorphic sectional curvature*, arXiv:1808.01854 (2018). ↑ [2](#), [4](#)
- [Hah78] Kyong T. Hahn, *Inequality between the Bergman metric and Caratheodory differential metric*, Proc. Amer. Math. Soc. **68** (1978), 193–193. ↑ [5](#), [17](#)
- [Kob59] Shoshichi Kobayashi, *Geometry of bounded domains*, Trans. Amer. Math. Soc. **92** (1959), 267–290. MR 0112162 ↑ [5](#)
- [Kob98] ———, *Hyperbolic complex spaces*, Grundlehren der Mathematischen Wissenschaften, vol. 318, Springer-Verlag, Berlin, 1998. ↑ [9](#), [17](#)
- [Kol07] János Kollár, *Lectures on resolution of singularities*, Princeton University Press, 2007. ↑ [16](#)
- [Lan86] Serge Lang, *Hyperbolic and Diophantine analysis*, Bull. Amer. Math. Soc. (N.S.) **14** (1986), no. 2, 159–205. MR 828820 ↑ [1](#)
- [Mok89] Ngaiming Mok, *Metric rigidity theorems on hermitian locally symmetric manifolds, series in Pure Mathematics*, vol. 6, World Scientific Publishing Company, 1989. ↑ [19](#)
- [Rou16] Erwan Rousseau, *Hyperbolicity, automorphic forms and Siegel modular varieties*, Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), no. 1, 249–255. ↑ [4](#)
- [Tai82] Yung-Sheng Tai, *On the Kodaira dimension of the moduli space of abelian varieties.*, Inventiones mathematicae **68** (1982), 425–440. ↑ [5](#)
- [Wei86] Rainer Weissauer, *Untervarietäten der Siegelschen Modulmannigfaltigkeiten von allgemeinem Typ*, Math. Ann. **275** (1986), no. 2, 207–220. MR 854007 ↑ [5](#)

- [WY16] Damir Wu and Shing-Tung Yau, *Negative holomorphic curvature and positive canonical bundle*, *Inventiones mathematicae* **204** (2016), no. 2, 595–604. ↑ 4

INSTITUT DE MATHÉMATIQUES DE TOULOUSE; UMR 5219, UNIVERSITÉ DE TOULOUSE; CNRS, UPS, 118 ROUTE DE NARBONNE, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `benoit.cadorel@univ-lorraine.fr`

Current address: Institut Élie Cartan de Lorraine, UMR 7502, Université de Lorraine, Site de Nancy, B.P. 70239, F-54506 Vandoeuvre-lès-Nancy Cedex

DIPARTIMENTO DI MATEMATICA “GUIDO CASTELNUOVO”, SAPIENZA UNIVERSITÀ DI ROMA, PIAZZALE ALDO MORO 5, I-00185 ROMA, ITALY

E-mail address: `diverio@mat.uniroma1.it`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE; UMR 5219, UNIVERSITÉ DE TOULOUSE; CNRS, UPS, 118 ROUTE DE NARBONNE, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `henri.guenancia@math.cnrs.fr`