



**HAL**  
open science

## Impact of galvanic vestibular stimulation on mood

Florane Pasquier, Pierre F Denise, Antoine Gauthier, Nicolas Bessot, Gaëlle Quarck

► **To cite this version:**

Florane Pasquier, Pierre F Denise, Antoine Gauthier, Nicolas Bessot, Gaëlle Quarck. Impact of galvanic vestibular stimulation on mood. ACAPS, Oct 2017, Dijon, France. hal-02135556

**HAL Id: hal-02135556**

**<https://hal.science/hal-02135556>**

Submitted on 21 May 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Impact of galvanic vestibular stimulation on mood

F. Pasquier, P. Denise, A. Gauthier, N. Bessot, G. Quarck  
Unité COMETE UMR 1075 UNICAEN/INSERM, 14032 Caen, France

## Introduction

Galvanic Vestibular Stimulation (GVS) is a variant of transcranial direct current (tDCS) used to treat depression. In GVS, the electrodes are attached to the mastoids process behind the ears in order to stimulate the vestibular system including vestibular cortex. In addition to its role in gaze stabilization and postural control, the vestibular system has an influence into dimensions of emotion processing, mental health and social cognition. Moreover, several studies demonstrated that vestibular stimulation has a modulating effect on mood, affective control, and especially in anxiety level.

### Objectives :

- Demonstrate the feasibility of our experimental protocol for clinical treatment (motion sickness).
- Test if the dose (duration) of stimulation has an impact on mood.

GVS Stimulation

Vestibular System



## Discussion

Our results confirm that GVS induces motion sickness symptoms. Vegetative disturbances rated by the participants appeared only for the double-dose session and provoked minor level of motion sickness symptoms.

Present study demonstrates also a significant decrease of anxiety level after a dose of GVS, in a student population. This result corroborates similar findings in the literature. It confirms that vestibular system consists of afferent and efferent connections with most of the brain structures involved in emotional processes, such as insula and prefrontal areas.

→ **Methodological perspectives**

→ **Chronobiological perspectives**

1

## Methodology

- **Participants :**  
22 students in sport sciences were recruited.  
Criteria of exclusion : vestibular disorders, psychiatric diagnoses.

2

- **Measures :**  
Motion sickness symptoms were assessed by Graybiel scale (Figure 1.A). To evaluate mood in particular anxiety, participants completed a Visual Analog Scale/VAS (Figure 1.B). Each test was completed before and after GVS.

A

Category	Pathognomonic 16 points	Major 8 points	Minor 4 points	Minimal 2 points	AQS* 1 point
Nausea syndrome	Vomiting or retching	Nausea? II, III	Nausea I	Epigastric discomfort	Epigastric awareness
Skin color		Pallor III	Pallor II	Pallor I	Flushing
Cold sweating		III	II	I	
Increased salivation		III	II	I	
Drowsiness		III	II	I	
Pain		III	II	I	Headache
Central nervous system					Dizziness: Eyes closed ≥ II Eyes open III

Levels of Severity Identified by Total Points Scored


Frank Sickness (S)	Severe Malaise (M III)	Moderate Malaise A (M IIA)	Moderate Malaise B (M IIB)	Slight Malaise (M I)
≥ 16 points	8-15 points	5-7 points	3-4 points	1-2 points

\*AQS = Additional qualifying symptoms. ?III = severe or marked, II = moderate, I = slight.



Figure 1 : Graybiel scale (A) and the visual analog scale for anxiety (B).

- **Experimental design :**  
3 randomized sessions (sham, dose, double-dose) of stimulation (NeuroConn®, Germany), the same day and the same hour for three consecutive weeks.  
The intensity was 1000 µA and the polarity was changed four or eight times according to the session.



## Results

