



**HAL**  
open science

# Rate constants of the reactions of O( 3 P) atoms with Br 2 and NO 2 over the temperature range 220-950 K

Yuri Bedjanian, Chaitanya Kalyan

► **To cite this version:**

Yuri Bedjanian, Chaitanya Kalyan. Rate constants of the reactions of O( 3 P) atoms with Br 2 and NO 2 over the temperature range 220-950 K. *International Journal of Chemical Kinetics*, 2019, 51 (7), pp.476-483. 10.1002/kin.21270 . hal-02136030

**HAL Id: hal-02136030**

**<https://hal.science/hal-02136030>**

Submitted on 12 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Rate Constants of the Reactions of O(<sup>3</sup>P) Atoms with Br<sub>2</sub> and NO<sub>2</sub> over the Temperature Range 220 – 950 K

YURI BEDJANIAN,\* CHAITANYA KALYAN

*Institut de Combustion, Aérodynamique, Réactivité et Environnement (ICARE), CNRS, 45071 Orléans Cedex 2, France*

**ABSTRACT:** The kinetics of the reactions of Br<sub>2</sub> and NO<sub>2</sub> with ground state oxygen atoms have been studied over a wide temperature range,  $T = 220 - 950$  K, using a low pressure flow tube reactor coupled with a quadrupole mass spectrometer:  $O + NO_2 \rightarrow NO + O_2$  (1) and  $O + Br_2 \rightarrow Br + BrO$  (2). The rate constant of reaction (1) was determined under pseudo-first order conditions, either monitoring the kinetics of O-atom or NO<sub>2</sub> consumption in excess of NO<sub>2</sub> or of the oxygen atoms, respectively:  $k_1 = (6.1 \pm 0.4) \times 10^{-12} \exp((155 \pm 18)/T)$  cm<sup>3</sup>molecule<sup>-1</sup>s<sup>-1</sup> (where the uncertainties represent precision at the  $2\sigma$  level, the estimated total uncertainty on  $k_1$  being 15% at all temperatures). The temperature dependence of  $k_1$ , found to be in excellent agreement with multiple previous low temperature data, was extended to 950K. The rate constant of reaction (2) determined under pseudo-first order conditions, monitoring the kinetics of Br<sub>2</sub> consumption in excess of O-atoms, showed upward curvature at low and high temperatures of the study and was fitted with the following three-parameter expression:  $k_2 = 9.85 \times 10^{-16} T^{1.41} \exp(543/T)$  cm<sup>3</sup>molecule<sup>-1</sup>s<sup>-1</sup> at  $T = (220 - 950)$  K, which is recommended from the present study with an independent of temperature conservative uncertainty of 15% on  $k_2$ .

**Keywords:** Oxygen atom, Br<sub>2</sub>, NO<sub>2</sub>, rate coefficient, temperature dependence.

\*Correspondence to: Yuri Bedjanian: Tel.: +33 238255474, e-mail: yuri.bedjanian@cnrs-orleans.fr  
Supporting Information is available in the online issue at [www.wileyonlinelibrary.com](http://www.wileyonlinelibrary.com).

## INTRODUCTION

The reaction of ground state atomic oxygen with nitrogen dioxide is of great atmospheric importance as a rate limiting step in catalytic cycle destroying odd oxygen ( $O+O_3$ ) in the middle stratosphere and affecting both the abundance and vertical profile of the stratospheric ozone [1]:



Besides the atmospheric importance, the reaction is often used in laboratory studies as a titration reaction of oxygen atoms and is included in  $NO_2$  sub-mechanism of kinetic models for combustion systems [2]. Hence, it is not surprising that kinetics of the reaction (1) has been studied very extensively in the past [3-14]. Currently, the rate constant of the  $O+NO_2$  reaction,  $k_1$ , can be considered as well established in the temperature range  $T = 220 - 425K$  [15]. At higher temperatures only three studies of the reaction are available [5,8,11]. However, the most extended measurements of  $k_1$  over the temperature range  $T = 297 - 1055K$  reported by Bemand *et al.* [8] are highly scattered and do not match the extrapolation of the Arrhenius expression for  $k_1$  based on the low temperature data [15]. In addition, the reaction rate constant at high temperatures has been recently challenged by appearance of a theoretical study [16], predicting U-shaped temperature dependence of  $k_1$  with rapid increase of the rate constant with temperature at  $T > 400 K$ . In this respect, the objective of the present work was to extend the measurements of  $k_1$  to high temperatures in order to compare them with previous experimental data and new theoretical predictions.

The reaction of oxygen atoms with  $Br_2$  was studied as a part of this work:



In fact, this reaction was used for detection of low concentrations of oxygen atoms *via* their chemical conversion to  $BrO$  radicals [17]. However, it is also of autonomous interest. Although this reaction is not of direct atmospheric importance, it is very often used in the laboratory as a source of  $BrO$  radicals, which is an important species in the atmospheric bromine/ozone chemistry [18]. The rate constant of the reaction (2),  $k_2$ , especially its temperature dependence, is highly uncertain. Thus, in four previous temperature dependent kinetic studies of the reaction [19-22], noticeable positive [19,20] and negative [22] dependence of  $k_2$  on temperature as well as independent of temperature  $k_2$  [21] were reported. In the present work, we have performed measurements of the reaction rate constant in an extended temperature range  $T = 220 - 950 K$ .

## EXPERIMENTAL


The rate constants of the title reactions were measured at total pressure of  $\approx 2$  Torr of He in a discharge flow reactor combined with a modulated molecular beam mass spectrometer [17,23,24]. Two flow reactors were used. The first one, used at low temperatures ( $T = 220 - 317$  K), consisted of a Pyrex tube (45 cm length and 2.4 cm i.d.) with a jacket for the thermostated liquid circulation (ethanol). The inner surface of the reactor as well as of the movable injector of O atoms was coated with halocarbon wax (HW) in order to minimize their heterogeneous loss. The second flow reactor, operating at high temperatures ( $T = 294 - 950$  K), consisted of an electrically heated Quartz tube (45 cm length and 2.5 cm i.d.) with water-cooled extremities (Fig. 1S, Supporting Information) [25]. Kinetic measurements were carried out in the part of the reactor where temperature gradient along the reaction zone (measured with a thermocouple inserted in the reactor through the movable injector [25]) was less than 1% (Fig. 1S, Supporting Information).

Oxygen atoms were generated from the microwave discharge in  $O_2/He$  mixtures. Two methods were used for the detection of oxygen atoms. At high concentrations of O atoms (experiments in excess of O atoms over  $Br_2$  or  $NO_2$ ), they were detected at their parent peak at  $m/z=16$  ( $O^+$ ). The absolute concentrations of the atoms were determined from the fraction of  $O_2$  dissociated in the microwave discharge ( $\Delta[O_2] = 2[O]$ ). At low concentrations of oxygen atoms (experiments in excess of  $NO_2$  over oxygen atoms), another method for their detection was preferred to the direct one at  $m/z = 16$  due to significant residual signal at this mass.  $Br_2$  was added at the end of the reactor (6 cm upstream of the sampling cone, Fig. 1S) in order to transform O atoms to BrO detected at  $m/z=95/97$  as  $BrO^+$ :



$$k_2 = 9.85 \times 10^{-16} T^{1.41} \exp(543/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \quad (T = 220-950 \text{ K, this work}).$$

The concentrations of  $Br_2$  ( $\sim 10^{14}$  molecule  $\text{cm}^{-3}$ ) were high enough to ensure rapid consumption of oxygen atoms. The impact of the recombination reaction of BrO radicals (3) and reaction (4) of BrO with NO (product of reaction (1)) on the measurements of [BrO] was insignificant due to relatively low concentrations of these species (initial concentration of oxygen atoms was  $\leq 4 \times 10^{11}$  molecule  $\text{cm}^{-3}$ ):



$$k_3 = 1.5 \times 10^{-12} \exp(230/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ over } T = 220-348 \text{ K [15].}$$



$$k_4 = 8.8 \times 10^{-12} \exp(260/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ over } T = 224\text{-}425\text{K [15].}$$

Reaction of BrO with NO<sub>2</sub> was also slow under experimental conditions of the study:



$$k_5 < 2.3 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } T \geq 220\text{K and } P = 2 \text{ Torr [15].}$$


Absolute concentrations of BrO radicals were determined using their chemical conversion to NO<sub>2</sub> through reaction (4) in excess of NO with subsequent detection of NO<sub>2</sub> formed ( $[\text{BrO}]_0 = [\text{NO}_2]_{\text{formed}}$ ). For stable species (NO<sub>2</sub>, NO, Br<sub>2</sub>, O<sub>2</sub>), the mass spectrometer was calibrated by introducing their known concentrations into the reactor from their calibrated mixtures with helium. The uncertainty on the absolute concentrations of the stable species was  $\leq 10\%$ . All species were detected by mass spectrometry at their parent peaks:  $m/z = 46$  (NO<sub>2</sub><sup>+</sup>), 30 (NO<sup>+</sup>), 158/160/162 (Br<sub>2</sub><sup>+</sup>), 95/97 (BrO<sup>+</sup>), 32 (O<sub>2</sub><sup>+</sup>), 16 (O<sup>+</sup>).

## RESULTS AND DISCUSSION

### Reaction O + NO<sub>2</sub>


*Measurements of  $k_1$ .* The reaction rate constant was determined under pseudo-first order conditions either from kinetics of O-atom consumption ( $[\text{O}]_0 \leq 4 \times 10^{11} \text{ molecule cm}^{-3}$ ) in excess of NO<sub>2</sub> or from NO<sub>2</sub> decays ( $[\text{NO}_2]_0 \leq 5 \times 10^{11} \text{ molecule cm}^{-3}$ ) monitored in excess of oxygen atoms. The ranges of concentrations of the respective excess reactants are shown in Table I. The flow velocity in the reactor was in the range (1950-2300) cm s<sup>-1</sup>. The concentrations of O and NO<sub>2</sub> were simultaneously measured as a function of reaction time. Generally, the consumption of excess reactant was negligible (within a few percent), although reaching up to 25 % in a few kinetic runs at lowest concentrations of the excess reactant: in all cases the average concentration of excess reactant along the reaction zone was used in calculation of the reaction rate constant. This simplified procedure did not significantly affect the calculated rate constants, which could also be derived from the simulation of the experimental NO<sub>2</sub> (or O-atom) kinetics using the observed temporal profiles of excess reactant. The values of  $k_1$  obtained by the two methods were in agreement within 4%.

Examples of O-atom and NO<sub>2</sub> decays observed at different concentrations of NO<sub>2</sub> and oxygen atoms, respectively, are shown in Figs. 1 and 2S (Supporting Information).




**Figure 1** Reaction O + NO<sub>2</sub>: example of the exponential decays of O atoms in excess of NO<sub>2</sub> at  $T = 950$  K.

The pseudo-first order rate constants,  $k_1' = k_1[\text{NO}_2] + k_w$ , determined from the slopes of the straight lines, like those in Figs. 1 and 2S, are shown in Fig. 2 as a function of NO<sub>2</sub> concentration. The intercepts in Fig. 2,  $k_w$ , were in the range (1 - 9) s<sup>-1</sup>, in good agreement with the rate of O-atom decay measured in separate experiments in the absence of NO<sub>2</sub> in the reactor. Diffusion corrections were applied to all the measured values of  $k_1'$  in order to take into account the axial and radial gradient of O-atom concentration in the flow tube:  $k_1' = k_{1' \text{ measured}} \times (1 + k_{1' \text{ measured}} \times D/v^2 + k_{1' \text{ measured}} \times r^2/48D)$ , where  $D$  is the molecular diffusion coefficient,  $v$  is the flow velocity and  $r$  is the radius of the flow tube [26]. The diffusion coefficient of O in He was calculated using  $D_0 = 770 \text{ Torr cm}^2 \text{ s}^{-1}$  at  $T = 298\text{K}$  [27] and assuming  $T^{1.75}$ -dependence on temperature. Corrections were generally less than 9 %, and only in three kinetic runs at  $T = 950$  K they were somewhat higher, up to 16%. Examples of the dependence of pseudo-first order rate constant  $k_1' = k_1[\text{O}]$  measured from NO<sub>2</sub> decays on concentration of excess reactant, oxygen atom, are shown in Fig. 3. Diffusion corrections on  $k_1'$  calculated in this case using the diffusion coefficient of NO<sub>2</sub> in He,  $D_0 = 550 \times (T/298)^{1.75} \text{ Torr cm}^2 \text{ s}^{-1}$  (estimated with Fuller's method) [28], were less than 8%.




**Figure 2** Reaction O + NO<sub>2</sub>: dependences of pseudo-first order rate constant of O-atom decay on concentration of NO<sub>2</sub> measured at  $T = 220, 298$  and  $950$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .



**Figure 3** Reaction O + NO<sub>2</sub>: dependences of pseudo-first order rate constant of NO<sub>2</sub> decay on concentration of oxygen atoms measured at  $T = 266$  and  $950$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .

The slopes of the straight lines in Figs. 2 and 3 provide the values of  $k_1$  at respective temperatures. All the results obtained for  $k_1$  in the present study are shown in Table I and Fig.

4. One can note very good agreement between the values for  $k_1$  obtained from kinetics of O-atom and  $\text{NO}_2$  consumption. The combined uncertainty on the measurements of  $k_1$  was estimated to be of nearly 15 %, by adding in quadrature (square root of the sum of squares) the statistical error ( $\leq 4$  %) and those on the measurements of the flows (5%), pressure (2%), temperature (1%), and absolute concentrations (10%) of  $\text{NO}_2$  and oxygen atoms.



**Figure 4** Reaction  $\text{O} + \text{NO}_2$ : summary of the measurements of the temperature dependence of the reaction rate constant. Error bars correspond to estimated 15 % uncertainty on the measurements of  $k_1$  in the present work. Uncertainties on  $k_1$  reported in previous studies are omitted for clarity.

*Comparison with previous data.* Figure 4 summarizes the results of temperature dependent measurements [5-14] of the rate constant of reaction (1). Westenberg and deHaas [5] have measured the reaction rate constant at  $T = 297$ - $543$  K in a fast-flow reactor combined with electron spin resonance (ESR) spectroscopy. Davis *et al.* [6], using flash photolysis-resonance fluorescence technique have reported temperature independent  $k_1 = (9.12 \pm 0.44) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  over the temperature range  $T = 230$ - $339$  K. Close results were obtained by Slinger *et al.* [7] who measured  $k_1 = 9.3$  and  $10.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  and  $T = 296$  and  $240$  K, respectively, using pulsed photolysis and the  $\text{O} + \text{NO} + \text{M}$  chemiluminescent emission for detection of O-atoms. In the only high temperature study of reaction (1), using atomic resonance fluorescence, Bemand *et al.* [8] measured  $k_1$  at 16 temperatures between 298 and 1055 K (average values of  $k_1$  at each temperature are shown in Fig. 4) and observed a slight negative temperature dependence of the rate constant. Small negative activation factor ( $E_a/R$ ) of the reaction (1) has been confirmed in all subsequent work:  $142 \pm 23$  K over the



temperature range 220-387 K, in a discharge flow study with chemiluminescent detection of O-atoms [9];  $202 \pm 27$  K over  $T = 199$ -357 K, using flash photolysis with chemiluminescent reaction of O-atoms with NO [10];  $209 \pm 35$  K over  $T = 220$ -412 K, using pulsed laser photolysis of NO<sub>2</sub> to produce oxygen atoms and time-resolved vacuum UV resonance fluorescence for their detection [12];  $273 \pm 18$  K over  $T = 221$ -425 K, with laser flash photolysis-resonance fluorescence technique [13] and  $165 \pm 10$  K over  $T = 216$ -413 K by discharge-flow resonance fluorescence technique [14]. The Arrhenius expression recommended by the JPL panel (black continuous line in Fig. 4) [15],  $k_1 = 5.1 \times 10^{-12} \exp(210/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , is based on the measurements reported in refs. [7,9,10,12,13]. A least-squares fit to  $k_1$  data from the present work (red line in Fig. 4) provides the following Arrhenius expression:

$$k_1 = (6.1 \pm 0.4) \times 10^{-12} \exp((155 \pm 18)/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 220-950 K and with  $2\sigma$  statistical uncertainties on the fit. Although this expression is somewhat different from recommended by JPL panel [15], the absolute values of  $k_1$  measured in the present study are in excellent agreement with the most recent measurements [9,10,12-14] of the rate constant and current recommendation [15] at  $T \leq 425$  K. At higher temperatures, present values of  $k_1$  are higher than those reported by Bemand *et al.* [8] by a factor of 1.3-1.9. One can also note that possible downward curvature of the Arrhenius plot at temperatures above 298 K discussed in ref. [14] is not confirmed by the present measurements.


In a recent theoretical study, the rate constant of the reaction (1) was computed over a temperature range 150–1500 K [16]. The values of  $k_1$  calculated at two different levels of theory (B3LYP/aug-cc-pVTZ and MP2/6-311+G(d,p)) are shown in Fig. 4. The authors concluded that computed values of  $k_1$  are in "an excellent agreement (within 5–10%) with all the available experimental data particularly at B3LYP method". As one can see from Fig. 4, this statement is valid only in a rather narrow temperature range. In fact, the temperature dependence of  $k_1$  predicted by simulations at the B3LYP level of theory does not reproduce the experimental results. Shiekh *et al.* [16] noted that the nonlinear behavior of  $k_1$  calculated with B3LYP method at higher temperatures may be due to the limitations of VPT2 model in this temperature range. On the other hand, the calculations with MP2 method seem to better reproduce temperature dependence of  $k_1$  although they are shifted by a factor of 1.4-1.7 with respect to the experimental data. Zuev and Starikovskii [11] in their shock tube study of thermal decomposition of N<sub>2</sub>O reported  $k_1 = (1.0 \pm 0.25) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $T = 1780$ -

2300 K. This value was derived from an analysis of concentration profiles of N<sub>2</sub>O, NO<sub>2</sub> and NO within a complex kinetic mechanism including nine reversible reactions. Although the indirect measurement of  $k_1$  [11] is by a factor of  $\approx 1.5$  higher than the extrapolation of the present data to these temperatures, the results from two studies are consistent in the range of reported uncertainties. On the other hand, our experimental data on  $k_1$  measured at  $T \leq 950\text{K}$  do not exclude the possible upward curvature in the temperature dependence of  $k_1$  at higher temperatures.

### **Reaction O + Br<sub>2</sub>**

*Measurements of  $k_2$ .* The rate constant of the reaction O+Br<sub>2</sub> was measured under pseudo-first order conditions from kinetics of Br<sub>2</sub> consumption ( $[\text{Br}]_0 \leq 3 \times 10^{11}$  molecule cm<sup>-3</sup>) in excess of oxygen atoms. The concentrations of O-atoms at different temperatures of the study are shown in Table II, the flow velocity in the reactor was in the range (2315-3000) cm s<sup>-1</sup>.

Typical example of Br<sub>2</sub> decays observed with different excess concentrations of oxygen atoms in the reactor is shown in Fig. 3S (Supporting Information). Figures 5 and 4S (Supporting Information) present examples of the pseudo-first order rate constant,  $k_2' = k_2[\text{O}]$ , as a function of concentration of oxygen atoms. The values of  $k_2'$  were corrected for axial and radial diffusion [26] of Br<sub>2</sub>. The diffusion coefficient of Br<sub>2</sub> in He,  $D_0 = 276 \times (T/298)^{1.75}$  Torr cm<sup>2</sup> s<sup>-1</sup>, used in the calculations was estimated with Fuller's method [28]. Corrections on  $k_2'$  were generally less than 10 % and reached up to 24% at lowest temperatures of the study. The slopes of the straight lines in Figs. 5 and 4S provide the values of  $k_2$  at respective temperatures. All the values of  $k_2$  measured in the present work are shown in Table II and Fig. 6.




**Figure 5** Reaction O + Br<sub>2</sub>: dependences of pseudo-first order rate constant of Br<sub>2</sub> consumption on concentration of oxygen atoms measured at T = 370, 655 and 775 K. Error bars represent maximum uncertainty of nearly 5 % on determination of k<sub>1</sub>'.

*Comparison with Previous Data.* The results of the present measurements of  $k_2$  are shown in Fig. 6 together with those from previous studies [19-22,29]. Clyne *et al.* [29] using atomic resonance fluorescence determined the rate constant of reaction (2) in a discharge flow system both from the rate of removal of O-atom in excess of Br<sub>2</sub> and from the rate of formation of Br in excess of oxygen atoms over Br<sub>2</sub>:  $k_2 = (1.4 \pm 0.2) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $T = 298 \text{ K}$ . Moin *et al.* [19] measured the rate constant of reaction (2) relative to that of O-atom reaction with Cl<sub>2</sub>,  $k_2/k_6 = 3.98 \times \exp(1310/T)$  at  $T = (306 - 425) \text{ K}$  (uncertainty factor of 3.4), using diffusion flame method combined with electron paramagnetic resonance detection of oxygen atoms:



With  $k_6 = 7.4 \times 10^{-12} \exp(-1650/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  [30], the reported by Moin *et al.* [19]  $k_2/k_6$  ratio leads to  $k_2 = 2.95 \times 10^{-11} \exp(-340/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ . These data, being highly uncertain and not suitable for quantitative comparison, are shown in Fig. 6 for completeness. Dodonov *et al.* [20], have also used diffusion flame method to measure the rate constant of reaction (2) over the temperature range 246-431 K:  $k_2 = 4.6 \times 10^{-11} \exp(-(210 \pm 75)/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ . Nicovich and Wine [21] employed laser flash photolysis-resonance fluorescence technique to measure the rate constant of reaction (2) and found virtually independent of temperature  $k_2 =$

$(1.76 \pm 0.8) \times 10^{-11} \exp((40 \pm 100)/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  in the temperature range 255 – 350 K. The rate constant data reported by Nicovich and Wine [21] are in excellent agreement with those of the present work.



**Figure 6** Reaction O + Br<sub>2</sub>: summary of the measurements of the reaction rate constant. Error bars represent the uncertainties reported in the respective studies: estimated total uncertainty of 15% in Nicovich and Wine [21] and present work; 1 $\sigma$  statistical uncertainty in Harwood *et al.* [22].

Finally, in a most recent study [22], the rate constant for the reaction of oxygen atoms with molecular bromine was determined in an indirect way from the measurements of [BrO] to [O<sub>3</sub>] ratio upon photolysis of O<sub>2</sub> in the presence of Br<sub>2</sub>:  $k_2 = (5.12 \pm 1.86) \times 10^{-11} \exp((989 \pm 91)/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $T = 222 - 298 \text{ K}$ :



One can note that at low temperatures the values of  $k_2$  measured in the present work differ considerably (up to a factor of 2 at  $T = 220 \text{ K}$ ) from those reported by Harwood *et al.* [22]. Three-parameter fit to the present data for  $k_2$  provides the following expression for the rate constant of the O+Br<sub>2</sub> reaction:

$$k_2 = 9.85 \times 10^{-16} T^{1.41} \exp(543/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } T = (220 - 950) \text{ K},$$

which is recommended from the present study with an independent of temperature conservative uncertainty of 15% on  $k_2$ . The information on the rate constant at elevated

temperatures may be useful for the future kinetic studies related to high-temperature bromine chemistry in near-vent volcanic plumes [31].

Both negative and U-shaped temperature dependence observed for reactions of oxygen atoms with  $\text{NO}_2$  and  $\text{Br}_2$ , respectively, indicate a complex mechanism of these reactions. In this respect, further theoretical high level calculations in comparison with the experimental observations would be very beneficial to understand the dynamics of the processes in these relatively simple chemical systems.

## CONCLUSION

In this work, using two (low and high temperature) flow reactors, we have measured the rate constants of the reactions of  $\text{O}(^3\text{P})$  atoms with  $\text{NO}_2$  and  $\text{Br}_2$  over an extended temperature range, 220 – 950 K. The temperature dependence of the rate constant of the important atmospheric reaction of oxygen atoms with  $\text{NO}_2$ , found to be in excellent agreement with multiple previous data at  $T \leq 425$  K, was extended to  $T = 950$  K. Both negative and U-shaped temperature dependence of the rate constant observed for reactions of oxygen atoms with  $\text{NO}_2$  and  $\text{Br}_2$ , respectively, seem to represent an interesting experimental basis for further theoretical developments. The high temperature data on the rate constant of the  $\text{O} + \text{Br}_2$  reaction may be useful for the laboratory kinetic studies related to high-temperature bromine chemistry in volcanic plumes.

Financial support from CNRS is gratefully acknowledged. C.K. is very grateful to région Centre (France) for financing his PhD grant.

## BIBLIOGRAPHY

1. Portmann, R. W.; Brown, S. S.; Gierczak, T.; Talukdar, R. K.; Burkholder, J. B.; Ravishankara, A. R. *Geophys. Res. Lett.* 1999, 26, 2387-2390.
2. Shrestha, K. P.; Seidel, L.; Zeuch, T.; Mauss, F. *Energy Fuels* 2018, 32, 10202-10217.
3. Klein, F. S.; Herron, J. T. *J. Chem. Phys.* 1964, 41, 1285-1290.
4. Smith, I. W. M. *Trans. Faraday Soc.* 1968, 64, 378-389.
5. Westenberg, A. A.; deHaas, N. J. *Chem. Phys.* 1969, 50, 707-719.
6. Davis, D. D.; Herron, J. T.; Huie, R. E. *J. Chem. Phys.* 1973, 58, 530-535.
7. Slinger, T. G.; Wood, B. J.; Black, G. *Int. J. Chem. Kinet.* 1973, 5, 615-620.
8. Bemand, P. P.; Clyne, M. A. A.; Watson, R. T. *J. Chem. Soc. Faraday Trans. 2* 1974, 70, 564-576.
9. Ongstad, A. P.; Birks, J. W. *J. Chem. Phys.* 1986, 85, 3359-3368.
10. Geers-Müller, R.; Stuhl, F. *Chem. Phys. Lett.* 1987, 135, 263-268.
11. Zuev, A. P.; Starikovskii, A. Y. *Khim. Fiz.* 1991, 10, 190-199.
12. Gierczak, T.; Burkholder, J. B.; Ravishankara, A. R. *J. Phys. Chem. A* 1999, 103, 877-883.
13. Estupiñán, E. G.; Nicovich, J. M.; Wine, P. H. *J. Phys. Chem. A* 2001, 105, 9697-9703.
14. Avallone, L. M. *J. Photochem. Photobio. A* 2003, 157, 231-236.
15. Burkholder, J. B.; Sander, S. P.; Abbatt, J.; Barker, J. R.; Huie, R. E.; Kolb, C. E.; Kurylo, M. J.; Orkin, V. L.; Wilmouth, D. M.; Wine, P. H. *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18*, JPL Publication 15-10, Jet Propulsion Laboratory, Pasadena, 2015 <http://jpldataeval.jpl.nasa.gov>, accessed January 2019.
16. Shiekh, B. A.; Kaur, D.; Seth, B.; Mahajan, S. *Chem. Phys. Lett.* 2016, 662, 244-249.
17. Morin, J.; Bedjanian, Y.; Romanias, M. N. *Int. J. Chem. Kinet.* 2017, 49, 53-60.
18. Bedjanian, Y.; Poulet, G. *Chem. Rev.* 2003, 103, 4639-4656.
19. Moin, F. B.; Yurkevich, Y. P.; Drogobyskii, V. M. *Dokl. Akad. Nauk SSSR* 1976, 226, 866-868.
20. Dodonov, A. F.; Zelenov, V. V.; Talroze, V. L. *Kinet. Catal.* 1981, 22, 245-247.
21. Nicovich, J. M.; Wine, P. H. *Int. J. Chem. Kinet.* 1990, 22, 379-397.

22. Harwood, M. H.; Rowley, D. M.; Cox, R. A.; Jones, R. L. *J. Phys. Chem. A* 1998, 102, 1790-1802.
23. Bedjanian, Y.; Morin, J. J. *Phys. Chem. A* 2017, 121, 1553-1562.
24. Morin, J.; Bedjanian, Y. *J. Phys. Chem. A* 2016, 120, 9063-9070.
25. Morin, J.; Romanias, M. N.; Bedjanian, Y. *Int. J. Chem. Kinet.* 2015, 47, 629-637.
26. Kaufman, F. J. *Phys. Chem.* 1984, 88, 4909-4917.
27. Ivanov, A. V.; Trakhtenberg, S.; Bertram, A. K.; Gershenzon, Y. M.; Molina, M. J. *J. Phys. Chem. A* 2007, 111, 1632-1637.
28. Tang, M. J.; Cox, R. A.; Kalberer, M. *Atmos. Chem. Phys.* 2014, 14, 9233-9247.
29. Clyne, M. A. A.; Monkhouse, P. B.; Townsend, L. W. *Int. J. Chem. Kinet.* 1976, 8, 425-449.
30. Wine, P. H.; Nicovich, J. M.; Ravishankara, A. R. *J. Phys. Chem.* 1985, 89, 3914-3918.
31. Roberts, T. J.; Martin, R. S.; Jourdain, L. *Atmos. Chem. Phys.* 2014, 14, 11201-11219.

## TABLE LEGENDS

**Table I** Reaction  $\text{O} + \text{NO}_2 \rightarrow \text{NO} + \text{O}_2$ : Experimental Conditions and Results of the Measurements of the Reaction Rate Constant

**Table II** Reaction  $\text{O} + \text{Br}_2 \rightarrow \text{Br} + \text{BrO}$ : Experimental Conditions and Results of the Measurements of the Reaction Rate Constant



## FIGURE LEGENDS

**Figure 1** Reaction  $O + NO_2$ : example of the exponential decays of O atoms in excess of  $NO_2$  at  $T = 950$  K.

**Figure 2** Reaction  $O + NO_2$ : dependences of pseudo-first order rate constant of O-atom decay on concentration of  $NO_2$  measured at  $T = 220, 298$  and  $950$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .

**Figure 3** Reaction  $O + NO_2$ : dependences of pseudo-first order rate constant of  $NO_2$  decay on concentration of oxygen atoms measured at  $T = 266$  and  $950$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .

**Figure 4** Reaction  $O + NO_2$ : summary of the measurements of the temperature dependence of the reaction rate constant. Error bars correspond to estimated 15 % uncertainty on the measurements of  $k_1$  in the present work. Uncertainties on  $k_1$  reported in previous studies are omitted for clarity.

**Figure 5** Reaction  $O + Br_2$ : dependences of pseudo-first order rate constant of  $Br_2$  consumption on concentration of oxygen atoms measured at  $T = 370, 655$  and  $775$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .

**Figure 6** Reaction  $O + Br_2$ : summary of the measurements of the reaction rate constant. Error bars represent the uncertainties reported in the respective studies: estimated total uncertainty of 15% in Nicovich and Wine [21] and present work;  $1\sigma$  statistical uncertainty in Harwood *et al.* [22].

## TABLES

**Table I** Reaction  $\text{O} + \text{NO}_2 \rightarrow \text{NO} + \text{O}_2$ : Experimental Conditions and Results of the Measurements of the Reaction Rate Constant

$T$ (K)	[Excess reactant] ( $10^{13}$ molecule $\text{cm}^{-3}$ ) <sup>a</sup>	$k_1$ ( $10^{-11}$ $\text{cm}^3$ molecule $^{-1}$ s $^{-1}$ ) <sup>b</sup>
220	0.10-4.07	$1.28 \pm 0.03$
246	0.28-4.47	$1.16 \pm 0.02$
266	0.27-4.74	$1.08 \pm 0.02$
294	0.10-4.37	$1.05 \pm 0.04$
298	0.17-5.09	$0.99 \pm 0.02$
340	0.27-4.40	$0.94 \pm 0.02$
403	0.35-4.50	$0.86 \pm 0.02$
496	0.21-4.19	$0.86 \pm 0.03$
653	0.18-3.23	$0.77 \pm 0.03$
950	0.17-4.55	$0.74 \pm 0.02$

<sup>a</sup> excess reactant: oxygen atom at  $T = 266, 294, 496$  and  $653$  K,  $\text{NO}_2$  at all other temperatures

<sup>b</sup> the errors correspond to  $2\sigma$  statistical uncertainty

**Table II** Reaction  $\text{O} + \text{Br}_2 \rightarrow \text{Br} + \text{BrO}$ : Experimental Conditions and Results of the Measurements of the Reaction Rate Constant

$T$ (K)	[O] ( $10^{13}$ molecule $\text{cm}^{-3}$ )	$k_2$ ( $10^{-11}$ $\text{cm}^3$ molecule $^{-1}$ s $^{-1}$ ) <sup>a</sup>
220	0.23-2.83	$2.28 \pm 0.08$
243	0.16-3.21	$2.17 \pm 0.06$
266	0.14-2.42	$2.11 \pm 0.04$
297	0.11-2.73	$1.87 \pm 0.06$
317	0.22-3.08	$1.81 \pm 0.04$
340	0.26-3.16	$1.72 \pm 0.08$
370	0.24-3.94	$1.74 \pm 0.06$
405	0.16-3.05	$1.78 \pm 0.04$
445	0.21-2.88	$1.84 \pm 0.10$
495	0.06-1.56	$1.89 \pm 0.06$
565	0.11-2.96	$1.96 \pm 0.10$
655	0.11-2.48	$2.02 \pm 0.08$
775	0.11-1.87	$2.52 \pm 0.08$
950	0.16-1.96	$2.69 \pm 0.22$

<sup>a</sup> the errors correspond to  $2\sigma$  statistical uncertainty

## Supporting Information

### Rate Constants of the Reactions of O(<sup>3</sup>P) Atoms with Br<sub>2</sub> and NO<sub>2</sub> over the Temperature Range 220 – 950 K.

YURI BEDJANIAN,\* CHAITANYA KALYAN


*Institut de Combustion, Aérodynamique, Réactivité et Environnement (ICARE), CNRS  
45071 Orléans Cedex 2, France*

Number of pages (including this one): 3


Number of Figures: 4

---


\*Correspondence to: Yuri Bedjanian: Tel.: +33 238255474, e-mail: [yuri.bedjanian@cnrs-orleans.fr](mailto:yuri.bedjanian@cnrs-orleans.fr)




**Figure 1S** Diagram of the flow reactor.



**Figure 2S** Reaction  $\text{O} + \text{NO}_2$ : example of the exponential decays of  $\text{NO}_2$  in excess of oxygen atoms at  $T = 294 \text{ K}$ .



**Figure 3S** Reaction  $O + Br_2$ : example of the exponential decays of  $Br_2$  in excess of oxygen atoms at  $T = 243$  K.



**Figure 4S** Reaction  $O + Br_2$ : dependences of pseudo-first order rate constant of  $Br_2$  consumption on concentration of oxygen atoms measured at  $T = 243, 297$  and  $340$  K. Error bars represent maximum uncertainty of nearly 5 % on determination of  $k_1'$ .