



HAL
open science

Spectroscopy of neutron-rich Sb isotopes using AGATA, VAMOS++ and EXOGAM

S. Biswas, A. Lemasson, M. Rejmund, Y.H. Kim, A. Navin, C. Michelagnoli,
I. Stefan, R. Banik, P. Bednarczyk, S. Bhattacharya, et al.

► **To cite this version:**

S. Biswas, A. Lemasson, M. Rejmund, Y.H. Kim, A. Navin, et al.. Spectroscopy of neutron-rich Sb isotopes using AGATA, VAMOS++ and EXOGAM. 63rd DAE-BRNS Symposium on Nuclear Physics, Dec 2018, Mumbai, India. pp.304-305. hal-02136308

HAL Id: hal-02136308

<https://hal.science/hal-02136308v1>

Submitted on 15 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spectroscopy of neutron-rich Sb isotopes using AGATA, VAMOS++ and EXOGAM

S. Biswas^{1,*}, A. Lemasson¹, M. Rejmund¹, Y. H. Kim¹, A. Navin¹, C. Michelagnoli¹, I. Stefan², R. Banik³, P. Bednarczyk⁴, S. Bhattacharya³, S. Bhattacharyya³, E. Clément¹, H. L. Crawford⁵, G. De France¹, P. Fallon⁵, J. Goupil¹, B. Jacquot¹, H. J. Li¹, J. Ljungvall⁶, A. O. Macchiavelli⁵, A. Maj⁴, L. Ménager¹, V. Morel¹, R. Palit⁷, R. M. Pérez-Vidal⁸, J. Ropert¹, and C. Schmitt¹

¹*GANIL, CEA/DRF-CNRS/IN2P3, Bd Henri Becquerel, BP 55027, F-14076 Caen Cedex 5, France*

²*Institut de Physique Nucléaire, IN2P3-CNRS,*

Université Paris-Saclay, F-91406 Orsay, Cedex, France

³*Variable Energy Cyclotron Centre, 1/AF Bidhan Nagar, Kolkata 700064, India*

⁴*Institute of Nuclear Physics PAN, 31-342 Kraków, Poland*

⁵*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

⁶*CSNSM, Université Paris-Saclay, CNRS/IN2P3, F-91405 Orsay, France*

⁷*Department of Nuclear and Atomic Physics,*

Tata Institute of Fundamental Research, Mumbai, 400005, India and

⁸*IFIC, Universitat de València, CSIC, E-46071 València, Spain*

Introduction

¹³²Sb is one of the rare neutron-rich unstable nuclei, with classical doubly magic shell closures, that is experimentally accessible with the recent developments in accelerators and detectors. The study of the excited states and the transition probabilities in nuclei in its vicinity, with few particles or/and holes added to the inert core, are crucial for understanding the shell model theory and the relevant *pp*, *pn* and *nn* interactions in this mass $A \sim 130$ region. Additionally, in this mass-region, there are presence of many seniority isomers in the range of ns to μ s. When the lifetime of the isomer is of the order of μ s's, an additional challenge appears in correlating the γ -ray transition feeding and depopulating these isomers. Therefore, these nuclei are often known either from their isomer decay or from their prompt γ spectroscopy. In the present work, the prompt-delayed spectroscopy of neutron-rich Sb isotopes have been carried out by utilising the unique combination of AGATA, VAMOS++ and EXOGAM detectors [1].

Experimental Details and Results

The neutron-rich ^{122–131}Sb fission fragments were populated using the inverse kinematics reaction, ⁹Be(²³⁸U, f) at 6.2 MeV/u. The experimental setup included the AGATA array, VAMOS++ spectrometer and the EXOGAM detectors. The prompt γ rays, emitted by the excited fission fragments, were detected using the AGATA γ -ray tracking array. The VAMOS++ spectrometer was used for the unambiguous identification of the neutron-rich fission fragments on an event by event basis. The delayed γ rays were detected by the EXOGAM detectors, placed at the focal plane of VAMOS++. More details regarding the experimental setup are given in Ref. [2]. This setup enabled the correlation of the prompt and delayed γ rays in the time range of 100 ns - 200 μ s.

A systematic analysis of the prompt and delayed transitions for all the neutron-rich ^{122–131}Sb isotopes have been carried out. The already known isomers in the odd-A ^{123–131}Sb isotopes have been confirmed in the present analysis. In addition, several new prompt and delayed transitions for these isotopes have been identified. New isomers, prompt and delayed γ rays have been identified in the even-A

*Electronic address: biswas@ganil.fr




FIG. 1: (a) The prompt γ -ray spectrum (γ_P) in coincidence with already known delayed 247 keV γ -ray (γ_D) in ^{127}Sb . (b) The γ_D in coincidence with newly identified 1195 keV prompt γ -ray in ^{128}Sb . The newly observed delayed transitions have been marked with an asterisk.

$^{122-130}\text{Sb}$ isotopes. An example of the prompt γ -ray spectrum (γ_P) with gate on an already known delayed 247 keV transition in ^{127}Sb and the delayed spectrum (γ_D) with gate on the newly identified prompt 1195 keV transition in ^{128}Sb have been shown in Fig. 1(a) and (b), respectively.

Shell model calculations using the restricted model space, consisting of neutron $d_{3/2}$, $s_{1/2}$, $h_{1/2}$ and proton $p_{1/2}$, $g_{9/2}$, and $g_{7/2}$ orbitals have been carried out using the interaction used in Ref. [3]. The interaction has been improved so as to have a better agreement for the binding energies, excitation energies and the transition probabilities. Also, a systematic comparison of the isomers in Sb

have been carried out with the corresponding isomers in $^{121-130}\text{Sn}$ isotopes, showing that the isomers in Sb dominantly arise due to a $\pi g_{7/2}$ particle coupled to the corresponding neutron configurations in the Sn isotopes. Though the energies of the excited states in the Sb isotopes seem to follow the seniority scheme, the transition probabilities do not. This could be probably due to the breaking of the neutron pairs in the $h_{11/2}$ orbital arising due to the $\pi g_{7/2}-\nu h_{11/2}$ interaction.

Acknowledgments

The authors would like to thank the entire AGATA Collaboration and the important technical contributions of the GANIL accelerator staff for the successful running of the experiment. PB and AM acknowledge support from National Science Centre under Contract No. 2013/08/M/ST2/00257 and the French LEA COPIGAL project. SB, RB, SB, SB and RP acknowledge support from CEFIPRA project No. 5604-4 and the LIA France-India agreement. HLC, PF and AOM acknowledge support from the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Contract No. DE-AC02-05CH11231 (LBNL).

References

- [1] S. Biswas, *et al.*, Manuscript in preparation.
- [2] Y. H. Kim, *et al.*, Eur. Phys. J. **24**, 39 (2017).
- [3] M. Rejmund, *et al.*, Phys. Lett. B **753**, 86 (2016).