



Rare earth elements sorption to iron oxyhydroxide: model development and application to groundwater

Haiyan Liu, Olivier Pourret, Huaming Guo, Jessica Bonhoure

► To cite this version:

Haiyan Liu, Olivier Pourret, Huaming Guo, Jessica Bonhoure. Rare earth elements sorption to iron oxyhydroxide: model development and application to groundwater. *Applied Geochemistry*, 2017, 10.1016/j.apgeochem.2017.10.020 . hal-02136360

HAL Id: hal-02136360

<https://hal.science/hal-02136360>

Submitted on 22 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rare earth elements sorption to iron oxyhydroxide: model development and application to groundwater

Haiyan Liu^{1, 2, 3}, Olivier Pourret^{3, *}, Huaming Guo^{1, 2}, Jessica Bonhoure³

¹State Key Laboratory of Biogeology and Environmental Geology, China University
of Geosciences (Beijing), Beijing 100083, P.R. China

²MOE Key Laboratory of Groundwater Circulation & Environment Evolution &
School of Water Resources and Environment, China University of Geosciences
(Beijing), Beijing 100083, P.R. China

³UniLaSalle, HydrISE, *60026 Beauvais Cedex*, France

*Corresponding author: olivier.pourret@unilasalle.fr, tel. +33 3 44 06 89 79.

Abstract: Iron oxyhydroxides are among the most important colloids that control rare earth elements (REE) concentrations and transport in natural hydrosystems. In this study, REE surface complexation to iron oxyhydroxides ($\text{Fe}(\text{OH})_3(\text{a})$) was described by using the Donnan diffuse layer model and a two-site (i.e. $\equiv\text{Fe}^{\text{s}}\text{OH}$ and $\equiv\text{Fe}^{\text{w}}\text{OH}$) model. The specific surface area and pH of zero charge were fixed as $100000 \text{ m}^2/\text{mol}$ and 8.0, respectively. The surface site density for weak and strong binding sites were fixed at 0.1 mol/mol Fe ($\equiv\text{Fe}^{\text{w}}\text{OH}$) and 0.001 mol/mol Fe ($\equiv\text{Fe}^{\text{s}}\text{OH}$) respectively. The two site types were used with $\text{pK}_{\text{a}1}^{\text{int}} = 7.29$ and $\text{pK}_{\text{a}2}^{\text{int}} = 8.93$. Using linear free energy relationship, the estimated equilibrium surface complexation constants ($\log K$) increased from light REE (LREE) to heavy REE (HREE). Results of REE modeling calculation using the determined $\log K$ revealed a good fit of experimental data, showing an order of sorption on iron oxyhydroxides: $\text{HREE} > \text{MREE} > \text{LREE}$ and preferential sorption of HREE at a lower pH. However, sorption edges only showed a slight change with ionic strength (0.1 to 0.7 mol/L) for the whole REE series. The generalized model was subsequently used to evaluate the impact of iron oxyhydroxides on REE speciation in groundwater. Application of the model to “model groundwater” showed that iron oxyhydroxide complexes of REE were significant in near neutral and weakly alkaline pH. This study contributes to putting forward a comprehensive database which would be useful for the application of surface complexation model to describe REE sorption by amorphous ferric hydroxides in nature.

Keywords: Rare earth elements; Iron oxyhydroxides; Sorption; Surface complexation modeling; Groundwater

1. Introduction

The great interest in determining the distribution and abundance of rare earth elements (REE) in aquatic systems has been shown by numerous research works generated over the past decades (Goldberg et al., 1963; Henderson, 1984; Dia et al., 2000; Tang and Johannesson, 2003; Gruau et al., 2004; Atwood, 2013; Noack et al., 2014; Gaillardet et al., 2014; Davranche et al., 2017). The solution and mineral properties of REE make them effectively robust proxies for low temperature geochemical reactions. It is generally recognized that REE systematic chemical properties usually lead to REE fractionation in geochemical systems (Henderson, 1984). REE form a coherent group and generally occur in the trivalent oxidation state. The effective ionic radii of the REE decrease with increasing atomic number (Shannon, 1976). This effect causes characteristic regular features of normalized REE patterns defining the CHArge and RAdius-Controlled process, CHARAC (Bau, 1996). If a low temperature geochemical system is characterized by a CHARAC patterns, REE of similar charge and radius should display a coherent behavior. This property, however, is no longer observed when chemical processes are driven by an external electronic configuration that produces a sub-partition (non-CHARAC processes). The CHARAC attribution of REE should thus generate smooth, rather

than irregular patterns (excepting redox-related Ce anomaly), which may indicate non-CHARAC behavior. The apparent subdivision of La and the fourteen *4f* REE elements into four groups or tetrads (I: La to Nd; II: Nd to Gd; III: Gd to Ho; IV: Ho to Lu) is referred as the Tetrad effect and a M-type lanthanide tetrad effect (Bau, 1996) is defined when subdivided into four concave-upward segments. The M-type tetrad effect has been shown on patterns of apparent REE distribution coefficients between iron oxyhydroxides and aqueous solutions (Bau, 1996) and contrasts to the complimentary W-type in the aqueous phase. In aquatic geochemical systems, REE fractionation takes place through complexation to reactive surface organic (Tang and Johannesson, 2003) and inorganic (Byrne and Li, 1995) ligands as well as surface complexation to aquifer minerals (Tang and Johannesson, 2005; Pourret and Davranche, 2013).

Despite the significant efforts that have been made toward the study of REE solution complexation with a variety of inorganic anions (carbonate, hydroxide, sulfate, fluoride, and chloride) and organic ligands (Wood, 1990; Tang and Johannesson, 2003; Pourret et al., 2007; Pourret and Martinez, 2009; Pourret et al., 2010). Far less is understood concerning REE surface complexation behavior on mineral surfaces. To date, few studies have been dedicated to REE sorption onto mineral surfaces (De Carlo et al., 1998; Bau, 1999; Ohta and Kawabe, 2001; Quinn et al., 2007; Schijf and Marshall, 2011). Early laboratory study of REE sorption onto Fe oxides showed that the light rare earth elements (LREE) were preferentially removed from solution over the heavy rare earth elements (HREE) (Koeppenkastrup and De

Carlo, 1992), although only a single pH value (i.e. pH 7.8) was considered. More recently, REE sorption on Fe (oxy)hydroxides in the absence of competing complexing ligands was investigated over a range of pH 4.0 to 9.0, as well as an ionic strength (IS) range of 0 to 0.7 mol/L (Ohta and Kawabe, 2001; Quinn et al., 2006, 2007). Further works of Bau (1999) and Ohta and Kawabe (2001), not only revealed the influences of pH and IS on REE sorption behaviour, but, in particular, non-CHARAC REE behaviors, such as lanthanide tetrad effect, were documented. Results of these experiments showed that the HREE exhibited a greater affinity for the solid phase than the LREE. Results from field experiments under acidic pH (pH<5.0) further indicated that LREE were preferentially enriched by hydrous Fe(III) oxides (Verplanck et al., 2004). Even if Mn oxides and humic substance play significant roles (Tang and Johannesson, 2003; Nakada et al., 2013; Pourret and Tuduri, 2017); previously cited studies demonstrated that Fe-oxyhydroxides may also control REE fractionation and partitioning in the natural environment.

However, although amorphous iron oxyhydroxides (e.g. $\text{Fe}(\text{OH})_3(\text{a})$) appear to be a most frequently studied solid for investigation of REE sorption, only a few works have accounted for this substrate (De Carlo et al., 1998; Bau, 1999; Ohta and Kawabe, 2001; Quinn et al., 2006). Poorly crystallized amorphous iron oxyhydroxides have a larger surface area and smaller site density than their crystalline counterparts (Henmi et al., 1980; Jambor and Dutrizac, 1998). This is evidenced by more extensive sorption of REE onto amorphous iron oxyhydroxides (Koeppenkastrop and De Carlo, 1992). The lack of data for REE binding by amorphous iron oxyhydroxides may be

110 attributed to the high variety of the equilibrium sorption constants, which complicates
111 modeling studies. Such that, for example, the estimated REE partition coefficients
112 (K_{Fe}) differ by a factor of approximately round 400 for individual REE binding
113 constants at fixed pH (Schijf and Marshall, 2011). However, in order to more
114 accurately predict REE behavior with respect to key environmental variables, such as
115 pH and ionic strength, a quantitative description of REE binding to amorphous iron
116 oxyhydroxides is urgently needed. Understanding of the fate and transport of REE is
117 of importance regarding addressing health-related (e.g. liver function) (Zhu et al.,
118 2007) and environmental issues (e.g. water pollution and farmland destruction) (Rim
119 et al., 2013), due to the increasingly extensive use of REE in various aspects entailing
120 an enhanced anthropogenic input (e.g. Gd anomaly) into environment (Bau and
121 Dulski, 1996a; Verplanck et al., 2010).

122 In this study, a surface complexation model (SCM) was proposed to quantify
123 REE sorption to amorphous iron oxyhydroxides. This SCM took into consideration
124 the generic HFO surface parameters provided by Dzombak and Morel (1990). The
125 REE-Fe(OH)₃(a) binding constants were extrapolated using the linear free energy
126 relationship (LFER) methodology and were further validated by reproducing
127 published experimental data within ionic strengths extending from 0.1 to 0.7
128 mol/L, and a pH range of 4 to 9 (De Carlo et al., 1998). The model was developed for
129 different IS and pH values (i.e. conditions of seawater and groundwater), and was
130 eventually used to discuss the importance of iron oxyhydroxides REE sorption in
131 natural groundwater.

2. Materials and methods



2.1. Surface complexation model description

The diffuse layer model is capable of describing the sorption of solutes onto hydrous oxides including those of iron, aluminum, manganese and silicon. As the central component of the generalized two-layer model, it has been used to compile a database of surface complexation reactions for hydrous ferric oxides (Dzombak and Morel, 1990), hydrous manganese oxides (Tonkin et al., 2004) and goethite (Mathur and Dzombak, 2006). The model is chosen principally because it is the simplest model which accounts qualitatively and quantitatively for all available sorption data for hydrous ferric oxides. Two binding sites, $\equiv\text{Fe}^{\text{s}}\text{OH}$ and $\equiv\text{Fe}^{\text{w}}\text{OH}$, (strong- and weak-affinity sites, respectively) were used in this SCM to describe sorption reactions with specific hydroxyl sites on the oxide surfaces. Surface equilibrium reactions (Eqs. (1) to (6)) are written as a combination of chemical free energy terms. The coulombic term represents the electrochemical sorption processes. This parameter was calculated from the Gouy-chapman electrical double layer theory, which considered a surface charge layer and a diffuse layer of counter charges.

$$P = \exp(-F\psi/RT)$$

where F stands for the molar Faraday constant (96485 C/mol); R is molar gas constant (8.314 J/mol); T is the absolute temperature (K), and ψ is the electrostatic potential.





151

152 Acidity constants (pK_{a1} and pK_{a2}) for iron oxyhydroxides, presented in Table 1,
153 were obtained from Dzombak and Morel (1990). Specific surface area (SSA) and site
154 density values required for the SCM model were those proposed by Parkhurst and
155 Appelo (2013). Total concentrations of surface sites (i.e. surface site densities) were
156 divided into strong and weak affinity fractions for REE binding (Table 1). Therefore,
157 REE sorption onto oxide surfaces was simulated by assuming two binding sites. It
158 should be noted that the site density can range from 0.001 to 0.1 mol/mol Fe, and
159 from 0.1 to 0.3 mol/mol Fe for strong and weak binding sites, respectively, due to
160 different estimation methods (Dzombak and Morel, 1990). A generic site density
161 value (0.001 mol/mol Fe and 0.1 mol/mol Fe fraction for strong and weak binding
162 sites, respectively) was chosen for $\text{Fe}(\text{OH})_3(\text{a})$ in the present model, following the
163 suggestion of Parkhurst and Appelo (2013). Modeling calculations were performed
164 using the hydrogeochemical code PHREEQC version 3.3.9 (Parkhurst and Appelo,
165 2013), following the method described by Pourret and Davranche (2013). The input
166 $\text{Fe}(\text{III})$ was considered to precipitate as $\text{Fe}(\text{OH})_3(\text{a})$ instantaneously with an
167 equilibrium constant $\log K=4.89$ (Nordstrom et al., 1990). The amounts of the binding
168 sites for these two-type of sites were determined by the amount of $\text{Fe}(\text{OH})_3(\text{a})$. The

Fe(OH)₃(a) was defined explicitly in keyword data block EQUILIBRIUM PHASES. The SSA was defined relative to the moles of Fe(OH)₃(a), in which the amount of specified binding sites changed as the SSA varied during batch-reaction simulation. Upon Fe(OH)₃(a) precipitation, the two types of oxide surface binding sites ($\equiv\text{Fe}^{\text{s}}\text{OH}$ and $\equiv\text{Fe}^{\text{w}}\text{OH}$) were assumed to be available for REE complexation. For SCM modeling, surface-complexed and diffuse layer species, were taken as the components of the system. REE surface complexation modeling was then performed in the presence of iron oxyhydroxides. Following the method described in Hummel et al. (2002) activity coefficients were calculated using the Davies (1962) approach.

2.2 Estimation of the REE-Fe(OH)₃(a) stability constants

Surface complexation constants were estimated using linear free energy relationship (LFER) by the method of Pourret and Davranche (2013), and as shown previously (Schindler et al., 1976; Dzombak and Morel, 1990; Tang and Johannesson, 2003; Tonkin et al., 2004; Mathur and Dzombak, 2006; Pourret et al., 2007). Specifically, due to the similarity of surface complexation for metal ions to the corresponding hydrolysis reaction in solution, the energy of metal ion complexation with hydroxyl ions in solution could be correlated with that of the binding of metal ions to deprotonated surface hydroxyl groups. Therefore, the first hydrolysis constants could be correlated with surface complexation constants (Mathur and Dzombak, 2006; Schott et al., 2009). As such, REE surface complexation constants could be obtained by extrapolation of aqueous hydrolysis reactions. The metal ion (including Ag⁺, Co²⁺, Ni⁺, Cd²⁺, Zn²⁺, Cu²⁺, Pb²⁺, Hg²⁺) complexation reaction parameters that were used for

verifying the fit of REE sorption data to iron oxyhydroxides were obtained from Dzombak and Morel (1990). The solution complexation reactions for these metals could be defined similarly to Eqs. (3) to (6). The LFER for metal cations was defined by the $\log K_{\text{Fe}}^{\text{s}}_{\text{OM}}$, $\log K_{\text{Fe}}^{\text{s}}_{\text{OMOH}}$, $\log K_{\text{Fe}}^{\text{w}}_{\text{OM}}$ and $\log K_{\text{Fe}}^{\text{w}}_{\text{OMOH}}$ and the hydrolysis (OH) constants. The correlation of $\log K$ with the first hydrolysis constant $\log K_{\text{MOH}}$ for these metal ions was given by equations (7) to (10). The REE first hydrolysis constant ($\log K_{\text{LnOH}}$) (Table 2) was obtained from Klungness and Byrne (2000) and was used for subsequent extrapolation of REE stability constants.

$$\text{Log } K_{\text{Fe}}^{\text{s}}_{\text{OM}} = 1.17 \log K_{\text{MOH}} - 7.89 \quad (7)$$

$$\text{Log } K_{\text{Fe}}^{\text{s}}_{\text{OMOH}} = 0.79 \log K_{\text{MOH}} - 7.69 \quad (8)$$

$$\text{Log } K_{\text{Fe}}^{\text{w}}_{\text{OM}} = 1.30 \log K_{\text{MOH}} - 7.89 \quad (9)$$

$$\text{Log } K_{\text{Fe}}^{\text{w}}_{\text{OMOH}} = 1.97 \log K_{\text{MOH}} - 16.56 \quad (10)$$

The extrapolated REE stability constants are shown in Table 2. Only the equilibrium stability constants $\log K_{\text{Fe}}^{\text{s}}_{\text{OLn}}$ and $\log K_{\text{Fe}}^{\text{w}}_{\text{OLn}}$ were considered as the amount of the REE binding through the reactions in Eqs. (4) and (6), with $\log K_{\text{Fe}}^{\text{s}}_{\text{OLnOH}}$ and $\log K_{\text{Fe}}^{\text{w}}_{\text{OLnOH}}$, was negligible.

Experimental REE data from literature (De Carlo et al., 1998) was subsequently used to validate the equilibrium surface complexation constants for the whole REE series in a reactor open to the atmosphere and at room temperature, in the presence of a $[\text{Ln}^{n+}]$ of 125 $\mu\text{g/L}$ for each of the 14 REE; $[\text{Fe(III)}] = 10 \text{ mg/L}$; and an IS range of 0.1 to 0.7 mol/L NaNO_3 . REE carbonate species were not considered in this study,

although De Carlo et al.'s (1998) experiment showed an influence of REE solution complexation with CO_3^{2-} . Modeling calculations were carried out with the Nagra/PSI database (Hummel et al., 2002). This database has been updated by the incorporation of the well-accepted stability constants at zero ionic strength and 25 °C for REE inorganic anion complexation including: (1) chloride (LnCl^{2+} , LnCl_2^+) (Luo and Byrne, 2001), (2) fluoride (LnF^{2+}) (Luo and Millero, 2004), (3) sulfate (LnSO_4^+) (Schijf and Byrne, 2004), (4) nitrate (LnNO_3^{2+}) (Millero, 1992), (4) hydroxide (LnOH^{2+} , Ln(OH)_2^+ , $\text{Ln(OH)}_3(\text{a})$) (Lee and Byrne, 1992; Klungness and Byrne, 2000), (5) carbonate (LnCO_3^+ and $\text{Ln(CO}_3)_2^-$) (Luo and Byrne, 2004). The specific log K values are presented in the supplementary information (Table S1). It should be pointed out that, to be consistent with the experimental settings (De Carlo et al., 1998), binding of the first hydrolysis species LnOH^{2+} and carbonate ions onto iron oxyhydroxides was not included in our simulations. All the REEs considered in the model in this study were trivalent, because oxidation of REEs like Ce(III) should occur after initial sorption onto mineral surface sites (Bau, 1999).

3. Results and discussion

3.1 REE- $\text{Fe(OH)}_3(\text{a})$ modeling with extrapolated stability constant

3.1.1 Modeling of REE complexation to iron oxyhydroxides

To test the validity of the extrapolated stability constants and the diffuse layer SCM, calculations herein were performed using data obtained from literature under the conditions described by De Carlo et al. (1998). The modeling results in this study

were compared to REE sorption experimental data recorded previously (De Carlo et al., 1998). As sorption behavior of the fourteen REE members were similar, subsequent results and discussions focus on La, Ce, Eu and Lu. Lanthanum, Eu and Lu represent light REE (LREE), middle REE (MREE), and heavy REE (HREE), respectively. Cerium was selected to highlight the peculiar behavior of the only redox sensitive lanthanide in the experimental conditions. As shown in figure 1, the calculated proportions of bound REE (i.e. La, Ce, Eu and Lu) increased with increasing pH (3 to 9). Relatively higher sorptive proportions were modeled with IS of 0.1 as compared to those of 0.3 and 0.7. for all REEs and pH region (Fig. 1). These results are consistent with experimental data sets therein (De Carlo et al., 1998), especially for LREE (Figs. 1a and b). For MREE (illustrated by Eu) and HREE (illustrated by Lu), however, although a relatively higher binding strength was predicted by the model under increasing pH (Fig. 1c and d), the model fit results were in agreement with experimental data within the pH range of 4 to 6.5. The discrepancy between modeled and experimental data, seemed to become significant with increasing pH and IS, and REE atomic number. The difference between modeled and experimental data was lower for LREE (e.g. less than 10% for La and Ce) than for MREE and HREE (between 20 and 30% for Eu and Lu, respectively). This may be attributed to REE carbonate complexation or to the presence of more than two binding sites on the mineral surfaces which were not accounted by the LFER model in De Carlo et al. (1998). Previous studies demonstrated that REE binding by carbonate in solution increased with REE atomic number (Lee and Byrne, 1992). Indeed, REE

carbonate complexation in solution was stronger than sorption of the HREE onto oxide surfaces (i.e. FeOOH and δ -MnO₂) when compared to the LREE (Koeppenkastrup and De Carlo, 1993). Kawabe et al. (1999) have examined the effect of REE carbonate complexation on REE sorption onto Fe oxyhydroxides. Their results further showed that distribution coefficients (K_d , commonly used in estimating the potential sorption of dissolved metals by a solid phase, Fe oxyhydroxide in our case) increased with the atomic number in carbonate-free system, whereas K_d declined across the REE series more significantly for HREE in the presence of carbonate. Therefore, the discrepancy observed in sorption between the HREE and the LREE indicated that larger concentrations of HREE were predicted to be sorbed by present model, which have not considered carbonate species, than those under the experimental conditions.

3.1.2 Effect of pH on REE sorption behavior

To a first approximation, REE sorption patterns displayed a similar behavior to that of trace-metal cations. As the pH becomes more alkaline, metal sorption onto reactive solids (e.g. iron oxyhydroxides) increased from 0 to 100% over a narrow pH range (Dzombak and Morel, 1990). Figure 1 showed that 5% of the LREE (i.e. La and Ce) were sorbed at pH<6.0, and >70% at pH=8.0 (i.e. 70% to 80% for La; 88% to 93% for Ce), whereas, sorbed fractions of MREE and HREE (i.e. Eu and Lu, respectively) ranged from about 10% at pH= 6.0 to near 100% at pH=8.0. This demonstrated pH edges shifted to lower values for HREE and MREE with respect to LREE. Therefore, the model fit suggested a preferential sorption of HREE and MREE

relative to LREE. HREE and MREE began to be sorbed at a lower pH than LREE (Fig. 2). Such that, under IS=0.1mol/L condition, sorption of Lu triggered at pH around 4, while Eu and La at pH 5 and 6, respectively (Fig. 2a). The similar characteristic was observed at IS=0.3 mol/L (Fig. 2b) and IS=0.7 mol/L (Fig. 2c), as well as has already been highlighted in De Carlo et al. (1998) and observed in Bau (1999). Indeed, when the amount of Fe(III) concentration in the SCM was decreased, the fractional sorption at a given pH was reduced (Fig. 3). In order for the proportion of sorbed REE to remain fixed as it was at low sorbent condition, pH edges shifted to the right. However, the shift appeared to be more pronounced for LREE than MREE and HREE (Fig. 3). Specifically, the pH where 50% of the total REE was sorbed, increases from 7.2 to 7.8, and from 6.2 to 6.5, and from 5.7 to 5.9 for La, Eu, Lu, respectively, when the Fe(III) input decreased by 50%. This was further evidenced for the more efficiency of sorption of HREE and MREE at lower pH as compared to LREE.

3.1.3 REE partitioning and fractionation

The distribution and fractionation of the REE sorbed by iron oxyhydroxides (at IS = 0.5 mol/L and within a pH range of 5.5-8.0) were further modeled using the SCM procedure described above. Figure 4 shows the proportion of sorbed REE as a function of REE atomic number. To further evaluate REE fractionation, Yb/Gd and La/Sm ratios were considered to fully quantify HREE and LREE enrichment, respectively. Calculated REE patterns showed HREE enrichments ($\%Yb_{sorb}/\%Gd_{sorb}$ ranging between 1.1 and 5.4; if $\%Yb_{sorb}/\%Gd_{sorb}>1$, REE patterns highlight HREE

295 enrichment, and vice versa) and LREE depletion ($\%La_{sorb}/\%Sm_{sorb}$ ranging between
296 0.08 and 0.91; if $\%La_{sorb}/\%Sm_{sorb} < 1$, REE patterns feature LREE enrichment, and
297 vice versa). Most importantly, this patterns exhibited (i) concave-upward Tetrad effect
298 with tetrads ranging from 1.00 to 1.32 (using the method proposed by Irber (1999))
299 and (ii) positive Ce anomalies ($Ce/Ce^* = \%Ce_{sorb}/(\%La_{sorb} \times \%Pr_{sorb})^{0.5}$) with Ce/Ce^*
300 ranging from 1.01 to 1.74 (average 1.54)) (Fig. 4). These features contrasted with
301 CHARAC REE patterns and strongly suggested that REE complexation behavior was
302 additionally controlled by their electron configuration and the complexing hydroxyl
303 group on the oxide surface, rather than by the REE ionic charge and radius (Bau,
304 1996). The REE fractionation patterns also coincided with SCM predictions,
305 indicating that the HREE had a stronger affinity for iron oxyhydroxides than the
306 LREE. Moreover, Ce was preferentially enriched on the iron oxyhydroxide surface
307 relative to neighboring La and Pr. The results are in good agreement with the
308 experimental datasets provided by Bau (1999) and Ohta and Kawabe (2001). At the IS
309 of 0.5 mol/L, Schijf and Marshall (2011) calculated REE distribution coefficients on
310 hydrous ferric oxides over the pH range 3.9-8.4. Their calculated REE patterns,
311 however, showed a less pronounced tetrad effect and no apparent positive Ce
312 anomalies. This is because of the weaker, and less pH dependent sorption of the REE
313 on iron oxyhydroxides, under their experimental conditions (Schijf and Marshall,
314 2011). However, the results from Verplanck et al. (2004) are strikingly similar to
315 those obtained in this study. The authors employed a lower pH (i.e. $pH < 5.1$) to model
316 REE partitioning relative to De Carlo et al. (1998) and Schijf and Marshall (2011).

The calculated complexation of REE with hydrous ferric oxides by Verplanck et al. (2004) showed a convex Tetrad curve and a positive Ce anomaly, as shown in Fig. 4. Moreover, the HREE fractionated more significantly with respect to the LREE in their study (i.e. Yb_n/Gd_n ranging from 4.3 to 7.1; for further information see Verplanck et al. (2004)). This further indicates that REE partitioning and fractionation on iron oxyhydroxides are more influenced by low pH conditions. The upward Tetrad effect curves were more apparent at acidic pH values (Fig. 4a). Indeed, as the pH decreased, enhanced protonation of ferric oxides surface occurs, leading to a decrease in the proportion of negatively charged groups being able to complex REE (Bethke, 2007).

One of the most striking features of the modeling results in this study was that Ce was preferentially sorbed by iron oxyhydroxides from the solution, with respect to La and Pr, at acidic pH (Fig. 4). The positive Ce anomaly began to decrease at a pH of 6, which may be due to a global increase of neighboring REE, since REE sorption increases as the pH rises above 6 (up to 100%). Therefore, the Ce sorption edge in iron oxyhydroxides was below pH 5, and lower than that of La. Although positive Ce anomaly may result from anomalous La enrichment, as interpreted by Bau and Dulski (1996b), further calculations showed that the observed accumulated Ce was not interfered by La (supplementary information Fig. S2). It should be noted that in low-temperature aquatic systems, Ce is the only REE that can occur in a stable tetravalent state (i.e. as Ce(IV)). This can be explained by the oxidation scavenging of Ce(III) from aqueous onto metal oxide surfaces, which leads to a decoupling of Ce from its neighbors, La and Pr (Bau, 1999). Nevertheless, sorption experiments

described by Nakada et al. (2013) showed that the sorbed Ce on ferrihydrite might not have been oxidized in the Ce/ferrihydrite system, which was evidenced by XANES analysis and the thermodynamic data. In consequence, although the redox properties of Ce(III) were not taken into account for REE modeling calculations in this study, the estimated REE sorption dataset are able to provide a better understanding of the mechanism of preferential accumulation of the Ce onto the iron oxyhydroxide surfaces.

3.2 Model validity and application

3.2.1 SCM of REE sorption to iron oxyhydroxides

In order to check the validity of the SCM of REE sorption to iron oxyhydroxides, additional calculations were performed using literature data from field studies (Pourret et al., 2010; Guo et al., 2010; Liu et al., 2016). Sampling location of the modeled groundwater extends from the recharge zone to discharge areas, which considers the variable aqueous chemistry (including pH, IS, ΣREE and Fe(III) concentrations) over a flow system. The considered pH ranges from 6.70 to 8.32, the metal loading (ΣREE) from 0.20 $\mu\text{g/L}$ to 4.25 $\mu\text{g/L}$ and the dissolved Fe(III) from 19.6 $\mu\text{g/L}$ to 740 $\mu\text{g/L}$. As shown in Fig. 5, the results from surface complexation modeling of the field data were similar to those obtained for the experimental data of De Carlo et al. (1998), showing an upward tetrad effect REE pattern and positive Ce anomalies (Ce/Ce* ranging from 1.13 to 1.61 (average 1.50)). This has been observed, in spite of the lower IS (e.g. $0.006 < \text{IS} < 0.019 \text{ mol/L}$) for these field samples with

respect to solutions used in De Carlo et al. (1998), as well as in Ohta and Kawabe (2001) and Bau (1999). In addition to the IS, a lower metal loading was observed with respect to the experimental conditions (e.g. 4.25 $\mu\text{g/L}$ for Pourret et al. (2010); 0.21 $\mu\text{g/L}$ for Guo et al. (2010)). The stronger REE binding (e.g. total sorbed REE ranging from 39% to 51%) for the Guo et al.'s (2010) sample resulted from the higher ferric oxide content and higher pH. However, under these conditions, the fractionation between individual REE occurred to a lesser extent (Fig. 5). As $\% \text{Yb}_{\text{sorb}}/\% \text{Gd}_{\text{sorb}}$ (i.e. 0.80) and $\% \text{La}_{\text{sorb}}/\% \text{Sm}_{\text{sorb}}$ (i.e. 1.04) ratios were close to 1, REE patterns tended to be flat. Eventually, modeling results are coherent with ultrafiltration experiments performed on sample PG3 (Pourret et al., 2010) which highlighted that large colloids comprising Fe oxides (>30 kDa) only controlled a small fraction of REE.

Apart from these discrepancies, the inclusion of carbonate species can be the major difference between the modeling and field studies. While CO_2 species were not considered in the modeling as shown in De Carlo et al.'s (2010), carbonate ligands are strong complexing agents for aqueous HREE at alkaline pH (Johannesson et al., 1996; Luo and Byrne, 2004). This may explain that a slightly higher sorption of the HREE by iron oxyhydroxides was calculated at alkaline pH by the model fit with respect to experimental data (Fig. 1d). When field samples were used (HCO_3^- ranging from 322 to 501 mg/L) for further modeling (Fig. 5), the calculated REE patterns changed to be less enriched in HREE (e.g., when pH increases), as evidenced by the generally lower $\% \text{Yb}_{\text{sorb}}/\% \text{Gd}_{\text{sorb}}$ ratios (1.04 to 2.32). However, these ratios were within the range of the De Carlo et al.'s (1998) modeling of experimental datasets, as shown

above (Fig. 4). To be more specific, REE species including Fe-oxyhydroxide complexes as function of atomic number are shown in Fig. 6. In the modeled groundwater of low Fe(III) concentration, REE chiefly occurred as carbonate complexes (LnCO_3^+ and $\text{Ln}(\text{CO}_3)_2^-$) (Fig. 6a and b), whereas as Fe(III) concentration increased, more REE were complexed with iron oxyhydroxides and carbonate ligands became the primarily competitor (Fig. 6c). The other REE species (including Ln^{3+} , LaSO_4^+ , LnOH^{2+} and $\text{Ln}(\text{OH})_2^+$) were not significant, although free metal ion (Ln^{3+}) had proportion greater than 10% for La and Ce in low pH groundwaters (Fig. 6a, b and c). This was best observed for Guo et al's (2010) groundwater (Fig. 6d), where dicarbonato complex ($\text{Ln}(\text{CO}_3)_2^-$) dominated the remaining aqueous REE and its proportion increased from LREE (i.e., La 22%) to HREE (i.e., Lu 48%). An increasing trend for Fe-oxyhydroxide complex fraction with increasing REE atomic number was also observed in Guo et al's (2010) sample (i.e., increasing from 38% for La to 47% for Lu) (Fig. 6d). This was attributed to the higher Fe(III) concentration and higher pH as compared to the other three samples. Therefore, although SCM estimated stability constants may be dependent on IS, pH, Fe(III) concentration and metal loading, the unique attributes of the REE series are useful for examining whole log K patterns. Furthermore, combine the well-accepted inorganic anion complexation with the SCM, the model is able to highlight the REE partitioning and fractionation between solution and iron oxide surfaces, as already highlighted in Fig. 6.

In particular, insight can be gained from SCM to understand lanthanide tetrad effect generated from REE sorption onto iron oxyhydroxide surfaces. Calculation of

the four tetrads following the method described by Irber (1999) showed that the value increases from the first tetrad (t_1) to the fourth tetrad (t_4) ($1.10 \leq T \leq 1.45$ for PG3 (Pourret et al., 2010); $1.01 \leq T \leq 1.05$ for No. well 48 (Guo et al., 2010); $1.09 \leq T_{\text{average}} \leq 1.28$ for Liu et al. (2016); $T = t_1 + t_2 + t_3 + t_4$). Moreover, the calculated tetrad values decreased with increasing pH. This has been featured in Figure 5. This aqueous-solid (i.e. iron oxyhydroxide) reaction has been considered as one of the most important processes responsible for the non-CHARAC REE behavior and REE abundance in aqueous solutions (Koeppenkastrop and De Carlo, 1992, 1993; Bau, 1996, 1999; Schijf and Marshallv, 2011; Verplanck et al., 2004). In addition to the tetrad effect, the modeling approach reproduced the stronger sorption proportion of Ce with respect to neighbors La and Pr. This was further evidenced by the largest positive Ce anomaly ($\text{Ce}/\text{Ce}^* 1.61$) observed by Pourret et al. (2010) and lowest positive Ce anomaly ($\text{Ce}/\text{Ce}^* 1.13$) in Guo et al. (2010). This may account in part for the negative Ce anomalies for normalized REE patterns in these field groundwater samples (0.58 for Pourret et al. (2010); 0.39 for Guo et al. (2010); average 0.94 for Liu et al. (2016)). The non-CHARAC (e.g. lanthanide tetrad effect) REE behavior and negative Ce anomaly are widely observed in natural waters (Elderfield et al., 1990; Bau, 1996). These are, for the most part, attributed to the surface complexation of the REE by iron oxyhydroxides. The complexation is, however, not only exclusively dependent on REE ionic charge and radius, but controlled by variations in the electron configurations of REE and by the type of complexing ligands. Furthermore, the model also suggests that the contribution of REE scavenging of iron oxyhydroxides towards

the lanthanide tetrad effect and the removal of Ce from solution should be most significant at acidic pH.

3.2.2 REE speciation calculation for world average groundwater

The developed model was used to evaluate REE speciation for world average groundwater, proposed as “model groundwater” by Tang and Johannesson (2003). The concentrations of major anions and REE for world average groundwater used in the model were obtained from Wood (1990), whereas the concentrations of cations and Fe were taken from Livingstone (1963). These parameters have been previously used by Lee and Byrne (1992), Johannesson et al. (1996) and Tang and Johannesson (2003) for investigation of REE complexation behavior in groundwater. Thus REE speciation calculation was performed keeping the solute compositions constant and changing pH. The dissolved ligands (i.e. carbonates, sulfates, hydroxyls) present in the modeled groundwater are similar to those considered by Wood (1990) at pH=7 and 20 °C (calculating using PHREEQC).

The results of these calculations for La, Eu and Lu are shown in Fig. 7. It is obviously observed from Fig. 7a, b and c, in the “model groundwater” with Fe(III) concentrations of 0.672 mg/L, the SCM predicted that REE: (i) occurred predominately as free metal ion (Ln^{3+}) and sulfate complex (LnSO_4^+) at pH between 4 and 6.5; (ii) were significantly associated with Fe oxide ($\text{Hfo}_2\text{Ln}^{2+}$) at pH between 6.5 and 8; (iii) chiefly as carbonate complex (LnCO_3^+ and $\text{Ln}(\text{CO}_3)_2^-$) at pH between 8 and 9.5, and (iv) were dominated by hydroxide complex (LnOH^{2+} and $\text{Ln}(\text{OH})_2^+$) at pH above 9.5; (v) negligibly as chloride complex (LnCl^{2+} and LnCl_2^+) and nitrate

448 complex (LnNO_3^{2+}) (below 1%) in the pH range. Generally speaking, the modeling
449 results for acidic and alkaline groundwaters were in good agreement with previous
450 investigations of natural groundwater (Wood, 1990; Lee and Byrne, 1992;
451 Johannesson, 2005). It should be noted, however, that these earlier studies did not
452 consider REE complexation with iron oxyhydroxides. The accordance of the present
453 SCM with previously purely inorganic speciation models for acid and alkaline
454 groundwaters should thus be verified by further field or laboratory investigations in
455 these pH conditions. However, it is also worth noting that the predominant
456 characteristic for the modeling calculations is that significant amounts of REE are
457 complexed with iron oxyhydroxide for groundwaters of near neutral and weakly
458 alkaline pH. Indeed, the model predicted that 20% to 30% of LREE (i.e. La) occurred
459 as iron-oxyhydroxide complexes ($\text{Hfo}_2\text{OLa}^{2+}$) in the pH range from 7.5 to 9.0 (Fig.
460 7a), whereas for MREE (i.e. Eu) and HREE (i.e. Lu), 20% to 50% ($\text{Hfo}_2\text{OEu}^{2+}$) and
461 20% to 80% ($\text{Hfo}_2\text{OLu}^{2+}$) were complexed with iron-oxyhydroxide in the pH ranges
462 from 6.5 to 9.0, and from 6.3 to 9.0, respectively. Furthermore, a strong competition
463 between REE surface complexation and solution complexation has also been observed
464 in the model, and that the competition changes with pH (Fig. 7). In particular, REE
465 carbonate complexes (LnCO_3^+ and $\text{Ln}(\text{CO}_3)_2^-$) were the most important inorganic
466 ligands competing with iron oxyhydroxide in near neutral and weakly alkaline
467 groundwaters, and thus this process was responsible for the mobility and transport of
468 REE in natural groundwater systems. This is well consistent with the modeling results
469 using field sampled groundwater, as discussed above (Fig. 6). In this model, the

influences of organic matter and manganese oxyhydroxides on REE speciation were not taken in account, although previously researches demonstrated they could played an important role in REE behavior (Tang and Johannesson, 2003; Nakada et al., 2013; Pourret et al., 2009; Pourret and Tuduri 2017). However, as was the case documented by Verplanck et al. (2004) and Quinn et al. (2007), the generalized proposed model is able to provide important suggestion for REE complexation with iron oxyhydroxides in Fe-rich groundwater-aquifer systems, unless other potential complexers (e.g. manganese or humics) are essentially present in aquifers and complex the REE.

4. Conclusion

Surface complexation modeling with high and low affinity sites was used to test REE sorption onto iron oxyhydroxides by considering a LFER to determine log K values. The extrapolated equilibrium surface complexation constants were further validated by using them to fit various experimental data sets (De Carlo et al., 1998). Results of SCM using the determined constants showed that the role of iron oxyhydroxide in complexing REE exhibited dependence upon pH, IS and the concentration of iron oxyhydroxide. The proportion of iron oxyhydroxide complexes showed upward lanthanide tetrad effect patterns as a function of atomic number, with stronger sorption of Ce than neighboring La and Pr over a wide variety of conditions (i.e. pH from 4.0 to 9.0; IS from 0 to 0.7 mol/L; [REE] from 0.14 µg/L to 125 µg/L; [Fe(III)] ranging from 0.0183 mg/L to 10 mg/L). Speciation calculations using the SCM suggest that iron oxyhydroxides are important complexers of REEs in groundwaters, and they effectively compete with dissolved inorganic ligands under

various conditions. For the first time, the generalized proposed model allows us to predict the impact of iron oxyhydroxides on REE in natural environment. Compared to previous modeling studies (e.g. Verplanck et al. (2004) and Schijf and Marshall (2011)), this type of SCM accounts for a wider range of pH and IS, and particularly can be a good application to natural groundwater. Furthermore, the increase of the determined constants with increasing atomic number provides insight into understanding of the discrepancy for HREE between modeled and experimental data. Further development of the modeling techniques should incorporate Mn oxides, humic substances and Fe oxides simultaneously into model, in order to obtain a comprehensive model, and to better understand the roles of particle colloids in REE mobility in natural systems.

Acknowledgement

We would like to acknowledge China Scholarship Council (CSC) for sponsoring the first author's study in France (CSC No. 201606400016). We thank Dr. Raul E. Martinez for comments on an early draft and post-editing the English style.

References

- Atwood, D.A. 2013. The rare earth elements: fundamentals and applications. John Wiley & Sons. pp.1-19.
- Bau, M., Dulski, P. 1996a. Distribution of yttrium and rare-earth elements in the Penge and Kuruman iron-formations, Transvaal Supergroup, South Africa. *Precambrian Research*, 79(1-2), 37-55.
- Bau, M., Dulski, P., 1996b. Anthropogenic origin of positive gadolinium anomalies in river waters. *Earth and Planetary Science Letters*, 143, 245-255.

515 Bau, M. 1996. Controls on the fractionation of isovalent trace elements in magmatic
 516 and aqueous systems: evidence from Y/Ho, Zr/Hf, and lanthanide tetrad
 517 effect. *Contributions to Mineralogy and Petrology*, 123(3), 323-333.

518 Bau, M. 1999. Scavenging of dissolved yttrium and rare earths by precipitating iron
 519 oxyhydroxide: experimental evidence for Ce oxidation, Y-Ho fractionation, and
 520 lanthanide tetrad effect. *Geochimica et Cosmochimica Acta*, 63(1), 67-77.

521 Bethke, C.M. 2007. *Geochemical and biogeochemical reaction modeling*. Cambridge
 522 University Press. Second edition.

523 Byrne, R.H., Li, B. 1995. Comparative complexation behavior of the rare
 524 earths. *Geochimica et Cosmochimica Acta*, 59(22), 4575-4589.

525 Davies, C.W., 1962. *Ion Association*. Butterworths, London.

526 Davranche, M., Gruau, G., Dia, A., Le Coz-Bouhnik, M., Marsac, R., Pédrot, M.,
 527 Pourret, O. 2017. Chapter 7. Rare Earth Elements in Wetlands. In *Trace
 528 Elements in Waterlogged Soils and Sediments*. Eds Rinklebe J., Knox A.S.,
 529 Paller M., Taylor & Francis Group/CRC Press, 135-162.

530 De Carlo, E.H., Wen, X.Y., Irving, M. 1998. The influence of redox reactions on the
 531 uptake of dissolved Ce by suspended Fe and Mn oxide particles. *Aquatic
 532 Geochemistry* 3, 357-389.

533 Dia, A., Gruau, G., Olivie-Lauquet, G., Riou, C., Molénat, J., Curmi, P. 2000. The
 534 distribution of rare earth elements in groundwaters: assessing the role of
 535 source-rock composition, redox changes and colloidal particles. *Geochimica et
 536 Cosmochimica Acta*, 64(24), 4131-4151.

537 Dzombak, D.A., Morel, F.M. 1990. *Surface complexation modeling: hydrous ferric
 538 oxide*. John Wiley & Sons.

539 Elderfield, H., Upstill-Goddard, R., Sholkovitz, E.R. 1990. The rare earth elements in
 540 rivers, estuaries, and coastal seas and their significance to the composition of
 541 ocean waters. *Geochimica et Cosmochimica Acta*, 54(4), 971-991.

542 Gaillardet, J., Viers, J. and Dupré, B. 2014. 7.7-Trace Elements in River Waters, in:
 543 Holland, H.D., Turekian, K.K. (Eds.), *Treatise on Geochemistry* (Second
 544 Edition). Elsevier, Oxford, pp. 195-235.

545 Goldberg, E.D., Koide, M., Schmitt, R.A., Smith, R.H. 1963. Rare-Earth distributions
546 in the marine environment. *Journal of Geophysical Research*, 68(14), 4209-4217.

547 Gruau, G., Dia, A., Olivie-Lauquet, G., Davranche, M., Pinay, G. 2004. Controls on
548 the distribution of rare earth elements in shallow groundwaters. *Water*
549 *Research*, 38(16), 3576-3586.

550 Guo, H.M., Zhang, B., Wang, G.C., Shen, Z.L., 2010. Geochemical controls on
551 arsenic and rare earth elements approximately along a groundwater flow path in
552 the shallow aquifer of the Hetao Basin, Inner Mongolia. *Chemical Geology*
553 270, 117-125.

554 Henderson, P. 1984. *Rare earth element geochemistry*, Elsevier, Amsterdam.

555 Henmi, T., Wells, N., Childs, C.W., Parfitt, R.L. 1980. Poorly-ordered iron-rich
556 precipitates from springs and streams on andesitic volcanoes. *Geochimica et*
557 *Cosmochimica Acta*, 44(2), 365-372.

558 Hummel, W., Berner, U., Curti, E., Pearson, F.J., Thoenen, T. 2002. Nagra/PSI
559 chemical thermodynamic data base 01/01. Universal Publishers, Parkland,
560 Florida.

561 Irber, W. 1999. The lanthanide tetrad effect and its correlation with K/Rb, Eu/Eu* ,
562 Sr/Eu, Y/Ho, and Zr/Hf of evolving peraluminous granite suites. *Geochimica et*
563 *Cosmochimica Acta*, 63(3), 489-508.

564 Jambor, J.L., Dutrizac, J.E. 1998. Occurrence and constitution of natural and synthetic
565 ferrihydrite, a widespread iron oxyhydroxide. *Chemical Reviews*, 98(7),
566 2549-2586.

567 Johannesson, K.H., Stetzenbach, K.J., Hodge, V.F., Lyons, W.B. 1996. Rare earth
568 element complexation behavior in circumneutral pH groundwaters: assessing the
569 role of carbonate and phosphate ions. *Earth and Planetary Science*
570 *Letters*, 139(1-2), 305-319.

571 Johannesson, K.H., 2005. *Rare earth elements in groundwater flow systems*. Springer,
572 Dordrecht. Book. Water Science and Technology Library. Volume 51

573 Kawabe, I., Ohta, A., Miura, N. 1999. Distribution coefficients of REE between Fe
574 oxyhydroxide precipitates and NaCl solutions affected by REE-carbonate
575 complexation. *Geochemical journal*, 33(3), 181-197.

576 Klungness, G.D., Byrne, R.H., 2000. Comparative hydrolysis behavior of the rare
577 earths and yttrium: the influence of temperature and ionic strength. *Polyhedron*
578 19, 99-107.

579 Koeppenkastrop, D., De Carlo, E.H.. 1992. Sorption of rare-earth elements from
580 seawater onto synthetic mineral particles: An experimental approach. *Chemical*
581 *Geology*, 95(3-4), 251-263.

582 Koeppenkastrop, D., De Carlo, E.H. 1993. Uptake of rare earth elements from
583 solution by metal oxides. *Environmental Science & Technology*, 27(9),
584 1796-1802.

585 Lee, J.H., Byrne, R.H., 1992. Examination of comparative rare earth element
586 complexation behavior using linear free-energy relationships. *Geochimica et*
587 *Cosmochimica Acta* 56, 1127-1137.

588 Liu, H.Y., Guo, H.M., Xing, L.N., Zhan, Y.H., Li, F.L., Shao, J.L., N, H., L, X., Li,
589 C.Q. 2016. Geochemical behaviors of rare earth elements in groundwater along a
590 flow path in the North China Plain. *Journal of Asian Earth Sciences*, 117, 33-51.

591 Livingstone, D.A. 1963. Chemical composition of rivers and lakes. US Government
592 Printing Office.

593 Luo, Y.R., Byrne, R.H. 2001. Yttrium and rare earth element complexation by
594 chloride ions at 25 C. *Journal of Solution Chemistry*, 30(9), 837-845.

595 Luo, Y., Millero, F.J. 2004. Effects of temperature and ionic strength on the stabilities
596 of the first and second fluoride complexes of yttrium and the rare earth
597 elements. *Geochimica et cosmochimica acta*, 68(21), 4301-4308.

598 Luo, Y.R., Byrne, R.H., 2004. Carbonate complexation of yttrium and the rare earth
599 elements in natural rivers. *Geochimica et Cosmochimica Acta* 68, 691-699.

600 Mathur, S.S., Dzombak, D.A., 2006. Chapter 16-surface complexation modeling:
601 goethite. In: Johannes, L. (Ed.), *Interface Science and Technology*. 11. Elsevier,
602 pp. 443-468.

- 603 Millero, F.J., 1992. Stability constants for the formation of rare earth inorganic
604 complexes as a function of ionic strength. *Geochimica et Cosmochimica Acta*
605 56, 3123-3132.
- 606 Nakada, R., Takahashi, Y., Tanimizu, M. 2013. Isotopic and speciation study on
607 cerium during its solid–water distribution with implication for Ce stable isotope
608 as a paleo-redox proxy. *Geochimica et Cosmochimica Acta*, 103, 49-62.
- 609 Noack, C.W., Dzombak, D.A., Karamalidis, A.K., 2014. Rare earth element
610 distributions and trends in natural waters with a focus on groundwater.
611 *Environmental Science & Technology* 48, 4317-4326.
- 612 Nordstrom, D.K., Plummer, L.N., Langmuir, D., Busenberg, E., May, H.M., Jones,
613 B.F., Parkhurst, D.L. 1990. Revised chemical equilibrium data for major
614 water-mineral reactions and their limitations.
- 615 Ohta, A., Kawabe, I. 2001. REE (III) adsorption onto Mn dioxide (δ -MnO₂) and Fe
616 oxyhydroxide: Ce (III) oxidation by δ -MnO₂. *Geochimica et Cosmochimica*
617 *Acta*, 65(5), 695-703.
- 618 Parkhurst, D.L., Appelo, C. A.J. 2013. Description of input and examples for
619 PHREEQC version 3-a computer program for speciation, batch-reaction,
620 one-dimensional transport, and inverse geochemical calculations. US geological
621 survey techniques and methods, book, 6, 497.
- 622 Pourret, O., Davranche, M., Gruau, G., Dia, A., 2007. Rare earth elements
623 complexation with humic acid. *Chemical Geology*, 243, 128-141.
- 624 Pourret, O., Martinez, R.E. 2009. Modeling lanthanide series binding sites on humic
625 acid. *Journal of colloid and interface science*, 330(1), 45-50.
- 626 Pourret, O., Gruau, G., Dia, A., Davranche, M., Molenat, J. 2010. Colloidal control on
627 the distribution of rare earth elements in shallow groundwaters. *Aquatic*
628 *Geochemistry*, 16(1), 31.
- 629 Pourret, O., Davranche, M. 2013. Rare earth element sorption onto hydrous
630 manganese oxide: A modeling study. *Journal of Colloid and Interface*
631 *Science*, 395, 18-23.

632 Pourret, O., Tuduri, J. 2017. Continental shelves as potential resource of rare earth
633 elements. *Scientific Reports*, 7.

634 Quinn, K.A., Byrne, R.H., Schijf, J. 2006. Sorption of yttrium and rare earth elements
635 by amorphous ferric hydroxide: influence of solution complexation with
636 carbonate. *Geochimica et Cosmochimica Acta*, 70(16), 4151-4165.

637 Quinn, K.A., Byrne, R.H., Schijf, J. 2007. Sorption of yttrium and rare earth elements
638 by amorphous ferric hydroxide: influence of temperature. *Environmental Science
639 & Technology*, 41(2), 541-546.

640 Rim, K.T., Koo, K.H., Park, J.S. 2013. Toxicological evaluations of rare earths and
641 their health impacts to workers: a literature review. *Safety and health at
642 work*, 4(1), 12-26.

643 Shannon, R.T. 1976. Revised effective ionic radii and systematic studies of
644 interatomic distances in halides and chalcogenides. *Acta crystallographica
645 section A: crystal physics, diffraction, theoretical and general
646 crystallography*, 32(5), 751-767.

647 Schindler, P.W., Fürst, B., Dick, R., Wolf, P.U. 1976. Ligand properties of surface
648 silanol groups. I. Surface complex formation with Fe^{3+} , Cu^{2+} , Cd^{2+} , and
649 Pb^{2+} . *Journal of Colloid and Interface Science*, 55(2), 469-475.

650 Schijf, J., Byrne, R.H. 2004. Determination of $\text{SO}_4 \beta_1$ for yttrium and the rare earth
651 elements at $I=0.66 \text{ m}$ and $t=25 \text{ }^\circ\text{C}$ -implications for YREE solution speciation in
652 sulfate-rich waters. *Geochimica et cosmochimica acta*, 68(13), 2825-2837.

653 Schijf, J., Marshall, K.S. 2011. YREE sorption on hydrous ferric oxide in 0.5 M NaCl
654 solutions: a model extension. *Marine Chemistry*, 123(1), 32-43.

655 Schott, J., Pokrovsky, O.S., Oelkers, E.H. 2009. The link between mineral
656 dissolution/precipitation kinetics and solution chemistry. *Reviews in mineralogy
657 and geochemistry*, 70(1), 207-258.

658 Tang, J., Johannesson, K.H. 2003. Speciation of rare earth elements in natural
659 terrestrial waters: assessing the role of dissolved organic matter from the
660 modeling approach. *Geochimica et Cosmochimica Acta*, 67(13), 2321-2339.

661 Tang, J., Johannesson, K.H. 2005. Adsorption of rare earth elements onto Carrizo
662 sand: experimental investigations and modeling with surface
663 complexation. *Geochimica et Cosmochimica Acta*, 69(22), 5247-5261.

664 Tonkin, J.W., Balistrieri, L.S., Murray, J.W. 2004. Modeling sorption of divalent
665 metal cations on hydrous manganese oxide using the diffuse double layer
666 model. *Applied Geochemistry*, 19(1), 29-53.

667 Verplanck, P.L., Nordstrom, D.K., Taylor, H.E., Kimball, B.A. 2004. Rare earth
668 element partitioning between iron oxyhydroxides and acid mine waters. *Applied*
669 *Geochemistry* 19, 1339-1354.

670 Verplanck, P.L., Furlong, E.T., Gray, J.L., Phillips, P.J., Wolf, R.E., Esposito, K.
671 2010. Evaluating the behavior of gadolinium and other rare earth elements
672 through large metropolitan sewage treatment plants. *Environmental Science &*
673 *Technology*, 44(10), 3876-3882.

674 Wood, S.A. 1990. The aqueous geochemistry of the rare-earth elements and yttrium:
675 1. Review of available low-temperature data for inorganic complexes and the
676 inorganic REE speciation of natural waters. *Chemical Geology*, 82, 159-186.

677 Zhu, W.F., Xu, S.Q., Shao, P.P., Zhang, H., Wu, D.S., Yang, W.J., Feng, J., Feng, L.
678 2005. Investigation on liver function among population in high background of
679 rare earth area in South China. *Biological trace element research*, 104(1), 1-7.

Table 1

Table1 SCM parameters for iron oxyhydroxide (The pK values are taken from Dzobmbak and Morel (1990); the site densities and specific surface area are from Parkhurst and Appelo (2013)).

pK _{a1}	pK _{a2}	Specific surface area (m ² /mol)	Total site density (mol/mol Fe)	Strong site density (≡Fe ^s OH) (mol/mol Fe)	Weak site density (≡Fe ^w OH) (mol/mol Fe)
7.29	8.93	100000	0.101	0.001	0.1

Table 2 Stability constants extrapolated from the LFER

REE	$\log K_{LnOH}$	$\log K_{Ln(OH)2}$	$\log K_{Ln(OH)3}$	$\log K_{=Fe^s OLn}$	$\log K_{=Fe^w OLn}$
La	5.19	-4.14	-13.9	1.68	-1.16
Ce	5.66	-3.6	-13.23	2.22	-0.54
Pr	5.68	-3.27	-12.63	2.25	-0.52
Nd	5.82	-3.04	-12.4	2.41	-0.33
Sm	6.16	-2.51	-11.91	2.81	0.11
Eu	6.24	-2.37	-11.41	2.90	0.21
Gd	6.17	-2.37	11.72	2.82	0.12
Tb	6.36	-2.18	-11.08	3.04	0.37
Dy	6.41	-2.1	-10.83	3.10	0.43
Ho	6.44	-2.07	-10.56	3.13	0.47
Er	6.48	-1.96	-10.35	3.18	0.52
Tm	6.61	-1.88	-10.18	3.33	0.69
Yb	6.76	-1.74	-9.85	3.51	0.89
Lu	6.73	-2.67	-9.85	3.47	0.85

Fig. 1




Figure 1 A comparison between the experimental and calculated proportion of sorbed La, Ce, Eu and Lu with the extrapolated stability constants under the experimental conditions documented by De Carlo et al. (1998). The dots correspond to the experimental data, and the solid lines correspond to the model fit ((a): La; (b): Ce; (c): Eu; (d): Lu).

Fig. 2




Figure 2 Proportion of REE ([REE]=125 $\mu\text{g/L}$) sorbed to iron oxyhydroxide (total [Fe(III)] = 10 mg/L) as a function of pH at various IS: (a) 0.1 mol/L, (b) 0.3 mol/L and (c) 0.7 mol/L.

Fig. 3




Figure 3 pH edges for REE sorption ((a): La; (b): Eu; (c): Lu; [REE]=125 $\mu\text{g/L}$;

IS=0.5 mol/L). Arrows indicate decreasing iron oxyhydroxide concentration.

Fig. 4




Figure 4 Patterns arising from the calculated proportion of sorbed REE under the experimental conditions described by De Carlo et al. (1998) (IS= 0.5 mol/L).

Fig. 5




Figure 5 Modeling calculation with literature data from the field studies including Pourret et al. (2010) (sample PG3), Guo et al. (2010) (No. well 48) and Liu et al. (2016) (sampling location at a distance of 0 km and 61 km from the recharge area, respectively; for detailed information see Liu et al. (2016)).


Fig. 6

Figure 6 Results of REE speciation calculation for groundwater samples ((a): Pourret et al., (2010) (sample PG3); (b) and (c): Liu et al., (2016) (sampling location at a distance of 0 km and 61 km from the recharge area, respectively); (d): Guo et al., (2010) (No. well 48)).

Fig. 7




Figure 7 Speciation of (a) La, (b) Eu, and (c) Lu in groundwater with [Fe(III)] of 0.672 mg/L as a function of pH (Concentrations of the major anions and REE from Wood (1990); concentrations of the major cations and Fe from Livingstone (1963)) It must be noted that the minor fluoride, chloride and nitrate complexes (below < 1%) are not presented for clarity.

Supplementary Information for
**Rare earth elements sorption to iron oxyhydroxide: model development and
application to groundwater**

Haiyan Liu^{1, 2, 3}, Olivier Pourret^{3,*}, Huaming Guo^{1, 2}, Jessica Bonhoure³

¹State Key Laboratory of Biogeology and Environmental Geology, China University
of Geosciences (Beijing), Beijing 100083, P.R. China

²MOE Key Laboratory of Groundwater Circulation & Environment Evolution &
School of Water Resources and Environment, China University of Geosciences
(Beijing), Beijing 100083, P.R. China

³UniLaSalle, HydrISE, *60026 Beauvais Cedex*, France

Contents of this file

Caption and Table S1

Caption and Figure S2

Table S1 Stability constants (log k) for REE inorganic anion complexation

	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
REE(OH) ²⁺	-8.81	-8.34	-8.32	-8.18	-7.84	-7.76	-7.83	-7.64	-7.59	-7.56	-7.52	-7.39	-7.24	-7.27
REE(OH) ₂ ⁺	-18.14	-17.6	-17.27	-17.04	-16.51	-16.37	-16.37	-16.18	-16.1	-16.07	-15.96	-15.88	-15.74	-16.67
REE(OH) ₃ (aq)	-27.9	-27.23	-26.63	-26.4	-25.91	-25.41	-2.28	-25.08	-24.83	-24.56	-24.35	-24.18	-23.85	-23.85
REE(CO ₃) ⁺	6.73	7.06	7.23	7.28	7.46	7.48	7.39	7.46	7.56	7.55	7.61	7.68	7.81	7.75
REE(CO ₃) ₂ ⁻	11.3	11.76	12.08	12.17	12.53	12.63	12.48	12.78	12.91	13	13.12	13.27	13.3	13.37
REE(NO ₃) ₂ ⁺	0.58	0.69	0.69	0.79	0.78	0.83	0.47	0.51	0.15	0.25	0.15	0.2	0.25	0.56
REE(SO ₄) ⁺	3.61	3.61	3.62	3.6	3.63	3.64	3.61	3.59	3.57	3.54	3.51	3.48	3.46	3.44
REEF ²⁺	3.63	3.85	3.86	3.82	4.14	4.26	4.23	4.36	4.39	4.28	4.27	4.28	4.37	4.24
REECl ²⁺	0.74	0.68	0.71	0.71	0.67	0.69	0.71	0.67	0.7	0.71	0.7	0.71	0.69	0.67
REECl ₂ ⁺	0.51	0.46	0.45	0.47	0.42	0.44	0.46	0.43	0.43	0.45	0.45	0.46	0.44	0.43




Figure S2 The calculated Ce/Ce* and Pr/Pr * for cases shown in figure 4 and 5. Field I: neither Ce nor La anomaly; field IIa: positive La anomaly, no Ce anomaly; field IIb: negative La anomaly, no Ce anomaly; field IIIa: positive Ce anomaly; field IIIb: negative Ce anomaly. For further explanation see Bau and Dulski. (1996). (The only sample located in area I is the case with IS=0.5 and pH=8, where nearly all of the REEs are sorbed onto solid phase (see figure 4 in the text))