



HAL
open science

OrthoMaM v10: Scaling-Up Orthologous Coding Sequence and Exon Alignments with More than One Hundred Mammalian Genomes

Celine Scornavacca, Khalid Belkhir, Jimmy Lopez, Rémy Dernas, Frédéric Delsuc, Emmanuel J.P. Douzery, Vincent Ranwez

► **To cite this version:**

Celine Scornavacca, Khalid Belkhir, Jimmy Lopez, Rémy Dernas, Frédéric Delsuc, et al.. OrthoMaM v10: Scaling-Up Orthologous Coding Sequence and Exon Alignments with More than One Hundred Mammalian Genomes. *Molecular Biology and Evolution*, 2019, 36 (4), pp.861-862. <10.1093/molbev/msz015>. <hal-02139915>

HAL Id: hal-02139915

<https://hal.science/hal-02139915v1>

Submitted on 26 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

OrthoMaM v10: Scaling-Up Orthologous Coding Sequence and Exon Alignments with More than One Hundred Mammalian Genomes

Celine Scornavacca,^{*1} Khalid Belkhir,¹ Jimmy Lopez,¹ Rémy Derrat,¹ Frédéric Delsuc,¹ Emmanuel J.P. Douzery,¹ and Vincent Ranwez²

¹Institut des Sciences de l'Evolution (ISEM), Univ. Montpellier, CNRS, EPHE, IRD, Montpellier, France

²AGAP, Univ. Montpellier, CIRAD, INRA, Montpellier SupAgro, Montpellier, France

***Corresponding author:** E-mail: celine.scornavacca@umontpellier.fr.

Associate editor: Koichiro Tamura

Abstract

We present version 10 of OrthoMaM, a database of orthologous mammalian markers. OrthoMaM is already 11 years old and since the outset it has kept on improving, providing alignments and phylogenetic trees of high-quality computed with state-of-the-art methods on up-to-date data. The main contribution of this version is the increase in the number of taxa: 116 mammalian genomes for 14,509 one-to-one orthologous genes. This has been made possible by the combination of genomic data deposited in Ensembl complemented by additional good-quality genomes only available in NCBI. Version 10 users will benefit from pipeline improvements and a completely redesigned web-interface.

Key words: orthologous sequences, mammals, coding sequences, phylogenomics, comparative genomics.

OrthoMaM is a database of high-quality orthologous sequence alignments and phylogenetic trees from mammalian genomes. It has been, for instance, used for developing new molecular markers, inferring mammalian phylogenies, simulating sequences for testing alignment filtering methods, and studying the evolution of base composition in protein-coding sequences. Previous versions of our database only included mammalian genomes from Ensembl (Ranwez et al. 2007; Douzery et al. 2014). With the progress of sequencing techniques, the number of genomes included in NCBI, but not yet in Ensembl, grows each year. This motivated us to totally rethink our database to include the annotated mammalian genomes available in NCBI only.

To identify the core set of orthologous sequences to be used, we rely on the OrthoMaM v8 pipeline described in Douzery et al. (2014). Briefly, we isolate 1-to-1 orthologous genes among pairs of our pillar placental species (*Homo-Mus*, *Homo-Canis*, and *Mus-Canis*) by using Ensembl v91 annotations (Zerbino et al. 2018). These clusters are then enriched by adding sequences of 69 additional mammals annotated in Ensembl v91 as 1-to-1 orthologues to the human gene, and turned into clusters of 1-to-1 orthologous CDSs by selecting the longest transcript of each gene. These proto-clusters are aligned using MAFFT at the amino acid level (Katoh and Standley 2013).

The CDSs of the 47 annotated mammal genomes available in NCBI (Rigden and Fernández 2017) but not in Ensembl as of March 2018 are then used to enrich the proto-alignments. For each proto-alignment, an HMM profile is created via *hmmbuild* using default parameters of the HMMER toolkit (Eddy 2011). Additionally, all HMM profiles are concatenated

and summarized using *hmmcompress* to construct an HMM database. Then, for each NCBI CDS C_i , *hmmsearch* is used on the HMM database to get the best hits among the proto-alignments—denoted $\text{bestAl}(C_i)$. For each proto-alignment A_j , the most similar sequences for each species S —denoted $\text{bestSeq}(S, A_j)$ —are detected via *hmmsearch*. Outputs from *hmmsearch* and *hmmsearch* are discarded if the first hit score is not substantially better than the second ($\text{hit}_2 < 0.9 \text{hit}_1$) and are combined in a best-reciprocal-hit fashion: Suppose that we are given a NCBI CDS sequence C_i for species S_k ; then denoting A_j the best proto-alignment for C_i , if the first sequence of $\text{bestSeq}(S_k, \text{bestAl}(C_i))$ equals C_i and none of the other sequences $C_z \neq C_i$ in $\text{bestSeq}(A_j)$ is such that $\text{bestAl}(S_k, C_z) = A_j$, then C_i is considered to belong to A_j . This ensures our orthology predictions for the NCBI CDSs to be robust. The enriched orthologous clusters have been thoughtfully aligned and filtered via the OMM_MACSE pipeline to construct high-quality codon alignments relying on MACSE v2 (Ranwez et al. 2018), MAFFT (Katoh and Standley 2013), and HMMcleaner (Philippe et al. 2017). Phylogenetic trees for each CDS alignment are constructed under maximum likelihood (ML) with RAXML (Stamatakis 2014) under the GTR+ Γ model (Yang 1994).

OrthoMaM v10 also includes an ad hoc nonorthologous sequence detection method. For a given marker, the distance between the most recent common ancestor (MRCA) of placental sequences and any sequence of the marker, denoted here S_i , depends mostly on two factors: the considered marker (some genes evolve faster) and the genome to which S_i belongs (some species evolve faster). We use linear regression to explain the patristic distance on inferred ML trees between a sequence

© The Author(s) 2019. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Open Access




Fig. 1. Screenshots from the OrthoMaM website version 10. Here, the database is queried for CDSs with a relative evolutionary rate between 0.5 and 2, a GC3 between 23% and 45%, a parameter of the gamma distribution ranging from 1 to 2, and an alignment length $>1,000$ characters (top left). The result returns 126 target CDSs, for which information can be downloaded (bottom left). The longest CDS (ABCA13) is visualized with the corresponding phylogenetic information (top right) and AA alignment (bottom right).

S_i and the placental MRCA based on those two factors. When the linear regression prediction departs from the observed patristic distance (standardized residual >3), we consider the sequence as spurious and remove it.

As in the previous versions of our database, we use exon positions in CDS alignments to infer exon orthology and alignments (see Douzery et al. 2014 for details). Phylogenetic trees for each exon alignment are reconstructed as for CDSs.

OrthoMaM v10 database is available at http://www.orthomam.univ-montp2.fr/orthomam_v10/. For each CDS and exon marker, we provide gene level information (gene name, GO annotation), full sequence traceability information (sequence identifier in Ensembl/NCBI, filtering details), nucleotide and amino acid alignments, phylogenetic trees, as well as several evolutionary indicators such as relative evolutionary rate and G + C content. The improved web-interface (see fig. 1) provides a user-friendly way to query the database based on alignment content (number of sequences, number of species that should be represented, and alignment length), evolutionary indicators, or sequence similarity using blast.

This new reactive web site, which has been completely redesigned to improve the user experience, the triplication of the number of species, the major update of the pipeline for orthology prediction and sequence filtering should make OrthoMaM v10 a central resource for anyone interested in mammalian comparative genomics.

Acknowledgments

Our project largely benefited from the Montpellier Bioinformatics Biodiversity cluster computing platform. This work was supported by the Agence Nationale de la Recherche “Investissements d’avenir/Bioinformatique” (ANR-10-BINF-

01-02 project Ancestrome), Labex CeMEB “Centre Méditerranéen de l’Environnement et de la Biodiversité” (ANR-10-LABX-0004), and the European Research Council (ERC-2015-CoG-683257 project ConvergeAnt). This is contribution ISEM 2018-292 of the Institut des Sciences de l’Evolution de Montpellier.

References

- Douzery EJP, Scornavacca C, Romiguier J, Belkhir K, Galtier N, Delsuc F, Ranwez V. 2014. OrthoMaM v8: a database of orthologous exons and coding sequences for comparative genomics in mammals. *Mol Biol Evol.* 31(7):1923–1928.
- Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput Biol.* 7(10):e1002195.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780.
- Philippe H, de Vienne DM, Ranwez V, Roure B, Baurain D, Delsuc F. 2017. Pitfalls in supermatrix phylogenomics. *Eur J Taxon.* 283:1–25.
- Ranwez V, Delsuc F, Ranwez S, Belkhir K, Tilak MK, Douzery EJP. 2007. OrthoMaM: a database of orthologous genomic markers for placental mammal phylogenetics. *BMC Evol Biol.* 7(1):241.
- Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. 2018. MACSE v2: toolkit for the alignment of coding sequences accounting for frameshifts and stop codons. *Mol Biol Evol.* 35(10):2582–2584.
- Rigden D. J., Fernández X. M. 2017. The 2018 Nucleic Acids Research database issue and the online molecular biology database collection. *Nucleic acids research.* 46(D1):D1–D7.
- Stamatakis A. 2014. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9):1312–1313.
- Yang Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *J Mol Evol.* 39(3):306–314.
- Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhaj J, Billis K, Cummins C, Gall A, Giron CG, et al. 2018. Ensembl 2018. *Nucleic Acids Res.* 46(D1):D754–D761.