

1

2 **Table 1.** Mass equilibrium defining linear equations used for the inverse analysis. Flow names in the LIM-MCMC were composed of the three-letter code of the compartment of origin followed by the three-letter code of the destination compartment. Abbreviations: gpp = gross primary
 3 production, res = respiration, los = loss for the considered system.
 4

5

Compartments	Mass balance equation	Equations common to the eight models
pic	Mass balance for picophytoplankton	$gpp-pic - (pic-res + pic-doc + pic-pro) = 0$
nan	Mass balance for nanophytoplankton	$gpp-nan - (nan-res + nan-doc + nan-det + nan-pro + nan-met1 + nan-met2 + nan-los) = 0$
mic	Mass balance for microphytoplankton	$gpp-mic - (mic-res + mic-doc + mic-det + mic-pro + mic-met1 + mic-met2 + mic-los) = 0$
pro	Mass balance for protozooplankton	$(pic-pro + nan-pro + mic-pro + bac-pro) - (pro-res + pro-doc + pro-det + pro-met1 + pro-los) = 0$
met1	Mass balance for metazooplankton 200-700 μm	$(nan-met1 + mic-met1 + pro-met1 + det-met1) - (met1-res + met1-doc + met1-det + met1-met2 + met1-los) = 0$
met2	Mass balance for metazooplankton > 700 μm	$(nan-met2 + mic-met2 + pro-met2 + met1-met2 + det-met2) - (met2-res + met2-doc + met2-det + met2-los) = 0$
bac	Mass balance for bacteria	$doc-bac - (bac-res + bac-doc + bac-pro) = 0$
doc	Mass balance for dissolved organic carbon	$(pic-doc + nan-doc + mic-doc + pro-doc + met1-doc + met2-doc + bac-doc + det-doc) - (doc-bac + doc-los) = 0$
det	Mass balance for detritus	$(nan-det + mic-det + pro-det + met1-det + met2-det) - (det-doc + det-met1 + det-met2 + det-los) = 0$

6

7 pic = picophytoplankton, nan = nanophytoplankton, mic = microphytoplankton, pro = protozooplankton, met1 = metazooplankton1,
 8 met2 = metazooplankton2, bac = bacteria, doc = dissolved organic carbon, det = detritus
 9

10 **Table 2.** Minimum and maximum values of flows (derived from *in situ* measurement for each seasons) used as inequalitiesfor the eight
 11 networks.

Process integrated as inequalities	Flux	Spring		Summer		Autumn		Winter	
		Channel model	Bay model	Channel model	Bay model	Channel model	Bay model	Channel model	Bay model
pic gross primary production	gpp-pic	54 - 77	46 - 69	36 - 53	92 - 128	202 - 268	10 - 68	28 - 38	62 - 82
nan gross primary production	gpp-nan	88 - 121	48 - 75	102 - 135	92 - 122	4 - 6	1 - 2	4 - 6	2 - 3
mic gross primary production	gpp-mic	687 - 904	522 - 698	208 - 276	172 - 227	133 - 175	326 - 562	75 - 140	139 - 206
pic grazing by protozooplankton	pic-pro	17- 22	31 - 46	33 - 36	41 - 93	168 - 204	68 - 81	18 - 28	37- 66
nan grazing by protozooplankton	nan-pro	73 - 90	31- 41	37 - 51	20 - 25	2 - 3	1 - 1.5	2 - 3	1.62 - 2.27
mic grazing by protozooplankton	mic-pro	425 - 468	210 - 478	80- 110	76 - 82	48 - 65	190 - 352	46 - 49	67 - 84
doc uptake by bacteria	doc-bac	90 - 117	20 - 59	46 - 61	40 - 92	42 - 85	63 - 67	90 - 135	244 - 274
bacterial grazing by protozoa	bac-pro	58 - 61	13 - 16	29 - 32	18 - 37	42 - 47	24 - 53	67 - 68	137 - 159
mic sinking	mic-los	86 - 521	29 - 55	12 - 19	36 - 44	31 - 35	6 - 7	39 - 57	24 - 54
nan sinking	nan-los	1.5- 44	1 - 1.5	0.02 - 0.03	0.4 - 0.6	2 - 3	0.3 - 0.5	0.20 - 0.31	0.71 - 0.91
mic and nan grazing by met1	mic+nan-met1	36 - 49	17- 26	32 - 48	5 - 8	9 - 11	19 - 29	7 - 8	18 - 22
mic and nan grazing by met2	mic+nan-met2	66 - 85	0.12 - 0.16	0.10 - 0.20	14 - 21	0.05 - 0.08	7 - 19	14 - 18	40 - 52
met1+met2 sinking	met1+met2-los	16*	21*	0.65*	0.62*	4.24*	0.91*	0.47*	0.48*
det sinking	det-los	2361**	657**	6849**	1611**	4126**	7083**	1238**	684**

12

13 All flows are considered as intervals between the minimum and maximum values estimated *in situ*, with the following exceptions:

14 * Flows considered as minimum

15 ** Flows considered as maximum

16

17 **Table 3.** Flows description and values ($\text{mg C m}^{-2} \text{ d}^{-1}$) estimated by the LIM-MCMC for the eight plankton food webs.

18

Description	Symbol	Interfered value ($\text{mg C m}^{-2} \text{ d}^{-1}$)							
		Spring		Summer		Autumn		Winter	
		Channel model	Bay model	Channel model	Bay model	Channel model	Bay model	Channel model	Bay model
Microphytoplankton gross primary production	gpp-mic	795	609	242	199	154	453	123	180
Nanophytoplankton gross primary production	gpp-nan	65	61	118	107	5	2	5	3
Picophytoplankton gross primary production	gpp-pic	105	56	44	109	236	117	33	73
Microphytoplankton respiration	mic-res	40	32	12	10	8	35	6	9
Microphytoplankton dissolved organic carbon exudation	mic-doc	69	54	21	17	13	49	11	16
Microphytoplankton det production	mic-det	35	143	77	41	34	76	0.2	0.3
Microphytoplankton grazing by protozooplankton	mic-pro	444	326	95	79	56	250	46	74
Microphytoplankton grazing by metazooplankton 1	mic-met1	33	15	20	3	10	24	6	18
Microphytoplankton grazing by metazooplankton 2	mic-met2	66	0.1	0.06	9	0.04	13	13	40
Microphytoplankton sinking	mic-los	109	40	15	40	33	6	40	24
Nanophytoplankton respiration	nan-res	4	4	6	6	0.3	0.1	0.2	0.1
Nanophytoplankton dissolved organic carbon exudation	nan-doc	6	6	10	9	0.5	0.2	0.4	0.3
Nanophytoplankton detritus production	nan-det	9	7	37	57	0.1	0.02	0.2	0.01
Nanophytoplankton grazing by protozooplankton	nan-pro	19	36	44	23	2	1	3	2
Nanophytoplankton grazing by metazooplankton 1	nan-met1	9	7	20	3	0.1	0.02	1	0.01
Nanophytoplankton grazing by metazooplankton 2	nan-met2	9	0.1	0.06	9	0.03	0.03	1	0.01
Nanophytoplankton sinking	nan-los	9	1	0.03	0.5	2	0.4	0.2	1
Picophytoplankton respiration	pic-res	6	4	3	7	12	6	2	4
Picophytoplankton dissolved organic carbon exudation	pic-doc	9	6	4	10	20	10	3	6
Picophytoplankton grazing by protozooplankton	pic-pro	90	46	36	92	203	101	28	62
Protozooplankton respiration	pro-res	181	127	63	67	89	113	39	89
Protozooplankton dissolved organic carbon excretion	pro-doc	95	66	33	35	47	59	38	89
Protozooplankton detritus production	pro-det	94	64	31	33	46	58	20	28
Protozooplankton consumption by met1	pro-met1	158	99	43	50	76	91	42	69

Protozooplankton consumption by metazooplankton 2	pro-met2	84	66	35	34	46	58	5	0.1
Metazooplankton 1 respiration	met1-res	92	80	54	45	43	63	19	36
Metazooplankton1 dissolved organic carbon excretion	met1-doc	47	41	28	23	22	33	18	36
Metazooplankton1 detritus production	met1-det	45	39	26	22	21	31	10	11
Metazooplankton1 consumption by metazooplankton 2	met1-met2	67	60	42	33	33	48	19	28
Metazooplankton1 sinking	met1-los	38	31	21	18	16	25	1	0.02
Metazooplankton 2 respiration	met2-res	96	68	45	43	37	57	13	22
Metazooplankton2 dissolved organic carbonexcretion	met2-doc	48	35	23	22	19	29	12	22
Metazooplankton 2 detritus production	met2-det	46	32	21	21	18	27	6	7
Metazooplankton 2 sinking	met2-los	96	69	45	44	38	58	11	17
Bacteria respiration	bac-res	26	13	14	20	19	16	27	55
Bacteria dissolved organic carbon production	bac-doc	32	21	17	30	23	22	40	82
Bacteria grazing by protozooplankton	bac-pro	58	14	29	26	42	28	67	137
Dissolved organic carbon uptake by bacteria	doc-bac	117	48	60	77	85	65	135	274
Dissolved organic carbon output	doc-los	236	223	103	96	81	167	0.5	0.01
Detritus dissolution	det-doc	46	43	26	25	21	31	13	22
Detritus consumption by metazooplankton1	det-met1	89	130	87	84	50	85	17	24
Detritus consumption by metazooplankton2	det-met2	61	78	56	44	53	33	5	0.1
Detritus sinking	det-los	34	34	23	20	16	25	1	0.01

19

20

21 **Table 4.** Constraints (inequalities) used for the eight networks based on literature.

Process	Concerned compartment	Bound	Description	Equation	Reference	
Respiration	pic, nan, mic	Lower	Autotroph respiration (r) is higher than 5% of gross primary production (gpp)	$5\% \text{ gpp} < r$	Vézina and Platt (1988)	
		Upper	Autotroph respiration (r) is no more than 30% of gross primary production (gpp)	$r < 30\% \text{ gpp}$		
	bac	Lower	Bacteria respiration (r') is higher than 20% of total uptake of doc (udoc)	$20\% \text{ udoc} < r'$	Vézina and Savenkoff (1999)	
		Lower	Zooplankton respiration (r'') is at least 20% of their total ingestion (ΣIng)	$20\% \Sigma \text{Ing} < r''$		
Dissolved organic carbon production	pic, nan, mic	Lower	Autotroph doc exudation (e) is higher than 10% of their net primary production (npp)	$10\% \text{ npp} < e$	Vézina and Savenkoff (1999)	
		Upper	Autotroph doc exudation (e) does not exceed 55% of their net primary production (npp)	$e < 55\% \text{ npp}$		
	pro, met1, met2	Lower	Zooplankton excretion (ex) of doc is at least 10% of their total ingestion (ΣIng)	$10\% \Sigma \text{Ing} < \text{ex}$	Vézina and Pace (1994)	
		Upper	Zooplankton excretion (ex) doesn't exceed their respiration (r'')	$\text{ex} < r''$		
	bac	Upper	Bacteria doc release (rdoc) is lower than their respiration (r')	$\text{rdoc} < r'$	Vézina and Pace (1994)	
	pro, met1, met2	Lower	The growth efficiency of zooplankton is at least 25% of their carbon uptake	$25\% \Sigma \text{Ing} < \text{Ing} - (r'' + \text{ex} + \text{det})$	Vézina and Pahlow (2003)	
Growth efficiency		Upper	The growth efficiency of zooplankton is no more than 50% of their carbon uptake	$\text{Ing} - (r'' + \text{ex} + \text{det}) < 50\% \Sigma \text{Ing}$		
		Lower	Growth efficiency of bacteria is at least 5 % of their total ingestion	$5\% \text{ udoc} < \text{udoc} - (r' + \text{rdoc})$	Vézina and Pahlow (2003)	
		Upper	Growth efficiency of bacteria is no more than 50% of their total ingestion	$\text{udoc} - (r' + \text{rdoc}) < 50\% \text{ udoc}$		
Assimilation efficiency	pro, met1, met2	Lower	Assimilation efficiency of zooplankton is at least 50 % of their total ingestion	$50\% \Sigma \text{Ing} < \text{Ing} - \text{det}$	Vézina et al. (2000)	
		Upper	Assimilation efficiency of zooplankton is no more than 90% of their total ingestion	$\text{Ing} - \text{det} < 90\% \Sigma \text{Ing}$		
Detritus production	pro, met1, met2	Lower	Zooplankton det production is at least 5% of their det production and respiration combined	$5\% (\text{zoo-det} + \text{zoo-res}) < \text{zoo-det}$	Steinberg et al. (2000)	
		Upper	Zooplankton det production is no more than 42% of their det production and respiration combined	$\text{Zoo-det} < 42\% (\text{zoo-det} + \text{zoo-res})$		
Detritus dissolution	det	Upper	Det dissolution does not exceed 10% of net primary production	$\text{det-doc} < 10\% \text{ npp}$	Pace et al. (1984)	

23 **Table 5.** Ecological indicators used for the Channel and the Bay models. Nomenclature of symbols used in formula calculation of network index
 24 equations: i: flow out of the system from i, i=1,... n; j: flow into the system to j, j=1... n; T_i : Total outflow from compartment i; T_j : Total inflow
 25 for compartment j; and T_{ij} : Flow from compartment i to compartment j. TST_c is the total flow that is recycled, Z: equals the sum of all
 26 exogenous inputs.
 27

Ecological indicator	Definition	Formula calculation	Unit	Reference
Total System Throughput (TST)	The sum of all the flows circulating through all compartments and a measure of the total system activity.	$TST = \sum_{i=1, j=1}^n T_{ij}$	$\text{mg C m}^{-2} \text{ d}^{-1}$	Kay et al. (1989)
Ascendency (A)	Ascendency represents the organized part of the ecosystem and merges the quantification of the system activity and the degree of specialization.	$A = \sum_{ij} (T_{ij}) * \log\left(\frac{T_{ij} * TST}{T_j * T_i}\right)$	$\text{mg C m}^{-2} \text{ d}^{-1} \times \text{bits}$	Ulanowicz and Wulff (1991) Ulanowicz (2000)
Development Capacity (DC)	Represents the maximum possible value of Ascendency that an ecosystem may reach.	$DC = -TST * \sum_{ij} \frac{T_{ij}}{TST} * \log\left(\frac{T_{ij}}{TST}\right)$	$\text{mg C m}^{-2} \text{ d}^{-1} \times \text{bits}$	Ulanowicz and Wulff (1991) Ulanowicz (2000)
Relative Ascendency (A/DC)	Defines the ecosystem degree of development. High A/DC ratios reflect a high degree of organization, highly specialized and less redundant pathways.	$\frac{A}{DC}$	%	Baird et al. (1991, 1998)
Average Path Length (APL)	It is the average number of compartments crossed by a unit of carbon from entering to exiting the ecosystem. Represents a measure of the ecosystem retention capacity.	$APL = \frac{TST - \sum_i \text{imports}}{\sum_i \text{imports}}$	no unit	Kay et al. (1989) Baird et al. (1991)
Finn Cycling Index (FCI)	Quantifies the fraction of all flows involved in recycling and can be considered as a measure of the retentiveness of the system.	$FCI = \frac{TST_c}{TST}$	%	Finn (1976)
Detritivory/Herbivory (D/H)	Indicates the importance of living trophic interactions compared to detritus chain.	$\frac{D}{H}$	No unit	Ulanowicz (1992)

28 **Table 6.** Results of the Cliff's δ test applied to ecological network indices calculated for the Channel and the Bay over the four seasons.
 29 Comparison was made between seasons (spring/summer, spring/autumn, spring/winter, summer/autumn, summer/winter and winter/autumn) for
 30 each ecological indices (TST, A, DC, A/DC, APL, FCI and D/H). Asterisk Between brackets represent the magnitude of the effect (***): large
 31 and (ns) negligible effects.

32
33

		Spring/Summer	Spring/Autumn	Spring/Winter	Summer/Autumn	Summer/Winter	Winter/Autumn
Channel (inshore station)	TST	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)
	A	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)
	DC	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)
	A/DC	0.54 (***)	0.54 (***)	-0.99 (***)	0.54 (***)	-0.99 (***)	0.99 (***)
	APL	-0.92 (***)	0.99 (***)	-1 (***)	0.99 (***)	-1 (***)	0.99 (***)
	FCI	-0.67 (***)	0.99 (***)	-1 (***)	0.99 (***)	-1 (***)	0.99 (***)
	D/H	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)
Bay (marine station)	TST	0.99 (***)	0.99 (***)	0.99 (***)	-0.99 (***)	0.99 (***)	0.99 (***)
	A	0.99 (***)	0.99 (***)	0.99 (***)	-0.99 (***)	0.87 (***)	0.99 (***)
	DC	0.99 (***)	0.99 (***)	0.99 (***)	-0.99 (***)	0.99 (***)	0.99 (***)
	A/DC	0.95 (***)	-0.01 (ns)	-0.94 (***)	-0.96 (***)	-1 (***)	-0.96 (***)
	APL	-0.62 (***)	-0.53 (***)	-1 (***)	0.12 (ns)	-1 (***)	-1 (***)
	FCI	-0.94 (***)	-0.60 (***)	-1 (***)	0.87 (***)	-1 (***)	-1 (***)
	D/H	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)	0.99 (***)

34

35