



HAL
open science

The Boltzmann equation with an external force on the torus: Incompressible Navier-Stokes-Fourier hydrodynamical limit

Marc Briant, Arnaud Debussche, Julien Vovelle

► **To cite this version:**

Marc Briant, Arnaud Debussche, Julien Vovelle. The Boltzmann equation with an external force on the torus: Incompressible Navier-Stokes-Fourier hydrodynamical limit. 2020. hal-02150286v2

HAL Id: hal-02150286

<https://hal.science/hal-02150286v2>

Preprint submitted on 26 Jun 2020 (v2), last revised 13 Nov 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE BOLTZMANN EQUATION WITH AN EXTERNAL FORCE ON THE TORUS: INCOMPRESSIBLE NAVIER-STOKES-FOURIER HYDRODYNAMICAL LIMIT

MARC BRIANT, ARNAUD DEBUSSCHE, JULIEN VOVELLE

ABSTRACT. We study the Boltzmann equation with external forces, not necessarily deriving from a potential, in the incompressible Navier-Stokes perturbative regime. On the torus, we establish Cauchy theories that are independent of the Knudsen number in Sobolev spaces. The existence is proved around a time-dependent Maxwellian that behaves like the global equilibrium both as time grows and as the Knudsen number decreases. We combine hypocoercive properties of linearized Boltzmann operators with linearization around a time-dependent Maxwellian that catches the fluctuations of the characteristics trajectories due to the presence of the force. This uniform theory is sufficiently robust to derive the incompressible Navier-Stokes-Fourier system with an external force from the Boltzmann equation. Neither smallness, nor time-decaying assumption is required for the external force, nor a gradient form, and we deal with general hard potential and cut-off Boltzmann kernels. As a by-product the latest general theories for unit Knudsen number when the force is sufficiently small and decays in time are recovered.

Keywords: Boltzmann equation with external force, Hydrodynamical limit, Incompressible Navier-Stokes equation, Hypocoercivity, Knudsen number.

CONTENTS

1. Introduction	1
2. Main results	6
3. Properties and estimates on the external operator and the Boltzmann operator	10
4. <i>A priori</i> estimates in Sobolev spaces	18
5. Proofs of the main results	30
References	32

1. INTRODUCTION

The Boltzmann equation is used to model rarefied gas dynamics when particles undergo elastic binary collisions, when one studies the gas from a mesoscopic point of view. It describes the time evolution of $f = f(t, x, v)$: the distribution of the particles constituting the gas in position x and velocity v . The equation can be derived from Newton's law under the assumption of rarefied gases [14] and it reads

$$(1.1) \quad \tau \partial_t F + v \cdot \nabla_x F = \frac{1}{\text{Kn}} Q(F, F).$$

The parameter Kn is a physical parameter, called the Knudsen number, that gauges the continuity of the gas. Physically speaking, a small Knudsen number indicates that fluid equations are more accurate to describe the gas. The parameter τ in (1.1) is a relaxation time.

For given ranges of the parameters τ and Kn , one can show that the physical observables - mass, momentum and energy - of the solution F converge are well approached by solutions of acoustic equations or Euler equation or incompressible Navier-Stokes equations, among others. We refer to [34, 33, 20] for a deep discussions on the matter. We will consider the regime

$$\tau = \text{Kn} = \varepsilon,$$

with $\varepsilon \rightarrow 0$. Describing by the decomposition $F = \mu + \varepsilon f$ the fluctuations of amplitude ε of the solution F around a global equilibrium μ , we expect an asymptotic description of f in terms of the incompressible Navier-Stokes-Fourier equations:

$$(1.2) \quad \begin{aligned} \partial_t u - \nu \Delta u + u \cdot \nabla u + \nabla p &= 0, \\ \nabla \cdot u &= 0, \\ \partial_t \theta - \kappa \Delta \theta + u \cdot \nabla \theta &= 0, \end{aligned}$$

together with the Boussinesq relation

$$(1.3) \quad \nabla(\rho + \theta) = 0.$$

It is interesting to mention that due to initial conservation laws for the Boltzmann equation, the Boussinesq equation actually imposes $\rho + \theta = 0$, which in turns gives (1.2) [20].

The resulting perturbative Boltzmann equation is

$$(1.4) \quad \partial_t f + \frac{1}{\varepsilon} v \cdot \nabla_x f = \frac{1}{\varepsilon^2} L[f] + \frac{1}{\varepsilon} \tilde{Q}(f, f)$$

where L is a linear operator.

Describing the evolution of the macroscopic parameters, the density, the momentum and the energy associated to f as ε tends to 0 has been the subject of numerous works starting from the *a priori* very weak convergence given by the Bardos-Golse-Levermore program [6] and using a wide range of tools from spectral theory in Fourier space [19, 11] to the setting of renormalized solutions [22, 23]. We point out [27, 12] in particular as they rely on two different manifestations of a very important property of the Boltzmann linear operator L : its hypocoercivity, which will play a central part in our study. Note that one may differentiate here the perturbative approach of References [11, 27, 12], for example, from the approach “in the large” of [5, 6, 8, 9, 10, 30, 21, 22, 23, 29, 2].

The present article focuses on the Boltzmann equation when the gas under consideration is evolving on the d -dimensional torus \mathbb{T}^d and is influenced by an external force $\vec{E}_t(x)$. We would like to derive the incompressible Navier-Stokes-Fourier hydrodynamical limit of the latter. In this setting, the Boltzmann equation reads, for (t, x, v) in $[0, T_{\max}) \times \mathbb{T}^d \times \mathbb{R}^d$,

$$(1.5) \quad \partial_t F(t, x, v) + \frac{1}{\varepsilon} v \cdot \nabla_x F(t, x, v) + \varepsilon \vec{E}_t(x) \cdot \nabla_v F(t, x, v) = \frac{1}{\varepsilon} Q(F, F)(t, x, v).$$

The bilinear operator $Q(g, h)$ is given under its symmetric form:

$$Q(g, h) = \frac{1}{2} \int_{\mathbb{R}^d \times \mathbb{S}^{d-1}} \Phi(|v - v_*|) b(\cos\theta) [h'g'_* + h'_*g' - hg_* - h_*g] dv_* d\sigma,$$

where f' , f_* , f'_* and f are the values taken by f at v' , v_* , v'_* and v respectively. Define:

$$\left\{ \begin{array}{l} v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma \\ v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma \end{array} \right., \text{ and } \cos\theta = \left\langle \frac{v - v_*}{|v - v_*|}, \sigma \right\rangle.$$

All along this paper we consider the Boltzmann equation with assumptions

(H1) *Hard potential* or *Maxwellian potential* ($\gamma = 0$), that is to say there is a constant $C_\Phi > 0$ such that

$$(1.6) \quad \Phi(z) = C_\Phi z^\gamma, \quad \gamma \in [0, 1].$$

(H2) Strong Grad's *angular cutoff* [24], expressed here by the fact that we assume the non-negative function b to be C^1 with the following controls

$$(1.7) \quad \forall z \in [-1, 1], \quad b(z), |b'(z)| \leq C_b.$$

Remark 1.1. We may relax (1.6) into $\Phi(z) \asymp z^\gamma$, in the sense that $C_\Phi^1 z^\gamma \leq \Phi(z) \leq C_\Phi^2 z^\gamma$ for all z , where C_Φ^1 and C_Φ^2 are two positive constants. We may also assume, instead of (1.7), that

$$(1.8) \quad \sup_{z \in (-1, 1)} b(z) < +\infty,$$

and that the non-degeneracy hypothesis

$$(1.9) \quad \inf_{\sigma_2, \sigma_3 \in \mathbb{S}^{d-1}} \int_{\sigma_3 \in \mathbb{S}^{d-1}} \min\{b(\sigma_1 \cdot \sigma_3), b(\sigma_2 \cdot \sigma_3)\} d\sigma_3 > 0,$$

is satisfied. Under (1.7), we can use [4] to get a spectral gap estimate on the linearized operator L (see (3.9)), while, under (1.8)-(1.9), this is [31] that can be applied.

There are two direct observations one can make comparing (1.5) to the standard Boltzmann equation (1.1). Firstly, the conservation of momentum and energy do not hold, and we are only left with the *a priori* mass conservation

$$(1.10) \quad \forall t \in [0, T_{\max}), \quad \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} F(t, x, v) dx dv = 0.$$

Secondly, the global equilibrium of the Boltzmann equation

$$(1.11) \quad \forall v \in \mathbb{R}^d, \quad \mu(v) = \frac{1}{(2\pi)^{d/2}} e^{-\frac{|v|^2}{2}},$$

which satisfies $Q(\mu, \mu) = 0$ is no longer a stationary solution to (1.5). However, as ε vanishes we expect the dynamics of the Boltzmann equation with external force to converge towards $\mu(v)$. We aim at constructing an existence and uniqueness theory in Sobolev spaces for solutions to (1.5) uniformly in ε . We shall look for solutions in a perturbative setting, mimicking the classical decomposition $F = \mu + \varepsilon f$, that will catch the hydrodynamical regime of the incompressible Navier-Stokes-Fourier with external force. More precisely, we intend to show that if, at initial time, F_0 is

sufficiently close to $\mu(v)$, then so is $F(t)$. Moreover the perturbations of the mass, momentum and energy:

$$(1.12) \quad \rho_\varepsilon(t, x) = \varepsilon^{-1} \int_{\mathbb{R}^d} [F(t, x, v) - \mu(v)] dv$$

$$(1.13) \quad u_\varepsilon(t, x) = \varepsilon^{-1} \int_{\mathbb{R}^d} v [F(t, x, v) - \mu(v)] dv$$

$$(1.14) \quad \theta_\varepsilon(t, x) = \varepsilon^{-1} \int_{\mathbb{R}^d} \frac{|v|^2 - d}{\sqrt{2d}} [F(t, x, v) - \mu(v)] dv$$

converge to (ρ, u, θ) , which are Leray solutions to the following Navier-Stokes-Fourier's system (1.15)-(1.16):

$$(1.15) \quad \begin{aligned} \partial_t u - \nu \Delta u + u \cdot \nabla u + \nabla p &= \frac{\vec{E}_t(x)}{2}, \\ \nabla \cdot u &= 0, \end{aligned}$$

$$(1.16) \quad \partial_t \theta - \kappa \Delta \theta + u \cdot \nabla \theta = 0,$$

together with the Boussinesq relation (1.3). Leray solutions of the latter means weak solutions integrated against test functions with null divergence. We show that (ρ, u, θ) are solutions in this weak sense, but, due to the estimates in Sobolev spaces with high indexes that we obtain (*cf.* Theorem 2.5), these solutions are classical, regular solutions close to the equilibrium state $(1, 0, 1)$.

The present hydrodynamical problem has not been addressed yet in the mathematical literature, and even the works with $\varepsilon = 1$ on the Boltzmann equation with an external force (1.5) in full generality are scarce. The main issue being that velocity derivatives can grow very rapidly. To our knowledge only [16] deals with general $\vec{E}_t(x)$: they solve the perturbative Cauchy theory around μ in Sobolev spaces for $\varepsilon = 1$ as long as the force $\vec{E}_t(x)$ is small and decreases to 0 as time increases (the latter assumption is removable if $d \geq 5$ or if one solely deals with linear terms).

There have been several studies for $\varepsilon = 1$ when the force comes from a potential $\vec{E}_t(x) = \nabla_x V_t(x)$. The latest result in this setting seems to be [28] and deals with large potential in an L^∞ framework, we refer to the references therein for the potential force framework. This framework is however irrelevant to derive Navier-Stokes-Fourier system with force because one can only construct Leray solutions from Boltzmann-type equations and such solutions do not see gradient terms.

At last we would like to present a related issue, still for $\varepsilon = 1$, that is when the force is nonlinear: $\vec{E}_t(x) = \vec{E}_t[f](x)$. This happens in electromagnetism for instance. Several results have been obtained in these settings in Sobolev space for perturbation of the global equilibrium. The advantage of this nonlinearity is a feedback that keeps the smallness of the force along the flow. We point out Vlasov-Poisson-Boltzmann equations [26, 17, 18, 36, 37] or Vlasov-Maxwell-Boltzmann equations [15], the strategies of which will prove themselves useful in our methods. See also [3] for a non-perturbative approach to those systems.

One of the main issue when dealing with the Boltzmann equation with an external force comes from the fact that the perturbative regime $F = \mu + f$ gives rise to a differential equation in f that includes the term $\vec{E}_t(x)(x) \cdot v f$. The latter generates a loss of weight in standard Sobolev estimates. The latest result we are aware of for

general non-potential forces comes from [16], where the authors work in the whole spatial domain \mathbb{R}^d close to μ and $\vec{E}_t(x)$ is assumed to be small and time-decaying like $(1+t)^{-\alpha}$ if $3 \leq d < 5$. Moreover their collision operator must satisfy the hard spheres assumption $\gamma = 1$ and $b(\cos \theta) = 1$. Working with a hard sphere kernel was mandatory in [16] to compensate the loss of weight in v , by means of the negative feedback of the linear Boltzmann operator, which generates a gain of $1 + |v|^\gamma$ (see (3.9)). Note however, that when only studying the semigroup generated by the linear part of the perturbative regime they do not need any time-decay for \vec{E}_t . It is important to understand that μ is no longer a stationary state when $\vec{E}_t(x) \neq 0$ so one hope that μ shall be stable when the force is very small or in the limit ε tends to 0 where formal Chapman-Enskog expansion easily shows that the first order term must be μ . To deal with non small force we propose a different regime.

The idea we have arises from the time-dependent norms proposed in [18, 37, 15], that compensate the increase of weight due to the nonlinearity of the external force. On the other hand, in a completely different setting, [1] linked the external force in a fractional Vlasov-Fokker-Planck equation to a new equilibrium that evolves with the external force. The new equilibrium can be explicitly written for the fractional Vlasov-Fokker-Planck equation whereas the non-local part of the Boltzmann equation seems to prevent such a direct treatment. However, we try to combine the two point of views described above : we cannot explicitly extract a new equilibrium for the Boltzmann equation with external force so we fake it by studying the equation around a Maxwellian distribution that depends on $\vec{E}_t(x)$. Such an approach sees the external force as a fluctuation of the classical characteristics of the Boltzmann equation rather than a direct interaction on the solution. We are therefore able to relax the hard sphere assumption, as the loss of weight generated by the external force is effectively compensated, although not by the non-positivity of the linear operator, but thanks to the negative feedback offered by the fluctuation of the Maxwellian. When $\vec{E}_t(x)$ is time-decaying, not only our strategy works for general hard potential kernels with angular cut-off, but it also enables to treat large forces. The core of the proof relies on the construction of twisted Sobolev norms, in the spirit of [32, 12], which pushes out the hypocoercivity of the Boltzmann linear equation. Namely, the commutator $[v \cdot \nabla_x, \nabla_v] = -\nabla_x$ offers a full negative feedback on x derivatives and one thus would like to work with functional of the form

$$\|f\|^2 = a \|f\|_{L_{x,v}^2}^2 + b \|\nabla_x f\|_{L_{x,v}^2}^2 + c \|\nabla_v f\|_{L_{x,v}^2}^2 + d \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2}$$

and equivalently in $H_{x,v}^s$ regularity. The mixed term in the twisted norm uses the commutator property and is sufficient in the classical case $\vec{E}_t = 0$. The presence of the external force, however, requires a more subtle use of this mixed part that will have to compensate much more terms arising from pure spatial derivatives. We shall see the interplay between the negative feedback offered by the commutator on one side and the one offered by the fluctuation on the other side. The main issue being the negative feedback coming only from the orthogonal part of the solution when dealing with pure x derivatives. Using commutator for fixed pure x -derivatives proved itself sufficient for the classical Boltzmann equation $\vec{E}_t = 0$ but in our case they have to be dealt with at the same time.

Unfortunately, when $\vec{E}_t(x)$ does not display a time-decaying property we can only use this strategy on fixed time intervals $[0, T_0]$, for any $T_0 > 0$, but not globally in

time. However, and of important note, in the hydrodynamical regime $\varepsilon \rightarrow 0$ our method provides solutions close to the global maxwellian $\mu(v)$.

All these thoughts make us look at the perturbative regime around
(1.17)

$$\forall(t, v) \in [0, +\infty) \times \mathbb{R}^d, \quad M(t, v) = \frac{e^{-\varepsilon^{1+e} \frac{A}{1+t}}}{(2\pi)^{d/2}} e^{-\frac{|v|^2}{2} (1 + \varepsilon^{1+e} \frac{a}{1+t})} = \mu e^{-\varepsilon^{1+e} \frac{A+a \frac{|v|^2}{2}}{1+t}}$$

where A and a stand for positive constant that we shall define in due time. Of core importance, e has to belong to $(0, 1)$. We refer to Remark 4.3 and Remark 4.7 to understand that when $e = 0$ then the fluctuation M is not close enough to μ to perform a relevant hydrodynamical limit whereas when $e = 1$ the fluctuation goes to fast towards μ compared to the variations of the characteristics.

We study the perturbative regime

$$(1.18) \quad \forall(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d, \quad F(t, x, v) = M(t, x) + \varepsilon M^{\frac{1}{2}} f(t, x, v)$$

which leads to the following perturbative equation

$$(1.19) \quad \partial_t f + \frac{1}{\varepsilon} v \cdot \nabla_x f + \varepsilon \vec{E}_t(x) \cdot \nabla_v f + \varepsilon \mathcal{E}(t, x, v) f = \frac{1}{\varepsilon^2} L[f] + \frac{1}{\varepsilon} \Gamma[f, f] - 2\mathcal{E}(t, x, v) M^{1/2}$$

where L and Γ are respectively the standard linear and bilinear perturbative Boltzmann operators around M

$$\begin{aligned} L[f] &= \frac{2}{\sqrt{M}} Q(M, \sqrt{M} f) \\ \Gamma[f, f] &= \frac{1}{\sqrt{M}} Q(\sqrt{M} f, \sqrt{M} f) \end{aligned}$$

and we defined the perturbative force term

$$(1.20) \quad \forall(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d, \quad \mathcal{E}(t, x, v) = \frac{1}{2} \left(\varepsilon^e \frac{A + a \frac{|v|^2}{2}}{(1+t)^2} - \left(1 + \varepsilon^{1+e} \frac{a}{1+t} \right) \vec{E}_t(x) \cdot v \right).$$

We conjecture that, by use of the maxwellian regularising properties of the compact part of L , one could directly solve the Cauchy problem around a global maxwellian $F = \mu + \varepsilon \sqrt{\mu} f$ for f in $H_{x,v}^s \left(e^{\varepsilon^2 \frac{A+a|v|^{1+0}}{1+t}} \right)$ when \vec{E}_t or ε are sufficiently small.

However, it would implies some technicalities we did not want to tackle in the present manuscript, where we are only interested in the limit when ε vanishes: working in $L_{x,v}^2$ framework makes usual properties of $L[f]$ directly applicable, thus our proofs only emphasizes the characteristics fluctuations. Moreover, the strategy we use enables to compensate a quadratic loss of weight $|v|^2$ (rather than the sole $|v|$ specific to the present problem), which may suit further investigations for more complex forces.

2. MAIN RESULTS

2.1. Notations. For $j = (j_1, \dots, j_d)$ and $l = (l_1, \dots, l_d)$ multi-indexes we define

$$\partial_l^j f = \frac{\partial^{|l|}}{\partial x_1^{l_1} \dots \partial x_d^{l_d}} \frac{\partial^{|j|}}{\partial v_1^{j_1} \dots \partial v_d^{j_d}} f.$$

And we define the multi-index: $\forall i \in \{1, \dots, d\}$, $\delta_i = (\delta_{ik})_{1 \leq k \leq d}$. For clarity purposes, and as it plays a central role for the linearized Boltzmann operator we shall use the shorthand notation

$$\forall \beta \in \mathbb{R}, \quad \|f\|_{L_\beta^2} = \left(\int_{\mathbb{T}^d \times \mathbb{R}^d} f(t)^2 \langle v \rangle^\beta dx dv \right)^{1/2}, \quad \langle v \rangle := (1 + |v|^2)^{1/2}.$$

Finally, we shall index by x , v or (x, v) the norms that will be used:

$$\|f\|_{L_x^2} = \left(\int_{\mathbb{T}^d} f(x, v)^2 dx \right)^{1/2}, \quad \|f\|_{L_v^2} = \left(\int_{\mathbb{R}^d} f(x, v)^2 dv \right)^{1/2},$$

$$\|f\|_{L_{x,v}^2} = \left(\int_{\mathbb{T}^d \times \mathbb{R}^d} f(x, v)^2 dx dv \right)^{1/2}.$$

The same notations apply for Sobolev spaces H_x^s (only x -derivatives), H_v^s (only v -derivatives) and $H_{x,v}^s$ (both derivatives).

In what follows any positive constant depending on a parameter α will be denoted C_α . Note that we will not keep track on the dependencies over d , γ or $b(\cos \theta)$.

2.2. Results on Cauchy theories. When \vec{E}_t is a given force we shall prove the following Cauchy problem. We recall Definition (1.17) of the fluctuation of a global Maxwellian

$$\forall (t, v) \in [0, +\infty) \times \mathbb{R}^d, \quad M(t, v) = \frac{e^{-\varepsilon^{1+e} \frac{A}{1+t}}}{(2\pi)^{d/2}} e^{-\frac{|v|^2}{2} (1 + \varepsilon^{1+e} \frac{a}{1+t})} \quad \text{with } e \in (0, 1).$$

We get a Cauchy theory under the perturbative regime around a given M .

Theorem 2.1. *Let the Boltzmann operator satisfies hypotheses (H1) – (H2) and let s be in \mathbb{N} . Further assume that \vec{E}_t verifies*

$$(2.1) \quad \left\| \vec{E}_t(x) \right\|_{L_t^\infty W_x^{s, \infty}} \leq C_E, \quad \forall t \geq 0, \quad \int_{\mathbb{T}^d} \vec{E}_t(x) dx = 0.$$

There exists $s_0 \in \mathbb{N}^$ such that for any $s \geq s_0$ the following holds. Let $T_0 > 0$ and $C_{\text{in}} \geq 0$. There exists $\varepsilon_{T_0, C_{\text{in}}, E, s} > 0$ such that if $\varepsilon = 1$ or $0 < \varepsilon < \varepsilon_{T_0, C_{\text{in}}, E, s}$, there exists a norm*

$$\|\cdot\|_{\mathcal{H}_{x,v}^s} \sim \sum_{|l| \leq s} \|\partial_l^0 \cdot\|_{L_{x,v}^2} + \varepsilon \sum_{\substack{|j|+|l| \leq s \\ |j| \geq 1}} \|\partial_l^j \cdot\|_{L_{x,v}^2}$$

and $A_{T_0, E, s}$, $a_{T_0, E, s}$, $\delta_{T_0, E, s}$, $C_{T_0, E, s} > 0$ such that if $F_{\text{in}} = M + \varepsilon \sqrt{M} f_{\text{in}}$ with

$$\|f_{\text{in}}\|_{\mathcal{H}_{x,v}^s} \leq \delta_{T_0, E, s} \quad \text{and} \quad \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} f_{\text{in}}(x, v) \sqrt{M}|_{t=0} dx dv \right| \leq C_{\text{in}} \varepsilon^e,$$

then there exists a unique solution $F = M + \varepsilon \sqrt{M} f$ on $[0, T_0)$ to the Boltzmann equation with external force (1.5) and it satisfies

$$\forall t \in [0, T_0), \quad \|f(t)\|_{\mathcal{H}_{x,v}^s} \leq \max \left\{ \|f_{\text{in}}\|_{\mathcal{H}_{x,v}^s}, C_{T_0, E, s} \right\}.$$

All the constants can be computed explicitly and are independent of ε .

Let us make a few comments about the theorem above.

Remark 2.2. • The $\mathcal{H}_{x,v}^s$ -norm is defined by (4.25) and the equivalence of norm is independent of ε .

- The hypothesis of cancellation of the first Fourier coefficient $\widehat{\vec{E}}_t(0)$ in (2.1) is not really necessary¹. Since the Boltzmann operator is commuting with translations in v , the change of variable

$$v' = v + \varepsilon w_t, \quad w_t := \int_0^t \widehat{\vec{E}}_s(0) ds$$

operates a reduction to the case where $\widehat{\vec{E}}_t(0) = 0$. In the case of a general forcing term satisfying only the bound by C_E in (2.1), the result of Theorem 2.1 holds true, except that the decomposition of F has to be modified into the following expansion:

$$F(t, x, v) = M(t, v - \varepsilon w_t) + \varepsilon \sqrt{M}(v - \varepsilon w_t) f(t, x, v - \varepsilon w_t),$$

where f is solution to (1.19) with a forcing term $\vec{E}_t' := \vec{E}_t - \widehat{\vec{E}}_t(0)$.

- We get a local existence result for $\varepsilon = 1$ for a non-small, non time-decreasing force and with more general kernels than considered formerly (only hard spheres has been obtained to our knowledge [16]). This is the first result of this kind we are aware of. Moreover, we get close-to-global Maxwellian μ existence for small C_E , recovering and extending on the torus the latest results (see Remark 2.4).
- We agree that in the case $\varepsilon = 1$ when C_E or T_0 are taken larger and larger, the fluctuation M is getting closer to 0 and our problem thus boils down to the perturbative study around a vacuum state. However, as proven in Corollary 2.3, in the regime of small epsilon we obtain a perturbative theory around the classical Maxwellian μ .
- We do not have to impose any decay in time on the force. On the contrary a polynomial decay is used, for instance, in [16]. As explained in the introduction, this comes from the use of a time-dependent Maxwellian as reference state. We think that our strategy is applicable if one looks at solution $F = \mu + \varepsilon \sqrt{\mu} h$ with h belonging to $H_{x,v}^s \left(e^{\varepsilon^{1+e} \frac{A+a|v|^{1+0}}{1+t}} \right)$ for small ε , by use of a gain of integrability of $K = L + \nu$. It would have been more technical to treat this case, and we thus decided to take a clearer approach which is sufficient to deal with the issue of the hydrodynamical limit.
- Note that the hypothesis

$$(2.2) \quad \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} f_{in}(x, v) \sqrt{M}|_{t=0} dx dv \right| \leq C_{in} \varepsilon^e$$

says that some global moments of f_{in} in both the space and velocity variables are small with ε . Actually, we may replace the right-hand side $C_{in} \varepsilon^e$ of (2.2) by any quantity that tends to 0 with ε . In the following Remark 2.6, we comment the incidence of (2.2) on the initial data for the solution to (1.15)-(1.3).

¹However, as such, it ensures a condition of quasiconservation of the total momentum in Navier-Stokes equation, which is used to obtain the estimate (4.17)

As a corollary of Theorem 2.1 we obtain a global perturbative Cauchy theory close to μ .

Corollary 2.3. *Under the assumptions of Theorem 2.1 there exists $\varepsilon_{T_0, E, s}$, $\delta_{T_0, E, s}$ and $C_{T_0, E, s}$ such that if $0 < \varepsilon \leq \varepsilon_{T_0, E, s}$ and $F_{\text{in}} = \mu + \varepsilon\sqrt{\mu} f_{\text{in}}$ with*

$$\|f_{\text{in}}\|_{\mathcal{H}_{x,v}^s} \leq \delta_{T_0, E, s} \quad \text{and} \quad \int_{\mathbb{T}^d \times \mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} f_{\text{in}}(x, v) \sqrt{\mu} \, dx dv = 0,$$

then there exists a unique solution $F = \mu + \varepsilon\sqrt{\mu} f$ on \mathbb{R}^+ to the Boltzmann equation with external force (1.5) and it satisfies

$$\forall t \in [0, T_0), \quad \|f(t)\|_{\mathcal{H}_{x,v}^s} \leq \max \left\{ \|f_{\text{in}}\|_{\mathcal{H}_{x,v}^s} + \varepsilon^e C_{T_0, E, s}, C_{T_0, E, s} \right\}.$$

Again, all the constants could be computed explicitly and are independent of ε .

Remark 2.4. *Two remarks are important at this point.*

- We emphasize that taking ε sufficiently small could be seen as requiring C_E to be sufficiently small in our estimates. The strenght of our result is that the resulting Incompressible Navier-Stokes limit can display non small force \vec{E}_t .
- We point out that $T_0 = +\infty$ is not reached in our study because of the negative return of our fluctuation that only works for finite T_0 (see Proposition 3.1). However, adapting our proofs when $\|\vec{E}_t\| \leq \frac{C_E}{(1+t)^\alpha}$ with $\alpha > 1$ around $M = \mu e^{\varepsilon^{1+e} \frac{A+a|v|^2}{(1+t)^{\alpha-1}}}$ gives that not only Proposition 3.1 is true for $T_0 = +\infty$ but also Corollary 2.3 holds globally in time and yields a polynomial time decay (see Remark 5.1): these are the results of [16] when $\varepsilon = 1$ which we recover on the torus.

2.3. Result on the hydrodynamical limit. Corollary 2.3 states that if one relabels the solution $f_\varepsilon = \varepsilon^{-1}(F - \mu)\mu^{-\frac{1}{2}}$ then we have uniform bounds on f_ε in Sobolev spaces and an existence theorem that does not depend on ε for the initial data. This yields the following convergence result.

Theorem 2.5. *Let $T_0 > 0$ and F_ε be the solution built in Corollary 2.3 on $[0, T_0]$ and define $f_\varepsilon = \varepsilon^{-1} \frac{F_\varepsilon - \mu}{\sqrt{\mu}}$. Then the sequence $(f_\varepsilon)_{\varepsilon > 0}$ converges (up to an extraction) weakly- $*$ in $L_{[0, T_0]}^\infty H_{x,v}^s$ towards an infinitesimal Maxwellian:*

$$(2.3) \quad \lim_{\varepsilon \rightarrow 0} f_\varepsilon(t, x, v) = \left[\rho(t, x) + v \cdot u(t, x) + \frac{|v|^2 - d}{2} \theta(t, x) \right] \sqrt{\mu},$$

where (ρ, u, θ) solves the incompressible Navier-Stokes-Fourier system in the sense of Leray with force (1.15) together with the Boussinesq equation (1.3).

Remark 2.6. *If the data are well prepared in the sense that f_{in} is of the form*

$$(2.4) \quad f_{\text{in}}(x, v) = \left[\rho_{\text{in}}(x) + v \cdot u_{\text{in}}(x) + \frac{|v|^2 - d}{2} \theta_{\text{in}}(x) \right] \sqrt{\mu},$$

with $\nabla \cdot u_{\text{in}} = 0$ and $\rho_{\text{in}} + \theta_{\text{in}} = 0$ and

$$(2.5) \quad \int_{\mathbb{T}^d} \rho_{\text{in}}(x) dx = 0, \quad \int_{\mathbb{T}^d} u_{\text{in}}(x) dx = 0$$

((2.5) is a consequence of (2.2)), then we expect (by adaptation of the arguments of the proof of [12, Theorem 2.5]) to have strong convergence in (2.3) in the space $C([0, T]; H_x^s L_v^2)$, and (ρ, u, θ) to be a regular solution to (1.15)-(1.3) on $[0, T_0]$ with initial datum $(\rho_{\text{in}}, u_{\text{in}}, \theta_{\text{in}})$.

3. PROPERTIES AND ESTIMATES ON THE EXTERNAL OPERATOR AND THE BOLTZMANN OPERATOR

In the present section we focus on the linear operator $L[f]$, the multiplicative $\mathcal{E}(t, x, v)$ and then on all the operators appearing in the perturbed equation (1.19).

3.1. Estimates on the external operator $\mathcal{E}(t, x, v)$. In this section, we will give some estimates on $\mathcal{E}(t, x, v)$ and fix the constants a , A and α for the rest of the manuscript. A consequence on the notations used in the paper is that we will drop the specific dependence of constants on the values a , A and α . Since $\vec{E}_t(x)$ is a datum of the problem on which no smallness assumption is required, we will also drop the possible dependency on E : $C_{a,A,\alpha,E} = C$, except in the Proposition below so that the reader can clearly see the improvement we can make when \vec{E}_t decreases in time, that is $C_E \sim (1+t)^{-\alpha}$ as mentioned in Remark 2.4.

Proposition 3.1. *Let \mathcal{E} be defined by (1.20) and let s be any integer. Under the assumption*

$$\left\| \vec{E}_t(x) \right\|_{L_t^\infty W_x^{s,\infty}} \leq C_E < +\infty,$$

for any T_0 and $\Lambda > 0$, there exists $A_{T_0,\Lambda}$ and $a_{T_0,\Lambda} > 0$ such that the following properties hold. Positivity:

$$(3.1) \quad \forall (t, x, v) \in [0, T_0) \times \mathbb{T}^d \times \mathbb{R}^d, \quad \mathcal{E}(t, x, v) \geq \varepsilon^e \Lambda (1 + |v|^2) - \frac{1}{4\varepsilon^e} \mathbf{1}_{\varepsilon < 1}.$$

Pure spatial derivative estimates: if $s \geq 1$

$$(3.2) \quad \exists C_{T_0,s,\Lambda} > 0, \forall |l| \geq 1, \forall (t, x, v) \in [0, T_0) \times \mathbb{T}^d \times \mathbb{R}^d, \quad |\partial_t^l \mathcal{E}(t, x, v)| \leq C_{T_0,s,\Lambda} |v|.$$

Second derivative estimates: if $s \geq 2$

$$(3.3) \quad \exists C_{T_0,s,\Lambda} > 0, \forall |j| \geq 1 \text{ and } |j| + |l| \geq 2, \forall (t, x, v) \in [0, T_0) \times \mathbb{T}^d \times \mathbb{R}^d, \quad |\partial_t^j \mathcal{E}| \leq C_{T_0,s,\Lambda}.$$

Higher order derivatives in v : if $s \geq 3$

$$(3.4) \quad \forall |j| \geq 2 \text{ and } |j| + |l| \geq 3, \quad |\partial_t^j \mathcal{E}(t, x, v)| = 0.$$

Proof of Proposition 3.1. The Proposition is rather straightforward. We recall:

$$\mathcal{E}(t, x, v) = \frac{\varepsilon^e (2A + a|v|^2)}{4(1+t)^2} - \frac{1}{2} \left(1 + \varepsilon^{1+e} \frac{a}{1+t} \right) \vec{E}_t(x) \cdot v$$

First a mere Cauchy-Schwarz inequality followed by Young inequality raises that for all t in $[0, T_0)$

$$\begin{aligned} \mathcal{E}(t, x, v) &\geq \varepsilon^e \frac{2A + a|v|^2}{4(1+t)^2} - \frac{C_E}{2}|v| - \frac{\varepsilon^{1+e} a C_E}{2(1+t)}|v| \\ &\geq \varepsilon^e \left[\frac{A}{2(1+t)^2} - \varepsilon^2 a C_E^2 \right] + \varepsilon^e |v|^2 \left[\frac{a}{8(1+t)^2} - \frac{C_E^2}{4} \right] - \frac{1}{4\varepsilon^e} \end{aligned}$$

So we can first choose $a = a_{T_0, \Lambda}$ sufficiently large so that

$$\frac{a}{8(1+T_0)^2} - \frac{C_E^2}{4} \geq \Lambda$$

and then choose $A = A_{T_0, \Lambda}$ sufficiently large so that

$$\frac{A}{2(1+T_0)^2} - a C_E^2 \geq \Lambda.$$

This yields the positivity property (3.1) because if $\varepsilon = 1$ we can make A larger so that $1/4$ is also absorbed. The rest of the estimates are direct computations once constant have been fixed. Note that the constants are independent of t since $\frac{1}{1+t} \leq 1$. \square

Remark 3.2. *Of important note is the fact that Λ shall be fixed later independently of ε : the value of Λ is determined in Proposition 4.5. We shall carefully keep track of the dependencies in Λ to ensure that no bad loop can occur.*

3.2. Known properties of the Boltzmann operator. We gather some well-known properties of the linear Boltzmann operator L (see [13, 14, 35, 25] for instance). For $1 \leq i \leq d$, let us set

$$\phi_0(v) = 1, \quad \phi_i(v) = v_i, \quad \phi_{d+1}(v) = \frac{1}{2}(|v|^2 - d).$$

The operator L (which is time-dependent) is a closed self-adjoint operator in L_v^2 with kernel

$$(3.5) \quad \text{Ker}(L) = \text{Span} \{1, v, |v|^2\} \sqrt{M} = \text{Span} \{\phi_0(v), \dots, \phi_{d+1}(v)\} \sqrt{M},$$

and $(\phi_0 \sqrt{M}, \cdot, \phi_{d+1} \sqrt{M})$ is an orthogonal basis of $\text{Ker}(L)$ in L_v^2 . We denote by π_L the orthogonal projection onto $\text{Ker}(L)$ in L_v^2 :

$$(3.6) \quad \pi_L(f) = \sum_{i=0}^{d+1} \left(\int_{\mathbb{R}^d} f(v_*) \bar{\phi}_i(v_*) \sqrt{M(t, v_*)} dv_* \right) \bar{\phi}_i(v) \sqrt{M(t, v)},$$

where we have used the normalized family

$$\bar{\phi}_0 = \phi_0, \quad \bar{\phi}_1 = \phi_1, \dots, \quad \bar{\phi}_d = \phi_d, \quad \bar{\phi}_{d+1} = \sqrt{\frac{d}{2}} \phi_{d+1}.$$

We set $\pi_L^\perp = \text{Id} - \pi_L$. The projection $\pi_L(f(x, \cdot))(v)$ of $f(x, v)$ onto the kernel of L is called the fluid part whereas $\pi_L^\perp(f)$ is called the microscopic part.

The operator L can be written under the following form

$$(3.7) \quad L = -\nu(v) + K,$$

where $\nu(v)$ is the collision frequency

$$\nu(v) = \int_{\mathbb{R}^d \times \mathbb{S}^{d-1}} b(\cos \theta) |v - v_*|^\gamma M_* d\sigma dv_*$$

and K is a bounded and compact operator in L_v^2 . We give a series of estimates on the operators above that have been proved in the case $a = A = 0$ in references we gave above. We solely empasize that the constants do not depend on t .

Proposition 3.3. *Let s be in \mathbb{N} , $|l| + |j| \leq s$ and f be in $H_{x,v}^s$.*

The collision frequency is strictly positive

$$(3.8) \quad \forall v \in \mathbb{R}^d, \quad 0 < \nu_{0,s,\Lambda} \langle v \rangle^\gamma \leq \nu(v) \leq \nu_{1,s,\Lambda} \langle v \rangle^\gamma.$$

The operator L acts on the v -variable and has a spectral gap $\lambda_{0,\Lambda} > 0$ in $L_{x,v}^2$

$$(3.9) \quad \langle \partial_l^0 L(f), \partial_l^0 f \rangle_{L_{x,v}^2} \leq -\lambda_{0,\Lambda} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2.$$

There exists $\lambda_{s,\Lambda}$, $C_{s,\Lambda} > 0$ such that, if $|j| \geq 1$, then

$$(3.10) \quad \langle \partial_l^j L[f], \partial_l^j f \rangle_{L_{x,v}^2} \leq -\lambda_{s,\Lambda} \|\partial_l^j f\|_{L_\gamma^2}^2 + C_{s,\Lambda} \|f\|_{H_{x,v}^{s-1}}^2.$$

At last we have the following estimates on scalar products: for $0 \leq |l| \leq s-1$ and any $\eta_0 > 0$ there exists C_{Λ,η_0} such that:

$$(3.11) \quad \langle \partial_{l+\delta_i}^0 L[f], \partial_{l+\delta_i}^{\delta_i} f \rangle_{L_{x,v}^2} = \frac{\lambda_{0,\Lambda} C_{\Lambda,\eta_0}}{\varepsilon} \|\pi_L^\perp(\partial_{l+\delta_i}^0 f)\|_{L_\gamma^2} + \varepsilon \lambda_{0,\Lambda} \eta_0 \|\partial_{l+\delta_i}^{\delta_i} f\|_{L_\gamma^2}^2.$$

Before getting into the proof of Proposition 3.3, let us emphasize that the multiplication by $\lambda_{0,\Lambda}$ in the scalar product estimate is of core importance for the case $\varepsilon = 1$.

Proof of Proposition 3.3. If we denote by $L_\mu = -\nu_\mu + K_\mu$ the linear operator when $A = a = 0$ then the results hold for L_μ , ν_μ and K_μ : see for instance [4, 31] for the spectral gap, [32, -(H1')+(H2')] page 13] for Sobolev estimates and [12, Appendix B.2.3 and B.2.5] for the scalar product.

The operator L only acts on the velocity variable thus the change of variable

$$v_* \mapsto \left(\sqrt{1 + \frac{\varepsilon^{1+e_a}}{1+t}} \right)^{-1} v_*$$

shows

$$(3.12) \quad L[f](v) = \left(1 + \frac{\varepsilon^{1+e_a}}{1+t} \right)^{-\frac{d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e_a} A}{1+t}} L_\mu[\tilde{f}] \left(v \sqrt{1 + \frac{\varepsilon^{1+e_a}}{1+t}} \right)$$

where $\tilde{f}(v) = f\left(\frac{v}{\sqrt{1 + \frac{\varepsilon^{1+e_a}}{1+t}}}\right)$. As $0 \leq (1+t)^{-1} \leq 1$, inequalities (3.8) – (3.9) – (3.10) directly follow from the case $a = A = 0$.

The scalar product (3.11) is a mere Cauchy-Schwarz inequality with Young inequality with constant η_0 . Let us show that the resulting constant is of the form $C_{\eta_0} \lambda_{0,\Lambda}$. Denoting $h = 1 + \frac{\varepsilon^{1+e_a}}{1+t}$, integrating (3.12) against f yields

$$\begin{aligned} \langle L[f], f \rangle_{L_{x,v}^2} &= h^{-\frac{2d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e_a} A}{1+t}} \langle L_\mu[\tilde{f}], \tilde{f} \rangle_{L_{x,v}^2} \leq -h^{-\frac{2d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e_a} A}{1+t}} \|\widetilde{\pi_{L_\mu}^\perp}(\tilde{f})\|_{L_\gamma^2}^2 \\ &\leq -\lambda_{0,0} h^{-\frac{d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e_a} A}{1+t}} \|\pi_L^\perp(f)\|_{L_\gamma^2}^2. \end{aligned}$$

We thus see that $\lambda_{0,\Lambda} = \lambda_{0,0} h^{-\frac{d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e}A}{1+t}}$ and $h^{-\frac{d+\gamma}{2}} e^{-\frac{\varepsilon^{1+e}A}{1+t}}$ is exactly the quantity appearing in (3.12) so the expected (3.11) follows. \square

We conclude the present section with estimates on the bilinear operator $\Gamma[f, g]$.

Proposition 3.4. *Let $s_0 = d + 1$ then for any $s \geq s_0$, for any $|j| + |l| \leq s$ and any $\eta_0, \eta'_0 > 0$, the following holds for f, g and h in $H_{x,v}^{|j|+|l|}$,*

$$\begin{aligned} \left| \langle \partial_l^0 \Gamma(g, h), f \rangle_{L_{x,v}^2} \right| &\leq \frac{\lambda_{0,\Lambda} \eta_0}{\varepsilon} \left\| \pi_L^\perp(f) \right\|_{L^2}^2 \\ &\quad + \varepsilon \lambda_{0,\Lambda} C_{s,\eta_0} \left(\|g\|_{H_x^s L_v^2}^2 \|h\|_{H_x^s L_v^2, \gamma}^2 + \|h\|_{H_x^s L_v^2}^2 \|g\|_{H_x^s L_v^2, \gamma}^2 \right) \\ \left| \langle \partial_l^j \Gamma(g, h), f \rangle_{L_{x,v}^2} \right| &\leq \frac{\eta'_0}{\varepsilon} \|f\|_{L_\gamma^2}^2 + \varepsilon C_{s,\Lambda, \eta'_0} \left(\|g\|_{H_{x,v}^s}^2 \|h\|_{H_\gamma^s}^2 + \|h\|_{H_{x,v}^s}^2 \|g\|_{H_\gamma^s}^2 \right). \end{aligned}$$

Proof of Proposition 3.4. As above, these estimates have been obtained when $A = a = 0$ and therefore the same arguments as before extend them to the general case. We refer the reader to [12, Appendix A.2] for constructive proofs in the case $A = a = 0$. We find the standard control

$$\begin{aligned} \left| \langle \partial_l^0 \Gamma(g, h), f \rangle_{L_{x,v}^2} \right| &\leq C_{s,\Lambda} \left(\|g\|_{H_x^s L_v^2} \|h\|_{H_x^s L_v^2, \gamma} + \|h\|_{H_x^s L_v^2} \|g\|_{H_x^s L_v^2, \gamma} \right) \left\| \pi_L^\perp(f) \right\|_{L_\gamma^2}, \\ \left| \langle \partial_l^j \Gamma(g, h), f \rangle_{L_{x,v}^2} \right| &\leq C_{s,\Lambda} \left(\|g\|_{H_{x,v}^s} \|h\|_{H_\gamma^s} + \|h\|_{H_{x,v}^s} \|g\|_{H_\gamma^s} \right) \|f\|_{L_\gamma^2}, \end{aligned}$$

that we complete with Young inequality. The dependency in $\lambda_{0,\Lambda}$ follows exactly from the same argument as in the proof of Proposition 3.3. \square

3.3. Estimates for each operator. The perturbative equation (1.19) that we shall study can be decomposed as the evolution by 6 different operators:

$$\begin{aligned} \partial_t f &= -\frac{1}{\varepsilon} v \cdot \nabla_x f - \varepsilon \vec{E}_t(x) \cdot \nabla_v f - \varepsilon \mathcal{E}(t, x, v) f + \frac{1}{\varepsilon^2} L[f] + \frac{1}{\varepsilon} \Gamma[f, f] - 2\mathcal{E}(t, x, v) M^{1/2} \\ &:= \sum_{i=1}^6 S_i(t, x, v). \end{aligned}$$

We prove a series of Lemmas to estimate each operator in Sobolev norms. To clarify the computations we shall use the convention that $\partial_l^j f = 0$ whenever the multi-indexes j or l contains one negative component. Thus any integration by parts can be computed.

Strategy. Our final aim is to get an estimate on the weighted norm (see (4.6))

$$(3.13) \quad f \mapsto \sum_{|l| \leq s} \left\| \partial_l^0 f \right\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|l|+|j| \leq s \\ |j| \geq 1}} \left\| \partial_l^j f \right\|_{L_{x,v}^2}^2.$$

Standard energy estimates will provide some gain and loss terms. The gain terms are due

- to the spectral gap estimates (3.9) and (3.10): they are

$$(3.14) \quad -\frac{\lambda_{0,\Lambda}}{\varepsilon^2} \sum_{|l| \leq s} \left\| \pi_L^\perp(\partial_l^0 f) \right\|_{L_{x,v}^2}^2, \quad -\lambda_{s,\Lambda} \sum_{\substack{|l|+|j| \leq s \\ |j| \geq 1}} \left\| \partial_l^j f \right\|_{L_{x,v}^2}^2;$$

- to the operator \mathcal{E} : associated to the derivative ∂_l^j , we have a gain (with weight $|v|^2$) which is $-\frac{\varepsilon^{1+e\Lambda}}{2} \|\partial_l^j f\|_{L_x^2}^2$ (see Lemma 3.7 below).

In a procedure that is standard for the derivation of hypocoercive estimates, we also introduce a correction by the twisted terms $\varepsilon \langle \partial_{l+\delta_i}^0 f, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2}$ in (3.13). Note those terms are pondered with a weight ε . Those terms will provide the gain term $-\|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2$ (see Lemma 3.6 below). Combining those latter terms with the terms of the second sum in (3.14), we obtain (up to the terms of order 0) a gain of almost a full $H_{x,v}^s$ -norm, having no weight ε . This is why the occurrence of a term $C\|f\|_{H_{x,v}^s}^2$ in the forthcoming estimates (Lemma 3.5 to 3.8) is admissible, a control on the size of the constant C being possibly necessary to ensure a good control when all estimates are gathered (which is done step by step in Proposition 4.1, Proposition 4.4, Proposition 4.5). In Proposition 4.5, we also study the evolution of the global moments of f^ε (in combination with a Poincaré-Wirtinger inequality), in order to recover the term of order 0 that is lacking in our estimates.

Lemma 3.5. *Let s be in \mathbb{N} and for f in $H_{x,v}^s$ define $S_1(t, x, v) = -\frac{1}{\varepsilon} v \cdot \nabla_x f$. Then for any $\eta_1 > 0$, there exists $C_{\eta_1} > 0$ such that for any multi-indexes l, j such that $|l| + |j| \leq s$,*

$$\left| \langle \partial_l^j S_1, \partial_l^j f \rangle_{L_{x,v}^2} \right| \leq \begin{cases} \frac{\eta_1}{\varepsilon^2} \|\partial_l^j f\|_{L_{x,v}^2}^2 + \frac{1}{\eta_1} \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 & \text{if } |j| \geq 1 \\ 0 & \text{if } j = 0. \end{cases}$$

We have moreover

$$\langle \partial_{l+\delta_i}^0 S_1, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} = -\frac{1}{2\varepsilon} \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2.$$

Proof of Lemma 3.5. By direct computations

$$\begin{aligned} \langle \partial_l^j (v \cdot \nabla_x f), \partial_l^j f \rangle_{L_{x,v}^2} &= \sum_{j_1+j_2=j} \sum_{k=1}^d \int_{\mathbb{T}^d \times \mathbb{R}^d} (\partial_0^{j_1} v_k) (\partial_{l+\delta_k}^{j_2} f) \partial_l^j f \, dx dv \\ &= \sum_{k=1}^d \int_{\mathbb{T}^d \times \mathbb{R}^d} v_k (\partial_{l+\delta_k}^j f) \partial_l^j f + \sum_{k=1}^d \int_{\mathbb{T}^d \times \mathbb{R}^d} \left(\partial_{l+\delta_k}^{j-\delta_k} f \right) \partial_l^j f \\ &= \sum_{k=1}^d \int_{\mathbb{T}^d \times \mathbb{R}^d} \left(\partial_{l+\delta_k}^{j-\delta_k} f \right) \partial_l^j f \, dx dv. \end{aligned}$$

We used the property $\partial_0^{j_1}(v_k) = 0$ if $(|j_1| \geq 2)$ or $(|j_1| = 1 \text{ and } j_1 \neq \delta_k)$. The first result then follows from Cauchy-Schwarz and Young inequalities: for any $\eta_1 > 0$,

$$\int_{\mathbb{T}^d \times \mathbb{R}^d} \left(\partial_{l+\delta_k}^{j-\delta_k} f \right) \partial_l^j f \, dx dv \leq \frac{\eta_1}{\varepsilon} \int_{\mathbb{T}^d \times \mathbb{R}^d} \left(\partial_{l+\delta_k}^{j-\delta_k} f \right)^2 \, dx dv + \frac{\varepsilon}{\eta_1} \int_{\mathbb{T}^d \times \mathbb{R}^d} (\partial_l^j f)^2 \, dx dv.$$

The second equality comes from direct integration by parts. \square

Lemma 3.6. *Let s be in \mathbb{N} and for f in $H_{x,v}^s$ define $S_2(t, x, v) = -\varepsilon \vec{E}_t(x) \cdot \nabla_v f$. Then for any $\eta_2 > 0$, there exists $C_{\eta_2} > 0$ such that for any multi-indexes l, j satisfying $|l| + |j| \leq s$, we have*

$$\left| \langle \partial_l^j S_2, \partial_l^j f \rangle_{L_{x,v}^2} \right| = 0 \quad \text{if } l = 0,$$

and

$$\begin{aligned} & \left| \langle \partial_l^j S_2, \partial_l^j f \rangle_{L_{x,v}^2} \right| \\ & \leq \varepsilon^{1+e} \eta_2 \Lambda \|\partial_l^j f\|_{L_{x,v}^2}^2 + \varepsilon^{1-e} \frac{C \eta_2}{\Lambda} \left(\sum_{1 \leq i, k \leq d} \|\partial_{l-\delta_i}^{j+\delta_k} f\|_{L_{x,v}^2}^2 + \|f\|_{H_{x,v}^{|j|+|l|-1}}^2 \right), \end{aligned}$$

if $|l| > 0$, where Λ is given by Proposition (3.1). We have moreover

$$\left| \langle \partial_{l+\delta_i}^0 S_2, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} \right| \leq \varepsilon^{2+e} \eta_2 \Lambda \|\partial_l^{\delta_i} f\|_{L_{x,v}^2}^2 + \frac{C \eta_2}{\varepsilon^e \Lambda} \sum_{k=1}^d \|\partial_l^{\delta_k} f\|_{L_{x,v}^2}^2 + \frac{C \eta_2}{\varepsilon^e} \|f\|_{H_{x,v}^{|l|}}^2.$$

Proof of Lemma 3.6. Here again direct computations give

$$\begin{aligned} \left| \langle \partial_l^j \left(\vec{E}_t(x) \cdot \nabla_v f \right), \partial_l^j f \rangle_{L_{x,v}^2} \right| &= \left| \sum_{k=1}^d \sum_{l_1+l_2=l} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 E_k(x) \partial_{l_2}^{j+\delta_k} f \partial_l^j f \, dx dv \right| \\ &= \left| \sum_{k=1}^d \sum_{\substack{l_1+l_2=l \\ |l_2| < |l|}} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 E_k(x) \partial_{l_2}^{j+\delta_k} f \partial_l^j f \, dx dv \right| \\ &\leq \left\| \vec{E}_t \right\|_{W_x^{s,\infty}} \sum_{k=1}^d \sum_{|l_2| < |l|} \int_{\mathbb{T}^d \times \mathbb{R}^d} \left| \partial_{l_2}^{j+\delta_k} f \right| \left| \partial_l^j f \right| \, dx dv \end{aligned}$$

and here again combining Cauchy-Schwarz and Young inequality with constant $\varepsilon^e \eta_2$ yields the expected result.

Let us look at the second estimate. We have

$$\langle \partial_{l+\delta_i}^0 S_2, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} = -\varepsilon \sum_{l_1+l_2=l+\delta_i} \sum_{k=1}^d \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 E_k \partial_{l_2}^{\delta_k} f \partial_l^{\delta_i} f \, dx dv.$$

The higher derivative appears when $l_2 = l + \delta_i$ and by integration by parts we see

$$\begin{aligned} \int_{\mathbb{T}^d \times \mathbb{R}^d} E_k \partial_{l+\delta_i}^{\delta_k} f \partial_l^{\delta_i} f \, dx dv &= - \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{\delta_i}^0 E_k \partial_l^{\delta_k} f \partial_l^{\delta_i} f - \int_{\mathbb{T}^d \times \mathbb{R}^d} E_k \partial_l^{\delta_k} f \partial_{l+\delta_i}^{\delta_i} f \\ &= - \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{\delta_i}^0 E_k \partial_l^{\delta_k} f \partial_l^{\delta_i} f - \int_{\mathbb{T}^d \times \mathbb{R}^d} E_k \partial_l^{\delta_i} f \partial_{l+\delta_i}^{\delta_k} f \end{aligned}$$

which implies

$$\int_{\mathbb{T}^d \times \mathbb{R}^d} E_k \partial_{l+\delta_i}^{\delta_k} f \partial_l^{\delta_i} f \, dx dv = -\frac{1}{2} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{\delta_i}^0 E_k \partial_l^{\delta_k} f \partial_l^{\delta_i} f \, dx dv$$

and therefore the result follows from Cauchy-Schwarz and Young inequalities with constant $\varepsilon^{1+e} \eta_2$. \square

Lemma 3.7. *Let s be in \mathbb{N} and for f in $H_{x,v}^s$ define $S_3(t, x, v) = -\varepsilon \mathcal{E}(t, x, v) f$, where we recall that \mathcal{E} is given by (1.20).*

Then there exists $C_3 > 0$ such that for any multi-indices l, j such that $|l| + |j| \leq s$,

$$\langle \partial_l^j S_3, \partial_l^j f \rangle_{L_{x,v}^2} \leq \begin{cases} -\frac{\varepsilon^{1+e} \Lambda}{2} \|\partial_l^j f\|_{L_{x,v}^2}^2 + \varepsilon^{1-e} \frac{\mathbf{1}_{\varepsilon < 1}}{4} \|\partial_l^j f\|_{L_{x,v}^2}^2 + \varepsilon^{1-e} \frac{C_3}{\Lambda} \|f\|_{H_{x,v}^{|j|+|l|-1}}^2 \\ -\varepsilon^{1+e} \Lambda \|f\|_{L_{x,v}^2}^2 + \frac{\varepsilon^{1-e}}{4} \mathbf{1}_{\varepsilon < 1} \|f\|_{L_{x,v}^2}^2 & \text{if } |j| + |l| = 0 \end{cases}$$

where Λ is given by Proposition (3.1). Moreover, for any $\eta_3 > 0$, there exists $C_{\eta_3} > 0$ such that

$$\begin{aligned} \left| \langle \partial_{l+\delta_i}^0 S_3, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} \right| &\leq \varepsilon^{2+e} \eta_3 \Lambda \left\| \partial_l^{\delta_i} f \right\|_{L_{x,v}^2}^2 + \varepsilon^{2-e} \Lambda C_{\eta_3} \left\| \partial_{l+\delta_i}^0 f \right\|_{L_{x,v}^2}^2 \\ &\quad + \varepsilon^{1-e} \frac{\mathbf{1}_{\varepsilon < 1}}{4} \left(\left\| \partial_{l+\delta_i}^0 f \right\|_{L_{x,v}^2}^2 + \left\| \partial_l^{\delta_i} f \right\|_{L_{x,v}^2}^2 \right) + \frac{C_{\eta_3}}{\Lambda \varepsilon^e} \|f\|_{H_{x,v}^{|l|}}^2. \end{aligned}$$

Proof of Lemma 3.7. The inequality for $|j| = |l| = 0$ is a direct consequence of Proposition 3.1 and more precisely (3.1). When $|j| + |l| > 0$ we compute

$$\langle \partial_l^j (\mathcal{E}f), \partial_l^j f \rangle_{L_{x,v}^2} = \sum_{j_1+j_2=j} \sum_{l_1+l_2=l} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^{j_1} \mathcal{E} \partial_{l_2}^{j_2} f \partial_l^j f \, dx dv.$$

Proposition 3.1 tells us that most of the derivatives of \mathcal{E} vanish: when $(|j_1| \geq 2$ and $|j_1| + |l_1| \geq 3)$. We therefore decompose the sum into three different parts:

$$\begin{aligned} \sum_{j_1+j_2=j} \sum_{l_1+l_2=l} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^{j_1} \mathcal{E} \partial_{l_2}^{j_2} f \partial_l^j f \, dx dv &= \int_{\mathbb{T}^d \times \mathbb{R}^d} \mathcal{E} (\partial_l^j f)^2 \\ &\quad + \sum_{\substack{l_1+l_2=l \\ |l_1| \geq 1}} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 \mathcal{E} \partial_{l_2}^j f \partial_l^j f \\ &\quad + \sum_{\substack{j_1+j_2=j \\ |l_1|=1}} \sum_{l_1+l_2=l} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^{j_1} \mathcal{E} \partial_{l_2}^{j_2} f \partial_l^j f \\ &\quad + \sum_{\substack{j_1+j_2=j \\ |l_1|=2}} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_0^{j_1} \mathcal{E} \partial_{l_2}^{j_2} f \partial_l^j f. \end{aligned}$$

Proposition 3.1 gives us the estimate of the first term, see (3.1). In the second and third terms $|\partial_{l_1}^{j_1} \mathcal{E}|$ is bounded by $C_s(1 + |v|)$, see (3.2) – (3.3). Finally, in the fourth term we have $|\partial_0^{j_1} \mathcal{E}|$ bounded by C_s . Hence using Cauchy-Schwarz and Young inequality with $\eta > 0$:

$$\begin{aligned} \langle \partial_l^j (\mathcal{E}f), \partial_l^j f \rangle_{L_{x,v}^2} &\geq \int_{\mathbb{T}^d \times \mathbb{R}^d} \varepsilon^e \Lambda (1 + |v|^2) (1 - 2\eta C_s) (\partial_l^j f)^2 \, dx dv \\ &\quad - \frac{\mathbf{1}_{\varepsilon < 1}}{4\varepsilon^e} \left\| \partial_l^j f \right\|_{L_{x,v}^2}^2 - \frac{C_{s,\Lambda,\eta}}{\varepsilon^e} \|f\|_{H_{x,v}^{|j|+|l|-1}}^2. \end{aligned}$$

We choose η sufficiently small and the result follows.

The second estimate is derived in the same spirit. We have

$$\langle \partial_{l+\delta_i}^0 S_2, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} = \sum_{l_1+l_2=l+\delta_i} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 \mathcal{E} \partial_{l_2}^0 f \partial_l^{\delta_i} f \, dx dv.$$

When $|l_1| \geq 1$ then $|l_2| \leq l$ and $|\partial_{l_1}^0 \mathcal{E}| \leq C_s \Lambda |v| \leq C_s \Lambda |v|$, by Proposition 3.1. Therefore using Cauchy-Schwarz and Young inequality with constant $\varepsilon^{1+e} \eta_3 > 0$ we

have

$$\left| \sum_{\substack{l_1+l_2=l+\delta_i \\ |l_1| \geq 1}} \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_{l_1}^0 \mathcal{E} \partial_{l_2}^0 f \partial_l^{\delta_i} f \, dx dv \right| \leq \int_{\mathbb{T}^d \times \mathbb{R}^d} \varepsilon^{1+e} \eta_3 \Lambda (1 + |v|^2) (\partial_l^{\delta_i} f)^2 \, dx dv \\ + \frac{\Lambda C_{s, \eta_3}}{\varepsilon^{1+e}} \|f\|_{H_{x,v}^{|l|}}^2.$$

At last, when $l_1 = 0$ we use Proposition 3.1 to bound $|\mathcal{E}| \leq \varepsilon^e C_s \Lambda (1 + |v|^2) + \frac{1-\varepsilon < 1}{4\varepsilon^e}$ and get with the Young inequality

$$\left| \int_{\mathbb{T}^d \times \mathbb{R}^d} \mathcal{E} \partial_{l+\delta_i}^0 f \partial_l^{\delta_i} f \, dx dv \right| \leq \varepsilon^{1+e} \Lambda \eta_3 \int_{\mathbb{T}^d \times \mathbb{R}^d} (1 + |v|^2) (\partial_l^{\delta_i} f)^2 \, dx dv \\ + \Lambda C_{s, \eta_3} \int_{\mathbb{T}^d \times \mathbb{R}^d} (1 + |v|^2) (\partial_{l+\delta_i}^0 f)^2 \, dx dv \\ + \frac{1-\varepsilon < 1}{4\varepsilon^e} \left(\|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 + \|\partial_l^{\delta_i} f\|_{L_{x,v}^2}^2 \right).$$

This concludes the proof. \square

It remains to estimate the last operator S_6 , which is a mere multiplicative operator.

Lemma 3.8. *Let s be in \mathbb{N} and for f in $H_{x,v}^s$ define $S_6(t, x, v) = -\mathcal{E}(t, x, v)M^{1/2}$, where we recall that \mathcal{E} is given by (1.20).*

Then for any $\eta_4 > 0$, there exists $C_{\Lambda, \eta_4} > 0$ such that for any multi-indexes l, j such that $|l| + |j| \leq s$, we have

$$\langle \partial_l^j S_6, \partial_l^j f \rangle_{L_{x,v}^2} \leq \begin{cases} \eta_4 \|\pi_L(\partial_l^0 f)\|_{L_{x,v}^2}^2 + C_{\Lambda, \eta_4} & \text{if } j = 0 \\ \frac{\eta_4}{\varepsilon^2} \|\partial_l^j f\|_{L_{x,v}^2}^2 + \varepsilon^2 C_{\Lambda, \eta_4} & \text{if } |j| \geq 1. \end{cases}$$

Moreover, we have

$$\left| \langle \partial_{l+\delta_i}^0 S_6, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} \right| \leq \frac{\eta_4 \lambda_{0, \Lambda}}{\varepsilon} \|\partial_l^{\delta_i} f\|_{L_{x,v}^2}^2 + \varepsilon C_{\Lambda, \eta_4}.$$

Proof of Lemma 3.8. We can use the estimates on \mathcal{E} derived in Proposition 3.1, that we multiply by the Maxwellian \sqrt{M} . Looking at the kernel of L given by (3.5) we see that $\mathcal{E}\sqrt{M}$ belongs to $\text{Ker}(L)$ and so does $\partial_l^0 \mathcal{E}\sqrt{M}$ for any multi-index l . Therefore by Cauchy-Schwarz inequality

$$\left| \langle \partial_l^0 (\mathcal{E}M^{1/2}), \partial_l^0 f \rangle_{L_{x,v}^2} \right| = \left| \langle \partial_l^0 (\mathcal{E}M^{1/2}), \pi_L(\partial_l^0 f) \rangle_{L_{x,v}^2} \right| \\ \leq \|\partial_l^0 (\mathcal{E}M^{1/2})\|_{L_{x,v}^2} \|\pi_L(\partial_l^0 f)\|_{L_{x,v}^2} \\ \leq C_{s, \Lambda} \|\pi_L(\partial_l^0 f)\|_{L_{x,v}^2}.$$

When there are v derivatives we still have $\partial_l^j (\mathcal{E}M^{1/2})$ that is a polynomial times a Maxwellian and therefore belongs to $L_{x,v}^2$. Thus

$$\left| \langle \partial_l^j (\mathcal{E}M^{1/2}), \partial_l^j f \rangle_{L_{x,v}^2} \right| \leq C_{s, \Lambda} \|\partial_l^j f\|_{L_{x,v}^2}.$$

Also for similar reasons

$$\left| \langle \partial_{l+\delta_i}^0 (\mathcal{E}M^{1/2}), \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} \right| \leq C_{s, \Lambda} \|\partial_l^{\delta_i} f\|_{L_{x,v}^2}.$$

Those three estimates yield the expected results using the Young inequality. \square

4. *A priori* ESTIMATES IN SOBOLEV SPACES

We provide here Sobolev estimates for the nonlinear perturbed equation (1.19). We shall work in twisted Sobolev norms that catch the hypocoercivity of the Boltzmann perturbed linear operator. Indeed, as shown by the estimates on the Boltzmann linear operator L , we do have a full negative feedback, and a gain of weight, as soon as ∂_l^j includes one velocity derivative. Unfortunately, the negative feedback offered by L on pure spatial derivative only controls the orthogonal part π_L^\perp . In the exact same spirit as [32, 12], a small portion of scalar product between spatial and velocity derivative is added to the standard Sobolev norm in order to take advantage of the commutator

$$[v \cdot \nabla_x, \nabla_v] = -\nabla_x.$$

We shall establish *a priori* estimates in Sobolev space to the perturbed equation (1.19) that we recall here

$$(4.1) \quad \begin{aligned} \partial_t f &= -\frac{1}{\varepsilon} v \cdot \nabla_x f - \varepsilon \vec{E}_t(x) \cdot \nabla_v f - \varepsilon \mathcal{E}(t, x, v) f + \frac{1}{\varepsilon^2} L[f] + \frac{1}{\varepsilon} \Gamma[f, f] - 2\mathcal{E}(t, x, v) M^{1/2} \\ &:= \sum_{i=1}^6 S_i(t, x, v). \end{aligned}$$

We gather the estimates derived in Section 3 to construct an equivalent Sobolev norm of f that can be controlled. As T_0 , a and $A > 0$ have been fixed in Proposition 3.1 we drop the dependencies on the subscripts. Also, as we shall always work with derivatives of order less than a given s , we drop the dependencies on s . Note however that a lot of different parameters are involved and so, to avoid any loop in their later choice, we will index the constants with these parameters, even if it complicates the reading: the important dependencies are Λ and η_i .

We shall address the velocity derivatives and the pure spatial derivatives at different orders in ε , in the spirit of [12]. In what follows we shall use the notation

$$(4.2) \quad \forall f \in H_{x,v}^s, \quad \|f\|_{\mathcal{H}_\varepsilon^s}^2 = \sum_{|l| \leq s} \|\partial_l^0 f\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|j|+|l| \leq s \\ |l| \geq 1}} \|\partial_l^j f\|_{L_{x,v}^2}^2.$$

4.1. Estimates for spatial derivatives. As mentioned at the beginning of the present section the pure x -derivatives in Sobolev spaces must be handled with the help of the transport operator. We thus define

$$(4.3) \quad \forall l \in \mathbb{N}^d, \forall 1 \leq i \leq d, \quad Q_{l,i}(f) = p \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 + q\varepsilon^2 \|\partial_l^{\delta_i} f\|_{L_{x,v}^2}^2 + \varepsilon r \langle \partial_{l+\delta_i}^0 f, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2}.$$

The numbers p , q and r are constants that we shall define later and select to ensure that $Q_{l,i}(f)$ is a norm equivalent to its standard Sobolev counterpart. Before getting a full Sobolev estimate, we first study the term $Q_{l,i}$. The crucial idea being that the terms $\|f\|_{L_{x,v}^2}^2$ arising from S_2 will be controlled by the fluctuation of the characteristics (*i.e.* the gain due to S_3), rather than by the negative feedback of the Boltzmann operator, whilst the source term S_6 will find itself controlled by the latter. In what follows our Propositions are divided into two different cases: $\varepsilon = 1$ and $\varepsilon < 1$. The

difference here is that, in the case $\varepsilon = 1$, we must keep the negative feedback brought by the fluctuation, whereas for small ε we can discard it (see Remark 4.2).

Proposition 4.1. *Let s be in \mathbb{N} and l be a multi-index such that $|l| = s$. There exist $0 < \varepsilon_s \leq 1$ for which we have the following results.*

Case $0 < \varepsilon \leq \varepsilon_s$. *There exists p, q, r and $C_0 > 0$, such that*

$$Q_{l,i}(\cdot) \sim \|\partial_{l+\delta_i}^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \|\partial_l^{\delta_i} \cdot\|_{L_{x,v}^2}^2$$

and if f is a solution to the perturbative equation (4.1), then

$$\begin{aligned} \forall t \in [0, T_0), \quad \frac{d}{dt} Q_{l,i}(f) &\leq -2 \left(\frac{1}{\varepsilon^2} \|\pi_L^\perp (\partial_{l+\delta_i}^0 f)\|_{L_\gamma^2}^2 + \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 + \|\partial_l^{\delta_i} f\|_{L_\gamma^2}^2 \right) \\ &\quad + \varepsilon^{1-e} C_0 \sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_\gamma^2}^2 + C_0 \varepsilon^2 \sum_{1 \leq j, k \leq d} \|\partial_{l-\delta_j}^{\delta_i+\delta_k} f\|_{L_\gamma^2}^2 \\ &\quad + C_0 \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

Case $\varepsilon = 1$. *There exists p, q, r and $C_0 > 0$ such that*

$$Q_{l,i}(\cdot) \sim \|\partial_{l+\delta_i}^0 \cdot\|_{L_{x,v}^2}^2 + \|\partial_l^{\delta_i} \cdot\|_{L_{x,v}^2}^2$$

and if f is a solution to the perturbative equation (4.1), then for any $\Lambda > 1$,

$$\begin{aligned} \forall t \in [0, T_0), \quad \frac{d}{dt} Q_{l,i}(f) &\leq -2\lambda_{s,\Lambda} \left(\|\pi_L^\perp (\partial_{l+\delta_i}^0 f)\|_{L_\gamma^2}^2 + \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 + \|\partial_l^{\delta_i} f\|_{L_\gamma^2}^2 \right) \\ &\quad - \Lambda \|\partial_{l+\delta_i}^0 f\|_{L_2^2}^2 - \Lambda \|\partial_l^{\delta_i} f\|_{L_2^2}^2 \\ &\quad + C_0 \sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_2^2}^2 + C_0 \sum_{1 \leq j, k \leq d} \|\partial_{l-\delta_j}^{\delta_i+\delta_k} f\|_{L_2^2}^2 \\ &\quad + C_{0,\Lambda} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

All the constants involved depend explicitly on T_0, s and E .

Proof of Proposition 4.1. We recall that f is solution to

$$\partial_t f = \sum_{j=1}^6 S_j(t, x, v)$$

which directly implies that

$$\frac{d}{dt} Q_{l,i}(f) = \sum_{j=1}^6 p \langle \partial_{l+\delta_i}^0 S_j, \partial_{l+\delta_i}^0 f \rangle_{L_{x,v}^2} + \varepsilon^2 q \langle \partial_l^{\delta_i} S_j, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2} + \varepsilon r \langle \partial_{l+\delta_i}^0 S_j, \partial_l^{\delta_i} f \rangle_{L_{x,v}^2}.$$

We therefore directly apply Propositions 3.3 and 3.4 to control S_4 and S_5 , whereas we use Lemmas 3.5, 3.6, 3.7 and 3.8 for the other terms. It yields

$$\begin{aligned}
(4.4) \quad \frac{1}{2} \frac{d}{dt} Q_{l,i}(f) &\leq \frac{C_{\eta_0} r + p\eta_0 - p}{\varepsilon^2} \lambda_{0,\Lambda} \|\pi_L^\perp (\partial_{l+\delta_i}^0 f)\|_{L_\gamma^2}^2 \\
&+ \left[\frac{q\varepsilon^2}{\eta_1} + \eta_4 p + \varepsilon^{1-e} \frac{1_{\varepsilon < 1}}{4} p + \varepsilon \frac{1_{\varepsilon < 1}}{4} r - r \right] \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 \\
&+ \left[q(\eta_1 + \eta'_0 + \eta_4 + \varepsilon^{3-e} \frac{1_{\varepsilon < 1}}{4}) + r(\eta_0 + \eta_4) \lambda_{0,\Lambda} + \varepsilon^{2-e} \frac{1_{\varepsilon < 1}}{4} r - q\lambda_{s,\Lambda} \right] \|\partial_l^{\delta_i} f\|_{L_\gamma^2}^2 \\
&+ \varepsilon^{1+e} \Lambda \left[C_{\eta_3} r + \eta_2 p - \frac{p}{2} \right] \|\partial_{l+\delta_i}^0 f\|_{L_2^2}^2 \\
&+ \varepsilon^{3+e} \Lambda \left[\eta_2 q + (\eta_2 + \eta_3) r - \frac{q}{2} \right] \|\partial_l^{\delta_i} f\|_{L_2^2}^2 \\
&+ \varepsilon^{1-e} \left[pC_{\eta_2} + d \frac{C_{\eta_2}}{\Lambda} r \right] \sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_{x,v}^2}^2 \\
&+ [\varepsilon^2 q C_{\eta_2}] \sum_{1 \leq j, k \leq d} \|\partial_{l-\delta_j}^{\delta_i+\delta_k} f\|_{L_{x,v}^2}^2 \\
&+ C_{p,q,r,\Lambda,\eta} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right].
\end{aligned}$$

We firstly emphasize that the twice indexed sums are a cruder estimate than the one we actually derived in the Lemmas: we added some terms that were formerly absent, but we think it provides a better reading. We secondly emphasize that the estimate on the bilinear term S_5 in Proposition 3.4 gives a control of the form $\|f\|_{H_{x,v}^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2$ which translates into a control of the form $\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2$ when multiplying by powers of ε . We lastly used $\|\pi_L(\partial_l^0 f)\|_{L_{x,v}^2} \leq \|\partial_l^0 f\|_{L_{x,v}^2}$ when applying Lemma 3.8.

In what follows, we recall that $0 < \varepsilon \leq 1$. We shall now choose the constants carefully, which is why we indexed all generic constant by their dependencies in order to avoid any loop.

Remark 4.2. *The choices are different for $\varepsilon = 1$ or any $\varepsilon < 1$ because the control of the specific term $\mathcal{T} := \sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_{x,v}^2}^2$ will be achieved in two different ways.*

For $\varepsilon = 1$, the negative feedback of the fluctuation $\|\partial_l^{\delta_i} f\|_{L_2^2}$ can control these terms, taking Λ sufficiently large. Such an approach does not work for general values of ε (because we control v -derivatives with a degenerate weight ε^2). In the general case therefore, the term \mathcal{T} will be absorbed by the negative feedback that the linear Boltzmann operator provides, and this latter approach requires a sufficiently small value of ε .

Note that this distinction is quite artificial, and is due to our choice of Sobolev norm with coefficient $\varepsilon^2 \|\partial_l^{\delta_i} \cdot\|_{L_{x,v}^2}^2$. Working with this weighted norm facilitates various computations and estimates, but [12] showed that a finer norm, which is not degenerating when ε tends to 0, can catch the hypo-coercivity of the Boltzmann linear operator. With more technicalities, we think we could use the latter norm to

avoid this splitting into two regimes, and always control the problematic term with the negative feedback generated by the fluctuations for any ε .

We start with the case $\varepsilon < 1$ and fix the following quantities:

- (1) $\Lambda = 1$;
- (2) q sufficiently large such that $-q\lambda_{s,1} \leq -5$;
- (3) η_1 small enough such that $q\eta_1 \leq 1$;
- (4) r sufficiently large such that $\frac{q}{\eta_1} - \frac{3r}{4} \leq -4$;
- (5) $\eta_0 = \eta'_0$ small enough such that
 - $2\eta_0 - \lambda_{s,1} \leq -\frac{\lambda_{s,1}}{2}$,
 - $2\eta_0 q + 2\eta_0 \lambda_{0,1} r \leq -1$;
- (6) η_2 small enough such that $\eta_2 \leq 1/8$ and $\eta_2 r \leq q/8$;
- (7) η_3 small enough such that $\eta_3 r \leq q/8$;
- (8) p sufficiently large such that
 - $C_{1,\eta_0} r + p\eta_0 - p \leq -1$,
 - $C_{\eta_3} r + \eta_2 p - \frac{p}{2} \leq 0$,
 - $r^2 \leq pq$ and $q \leq p$ so that $Q_{l,i}(\cdot) \sim \|\partial_{l+\delta_i}^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \|\partial_l^{\delta_i} \cdot\|_{L_{x,v}^2}^2$
- (9) $\eta_4 \leq \eta_0$ small enough such that $\eta_4 p \leq 1$ - note that point this allows to use point (5) above;
- (10) At last we need ε small enough such that $\varepsilon^{1-e} \frac{p}{4} \leq 1$, $\varepsilon^{3-e} \frac{q}{4} \leq 1$, $\varepsilon^{2-e} \frac{r}{4} \leq 1$.

Such a choice yields exactly the expected result for $0 < \varepsilon < 1$.

Remark 4.3. *One clearly sees here that, if e were too large, $e \geq 1$, then one could not make $\varepsilon^{1-e} \frac{p}{4}$ small as desired. In other terms, in that case of large coefficient e , the amplitude of the evolution of the v characteristics would not be compensated by the gain due, via hypocoercive estimates, to the free transport in x .*

Now let us deal with the particular case $\varepsilon = 1$. The crucial step will be to fix the constants p , r and η_2 to ensure that the term $\sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_{x,v}^2}^2$ has a multiplicative constant independent of Λ . Then Λ should precisely be chosen large enough to absorb these contributions. In order to achieve this goal we transform (4.4) by estimating

$$q\eta_1 \|\partial_l^{\delta_i} f\|_{L_x^2}^2 \leq Cq\eta_1 \|\partial_l^{\delta_i} f\|_{L_x^2}^2 \quad \text{and} \quad 2r\eta_0 \lambda_{0,\Lambda} \|\partial_l^{\delta_i} f\|_{L_x^2}^2 \leq Cr\eta_0 \|\partial_l^{\delta_i} f\|_{L_x^2}^2,$$

since $\lambda_{0,\Lambda} \leq \lambda_{0,0}$ from the proof of Proposition 3.3. We infer, for $\Lambda \geq 1$, the estimate

$$\begin{aligned} \frac{d}{dt} Q_{l,i}(f) &\leq [C_{\eta_0} r + p\eta_0 - p] \lambda_{0,\Lambda} \|\pi_L^\perp (\partial_{l+\delta_i}^0 f)\|_{L_x^2}^2 + \left[\frac{q}{\eta_1} + \eta_4 p - r \right] \|\partial_{l+\delta_i}^0 f\|_{L_{x,v}^2}^2 \\ &\quad + [2q\eta'_0 - q\lambda_{s,\Lambda}] \|\partial_l^{\delta_i} f\|_{L_x^2}^2 \\ &\quad + \Lambda \left[C_{\eta_3} r + \eta_2 p - \frac{p}{2} \right] \|\partial_{l+\delta_i}^0 f\|_{L_x^2}^2 \\ &\quad + \Lambda \left[(C\eta_1 + \eta_2 + \eta_4)q + (C\eta_0 + \eta_2 + \eta_3 + \eta_4)r - \frac{q}{2} \right] \|\partial_l^{\delta_i} f\|_{L_x^2}^2 \\ &\quad + [pC_{\eta_2} + dC_{\eta_2} r] \sum_{1 \leq j, k \leq d} \|\partial_{l+\delta_i-\delta_j}^{\delta_k} f\|_{L_{x,v}^2}^2 + qC_{\eta_2} \sum_{1 \leq j, k \leq d} \|\partial_{l-\delta_j}^{\delta_i+\delta_k} f\|_{L_{x,v}^2}^2 \\ &\quad + C_{p,q,r,\Lambda,\eta} \left[\|f\|_{\mathcal{H}_x^{s+1}}^2 \|f\|_{H_x^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

We can now choose our different constants in the following way:

- (1) $q = 8$;
- (2) η_1 small enough such that $Cq\eta_1 \leq 1$;
- (3) r sufficiently large such that $\frac{q}{\eta_1} - r \leq -3$;
- (4) η_0, η_2 and η_3 small enough such that
 - $\eta_0 \leq \frac{1}{4}$,
 - $C\eta_1 + 2\eta_2 \leq 1$,
 - $(C\eta_0 + 2\eta_2 + \eta_3)r \leq 1$,
 - $\eta_2 \leq \frac{1}{4}$
- (5) $\eta'_0 = \eta'_0(\Lambda)$ small enough such that $3\eta'_0 \leq -\lambda_{s,\Lambda}$
- (6) p sufficiently large such that
 - $C_{\eta_0}r + 2p\eta_0 - p \leq -1$,
 - $C_{\eta_3}r + \eta_2p - \frac{p}{2} \leq -1$,
 - $r^2 \leq pq$ and $q \leq p$ so that $Q_{l,i}(\cdot) \sim \|\partial_{l+\delta_i}^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \|\partial_l^{\delta_i} \cdot\|_{L_{x,v}^2}^2$.
- (7) At last $\eta_4 \leq \eta_2$ small enough such that $\eta_4p \leq 1$ - note that point this allows to use point (5) above;

These choices yield the expected result, emphasizing that we manage to choose p , q , r and C_{η_2} independently of Λ . \square

4.2. Estimates for velocity derivatives. We now turn to the terms that include velocity derivatives for which the linear Boltzmann operator provides a full negative feedback (this is the second term in (3.14)).

Proposition 4.4. *Let s be in \mathbb{N}^* and l and j be multi-indexes with $|l| + |j| = s + 1$ with $|j| \geq 2$. There exist $0 < \varepsilon_s \leq 1$*

Case $0 < \varepsilon \leq \varepsilon_s$. *There exists $\lambda_{s,1}, C_1 > 0$ such that if f is a solution to the perturbative equation (4.1) then*

$$\begin{aligned} \forall t \in [0, T_0), \quad \frac{d}{dt} \|\partial_l^j f\|_{L_{x,v}^2}^2 &\leq -\frac{\lambda_{s,1}}{\varepsilon^2} \|\partial_l^j f\|_{L_\gamma^2}^2 + C_1 \sum_{1 \leq i, k \leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_\gamma^2}^2 \\ &\quad + C_1 \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 \\ &\quad + \frac{C_1}{\varepsilon^2} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

Case $\varepsilon = 1$. *There exists $C_1 > 0$ such that if f is a solution to the perturbative equation (4.1) then for any $\Lambda \geq 1$,*

$$\begin{aligned} \forall t \in [0, T_0), \quad \frac{d}{dt} \|\partial_l^j f\|_{L_{x,v}^2}^2 &\leq -\lambda_{s,\Lambda} \|\partial_l^j f\|_{L_\gamma^2}^2 + C_1 \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 \\ &\quad - \Lambda \|\partial_l^j f\|_{L_2^2}^2 + C_1 \sum_{1 \leq i, k \leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_2^2}^2 \\ &\quad + C_{1,\Lambda} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

All the constants depend explicitly on s , E and T_0 .

Proof of Proposition 4.4. As for $Q_{l,i}$, we recall that f is solution to

$$\partial_t f = \sum_{j=1}^6 S_j(t, x, v)$$

so we directly apply Propositions 3.3 and 3.4 to control S_4 and S_5 , whereas we use Lemmas 3.5, 3.6, 3.7 and 3.8 for the other terms. We have

$$(4.5) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|\partial_l^j f\|_{L_{x,v}^2}^2 &\leq \frac{\eta_0 + \eta'_0 + \eta_1 + \eta_4 + \varepsilon^{3-e} \frac{1-\varepsilon < 1}{4} - \lambda_{s,\Lambda}}{\varepsilon^2} \|\partial_l^j f\|_{L_{x,v}^2}^2 \\ &\quad + [\varepsilon^{1+e} \eta_2 \Lambda - \varepsilon^{1+e} \Lambda] \|\partial_l^j f\|_{L_{x,v}^2}^2 \\ &\quad + \frac{1}{\eta_1} \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 + \varepsilon^{1-e} C_{\eta_2} \sum_{1 \leq i, k \leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_{x,v}^2}^2 \\ &\quad + \frac{C_{1,\Lambda}}{\varepsilon^2} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

Here again our choice of constant will differ if $\varepsilon = 1$. First let us consider the general case $0 < \varepsilon < 1$. We take

- (1) $\Lambda = 1$;
- (2) $\eta_0 = \eta'_0 = \eta_1 = \eta_4$ small enough such that
 - $\eta_0 + \eta'_0 + \eta_1 - \lambda_{s,1} \leq -\lambda_{s,1}/2$,
 - $\eta_2 \leq 1/2$;
- (3) ε sufficiently small such that $\frac{\varepsilon^{3-e}}{4} \leq \frac{\lambda_{s,1}}{4}$.

and these choices lead to the expected estimate.

The particular case $\varepsilon = 1$ is dealt with differently and we modify (4.5) by estimating

$$\eta_1 \|\partial_l^j f\|_{L_{x,v}^2}^2 \leq \eta_1 \|\partial_l^j f\|_{L_{x,v}^2}^2,$$

to obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\partial_l^j f\|_{L_{x,v}^2}^2 &\leq [\eta_0 + \eta'_0 - \lambda_{s,\Lambda}] \|\partial_l^j f\|_{L_{x,v}^2}^2 + [2\eta_1 + \eta_2 \Lambda - \Lambda] \|\partial_l^j f\|_{L_{x,v}^2}^2 \\ &\quad + \frac{1}{\eta_1} \sum_{1 \leq k \leq d} \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 + C_{\eta_2} \sum_{1 \leq i, k \leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_{x,v}^2}^2 \\ &\quad + C_{1,\Lambda,\eta} \left[\|f\|_{\mathcal{H}_\varepsilon^{s+1}}^2 \|f\|_{H_\gamma^{s+1}}^2 + \|f\|_{H_{x,v}^s}^2 + 1 \right]. \end{aligned}$$

Taking $\Lambda \geq 1$ we can choose the constants η_1 and η_2 independently of Λ in the same manner as for the general case and obtain the expected result. \square

4.3. Estimates for the full Sobolev norm. We now gather the previous estimates to establish a full control over the twisted Sobolev norm. We start with the Sobolev norm corresponding to a fixed total number of derivatives, and then, layer by layer, deal with the norm that estimates all the derivatives.

Proposition 4.5. *Let $s \geq s_0$ (where s_0 is given in Proposition 3.4) and $1 \leq s' \leq s$. Then there exists $\Lambda > 0$, $\lambda_{s'} > 0$, $C_{s'} > 0$ and $0 < \varepsilon_s s' \leq 1$ such that for $(\varepsilon = 1$ or*

$0 < \varepsilon \leq \varepsilon_{s'}$) there exists a functional $F_{s'}$ such that

$$F_{s'} \sim \sum_{|l|=s'} \|\partial_l^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|l|+|j|=s' \\ |j| \geq 1}} \|\partial_l^j \cdot\|_{L_{x,v}^2}^2$$

and if f is a solution to the perturbative equation (4.1) then for any $\Lambda > 0$, for all $t \in [0, T_0)$,

$$\begin{aligned} \frac{d}{dt} F_{s'}(f) \leq & -\lambda \left(\frac{1}{\varepsilon^2} \sum_{|l|=s'} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2 + \sum_{\substack{|l|+|j|=s' \\ |j| \geq 1}} \|\partial_l^j f\|_{L_\gamma^2}^2 + \sum_{|l|=s'} \|\partial_l^0 f\|_{L_{x,v}^2}^2 \right) \\ & + C_{s'} \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + \|f\|_{H_{x,v}^{s'-1}}^2 + 1 \right]. \end{aligned}$$

All the constants depend on s , E and T_0 .

Proof of Proposition 4.5. We present the proof in two different cases: ε sufficiently small first and then $\varepsilon = 1$. The technicalities are identical but the absorption mechanisms are different as explained in Remark 4.2. Consider some constant $B_{j,l} > 0$ to be fixed later, and define the functional

$$F_{s'}(f) = \sum_{|l|=s'} \sum_{i=1}^d Q_{l,i}(f) + \varepsilon^2 \sum_{\substack{|l|+|j|=s' \\ |j| \geq 2}} B_{j,l} \|\partial_l^j f\|_{L_{x,v}^2}^2.$$

By Proposition 4.1, we know that, for any $B > 0$:

$$F_s \sim \sum_{|l|=s} \|\partial_l^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|l|+|j|=s \\ |j| \geq 1}} \|\partial_l^j \cdot\|_{L_{x,v}^2}^2.$$

Case ε sufficiently small. Using directly Proposition 4.1 and Proposition 4.4 we see that, for $B_{j,l} = B$ sufficiently large, we have a constant $C_B > 0$ independent of ε such that

$$\begin{aligned} \frac{d}{dt} F_{s'}(f) \leq & - \left(\frac{1}{\varepsilon^2} \sum_{|l|=s'} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2 + \sum_{\substack{|l|+|j|=s' \\ |j| \geq 1}} \|\partial_l^j f\|_{L_\gamma^2}^2 + \sum_{|l|=s'} \|\partial_l^0 f\|_{L_{x,v}^2}^2 \right) \\ & + \varepsilon C_B \sum_{|l|+|j|=s'} \sum_{1 \leq i, k \leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_\gamma^2}^2 + \left\| \partial_{l+\delta_j}^{j-\delta_k} f \right\|_{L_{x,v}^2}^2 \\ & + C_{1,B} \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + \|f\|_{H_{x,v}^{s'-1}}^2 + 1 \right]. \end{aligned}$$

Therefore taking ε sufficiently small allows to absorb the second line with the negative first term. This is the expected result.

Case $\varepsilon = 1$. The proof is exactly the same. Gathering Proposition 4.1 and Proposition 4.4 with $B_{j,l} < 1$, we infer that there exists $C > 0$ independent of Λ and

$B_{j,l}$ such that

$$\begin{aligned}
\frac{d}{dt} F_{s'}(f) &\leq -\lambda_{s',\Lambda} \left(\sum_{|l|=s'} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2 + \sum_{|l|=s'} \|\partial_l^0 f\|_{L_{x,v}^2}^2 \right. \\
&\quad \left. + \sum_{\substack{|l|+|j|=s' \\ |j|\geq 2}} B_{j,l} \|\partial_l^j f\|_{L_\gamma^2}^2 + \sum_{|l|=s'-1} \|\partial_l^{\delta_i} f\|_{L_\gamma^2}^2 \right) \\
&\quad + C_1 \sum_{\substack{|j|+|l|=s' \\ |j|\geq 2}} B_{j,l} \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_\gamma^2}^2 \\
&\quad - \Lambda \sum_{|j|+|l|=s'} \|\partial_l^j f\|_2^2 \\
&\quad + C \left(\sum_{|l|+|j|=s'} \sum_{1\leq i,k\leq d} \left\| \partial_{l-\delta_i}^{j+\delta_k} f \right\|_{L_2^2}^2 + \left\| \partial_{l+\delta_j}^{j-\delta_k} f \right\|_{L_2^2}^2 \right) \\
&\quad + C \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + \|f\|_{H_{x,v}^{s'-1}}^2 + 1 \right].
\end{aligned}$$

We can then choose $B_{j,l} = B_{j,l}(\Lambda)$ sufficiently small hierarchically to absorb

$$C_1 \sum_{\substack{|j|+|l|=s' \\ |j|\geq 2}} B_{j,l} \sum_{k=1}^d \left\| \partial_{l+\delta_k}^{j-\delta_k} f \right\|_{L_\gamma^2}^2$$

inside the full negative terms on the first line of the estimate. As C is independent of Λ (due to the fact that we could choose the $B_{j,l} < 1$) we can at last fix Λ sufficiently large such that the L_2^2 -norms give a negative contribution. This concludes the proof. \square

We finally have all the tools to perform a full $H_{x,v}^s$ estimate. The following proposition indeed shows that our choice of perturbative regime compensate the modification of the characteristics due to the presence of the external force even on the fluid part of the solution $\pi_L(f)$.

Proposition 4.6. *Let $s \geq s_0$ (where s_0 is given in Proposition 3.4). Then there exists $\Lambda > 0$, $C > 0$ and $0 < \varepsilon_s \leq 1$ such that, for $\varepsilon = 1$ or $0 < \varepsilon \leq \varepsilon_s$, there exists a functional*

$$(4.6) \quad \|\cdot\|_{\mathcal{H}_{x,v}^s}^2 \sim \sum_{|l|\leq s} \|\partial_l^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|l|+|j|\leq s \\ |j|\geq 1}} \|\partial_l^j \cdot\|_{L_{x,v}^2}^2$$

such that, if f is a solution to the perturbative equation (4.1) with initial data f_{in} satisfying

$$\left| \int_{\mathbb{T}^d \times \mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} f_{\text{in}}(x, v) \sqrt{M}|_{t=0} dx dv \right| \leq C_{\text{in}} \varepsilon^e,$$

then

$$(4.7) \quad \forall t \in [0, T_0), \quad \frac{d}{dt} \|f\|_{\mathcal{H}_{x,v}^s}^2 \leq -(\lambda - C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2) \|f\|_{H_\gamma^s}^2 + C_s.$$

All the constants depend on s , E and T_0 , but are independent of ε .

Proof of Proposition 4.6. As we shall see, the proof follows directly from Proposition 4.5 apart from $s = 1$ which has to be treated more carefully due to the lack of full negative return on the $L_{x,v}^2$ -norm. Fixing s in \mathbb{N}^* we use Proposition 4.5 to construct

$$\begin{aligned} & \frac{d}{dt} \sum_{s'=2}^s \alpha_{s'} F_s(f) \\ & \leq -\lambda \sum_{s'=1}^s \left(\frac{1}{\varepsilon^2} \sum_{|l|=s'} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2 + \sum_{\substack{|l|+|j|=s' \\ |j| \geq 1}} \|\partial_l^j f\|_{L_\gamma^2}^2 + \sum_{|l|=s} \|\partial_l^0 f\|_{L_{x,v}^2}^2 \right) \\ & \quad + C \sum_{1 \leq s' \leq s} \alpha_{s'} \|f\|_{H_{x,v}^{s'-1}}^2 + C_s \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + 1 \right], \end{aligned}$$

where $\alpha_{s'}$ are constants that we choose sufficiently small hierarchically (from s to 1) in order to absorb the first positive term at rank s' on the right-hand side by the negative feedback at rank $s' - 1$. We infer for all t in $[0, T_0)$,

$$(4.8) \quad \begin{aligned} & \frac{d}{dt} \sum_{s'=2}^s \alpha_{s'} F_s(f) \\ & \leq -\lambda \sum_{s'=1}^s \left(\frac{1}{\varepsilon^2} \sum_{|l|=s'} \|\pi_L^\perp(\partial_l^0 f)\|_{L_\gamma^2}^2 + \sum_{\substack{|l|+|j|=s' \\ |j| \geq 1}} \|\partial_l^j f\|_{L_\gamma^2}^2 + \sum_{|l|=s} \|\partial_l^0 f\|_{L_{x,v}^2}^2 \right) \\ & \quad + \|f\|_{H_{x,v}^1}^2 + C_s \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + 1 \right] \end{aligned}$$

and we have from Proposition 4.5:

$$\sum_{1 \leq s' \leq s} \alpha_{s'} F_s(f) \sim \sum_{1 \leq s' \leq s} \left(\sum_{|l| \leq s'} \|\partial_l^0 \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \sum_{\substack{|l|+|j| \leq s' \\ |j| \geq 1}} \|\partial_l^j \cdot\|_{L_{x,v}^2}^2 \right)$$

so that

$$(4.9) \quad \frac{d}{dt} \sum_{s'=2}^s \alpha_{s'} F_s(f) \leq - \left(\frac{\lambda}{2} - C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2 \right) \|f\|_{H_\gamma^s}^2 + C_s \|f\|_{H_{x,v}^1}^2 + C_s.$$

Control of fluid part by spatial derivatives. A key property of our proof will be to recover the full coercivity on the $L_{x,v}^2$ -norm by controlling $\|\pi_L(f)\|_{L_{x,v}^2}^2$ by $\|\nabla_x f\|_{L_{x,v}^2}^2$ which is fully coercive. First, as the eigenfunction of L are polynomials times Maxwellian, see (3.5) and (3.6), we easily obtain (we refer to [12, equation (3.3)] for a direct proof) that there exists $c_\pi, C_\pi > 0$ such that

$$(4.10) \quad \forall 0 \leq |j| + |l| \leq s, \quad c_\pi \|\partial_l^j \pi_L^\perp(f)\|_{L_{x,v}^2}^2 \leq \|\pi_L(\partial_l^0 f)\|_{L_\gamma^2}^2 \leq C_\pi \|\pi_L(\partial_l^0 f)\|_{L_{x,v}^2}^2.$$

So the only part left to estimate is $\|\pi_L(f)\|_{L_{x,v}^2}^2$. Coming back to (3.6), we have, for a constant C depending on a, A, e and T_0 ,

$$\|\pi_L(f)\|_{L_{x,v}^2}^2 \leq C \sum_{i=0}^{d+1} \int_{\mathbb{T}^d} |\langle f, \bar{\phi}_i \sqrt{M} \rangle|^2 dx.$$

We use the Poincaré-Wirtinger inequality to obtain, for possibly a different constant C ,

$$\|\pi_L(f)\|_{L_{x,v}^2}^2 \leq C \sum_{i=0}^{d+1} \left[\int_{\mathbb{T}^d} |\nabla_x \langle f, \bar{\phi}_i \sqrt{M} \rangle|^2 dx + \left| \int_{\mathbb{T}^d} \langle f, \bar{\phi}_i \sqrt{M} \rangle dx \right|^2 \right],$$

which gives in turn

$$(4.11) \quad \|\pi_L(f)\|_{L_{x,v}^2}^2 \leq C \|\nabla_x(\pi_L(f))\|_{L_{x,v}^2}^2 + C \int_{\mathbb{R}^d} \left| \int_{\mathbb{T}^d} \pi_L(f) dx \right|^2 dv.$$

For the classical Boltzmann equation the preservation of mass, momentum and energy gives the cancellation $\int_{\mathbb{T}^d} \pi_L(f) dx = 0$, which is no longer satisfied here however.

We come back to the original equation (1.5) on $F = M + \varepsilon \sqrt{M} f$:

$$\partial_t F + \frac{1}{\varepsilon} v \cdot \nabla_x F + \varepsilon \vec{E}_t(x) \cdot \nabla_v F = \frac{1}{\varepsilon} Q(F, F).$$

We multiply this equation by 1, v and $|v|^2$ and we integrate over $\mathbb{T}^d \times \mathbb{R}^d$. This yields

$$(4.12) \quad \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} F(t, x_*, v_*) dx_* dv_* = 0,$$

$$(4.13) \quad \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} v_* F(t, x_*, v_*) dx_* dv_* = \varepsilon \int_{\mathbb{T}^d \times \mathbb{R}^d} \vec{E}_t(x_*) F(t, x_*, v_*) dx_* dv_*,$$

$$(4.14) \quad \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} |v_*|^2 F(t, x_*, v_*) dx_* dv_* = 2\varepsilon \int_{\mathbb{T}^d \times \mathbb{R}^d} \vec{E}_t(x_*) \cdot v_* F(t, x_*, v_*) dx_* dv_*.$$

Note that we used that $Q(f, g)$ is orthogonal to $\text{Ker}(L)$ in $L_{x,v}^2$. We insert the relation $F = M + \varepsilon \sqrt{M} f$ in the previous estimates. We compute and bound for $\delta = 0$ or $\delta = 2$,

$$(4.15) \quad \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} |v_*|^\delta (M(0, x_*, v_*) - M(t, x_*, v_*)) \right| = \frac{e^{-\varepsilon^{1+e} \frac{A}{1+t}}}{(1 + \varepsilon^{1+e} \frac{a}{1+t})^{\frac{2d+\delta}{4}}} - \frac{e^{-\varepsilon^{1+e} A}}{(1 + \varepsilon^{1+e} a)^{\frac{2d+\delta}{4}}} \leq \varepsilon^{1+e} C_{a,A}.$$

From (4.12), (4.15) and the smallness hypothesis (2.2), we deduce that

$$(4.16) \quad \sup_{t \in [0, T_0]} \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} f_*(t) \sqrt{M}_*(t) dx_* dv_* \right| \leq C \varepsilon^e.$$

We use the cancellation condition in (2.1) and (2.2) to get

$$(4.17) \quad \sup_{t \in [0, T_0]} \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} v_* f_*(t) \sqrt{M}_*(t) dx_* dv_* \right| \leq C \varepsilon^e + \varepsilon T_0 C_E \sup_{t \in [0, T_0]} \int_{\mathbb{T}^d} \left| \int_{\mathbb{R}^d} v_* f_*(t) \sqrt{M}_*(t) dv_* \right| dx.$$

We use the bounds

$$\begin{aligned} \varepsilon T_0 C_E \sup_{t \in [0, T_0]} \int_{\mathbb{T}^d} \left| \int_{\mathbb{R}^d} v_* f_*(t) \sqrt{M_*(t)} dv_* \right| dx \\ \leq \varepsilon T_0 C_E \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2} \leq C \varepsilon^e + \varepsilon^{2(1-e)} \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2 \end{aligned}$$

to obtain

$$(4.18) \quad \sup_{t \in [0, T_0]} \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} v_* f_*(t) \sqrt{M_*(t)} dx_* dv_* \right| \leq C \varepsilon^e + \varepsilon^{2(1-e)} \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2.$$

A similar procedure gives

$$(4.19) \quad \sup_{t \in [0, T_0]} \left| \int_{\mathbb{T}^d \times \mathbb{R}^d} |v_*|^2 f_*(t) \sqrt{M_*(t)} dx_* dv_* \right| \leq C \varepsilon^e + \varepsilon^{2(1-e)} \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2.$$

The identities (4.16)-(4.18)-(4.19) above imply

$$(4.20) \quad \sup_{t \in [0, T_0]} \int_{\mathbb{R}^d} \left| \int_{\mathbb{T}^d} \pi_L(f(t)) dx \right|^2 dv \leq C \varepsilon^{2e} + 2\varepsilon^{2(1-e)} \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2.$$

We combine (4.20) with (4.11) to obtain

$$\sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2 \leq C \|\nabla_x(\pi_L(f(t)))\|_{L_{x,v}^2}^2 + C \varepsilon^{2e} + 2\varepsilon^{2(1-e)} \sup_{t \in [0, T_0]} \|\pi_L(f(t))\|_{L_{x,v}^2}^2.$$

For ε small enough, this gives

$$(4.21) \quad \|\pi_L(f)\|_{L_{x,v}^2}^2 \leq C_\pi C \|\nabla_x(\pi_L(f(t)))\|_{L_{x,v}^2}^2 + C_\pi C \varepsilon^{2e} \leq C'_\pi \left(\|\nabla_x(f(t))\|_{L_{x,v}^2}^2 + 1 \right),$$

where C_π was defined in (4.10).

Evolution of the H^1 -norm. Let us now look at the evolution of the full $H_{x,v}^1$ -norm. We take p , q and r and define

$$Q_1(f) = p \|\nabla_x f\|_{L_{x,v}^2}^2 + q \varepsilon^2 \|\nabla_v f\|_{L_{x,v}^2}^2 + \varepsilon r \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2}.$$

Using the estimates given in Section 3 exactly as for (4.4) but keeping explicit the dependencies $C_\Lambda = \frac{C}{\Lambda}$ we find

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} Q_1(f) &\leq \frac{C_{\eta_0} r + p\eta_0 - p}{\varepsilon^2} \lambda_{0,\Lambda} \|\pi_L^\perp(\nabla_x f)\|_{L_\gamma^2}^2 \\
&\quad + \left[\frac{q\varepsilon^2}{\eta_1} + \eta_4 p + \varepsilon^{1-e} \frac{1_{\varepsilon < 1}}{4} p + \varepsilon \frac{1_{\varepsilon < 1}}{4} r - r \right] \|\nabla_x f\|_{L_{x,v}^2}^2 \\
&\quad + \left[q(\eta_1 + \eta'_0 + \eta_4 + \varepsilon^{3-e} \frac{1_{\varepsilon < 1}}{4}) + r(\eta_0 + \eta_4) \lambda_{0,\Lambda} + \varepsilon^{2-e} \frac{1_{\varepsilon < 1}}{4} r - q\lambda_{s,\Lambda} \right] \|\nabla_v f\|_{L_\gamma^2}^2 \\
&\quad + \varepsilon^{1+e} \Lambda \left[C_{\eta_3} r + \eta_2 p - \frac{p}{2} \right] \|\nabla_x f\|_{L_2^2}^2 \\
&\quad + \varepsilon^{3+e} \Lambda \left[\eta_2 q + (\eta_2 + \eta_3) r - \frac{q}{2} \right] \|\nabla_v f\|_{L_2^2}^2 \\
&\quad + \varepsilon^{1-e} \left[pC_{\eta_2} + d \frac{C_{\eta_2}}{\Lambda} r \right] \|\nabla_v f\|_{L_{x,v}^2}^2 \\
&\quad + \left[\mathbf{q} \frac{C_1 C'_\pi}{\Lambda} + 3\mathbf{p} \varepsilon^{1-e} \frac{C_2 C'_\pi}{\Lambda} + 2\mathbf{r} \varepsilon^{1-e} \frac{C_3 C'_\pi}{\Lambda} \right] \|\nabla_{\mathbf{x}f}\|_{L_{\mathbf{x},v}^2}^2 \\
&\quad + C_{p,q,r,\Lambda,\eta} \left[\|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + \|\pi_L^\perp f\|_{L_{x,v}^2}^2 + 1 \right].
\end{aligned}$$

We solely decomposed the $\|f\|_{L_{x,v}^2}^2$ appearing in (4.4) into $\pi_L(f)$ - which we controlled thanks to (4.21) - and $\pi_L^\perp(f)$. To clarify we emphasized in bold the newly added terms. For $\varepsilon < 1$ we see that we can make exactly the same choices for the constants as in Proposition 4.1 with the following two modifications which have no impact on the proof:

- (8) p sufficiently large such that $C_{1,\eta_0} r + qC_1 + p\eta_0 - p\lambda_{0,1} \leq -2$ - instead of $C_{1,\eta_0} r + p\eta_0 - p \leq -1$
- (10) ε small enough as before plus $3p\varepsilon^{1-e} \frac{C_2 C'_\pi}{\Lambda} + 2r\varepsilon^{1-e} \frac{C_3 C'_\pi}{\Lambda} \leq \lambda_{0,1}$.

In the case $\varepsilon = 1$ it is easier since the choices made in Proposition 4.1 leave $\Lambda \geq 1$ free so we can fix all the constant in the same way - recall that p , q and r are independant of Λ - and then choose Λ sufficiently large such that

$$\mathbf{q} \frac{C_1 C'_\pi}{\Lambda} + 3\mathbf{p} \varepsilon^{1-e} \frac{C_2 C'_\pi}{\Lambda} + 2\mathbf{r} \varepsilon^{1-e} \frac{C_3 C'_\pi}{\Lambda} \leq -\frac{1}{2} \left[\frac{q\varepsilon^2}{\eta_1} + \eta_4 p + \varepsilon^{1-e} \frac{1_{\varepsilon < 1}}{4} p + \varepsilon \frac{1_{\varepsilon < 1}}{4} r - r \right].$$

In all cases we can find p , q and r such that $Q_1 \sim \|\nabla_x \cdot\|_{L_{x,v}^2}^2 + \varepsilon^2 \|\nabla_v \cdot\|_{L_{x,v}^2}^2$ and

$$\begin{aligned}
(4.22) \quad \frac{1}{2} \frac{d}{dt} Q_1(f) &\leq -\lambda \left[\frac{1}{\varepsilon^2} \|\pi_L^\perp(\nabla_x f)\|_{L_\gamma^2}^2 + \|\nabla_v f\|_{L_\gamma^2}^2 + \|\nabla_x f\|_{L_{x,v}^2}^2 \right] \\
&\quad + C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{\mathcal{H}_\gamma^s}^2 + C_s \|\pi_L^\perp(f)\|_{L_{x,v}^2}^2 + C_s.
\end{aligned}$$

At last, the evolution of the full $L_{x,v}^2$ -norm is derived as before using the estimates given in Section 3. One bounds

$$\begin{aligned}
\frac{d}{dt} \|f\|_{L_{x,v}^2}^2 &\leq -\frac{\lambda_{0,\Lambda}}{\varepsilon^2} (1 - 2\eta_0) \|\pi_L^\perp(f)\|_{L_\gamma^2}^2 - \varepsilon^{1+e} \Lambda \|f\|_{L_2^2}^2 + \varepsilon^{1-e} \frac{1_{\varepsilon < 1}}{4} \|f\|_{L_{x,v}^2}^2 \\
&\quad + \lambda_{0,\Lambda} \|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + 2C_{\Lambda,\eta_0}.
\end{aligned}$$

Once we fix η_0 sufficiently small and use the orthogonal projection together with the control of $\pi_L(f)$ by the spatial derivatives (4.21) the above implies

$$(4.23) \quad \begin{aligned} \frac{d}{dt} \|f\|_{L_{x,v}^2}^2 &\leq - \left[\frac{\lambda}{\varepsilon^2} - \varepsilon^{1-e} \frac{\mathbf{1}_{\varepsilon < 1}}{4} \right] \|\pi_L^\perp(f)\|_{L_\gamma^2}^2 - \varepsilon^{1+e} \Lambda \|f\|_{L^2}^2 + \varepsilon^{1-e} \frac{\mathbf{1}_{\varepsilon < 1}}{4} \|\nabla_x f\|_{L_{x,v}^2}^2 \\ &\quad + C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + C_s. \end{aligned}$$

We add $\alpha(4.22) + (4.23)$ with α sufficiently small so that $\alpha C_s \|\pi_L^\perp(f)\|_{L_{x,v}^2}^2$ from (4.22) is absorbed by $\frac{\lambda}{2\varepsilon^2} \|\pi_L^\perp(f)\|_{L_{x,v}^2}^2$ from (4.23). Then we can make $\varepsilon = 1$ or ε sufficiently small and obtain

$$(4.24) \quad \frac{1}{2} \frac{d}{dt} \left(\|f\|_{L_{x,v}^2}^2 + \alpha Q_1(f) \right) \leq -\lambda' \|f\|_{H_\gamma^1}^2 + C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2 \|f\|_{H_\gamma^s}^2 + C_s.$$

Conclusion of the proof. Taking a small parameter η , the linear combination $\eta(4.9) + (4.24)$ shows the existence of $\lambda > 0$ and $C_s > 0$ such that

$$\frac{1}{2} \frac{d}{dt} \left(\|f\|_{L_{x,v}^2}^2 + Q_1(f) + \eta \sum_{s'=2}^s F_{s'}(f) \right) \leq -(\lambda - C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2) \|f\|_{H_\gamma^s}^2 + C_s$$

which concludes the proof by defining

$$(4.25) \quad \|f\|_{\mathcal{H}_{x,v}^s}^2 = \|f\|_{L_{x,v}^2}^2 + Q_1(f) + \eta \sum_{s'=2}^s F_{s'}(f).$$

□

Remark 4.7. We see in the derivation of the estimate (4.15), that we need $e > 0$, to ensure that, although not conserved, the global quantities associated to the mass, momentum and energy of the perturbation f are controlled in a suitable way.

5. PROOFS OF THE MAIN RESULTS

At last, we have all the tools to give the proof of the main results.

5.1. Results on the Cauchy theory for the Boltzmann equation.

Proof of Theorem 2.1. The proof of existence follows from a standard iterative scheme:

$$\begin{aligned} \partial_t f_{n+1} + \frac{1}{\varepsilon} v \cdot \nabla_x f_{n+1} + \varepsilon \vec{E}_t(x) \cdot \nabla_v f_{n+1} + \varepsilon \mathcal{E}(t, x, v) f_{n+1} \\ = \frac{1}{\varepsilon^2} L[f_{n+1}] + \frac{1}{\varepsilon} \Gamma[f_n, f_{n+1}] - 2\mathcal{E}(t, x, v) M^{1/2}. \end{aligned}$$

A detailed procedure is given in [12, Section 6.1] for $\vec{E}_t = 0$. This proof is directly applicable here, in combination with our estimates of S_2 and S_3 (terms involving \vec{E}_t). We obtain a uniform bound on a sequence of approximations $(f_n)_{n \in \mathbb{N}}$ in $L_t^\infty H_{x,v}^s \cap L_t^1 H_\gamma^s$, and therefore the strong convergence towards f in less regular Sobolev spaces by Rellich's theorem. The uniqueness of the solution is also standard when $\vec{E}_t = 0$. When \vec{E}_t is non-trivial, uniqueness directly follows from our *a priori* estimate method applied to the difference $f - g$ of two solutions: the linear parts are estimated

in exactly the same way and the bilinear term is controlled when $\|f\|_{H_{x,v}^s}^2 + \|g\|_{H_{x,v}^s}^2$ is small enough, which is why we obtain the uniqueness only in a perturbative regime.

We infer from our *a priori* estimates described in Proposition 4.6 that

$$\forall t \in [0, T_0), \quad \frac{d}{dt} \|f\|_{\mathcal{H}_{x,v}^s}^2 \leq -(\lambda - C_s \|f\|_{\mathcal{H}_\varepsilon^s}^2) \|f\|_{H_\gamma^s}^2 + C_s.$$

Coming back to the definition of the $\mathcal{H}_\varepsilon^s$ -norm given by (4.2) we see that it is uniformly equivalent to the $\mathcal{H}_{x,v}^s$ -norm. Therefore we have

$$\forall t \in [0, T_0), \quad \frac{d}{dt} \|f\|_{\mathcal{H}_{x,v}^s}^2 \leq -(\lambda - C_s \|f\|_{\mathcal{H}_{x,v}^s}^2) \|f\|_{H_\gamma^s}^2 + C_s.$$

As $\|\cdot\|_{\mathcal{H}_{x,v}^s} \leq \|\cdot\|_{H_\gamma^s}$ we directly apply Grönwall lemma which implies that

$$\forall t \in [0, T_0), \quad \|f(t)\|_{\mathcal{H}_{x,v}^s} \leq \max \left\{ \|f_{\text{in}}\|_{\mathcal{H}_{x,v}^s}, C_{T_0, E, s} \right\}$$

as long as $\|f\|_{\mathcal{H}_{x,v}^s}^2$ ($t = 0$) is sufficiently small.

Remark 5.1. *In the specific case where $\|\vec{E}_t\| \leq \frac{C_E}{(1+t)^\alpha}$ with $\alpha > 1$ the second constant C_s behaves like $C_s/(1+t)^\alpha$ (C_E is merely replaced by $C_E/(1+t)^\alpha$ in the computations). Grönwall lemma then gives a polynomial time decay for f for sufficiently small ε .*

□

The corollary follows at once.

Proof of Corollary 2.3. The corollary is a direct consequence of Theorem 2.1 and our definition of fluctuation M . Indeed, direct computations show that

$$\exists C_s > 0, \forall t \geq 0, \quad \|\mu - M\|_{H_{x,v}^s} \leq \varepsilon^{1+e} C_s.$$

Therefore, using the notations of Theorem 2.1

$$\left(\left\| \frac{F_{\text{in}} - \mu}{\varepsilon \sqrt{\mu}} \right\|_{H_{x,v}^s} \leq \frac{\delta_{T_0, E, s}}{2} \right) \Rightarrow \left(\left\| \frac{F_{\text{in}} - M}{\varepsilon \sqrt{M}} \right\|_{H_{x,v}^s} \leq \frac{\delta_{T_0, E, s}}{2} + \varepsilon^e C_s \right)$$

which raises the expected corollary for ε sufficiently small because $e > 0$: we construct a solution provided by Theorem 2.1 and this solution remains close to μ by an additive constant $\varepsilon^e C_s$. □

5.2. Results on the Hydrodynamical limit. We are left with the computation of the limit equations and convergence issues.

Proof of Theorem 2.5. Let $T_0 > 0$ and F_ε be the solution built in Corollary 2.3 on $[0, T_0]$ and define $f_\varepsilon = \varepsilon^{-1} \frac{F_\varepsilon - \mu}{\sqrt{\mu}}$. The sequence $(f_\varepsilon)_{\varepsilon > 0}$ is uniformly bounded in $L_{[0, T_0]}^\infty H_{x,v}^s$ and solves

$$(5.1) \quad BE_\varepsilon(f_\varepsilon) = \varepsilon^2 \mathcal{E}(t, x, v) \mu^{1/2} + \varepsilon^3 \left[\vec{E}_t(x) \cdot \nabla_v f + \mathcal{E}(t, x, v) f \right]$$

where $BE_\varepsilon(f)$ is the standard Boltzmann equation operator given by

$$BE_\varepsilon(f) = \varepsilon^2 \partial_t f + \varepsilon v \cdot \nabla_x f - L[f] - \varepsilon \Gamma[f, f]$$

and we recall that

$$\mathcal{E}(t, x, v) = \varepsilon^e \frac{2A + a|v|^2}{4(1+t)^2} - \frac{1}{2} \left(1 + \varepsilon^{1+e} \frac{a}{1+t} \right) \vec{E}_t(x) \cdot v.$$

Having $(f_\varepsilon)_{\varepsilon>0}$ uniformly bounded in $L^\infty_{[0, T_0]} H^s_{x,v}$ means that up to a subsequence f_ε converges weakly-* in this space towards \bar{f} . The choice of $s \geq s_0$ allows us to take the weak-* limit in $BE_\varepsilon(f_\varepsilon)$ as it is now standard in the field [11, 33], [12, Section 8] and obtain

$$\lim_{\varepsilon \rightarrow 0} BE_\varepsilon(\bar{f}) = L[\bar{f}].$$

The right-hand side of (5.1) is linear in f_ε and it therefore converges weakly-* towards 0. In the limit one must have

$$L[\bar{f}] = 0 \quad \text{so} \quad \bar{f}(t, x, v) = \left[\rho(t, x) + v \cdot u(t, x) + \frac{|v|^2 - d}{2} \theta(t, x) \right] \sqrt{\mu(v)}.$$

The fluid equations are obtained by integrating in velocity (5.1) against $\sqrt{\mu}$, $v\sqrt{\mu}$ and $\frac{|v|^2 - (d+2)}{2} \sqrt{\mu}$. The computations on the Boltzmann equation part $BE_\varepsilon(f_\varepsilon)$ have been done and proven rigorously for weaker convergences [7, 20] and one obtain that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_{\mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ \frac{|v|^2 - (d+2)}{2} \end{pmatrix} \sqrt{\mu} BE_\varepsilon(f_\varepsilon) dv = \begin{pmatrix} \nabla_x \cdot u(t, x) \\ \nabla_x(\rho + \theta) \\ * \end{pmatrix}$$

and so looking at the right-hand side of (5.1) we see that in the limit

$$\nabla_x \cdot u(t, x) = 0 \quad \text{and} \quad \nabla_x(\rho + \theta) = 0$$

which are the incompressibility and Boussinesq relation.

Looking at the order ε^2 in $BE_\varepsilon(f_\varepsilon)$ yields the Navier-Stokes-Fourier system in the Leray sense [7, 20] - that is integrated against test functions with null divergence. It only remains to see what the right-hand side of (5.1) becomes in the limit at order ε^2 :

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \int_{\mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ \frac{|v|^2 - (d+2)}{2} \end{pmatrix} \left[\varepsilon^2 \mathcal{E}(t, x, v) \mu + \varepsilon^3 \left[\vec{E}_t(x) \cdot \nabla_v f + \mathcal{E}(t, x, v) f \right] \sqrt{\mu} \right] dv \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} \begin{pmatrix} 1 \\ v \\ \frac{|v|^2 - (d+2)}{2} \end{pmatrix} \frac{\vec{E}_t(x)}{2} \cdot v \mu dv = \begin{pmatrix} 0 \\ \frac{1}{2} \vec{E}_t(x) \\ 0 \end{pmatrix}. \end{aligned}$$

Therefore, taking the limit of the hydrodynamic quantities when ε goes to 0 of ε^{-2} (5.1) yields that (ρ, u, θ) is a Leray solution to the incompressible Navier-Stokes equation with a force (1.15) together with the Boussinesq equation (1.3). \square

REFERENCES

- [1] ACEVES-SANCHEZ, P., AND CESBRON, L. Fractional diffusion limit for a fractional Vlasov-Fokker-Planck equation. *SIAM J. Math. Anal.* 51, 1 (2019), 469–488.
- [2] ARSÉNIO, D. From Boltzmann's Equation to the Incompressible Navier–Stokes–Fourier System with Long-Range Interactions. *Arch. Ration. Mech. Anal.* 206, 2 (2012), 367–488.

- [3] ARSÉNIO, D., AND SAINT-RAYMOND, L. *From the Vlasov–Maxwell–Boltzmann System to Incompressible Viscous Electro-magneto-hydrodynamics*, vol. 1 of *EMS Monographs in Mathematics*. European Mathematical Society (EMS), Zürich, 2019.
- [4] BARANGER, C., AND MOUHOT, C. Explicit spectral gap estimates for the linearized Boltzmann and Landau operators with hard potentials. *Rev. Mat. Iberoamericana* 21, 3 (2005), 819–841.
- [5] BARDOS, C., GOLSE, F., AND LEVERMORE, D. Sur les limites asymptotiques de la théorie cinétique conduisant à la dynamique des fluides incompressibles. *C. R. Acad. Sci. Paris Sér. I Math.* 309, 11 (1989), 727–732.
- [6] BARDOS, C., GOLSE, F., AND LEVERMORE, D. Fluid dynamic limits of kinetic equations. I. Formal derivations. *J. Statist. Phys.* 63, 1-2 (1991), 323–344.
- [7] BARDOS, C., GOLSE, F., AND LEVERMORE, D. Fluid dynamic limits of kinetic equations. I. Formal derivations. *J. Statist. Phys.* 63, 1-2 (1991), 323–344.
- [8] BARDOS, C., GOLSE, F., AND LEVERMORE, D. Fluid dynamic limits of kinetic equations. II. Convergence proofs for the Boltzmann equation. *Comm. Pure Appl. Math.* 46, 5 (1993), 667–753.
- [9] BARDOS, C., GOLSE, F., AND LEVERMORE, D. Acoustic and Stokes limits for the Boltzmann equation. *C. R. Acad. Sci. Paris Sér. I Math.* 327, 3 (1998), 323–328.
- [10] BARDOS, C., GOLSE, F., AND LEVERMORE, D. The acoustic limit for the Boltzmann equation. *Arch. Ration. Mech. Anal.* 153, 3 (2000), 177–204.
- [11] BARDOS, C., AND UKAI, S. The classical incompressible Navier-Stokes limit of the Boltzmann equation. *Math. Models Methods Appl. Sci.* 1, 2 (1991), 235–257.
- [12] BRIANT, M. From the Boltzmann equation to the incompressible Navier-Stokes equations on the torus: a quantitative error estimate. *J. Differential Equations* 259, 11 (2015), 6072–6141.
- [13] CERCIGNANI, C. *The Boltzmann equation and its applications*, vol. 67 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1988.
- [14] CERCIGNANI, C., ILLNER, R., AND PULVIRENTI, M. *The mathematical theory of dilute gases*, vol. 106 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1994.
- [15] DUAN, R., LIU, S., YANG, T., AND ZHAO, H. Stability of the nonrelativistic Vlasov-Maxwell-Boltzmann system for angular non-cutoff potentials. *Kinet. Relat. Models* 6, 1 (2013), 159–204.
- [16] DUAN, R., UKAI, S., YANG, T., AND ZHAO, H. Optimal decay estimates on the linearized Boltzmann equation with time dependent force and their applications. *Comm. Math. Phys.* 277, 1 (2008), 189–236.
- [17] DUAN, R., YANG, T., AND ZHAO, H. The Vlasov-Poisson-Boltzmann system in the whole space: the hard potential case. *J. Differential Equations* 252, 12 (2012), 6356–6386.
- [18] DUAN, R., YANG, T., AND ZHAO, H. The Vlasov-Poisson-Boltzmann system for soft potentials. *Math. Models Methods Appl. Sci.* 23, 6 (2013), 979–1028.
- [19] ELLIS, R. S., AND PINSKY, M. A. The first and second fluid approximations to the linearized Boltzmann equation. *J. Math. Pures Appl. (9)* 54 (1975), 125–156.
- [20] GOLSE, F. Fluid dynamic limits of the kinetic theory of gases. In *From particle systems to partial differential equations*, vol. 75 of *Springer Proc. Math. Stat.* Springer, Heidelberg, 2014, pp. 3–91.
- [21] GOLSE, F., AND LEVERMORE, C. D. Stokes-Fourier and acoustic limits for the Boltzmann equation: convergence proofs. *Comm. Pure Appl. Math.* 55, 3 (2002), 336–393.
- [22] GOLSE, F., AND SAINT-RAYMOND, L. The Navier-Stokes limit of the Boltzmann equation for bounded collision kernels. *Invent. Math.* 155, 1 (2004), 81–161.
- [23] GOLSE, F., AND SAINT-RAYMOND, L. The incompressible Navier-Stokes limit of the Boltzmann equation for hard cutoff potentials. *J. Math. Pures Appl. (9)* 91, 5 (2009), 508–552.
- [24] GRAD, H. Principles of the kinetic theory of gases. In *Handbuch der Physik (herausgegeben von S. Flügge)*, Bd. 12, *Thermodynamik der Gase*. Springer-Verlag, Berlin, 1958, pp. 205–294.
- [25] GUALDANI, M. P., MISCHLER, S., AND MOUHOT, C. Factorization of non-symmetric operators and exponential H -theorem. *Mém. Soc. Math. Fr. (N.S.)*, 153 (2017), 137.
- [26] GUO, Y. The Vlasov-Poisson-Boltzmann system near Maxwellians. *Comm. Pure Appl. Math.* 55, 9 (2002), 1104–1135.
- [27] GUO, Y. Boltzmann diffusive limit beyond the Navier-Stokes approximation. *Comm. Pure Appl. Math.* 59, 5 (2006), 626–687.

- [28] KIM, C. Boltzmann equation with a large potential in a periodic box. *Comm. Partial Differential Equations* 39, 8 (2014), 1393–1423.
- [29] LEVERMORE, D., AND MASMOUDI, N. From the Boltzmann equation to an incompressible Navier-Stokes-Fourier system. *Arch. Ration. Mech. Anal.* 196, 3 (2010), 753–809.
- [30] LIONS, P.-L., AND MASMOUDI, N. From the Boltzmann equations to the equations of incompressible fluid mechanics. I, II. *Arch. Ration. Mech. Anal.* 158, 3 (2001), 173–193, 195–211.
- [31] MOUHOT, C. Explicit coercivity estimates for the linearized Boltzmann and Landau operators. *Comm. Partial Differential Equations* 31, 7-9 (2006), 1321–1348.
- [32] MOUHOT, C., AND NEUMANN, L. Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus. *Nonlinearity* 19, 4 (2006), 969–998.
- [33] SAINT-RAYMOND, L. *Hydrodynamic limits of the Boltzmann equation*, vol. 1971 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
- [34] VILLANI, C. Limites hydrodynamiques de l'équation de Boltzmann (d'après C. Bardos, F. Golse, C. D. Levermore, P.-L. Lions, N. Masmoudi, L. Saint-Raymond). *Astérisque*, 282 (2002). Séminaire Bourbaki, Vol. 2000/2001.
- [35] VILLANI, C. A review of mathematical topics in collisional kinetic theory. In *Handbook of mathematical fluid dynamics, Vol. I*. North-Holland, Amsterdam, 2002, pp. 71–305.
- [36] XIAO, Q., XIONG, L., AND ZHAO, H. The Vlasov-Poisson-Boltzmann system with angular cutoff for soft potentials. *J. Differential Equations* 255, 6 (2013), 1196–1232.
- [37] XIAO, Q., XIONG, L., AND ZHAO, H. The Vlasov-Poisson-Boltzmann system for the whole range of cutoff soft potentials. *J. Funct. Anal.* 272, 1 (2017), 166–226.

MARC BRIANT

UNIVERSITÉ DE PARIS,
MAP5, CNRS UMR 8145
F-75006 PARIS, FRANCE

E-MAIL: briant.maths@gmail.com

ARNAUD DEBUSSCHE

ENS RENNES,
AVENUE ROBERT SCHUMANN,
35170 BRUZ, FRANCE

E-MAIL: arnaud.debussche@ens-rennes.fr

JULIEN VOVELLE

UMPA UMR 5669 CNRS,
ENS DE LYON SITE MONOD,
46, ALLE D'ITALIE,
9364 LYON CEDEX 07, FRANCE

E-MAIL: julien.vovelle@ens-lyon.fr