



HAL
open science

Model Predictive Control, Cost Controllability, and Homogeneity *

Jean-Michel Coron, Lars Grüne, Karl Worthmann

► **To cite this version:**

Jean-Michel Coron, Lars Grüne, Karl Worthmann. Model Predictive Control, Cost Controllability, and Homogeneity *. 2019. hal-02151402v1

HAL Id: hal-02151402

<https://hal.science/hal-02151402v1>

Preprint submitted on 8 Jun 2019 (v1), last revised 9 Dec 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Model Predictive Control, Cost Controllability, and Homogeneity*

Jean-Michel Coron[†] Lars Grüne[‡] Karl Worthmann[§]

June 4, 2019

We are concerned with the design of Model Predictive Control (MPC) schemes such that asymptotic stability of the resulting closed loop is guaranteed – even if the linearization at the desired set point fails to be stabilizable. Therefore, we propose to construct the stage cost based on the homogeneous approximation and rigorously show that applying MPC yields an asymptotically stable closed-loop behavior if the homogeneous approximation is asymptotically null controllable. To this end, we verify cost controllability – a condition relating the current state, the stage cost, and the growth of the value function w.r.t. time – for this class of systems in order to provide stability and performance guarantees for the proposed MPC scheme without stabilizing terminal costs or constraints.

1 Introduction

Model Predictive Control (MPC) is nowadays a well-established and widely applied control technique for linear and nonlinear systems, see, e.g. the textbooks [15, 21, 27] and the survey papers [12, 22]. One of the main reasons for its success is the simplicity of the underlying idea: measure the current state of the system in consideration, optimize its

*Jean-Michel Coron acknowledges funding from the Miller Institute and from the Agence Nationale de La Recherche (ANR), grant ANR Finite4SoS. He also thanks the Miller Institute and UC Berkeley for their hospitality. Karl Worthmann gratefully acknowledges funding from the German Research Foundation (Deutsche Forschungsgemeinschaft, grant WO 2056/6-1).

[†]Sorbonne Université, CNRS, INRIA, Laboratoire Jacques-Louis Lions, équipe Cage, Paris, France (*coron@ann.jussieu.fr*).

[‡]Mathematical Institute, Faculty of Mathematics, Physics and Computer Science, University of Bayreuth, Germany (*lars.gruene@uni-bayreuth.de*).

[§]Institute for Mathematics, Faculty of Mathematics and Natural Sciences, Technische Universität Ilmenau, Ilmenau, Germany (*karl.worthmann@tu-ilmenau.de*).

behaviour on a finite prediction window, and implement the first piece of the computed control function before repeating the procedure ad infinitum. However, the stability analysis (or the construction of stabilizing terminal costs and constraints) is often based on the (explicit or implicit) assumption that the linearization at the desired set point is stabilizable [8]. Otherwise, the origin may not be asymptotically stable w.r.t. the MPC closed loop if, e.g. a quadratic stage cost is used – independently of the length of the optimization horizon as rigorously shown in [26] for the mobile robot example, i.e. a finite-time controllable system.

We propose to design the stage cost based on the homogeneous approximation [18, 2] and [10, Section 12.3]. Homogeneity – which is a generalization of linearity to nonlinear systems – is a property, which was extensively used to construct stabilizing (homogeneous) control laws [19, 11, 17, 24, 25]. We show that, if the homogeneous approximation is stabilizable, the desired set point is asymptotically stable w.r.t. the MPC closed-loop without terminal costs or constraints. Here, asymptotic stability is understood in a global (homogeneous system with degree zero), semi-global (negative degree of homogeneity), or local (homogeneous approximation) sense. All results are motivated and illustrated by examples.

To this end, we verify *cost controllability* – a condition relating the growth of the value function w.r.t. time with the stage cost. Then, stability guarantees or a desired performance bound for the proposed MPC scheme can be deduced if the prediction horizon is sufficiently large using techniques originally developed in a discrete-time setting, see [13, 16], adopted to the cost-controllability condition [31, 32], and transferred to the continuous-time setting [28, 35]. While we first show our results for homogeneous systems, we make use of homogeneous Lyapunov functions (see, in particular, [29, 11, 14, 4] and [10, Section 12.3]) in order to show that our results are locally preserved for (general) nonlinear systems, i.e. that cost controllability and local asymptotic stability of the origin for the MPC closed loop hold if the homogeneous approximation at the origin is stabilizable.

The paper is organized as follows: In section 2, cost controllability and its relationship with MPC are recalled and adapted to our setting. Then, we introduce homogeneous control systems in accordance to [14] and [5] and give an example that MPC with quadratic costs may fail for finite-time stabilizable systems with zero degree of homogeneity. In section 3, we establish cost controllability and asymptotic stability of the origin for the MPC closed loop for control systems with zero degree of homogeneity and the local counterparts of these results if the degree of homogeneity is negative. Then, in section 4, we extend our results to systems with globally asymptotically null controllable homogeneous approximation before conclusions are drawn in section 5.

Notation: We call a continuous function $\rho : [0, \infty) \rightarrow [0, \infty)$ of class \mathcal{K}_∞ if it maps zero to zero, is strictly monotonically increasing, and unbounded. For further details on comparison functions, we refer to the compendium [20] and the references therein.

2 Problem Formulation

We consider the control system governed by the autonomous ordinary differential equation

$$\dot{x}(t) = f(x(t), u(t)) \quad (1)$$

with state $x(t) \in \mathbb{R}^n$ and control $u(t) \in \mathbb{R}^m$ at time $t \geq 0$. We assume continuity of the map $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$ and local Lipschitz continuity w.r.t. its first argument on $\mathbb{R}^n \setminus \{0\}$ such that, for each initial value x^0 and each control function $u \in \mathcal{L}_{\text{loc}}^\infty([0, \infty), \mathbb{R}^m)$, the solution trajectory $x(\cdot; x^0, u)$ uniquely exists (on its maximal interval I_{x^0} of existence).

2.1 Model Predictive Control

Let $f(0, 0) = 0$ hold, i.e. the origin $0 \in \mathbb{R}^n$ is a (controlled) equilibrium. Our goal is the stabilization of system eq. (1) at the origin. As a preliminary condition, we require a stage cost $\ell : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}_{\geq 0}$ satisfying

$$\rho(\|x\|) \leq \ell^*(x) := \inf_{u \in \mathbb{R}^m} \ell(x, u) \leq \varrho(\|x\|) \quad \forall x \in \mathbb{R}^n, \quad (2)$$

for \mathcal{K}_∞ -functions ρ and ϱ .

We briefly introduce model predictive control. The basic idea is simple: For the current state $\hat{x} \in \mathbb{R}^n$, optimize a cost functional on a finite prediction horizon $[0, T]$ w.r.t. the system dynamics eq. (1), the initial condition $x(0) = \hat{x}$, and, if present, state and control constraints. Then, the first portion $u^*(t)$, $t \in [0, \delta)$, of the computed optimal control function $u^* = u^*(\cdot; \hat{x}) \in \mathcal{L}^\infty([0, T], \mathbb{R}^m)$ is implemented before the prediction window is shifted forward in time. The procedure is repeated ad infinitum. Assuming existence of a minimizer for simplicity of exposition, the MPC scheme reads as displayed in algorithm 1.

Algorithm 1 Model Predictive Control

Set $t = 0$ and let the prediction horizon T and the time shift $\delta \in (0, T)$ be given.

1. Measure the current state $\hat{x} := x(t)$
2. Solve the optimal control problem

$$\min_{u \in \mathcal{L}^\infty([0, T], \mathbb{R}^m)} \int_0^T \ell(x(s; \hat{x}, u), u(s)) ds \quad \text{subject to} \quad x(0; \hat{x}, u) = \hat{x} \text{ and eq. (1)}$$

and denote its solution by $u^* = u^*(\cdot; \hat{x})$.

3. Implement $u^*(s)|_{s \in [0, \delta)}$ at the plant, increment t by δ , and goto Step 1.
-

We define the value function $V_t : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ by

$$V_t(\hat{x}) := \inf_{u \in \mathcal{L}^\infty([0, t], \mathbb{R}^m)} \int_0^t \ell(x(s; \hat{x}, u), u(s)) ds \quad \text{subject to} \quad x(0; \hat{x}, u) = \hat{x} \text{ and eq. (1),}$$

with the convention that $V_t(\hat{x}) = \infty$ if there exists no $u \in \mathcal{L}^\infty([0, t], \mathbb{R}^m)$ such that $x(t; \hat{x}, u)$, $t \in [0, T]$, is defined. This definition allows us to state the following condition relating the stage cost ℓ at the current state \hat{x} and the value function.

Assumption 1 (Cost controllability on a set K). *Consider a set $K \subseteq \mathbb{R}^n$ containing 0 in its interior. We assume that there exists a monotonically increasing, bounded function $B_K : [0, \infty) \rightarrow \mathbb{R}_{\geq 0}$, which satisfies the inequality*

$$V_t(x) \leq B_K(t) \ell^*(x) \quad \forall (t, x) \in \mathbb{R}_{\geq 0} \times K. \quad (3)$$

For the formulation of the following theorem, which is an extension of a result from [28], we moreover need the sublevel set

$$V_T^{-1}([0, L]) := \{x \in \mathbb{R}^n \mid V_T(x) \leq L\}.$$

Theorem 1. *Consider the MPC scheme given in algorithm 1 and let Condition eq. (2) and assumption 1 for a set $K \subseteq \mathbb{R}^n$ be satisfied for the stage cost ℓ and the system dynamics eq. (1). Then, for given $\delta > 0$, there exists a sufficiently large prediction horizon $T > \delta$ such that the origin is locally asymptotically stable for the MPC closed loop. Moreover, the domain of attraction contains all level sets $V_T^{-1}([0, L]) \subseteq K$.*

If $K = \mathbb{R}^n$, the origin is globally asymptotically stable for the MPC closed loop. Moreover, if the assumptions hold for each compact set $K \subset \mathbb{R}^n$ containing 0 in its interior (with B_K possibly depending on K), then the origin is semiglobally asymptotically stable for the MPC closed loop, i.e., for each compact set $\hat{K} \subset \mathbb{R}^n$ and each $\delta > 0$ there is $T > 0$, such that the origin is locally asymptotically stable for the MPC closed loop and \hat{K} is contained in the domain of attraction.

Proof. The assertion for $K = \mathbb{R}^n$ was already shown in [28]. To prove the non-global version, i.e. $K \subsetneq \mathbb{R}^n$, we proceed similarly to the discrete time counterpart given in [15, Theorem 6.35] (see also [6] for related results). Hence we only give a sketch.

First one shows using the same techniques as in [28] that for sufficiently large T the optimal value function V_T decreases along solutions in K . This implies that each level set $V^{-1}([0, L]) \subset K$ is forward invariant. On this level set asymptotic stability with domain of attraction $V^{-1}([0, L])$ follows with usual Lyapunov function arguments. The second statement follows by estimating the size of the level sets $V_T^{-1}([0, L])$ exactly as in the proof of [15, Theorem 6.35]. \square

Monotonicity of the (growth) function B_K in assumption 1 can be assumed w.l.o.g. since the left hand side of Inequality eq. (3) is monotonically increasing due to the absence of a terminal cost. Furthermore, the condition $\limsup_{t \rightarrow \infty} B_K(t)/t < 1$ suffices to ensure the existence of a bounded (growth) function B_K , see [26, Proposition 6]. Using the abbreviation $\gamma := \limsup_{t \rightarrow \infty} B_K(t)$, assumption 1 ensures the inequality

$$\sup_{\hat{x} \in K \setminus \{0\}} \left(\frac{V_t(\hat{x})}{\ell^*(\hat{x})} \right) \leq \gamma \quad \forall t \geq 0.$$

In particular, this implies that the value function V_t cannot contain terms of lower order than the *minimized* stage cost $\ell^*(\hat{x}) = \inf_{u \in \mathbb{R}^m} \ell(\hat{x}, u)$. Otherwise assumption 1 cannot hold in a neighbourhood of the origin and, thus, on the set K .

2.2 Homogeneous Systems and quadratic stage costs

The following definition of homogeneity is taken from [14] with a slightly adapted notation.

Definition 1 (Homogeneous System).

Let $r = (r_1, \dots, r_n) \in (0, +\infty)^n$, let $s = (s_1, \dots, s_m) \in (0, +\infty)^m$ and let $\tau \in (-\min_i r_i, \infty)$. System eq. (1) is said to be (r, s, τ) -homogeneous if

$$f(\Lambda_\alpha x, \Delta_\alpha u) = \alpha^\tau \Lambda_\alpha f(x, u) \quad \forall (x, u) \in \mathbb{R}^n \times \mathbb{R}^m \text{ and } \alpha \geq 0 \quad (4)$$

is satisfied, where the matrices $\Lambda_\alpha \in \mathbb{R}^{n \times n}$ and $\Delta_\alpha \in \mathbb{R}^{m \times m}$ are defined as

$$\Lambda_\alpha = \begin{pmatrix} \alpha^{r_1} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & \alpha^{r_n} \end{pmatrix} \quad \text{and} \quad \Delta_\alpha = \begin{pmatrix} \alpha^{s_1} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & \alpha^{s_m} \end{pmatrix}.$$

Homogeneity can be considered as a generalization of linearity. For linear systems, i.e. $\dot{x}(t) = Ax(t) + Bu(t)$, homogeneity is trivially satisfied with $\tau = 0$ and $r_i = 1 = s_j$ for $(i, j) \in \{1, \dots, n\} \times \{1, \dots, m\}$. Using definition 1, one easily obtains the identity

$$x(t; \Lambda_\alpha x^0, \Delta_\alpha u(\alpha^\tau \cdot)) = \Lambda_\alpha x(\alpha^\tau t; x^0, u) \quad (5)$$

for all $t \in I_{x^0} \cap [0, \infty)$. For $\tau = 0$, the Identity eq. (5) simplifies to $x(t; \Lambda_\alpha x^0, \Delta_\alpha u) = \Lambda_\alpha x(t; x^0, u)$.

We consider the following example in order to show that the design of stage cost in MPC matters.

Example 1. Let the system dynamics eq. (1) be given by

$$\dot{x}(t) = f(x(t), u(t)) := \begin{pmatrix} u_1(t) \\ x_3(t)u_1(t) \\ u_2(t) \end{pmatrix} = \begin{pmatrix} 1 \\ x_3(t) \\ 0 \end{pmatrix} u_1(t) + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} u_2(t). \quad (6)$$

example 1 is governed by homogeneous system dynamics since Condition eq. (4) holds with $\tau = 0$, $r_2 = 2$, and $r_1 = r_3 = s_1 = s_2 = 1$. Moreover, the system is null controllable in finite time, which means that starting from any given state one can steer the control system to $0 \in \mathbb{R}^3$ in finite time (even in arbitrary small positive time). This property follows from the Rashevski-Chow Theorem [10, Theorem 3.18]. It can also be checked directly as follows. Ensure that the x_3 -component is not equal to 0 by using u_2 . Then take $u_2 = 0$ (so x_3 does not change) and use u_1 to steer the x_2 -component to 0. Then take $u_1 = 0$ (so x_2 remains equal to 0) and set the x_3 -component equal to 0 using u_2 . Finally, take $u_2 = 0$ (so both x_2 and x_3 remain equal to 0) and use u_1 to steer the x_1 -component to 0. It follows from this method to steer the control system eq. (6) to $0 \in \mathbb{R}^3$ that the value function V_t is uniformly bounded w.r.t. $t \in \mathbb{R}_{\geq 0}$ on compact sets of \mathbb{R}^3 . Nevertheless, one can show that there does not exist a (finite) prediction horizon such that the origin is asymptotically stable w.r.t. the MPC closed loop.

Proposition 1. *We consider the MPC scheme presented in algorithm 1 for the combination of System eq. (6) and the quadratic stage cost $\ell(x, u) = x^\top Qx + u^\top Ru$ with positive definite matrices $Q \in \mathbb{R}^{3 \times 3}$ and $R \in \mathbb{R}^{2 \times 2}$. Then, for arbitrary but fixed prediction horizon T , the origin is not (locally) asymptotically for the MPC closed loop.*

Proof. Firstly, it can be shown that, for every $\varepsilon > 0$, there exists an initial value $x^0 \in \{x \in \mathbb{R}^3 \mid \|x\| \leq \varepsilon\}$ such that $u^* \equiv 0$ is the unique optimal control. The proof is almost literally the same as its analogon for the nonholonomic robot presented in [26, Subsection 4.1] if Condition (13) is replaced by $(Qx^0)_3 = 0 = (Qx^0)_1 + (Qx^0)_2\theta^0$ in view of the slightly different adjoint equation. Hence, the closed-loop trajectory emanating at x^0 stays at x^0 forever and does, thus, not converge to the origin, which shows the assertion. \square

We note that this implies that assumption 1 must be violated. In the following, our goal is to construct a stage cost such that asymptotic stability of the origin w.r.t. the MPC closed loop is ensured for a sufficiently long prediction horizon T for a class of systems including both example 1 and the mentioned nonholonomic robot example provided that the homogeneous (approximation of the) system is globally asymptotically null controllable. Moreover, we want to keep the construction simple by solely replacing the quadratic performance measure by one based on the *dilated norm* $N : \mathbb{R}^n \rightarrow [0, \infty)$, i.e.

$$N(x) := \left(\sum_{i=1}^n x_i^{\frac{d}{r_i}} \right)^{\frac{1}{d}} \quad \text{with} \quad d := 2 \prod_{i=1}^n r_i. \quad (7)$$

Note that the *dilated norm* satisfies the equality

$$N(\Lambda_\alpha x) = \sqrt[d]{\sum_{i=1}^n (\alpha^{r_i} x_i)^{\frac{d}{r_i}}} = \sqrt[d]{\alpha^d \sum_{i=1}^n x_i^{\frac{d}{r_i}}} = \alpha N(x) \quad \forall (\alpha, x) \in \mathbb{R}_{\geq 0} \times \mathbb{R}^n \quad (8)$$

and that it is positive on $\mathbb{R}^n \setminus \{0\}$. However it is not a norm unless $\Lambda_\alpha = \alpha I$.

3 Cost Controllability for Homogeneous Systems

We consider homogeneous systems and propose to define the stage cost $\ell : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}_{\geq 0}$ by

$$\ell(x, u) := N(x)^d + \sum_{j=1}^m u_j^{\frac{d}{s_j}} = \sum_{i=1}^n x_i^{\frac{d}{r_i}} + \sum_{j=1}^m u_j^{\frac{d}{s_j}}. \quad (9)$$

The stage cost is based on the *dilated norm* and the coefficients $r_i, i \in \{1, \dots, n\}$, and $s_j, j \in \{1, \dots, m\}$, from definition 1. Note that Definition eq. (9) leads to quadratic stage cost for linear systems $\dot{x}(t) = Ax(t) + Bu(t)$. Moreover, the definition is compatible with homogeneity analogously to eq. (8).

Proposition 2. *The stage costs ℓ are homogeneous, i.e.*

$$\ell(x(s; \Lambda_\alpha x^0, \Delta_\alpha u(\alpha^\tau \cdot)), \Delta_\alpha u(\alpha^\tau s)) = \alpha^d \ell(x(\alpha^\tau s; x^0, u), u(\alpha^\tau s)) \quad (10)$$

holds for $\alpha \geq 0$. For $\tau = 0$, equality eq. (10) simplifies to

$$\ell(x(s; \Lambda_\alpha x^0, \Delta_\alpha u), \Delta_\alpha u(s)) = \alpha^d \ell(x(s; x^0, u), u(s)). \quad (11)$$

Proof. A direct calculation shows

$$\begin{aligned} \ell(x(s; \Lambda_\alpha x^0, \Delta_\alpha u(\alpha^\tau \cdot)), \Delta_\alpha u(t)) &= \underbrace{N(x(s; \Lambda_\alpha x^0, \Delta_\alpha u(\alpha^\tau \cdot)))^d}_{\stackrel{\text{eq. (5)}}{=} \Lambda_\alpha x(\alpha^\tau s; x^0, u)} + \sum_{j=1}^m (\alpha^{s_j} u_j(t))^{\frac{d}{s_j}} \\ &\stackrel{\text{eq. (8)}}{=} \alpha^d N(x(\alpha^\tau s; x^0, u))^d + \alpha^d \sum_{j=1}^m u_j(t)^{\frac{d}{s_j}} \\ &= \alpha^d \ell(x(\alpha^\tau s; x^0, u), u(t)), \end{aligned}$$

i.e. the assertion if t is replaced by $\alpha^\tau s$. \square

Our goal in this section is to verify cost controllability for homogeneous, globally asymptotically null controllable systems.

Definition 2. *The system eq. (1) is said to be globally asymptotically null controllable if, for each $x^0 \in \mathbb{R}^n$, there exists a control function $u_{x^0} \in \mathcal{L}_{\text{loc}}^\infty([0, \infty), \mathbb{R}^m)$ such that $\lim_{t \rightarrow \infty} \|x(t; x^0, u_{x^0})\| = 0$ holds.*

Note that a locally asymptotically null controllable homogeneous system is always globally asymptotically null controllable.

Inequality eq. (3) is equivalent to

$$V(t) \leq B_K(t) N(x^0)^d \quad \forall (t, x^0) \in \mathbb{R}_{\geq 0} \times K, \quad (12)$$

using the stage cost eq. (9). Under this condition, asymptotic stability of the origin w.r.t. the MPC closed loop is ensured for a sufficiently large prediction horizon T using theorem 1 since Condition eq. (2) is satisfied.

Before we proceed, we consider a simple example in order to illustrate and motivate our results.

Example 2. *Let the system dynamics eq. (1) be given by*

$$\dot{x}(t) = |x(t)|^k \text{sign}(x) + u \quad \text{with } k \in (0, +\infty). \quad (13)$$

Since we have

$$\begin{aligned} f(\Lambda_\alpha x, \Delta_\alpha u) &= |\alpha^r x|^k \text{sign}(x) + \alpha^s u = |\alpha^r x|^k \text{sign}(x) + \alpha^s u, \\ \alpha^\tau \Lambda_\alpha f(x, u) &= \alpha^{\tau+r} (|x|^k \text{sign}(x) + u), \end{aligned}$$

we get the (r, s, τ) -homogeneity of eq. (13) if and only if $\tau + r = rk$ and $\tau + r = s$. This leads to the following three cases:

- $k = 1$: Degree of homogeneity zero, i.e. $\tau = 0$ with coefficients $r = s$
- $k \in (0, 1)$: Negative degree of homogeneity, i.e. $\tau < 0$ with coefficients $r = s/k$
- $k > 1$: Positive degree of homogeneity, i.e. $\tau > 0$ with $r = s/k$

Next, we compute the infinite horizon optimal value function using the Hamilton-Jacobi-Equation

$$\min_{u \in \mathbb{R}^m} \left(\langle \nabla V(x), f(x, u) \rangle + \ell(x, u) \right) = 0 \quad (14)$$

with $m = 1$. To this end, we quadratically penalize the control effort to simplify the following calculations by using the stage cost $\ell : \mathbb{R}^n \times \mathbb{R}^m$, $n = m = 1$, defined by $\ell(x, u) = (x^2)^d + u^2$, which yields $u^* = -\frac{1}{2}V'(x)$. Then, plugging this expression into eq. (14), i.e. $V'(x) (|x|^k \text{sign}(x) + u^*) + \ell(x, u^*) = 0$, leads to $V'(x)|x|^k \text{sign}(x) - \frac{1}{4}V'(x)^2 + (x^2)^d = 0$, or (only the root having the sign of x has to be taken into account since V is increasing on $[0, +\infty)$ and decreasing on $(-\infty, 0]$)

$$V'(x) = 2|x|^k \text{sign}(x) + \text{sign}(x) \sqrt{4|x|^{2k} + 4(x^2)^d} \stackrel{d=k}{=} 2(1 + \sqrt{2})|x|^k \text{sign}(x).$$

Overall, we get the value function

$$V(x) = \frac{2(1 + \sqrt{2})}{k + 1} |x|^{k+1} \quad \text{and, thus,} \quad \frac{V(x)}{\ell^*(x)} = \frac{2(1 + \sqrt{2})}{k + 1} |x|^{1-k}.$$

In conclusion, we get for

- $k = 1$: The growth bound is uniformly upper bounded by $1 + \sqrt{2}$, i.e. cost controllability in the sense of assumption 1 with $K = \mathbb{R}^n = \mathbb{R}$.
- $k \in (0, 1)$: The growth bound is uniformly upper bounded on each compact set K by $\frac{2(1+\sqrt{2})}{k+1} \max_{x \in K} \{|x|^{1-k}\}$, i.e. cost controllability in the sense of assumption 1.
- $k > 1$: the quotient $V(x)/\ell^*(x)$ is uniformly upper bounded on $\mathbb{R} \setminus (-\varepsilon, \varepsilon)$ for arbitrary but fixed $\varepsilon > 0$ by $\frac{2(1+\sqrt{2})}{k+1} \max_{x \in \mathbb{R} \setminus (-\varepsilon, \varepsilon)} \{|x|^{1-k}\}$.

Moreover, note that the choice $d = 1$, i.e. purely quadratic costs, implies for $k \in (0, 1)$ that the value function still exhibits a term of order $1 + k$ but $\ell^*(x) = x^2$ holds. Hence, there does not exist a uniform bound for the quotient $V(x)/\ell^*(x)$ for $x \rightarrow 0$, $x \neq 0$.

Remark 1. In general, the degree of homogeneity ($\tau < 0$, $\tau = 0$, and $\tau > 0$) is not unique, which can be seen from the (driftless) system example 1. Here, Condition eq. (4) is also satisfied for the combinations

$$\tau = 0.5, r_1 = r_3 = 1, r_2 = 2, \text{ and } s_1 = s_2 = 1.5$$

and

$$\tau = -0.5, r_1 = r_3 = 1, r_2 = 2, \text{ and } s_1 = s_2 = 0.5.$$

We emphasize that the relative weighting of the control in comparison to the penalization of deviations w.r.t. the state depends on the choice, which explains the different behavior of the MPC closed loop. For control affine systems, we set $s_j = 2$ whenever possible in view of the Hamilton-Jacobi-Equation.

3.1 Degree of homogeneity zero

In this subsection, we derive cost controllability and, then, conclude asymptotic stability for globally asymptotically null controllable systems with degree of homogeneity $\tau = 0$.

Theorem 2 (Cost Controllability). *Let the system eq. (1) be globally asymptotically null controllable and $(r, s, \tau = 0)$ -homogeneous. In addition, let the stage cost be defined by eq. (9). Then, cost controllability, i.e. inequality eq. (12), holds. In particular, given $\delta > 0$, for any sufficiently large $T > 0$ the origin is globally asymptotically stable for the MPC closed loop.*

Proof. First, we show inequality eq. (12) only for initial values x^0 contained in the compact set

$$\mathcal{N}_{c_1, c_2} := \{x \in \mathbb{R}^n \mid c_1 \leq N(x)^d \leq c_2\}$$

with $0 < c_1 < c_2 < \infty$. Later on, we show that the same growth function B can be used for arbitrary initial values $x^0 \in \mathbb{R}^n$.

Global asymptotical null controllability implies that, for each $x^0 \in \mathcal{N}_{c_1, c_2}$, there exists a control function $u_{x^0} \in \mathcal{L}_{\text{loc}}^\infty([0, \infty), \mathbb{R}^m)$ such that $\lim_{t \rightarrow \infty} x(t; x^0, u_{x^0}) = 0$ holds. Hence, continuity of the solution trajectory $x(\cdot; x^0, u_{x^0})$ implies the existence of a time instant $T_{x^0} \in (0, \infty)$ such that

$$N(x(T_{x^0}; x^0, u_{x^0}))^d = \alpha \frac{c_1 + c_2}{2}$$

where $\alpha \in (0, 1)$ is chosen such that $\alpha(c_1 + c_2)/2 < c_1$ holds, i.e. $x(T_{x^0}; x^0, u_{x^0})$ is contained in the sublevel set $\{x \in \mathbb{R}^n \mid N(x)^d < c_1\}$. Due to continuity and compactness of the time interval $[0, T_{x^0}]$, there exists $\delta_{x^0} > 0$ sufficiently small such that the chain of inequalities

$$\alpha c_1 < N(x(T_{x^0}; \bar{x}, u_{x^0}))^d < \alpha c_2 \quad (15)$$

holds for all $\bar{x} \in \mathcal{B}_{\delta_{x^0}}(x^0)$. Since $\bigcup_{x^0 \in \mathcal{N}_{c_1, c_2}} \mathcal{B}_{\delta_{x^0}}(x^0)$ is an open cover of the compact set \mathcal{N}_{c_1, c_2} , there exists a finite subcover and, thus, there exist finitely many *initial values* x_1^0, \dots, x_k^0 , $k \in \mathbb{N}$, satisfying

$$\mathcal{N}_{c_1, c_2} \subset \bigcup_{i=1}^k \mathcal{B}_{\delta_{x_i^0}}(x_i^0).$$

Let us set $T^* := \min_{i=1, \dots, k} T_{x_i^0} \in (0, \infty)$. Then, the following function is obviously a growth function B on the finite interval $[0, T^*]$ for all $x^0 \in \mathcal{N}_{c_1, c_2}$:

$$B(t) := \max_{i=1, \dots, k} \sup_{\bar{x} \in \mathcal{B}_{\delta_{x_i^0}}(x_i^0) \cap \mathcal{N}_{c_1, c_2}} N(\bar{x})^{-d} \int_0^t \ell(x(s; \bar{x}, u_{x_i^0}), u_{x_i^0}(s)) ds \quad \forall t \in [0, T^*),$$

$$B(T^*) := \max_{i=1, \dots, k} \sup_{\bar{x} \in \mathcal{B}_{\delta_{x_i^0}}(x_i^0) \cap \mathcal{N}_{c_1, c_2}} N(\bar{x})^{-d} \int_0^{T_{x_i^0}} \ell(x(s; \bar{x}, u_{x_i^0}), u_{x_i^0}(s)) ds. \quad (16)$$

Note that the preceding line of argumentation implicitly implies well-posedness and boundedness of B on $[0, T^*]$ in view of the fact that $N(\bar{x})$ is uniformly bounded from

below by c_1 .¹ This construction is quite conservative and does not rely on homogeneity. But it is essential to have a bounded B on a compact time interval of positive length for the following line of argumentation.

Before we proceed, let us present some preliminary considerations. Let $\tilde{x}^0 \notin \mathcal{N}_{c_1, c_2}$ be arbitrary but fixed. Then, if $\tilde{x}^0 \neq 0$, there exists a scaling factor $c \in (0, \infty)$ such that $\Lambda_c \tilde{x}^0 \in \mathcal{N}_{c_1, c_2}$. This implies the existence of an index i^* , $i^* \in \{1, 2, \dots, k\}$, such that $\Lambda_c \tilde{x}^0 \in \mathcal{B}_{\delta_{x_{i^*}^0}}(x_{i^*}^0)$. Hence, using $\tilde{x}^0 = \Lambda_{c^{-1}}(\Lambda_c \tilde{x}^0)$, we get

$$\ell(x(s; \tilde{x}^0, \Delta_{c^{-1}} u_{x_{i^*}^0}), \Delta_{c^{-1}} u_{x_{i^*}^0}(s)) \stackrel{eq. (11)}{=} c^{-d} \ell(x(s; \Lambda_c \tilde{x}^0, u_{x_{i^*}^0}), u_{x_{i^*}^0}(s)) \quad (17)$$

and, for $\ell^*(\tilde{x}^0) := \inf_{u \in \mathbb{R}^m} \ell(\tilde{x}^0, u)$,

$$\ell^*(\tilde{x}^0) = N(\tilde{x}^0)^d = N(\Lambda_{c^{-1}} \Lambda_c \tilde{x}^0)^d \stackrel{eq. (8)}{=} c^{-d} N(\Lambda_c \tilde{x}^0)^d = c^{-d} \ell^*(\Lambda_c \tilde{x}^0). \quad (18)$$

Hence, showing the desired inequality (see assumption 1) for $x^0 \in \mathcal{N}_{c_1, c_2}$ suffices.

The Identity section 3.1 is also the key ingredient to show the assertion for $t > T^*$ based on the following contraction argument. For each $x^0 \in \mathcal{N}_{c_1, c_2}$, there exists an index i^\sharp , $i^\sharp \in \{1, \dots, k\}$, and a time instant $t_{x^0} \in [T^*, \max_{i=1, \dots, k} T_{x_i^0}]$ such that $x(t_{x^0}; x^0, u_{x_{i^\sharp}^0}) \in \alpha \mathcal{N}_{c_1, c_2}$ and

$$\int_0^t \ell(x(s; x^0, u_{x_{i^\sharp}^0}), u_{x_{i^\sharp}^0}(s)) ds \leq B(\min\{t, T^*\}) N(x^0)^d \quad \forall t \in [0, t_{x^0}]$$

hold. Next, using the Identity section 3.1 for $\tilde{x}^0 = x(t_{x^0}; x^0, u_{x_{i^\sharp}^0})$ with $c = \alpha^{-1} > 1$ allows us to repeat the presented line of reasoning to show that the stage cost are scaled with α^d . In addition,

$$x(T_{x_{i^*}^0}; \tilde{x}^0, \Delta_\alpha u_{x_{i^*}^0}) = \underbrace{\Lambda_\alpha x(T_{x_{i^*}^0}; \Lambda_{\alpha^{-1}} \tilde{x}^0, u_{x_{i^*}^0})}_{\in \alpha \mathcal{N}_{c_1, c_2}} \in \alpha^2 \mathcal{N}_{c_1, c_2}.$$

Hence, we get the desired growth condition if we periodically extend the growth function B on (T^*, ∞) by setting

$$B(t) := \sum_{i=0}^{\lfloor t/T^* \rfloor - 1} \alpha^{id} B(T^*) + \alpha^{d\lfloor t/T^* \rfloor} B(t - T^* \lfloor t/T^* \rfloor) \quad \forall t > T^*,$$

which is bounded by $\sum_{i=0}^{\infty} \alpha^{id} B(T^*) = (1 - \alpha^d)^{-1} B(T^*) \in (0, \infty)$. This completes the proof of the first statement. Asymptotic stability then follows from theorem 1. \square

¹Every trajectory $x(\cdot; x_i^0, u_{x_i^0})$ is bounded on $[0, T^*]$. In addition $u_{x_i^0}$ is essentially bounded. Hence, the stage costs are also bounded. Then, continuity of the solution w.r.t. the initial value and continuity of the stage cost imply the assertion.

3.2 Degree of homogeneity not equal to zero

We begin with globally asymptotically null controllable systems with negative degree of homogeneity, i.e. $\tau < 0$. Here, we show semiglobal asymptotic stability of the origin w.r.t. the MPC closed loop, i.e. local asymptotic stability of the origin with an arbitrary but fixed compact set being the domain of attraction, cf. theorem 1. We refer to example 2, which shows the necessity of this restriction for the derivation of cost controllability.

In proposition 3 we extend theorem 2 to homogeneous systems with negative degree.

Proposition 3 (Cost Controllability for $\tau < 0$). *Let the system eq. (1) be globally asymptotically null controllable and (r, s, τ) -homogeneous with $\tau < 0$. In addition, let the stage cost be defined by eq. (9). Then, eq. (3), i.e. cost controllability, holds on any compact set $K \subset \mathbb{R}^n$ with $0 \in \text{int } K$. In particular, given $\delta > 0$, for any sufficiently large $T > 0$ the origin is semiglobally asymptotically stable for the MPC closed loop.*

Proof. Since the set K is compact there exists $c_2 \in (0, \infty)$ such that

$$K \subseteq \{x^0 \in \mathbb{R}^n \mid N(x^0)^d \leq c_2\}$$

holds. We show that Inequality eq. (12) holds for all x^0 contained in this sublevel set. To this end, we start analogously to the verification of theorem 2 with $c_1 \in (0, c_2)$ but define the growth function $B : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ differently. Let \tilde{B} denote the growth function defined analogously as in the proof of theorem 2 and α , $\alpha \in (0, 1)$, be an arbitrarily chosen but fixed contraction parameter. Then, the adapted growth function B is given by

$$B(t) := \tilde{B}(\alpha^\tau t) \quad \text{for all } t \geq 0. \quad (19)$$

Note that $\alpha^\tau > 1$ holds since $\alpha \in (0, 1)$ and $\tau < 0$. Since \tilde{B} is monotonically increasing, $B(t) \geq \tilde{B}(t)$ holds for all $t \geq 0$.²

Next, we show that Inequality eq. (12) holds for all $t \geq 0$ for $x^0 \in \mathcal{N}_{c_1, c_2}$. For given initial state $x^0 \in \mathcal{N}_{c_1, c_2}$ there exists an index $i^\# \in \{1, 2, \dots, k\}$ such that $\tilde{x}^0 := x(T_{x_{i^\#}^0}; x^0, u_{x_{i^\#}^0}) \in \alpha \mathcal{N}_{c_1, c_2}$ holds. In addition, there exists an index $i^* \in \{1, 2, \dots, k\}$ such that $\Lambda_{\alpha^{-1}} \tilde{x}^0 \in \mathcal{B}_{\delta_{x_{i^*}^0}}(x_{i^*}^0)$ holds and the analogon of Identity section 3.1 for $\tau < 0$ is

$$\ell(x(s; \tilde{x}^0, \Delta_{c^{-1} u_{x_{i^*}^0}}(c^{-\tau} \cdot)), \Delta_{c^{-1} u_{x_{i^*}^0}}(c^{-\tau} s)) \stackrel{\text{eq. (10)}}{=} c^{-d} \ell(x(c^{-\tau} s; \Lambda_c \tilde{x}^0, u_{x_{i^*}^0}), u_{x_{i^*}^0}(c^{-\tau} s)).$$

Next, we use the scaling factor $c := \alpha^{-1} > 1$ for \tilde{x}^0 . Note that Equation eq. (18) yields

²This preliminary step is important to uniformly (independent of the initial value) reparametrize the time argument in the former growth bound by the factor α^τ

$\ell^*(\tilde{x}^0) = \alpha^d \ell^*(\Lambda_{\alpha^{-1}} \tilde{x}^0)$. Overall, for $t \leq \alpha^{-\tau} T_{x_{i^*}^0}$, we get

$$\begin{aligned}
\frac{V_t(\tilde{x}^0)}{\ell^*(\tilde{x}^0)} &= \alpha^{-d} \frac{V_t(\tilde{x}^0)}{\ell^*(\Lambda_{\alpha^{-1}} \tilde{x}^0)} \\
&\leq \frac{\alpha^{-d}}{\ell^*(\Lambda_{\alpha^{-1}} \tilde{x}^0)} \int_0^t \underbrace{\ell(x(s; \tilde{x}^0, \Delta_{\alpha} u_{x_{i^*}^0}(\alpha^\tau \cdot)), \Delta_{\alpha} u_{x_{i^*}^0}(\alpha^\tau s))}_{=\ell(x(\alpha^\tau s; \Lambda_{\alpha^{-1}} \tilde{x}^0, u_{x_{i^*}^0}(\alpha^\tau s)), u_{x_{i^*}^0}(\alpha^\tau s))} ds \\
&= \alpha^{-d} \alpha^{-\tau} \frac{\int_0^{\alpha^\tau t} \ell(x(\tilde{s}; \Lambda_{\alpha^{-1}} \tilde{x}^0, u_{x_{i^*}^0}(\tilde{s})), u_{x_{i^*}^0}(\tilde{s})) d\tilde{s}}{\ell^*(\Lambda_{\alpha^{-1}} \tilde{x}^0)} \\
&\leq \alpha^{-d} \alpha^{-\tau} B(\min\{t, \alpha^{-\tau} T^*\}) \leq \alpha^{-d} B(\min\{t, \alpha^{-\tau} T^*\})
\end{aligned}$$

where we have used that $\alpha^\tau > 1$ and $\alpha^{-\tau} < 1$ hold. Furthermore, for a scaling factor $c > 1$ (e.g. α^{-1}), we get

$$\begin{aligned}
x(c^\tau T_{x_{i^*}^0}; \tilde{x}^0, \Delta_{c^{-1}} u_{x_{i^*}^0}(c^{-\tau} \cdot)) &= x(c^\tau T_{x_{i^*}^0}; \Lambda_{c^{-1}} \Lambda_c \tilde{x}^0, \Delta_{c^{-1}} u_{x_{i^*}^0}(c^{-\tau} \cdot)) \\
&\stackrel{\text{eq. (5)}}{=} \underbrace{\Lambda_{c^{-1}} x(T_{x_{i^*}^0}; \Lambda_c \tilde{x}^0, u_{x_{i^*}^0})}_{\in \alpha \mathcal{N}_{c_1, c_2}}
\end{aligned}$$

where $c^\tau < 1$ and $c^{-1} < 1$. This shows that a contraction is attained already at time $\alpha^{-\tau} T_{x_{i^*}^0}$ ($c = \alpha^{-1}$). For general initial value, the index i^* has to be suitably chosen. Hence, an iterative application of this line of reasoning shows the claimed inequality not only for $x^0 \in \alpha^i \mathcal{N}_{c_1, c_2}$, $i \in \mathbb{N} \cup \{0\}$, but indeed for all $x^0 \in K$.

Semiglobal asymptotic stability then follows from theorem 1. \square

For systems with positive degree of homogeneity, we expect global practical asymptotic stability in view of example 2. To this end, an adaptation of [15, Theorem 6.37] to the continuous-time setting seems to be a valid option to infer global practical asymptotic stability from a suitably adapted cost controllability condition and, thus, to prove this conjecture.

4 Systems with globally asymptotically null controllable homogeneous approximation

In this section, we consider systems with homogeneous approximation at the origin in the following sense [2].

Definition 3. Consider two controlled vector field $f, h : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$, where h is homogeneous with parameters $r_i > 0$, $s_j > 0$ and $\tau \in (-\min_i r_i, \infty)$. Then h is called a homogeneous approximation of f near the origin, if the relation

$$f(x, u) = h(x, u) + R(x, u)$$

holds with

$$|R_i(\Lambda_\alpha x, \Delta_\alpha u)| \leq M\alpha^{r_i+\tau+\eta} \quad (20)$$

for all x, u with $\|x\| \leq \rho$, $\|u\| \leq \rho$ and all $\alpha \in (0, 1]$, where ρ, M, η are positive constants.

In case the approximation h is globally asymptotically null controllable, we will establish cost controllability for the system governed by f if the stage cost eq. (9) are employed in the sense of assumption 1, where K is a neighborhood of the origin. As a consequence, by means of theorem 1 we obtain local asymptotic stability of the origin for the MPC closed loop.

Our main tool for deriving these results are homogeneous control Lyapunov functions, whose existence was established in [14]. We first show preliminary results in section 4.1, i.e. in particular robustness of homogeneous Lyapunov functions. Then, we generalize theorem 2 ($\tau = 0$) and its extension proposition 3 ($\tau < 0$) before our findings are illustrated by means of the nonholonomic robot example, a variant of Brockett's famous nonholonomic integrator.

4.1 Robustness of Homogeneous Lyapunov Functions

In this section we assume that approximation h is globally asymptotically null controllable. Then [14, Theorem 3.5] establishes the existence of a homogeneous control Lyapunov function $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$, which

- 1) is continuous on \mathbb{R}^n and Lipschitz on $\mathbb{R}^n \setminus \{0\}$,
- 2) obeys the equality $V(\Lambda_\alpha x) = \alpha^{2k}V(x)$, and
- 3) satisfies the inequality $\min_{v \in \text{co}h(x, W_x)} DV(x; v) \leq -\mu N(x)^\tau V(x)$,

where $k := \min_{i=1, \dots, n} r_i$, $\mu > 0$, $W_x = \Delta_{N(x)}U$ for a suitable compact set $U \subset \mathbb{R}^m$, and $DV(x; v)$ denotes the lower Dini derivative

$$DV(x; v) := \liminf_{t \searrow 0, v' \rightarrow v} \frac{1}{t} (V(x + tv') - V(x)).$$

Next, we show an auxiliary result. Namely, that a suitably scaled version of the homogeneous Lyapunov function V is locally Lipschitz.

Lemma 1. *Let p be defined by $\max_{i=1, \dots, n} r_i$ and let the homogeneous Lyapunov function $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ satisfy Properties 1) and 2). Then, for each compact set $K \subset \mathbb{R}^n$, there is $L > 0$ such that $\hat{V} := V^{p/(2k)}$ is Lipschitz continuous on K with Lipschitz constant L .*

Proof. A straightforward computation yields that \hat{V} satisfies the (in)equalities

$$\hat{V}(\Lambda_\alpha x) = \alpha^p \hat{V}(x) \quad (21)$$

and

$$\min_{v \in \text{co}h(x, W_x)} D\hat{V}(x; v) \leq -\sigma N(x)^\tau \hat{V}(x) \quad (22)$$

with $\sigma := p\mu/(2k) > 0$. Moreover, Property 1) is preserved. Based on this observation, we can prove the improved Lipschitz property of \hat{V} .

Throughout the proof we use the 1-norm $\|\cdot\| = \|\cdot\|_1$. Because of the homogeneity, it suffices to show that \hat{V} is Lipschitz on the compact set $K := \{x \in \mathbb{R}^n \mid N(x) \leq 1\}$. To this end, first observe that the Lipschitz continuity on $\mathbb{R}^n \setminus \{0\}$ implies the existence of a Lipschitz constant $L_1 > 0$ on the set $K_1 := \{x \in \mathbb{R}^n \mid 1/2 \leq N(x) \leq 1\}$. Moreover, by continuity of \hat{V} there exists a constant $C_1 > 0$ such that for all $\hat{x} \in \mathbb{R}^n$ with $N(\hat{x}) = 1$ and all $\hat{y} \in \mathbb{R}^n$ with $N(\hat{y}) \leq 1/2$ the inequality $|\hat{V}(\hat{x}) - \hat{V}(\hat{y})| \leq C_1$ holds. Together with the fact that there is a constant $C_2 > 0$ with $\|\hat{x} - \hat{y}\| \geq C_2$ for all such \hat{x} and \hat{y} , for $L_2 = C_1/C_2$ we obtain $|\hat{V}(\hat{x}) - \hat{V}(\hat{y})| \leq C_1 \leq C_1/C_2 \|\hat{x} - \hat{y}\| = L_2 \|\hat{x} - \hat{y}\|$. This yields that, for all $\hat{x} \in \mathbb{R}^n$ with $N(\hat{x}) = 1$ and all $\hat{y} \in K$, the inequality

$$|\hat{V}(\hat{x}) - \hat{V}(\hat{y})| \leq L \|\hat{x} - \hat{y}\| \quad (23)$$

holds with $L := \max\{L_1, L_2\}$.

Now consider two arbitrary points $x, y \in K$. Without loss of generality assume $N(x) \geq N(y)$. Let $\alpha := N(x) \leq 1$ and define $\hat{x} := \Lambda_\alpha^{-1}x$ and $\hat{y} := \Lambda_\alpha^{-1}y$. Then we obtain

$$\|\hat{x} - \hat{y}\| = \sum_{i=1}^n |\hat{x}_i - \hat{y}_i| = \sum_{i=1}^n \alpha^{-r_i} |x_i - y_i| \leq \alpha^{-p} \sum_{i=1}^n |x_i - y_i| = \alpha^{-p} \|x - y\|.$$

Observing that $N(\hat{x}) = 1$ and $N(\hat{y}) \leq 1$, we can apply eq. (23) in order to conclude

$$\begin{aligned} |\hat{V}(x) - \hat{V}(y)| &= |\hat{V}(\Lambda_\alpha \hat{x}) - \hat{V}(\Lambda_\alpha \hat{y})| = \alpha^p |\hat{V}(\hat{x}) - \hat{V}(\hat{y})| \\ &\leq \alpha^p L \|\hat{x} - \hat{y}\| \leq \alpha^p L \alpha^{-p} \|x - y\| = L \|x - y\|, \end{aligned}$$

which shows the claim. \square

Based on the Lipschitz continuity of the suitably scaled homogeneous Lyapunov function, we can now derive a robustness statement for the Dini derivative of the Lyapunov function along the vector field f that is homogeneously approximated by h .

Proposition 4. *Consider a vector field f with homogeneous approximation h according to definition 3 and a homogeneous control Lyapunov function $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ satisfying Properties 2) and 3). Then, for each $\delta \in (0, 1)$, there exists $\varepsilon > 0$ such that the inequality*

$$\min_{v \in \text{co} f(x, W_x)} DV(x; v) \leq -\delta \sigma N(x)^T V(x). \quad (24)$$

holds for all $x \in \mathbb{R}^n$ with $N(x) \leq \varepsilon$.

Proof. At the beginning of the proof of lemma 1 we made the observation that scaling of the homogeneous Lyapunov function does not change Properties 2) and 3). Analogously, it can be seen that once Inequality eq. (24) is established for $\hat{V} := V^{p/(2k)}$, a straightforward computation shows that it also holds (with $\delta\sigma$ replaced by $\delta\sigma q$) for \hat{V}^q , $q > 0$, and, thus, in particular for V . Hence, it is sufficient to prove Inequality eq. (24)

for \hat{V} . In what follows, we again write V instead of \hat{V} in order to simplify the notation within the remainder of the proof.

From the form of f it follows that for all $\alpha \in (0, 1]$ and each $v_h \in \text{coh}(\Lambda_\alpha x, W_{\Lambda_\alpha x})$ there exists $v \in \text{cof}(\Lambda_\alpha x, W_{\Lambda_\alpha x})$ with $v = v_h + v_R(v_h, x, \alpha)$, where

$$|[v_R(v_h, x, \alpha)]_i| \leq M\alpha^{r_i + \tau + \eta}, \quad (25)$$

provided x is sufficiently close to 0. We choose $\nu > 0$ small enough such that eq. (25) holds whenever $N(x) \leq \nu$. Inequality eq. (25) implies that $v_R(v_h, x, \alpha)$ can be written as

$$v_R(v_h, x, \alpha) = \alpha^{\tau + \eta} \Lambda_\alpha \hat{v}_R$$

for a vector $\hat{v}_R = \hat{v}_R(v_h, x, \alpha) \in \mathbb{R}^n$ with $|\hat{v}_R|_i \leq M$. Any sequence $v' \rightarrow v$ can thus be written as $v' = v'_h + \alpha^{\tau + \eta} \Lambda_\alpha \hat{v}_R$ with $v'_h \rightarrow v_h$. From the Lipschitz continuity of V we obtain

$$\begin{aligned} |V(\Lambda_\alpha x + tv') - V(\Lambda_\alpha x + tv'_h)| &\leq \alpha^p |V(x + t\Lambda_\alpha^{-1}v') - V(x + t\Lambda_\alpha^{-1}v'_h)| \\ &\leq \alpha^p tL \|\Lambda_\alpha^{-1} \alpha^{\tau + \eta} \Lambda_\alpha \hat{v}_R\| = \alpha^{p + \tau + \eta} tL \|\hat{v}_R\|. \end{aligned}$$

For the Dini-derivative this implies

$$\min_{v \in \text{cof}(\Lambda_\alpha x, W_{\Lambda_\alpha x})} DV(\Lambda_\alpha x; v) \leq \min_{v_h \in \text{coh}(\Lambda_\alpha x, W_{\Lambda_\alpha x})} DV(\Lambda_\alpha x; v) + \alpha^{p + \tau + \eta} M_R.$$

with M_R being a bound on $\|\hat{v}_R\|$.

Now let $m_V := \min\{V(x) \mid N(x) = \nu\}$. Then eq. (21) implies

$$\alpha^{p + \tau + \eta} M_R \leq (1 - \delta) \sigma \alpha^{p + \tau} V(x) \nu^\tau$$

for all $x \in \mathbb{R}^n$ with $N(x) = \nu$, whenever $\alpha \leq \alpha_\delta := ((1 - \delta) \sigma m_V \nu^\tau / M_R)^{1/\eta}$. Given $x \in \mathbb{R}^n$ with $N(x) \leq \nu$, setting $\alpha := N(x) / \nu \in (0, 1]$ and $\hat{x} := \Lambda_\alpha^{-1}(x)$ we obtain $N(\hat{x}) = \nu$. Hence, for $\varepsilon = \alpha_\delta \nu$ and $N(x) \leq \varepsilon$ (implying $\alpha \leq \alpha_\delta$) we obtain

$$\begin{aligned} \min_{v \in \text{cof}(x, W_x)} DV(x; v) &= \min_{v \in \text{cof}(\Lambda_\alpha \hat{x}, W_{\Lambda_\alpha \hat{x}})} DV(\Lambda_\alpha \hat{x}; v) \\ &\leq \min_{v_h \in \text{coh}(\Lambda_\alpha \hat{x}, W_{\Lambda_\alpha \hat{x}})} DV(\Lambda_\alpha \hat{x}; v) + \alpha^{p + \tau + \eta} M_R \\ &\leq -\sigma \underbrace{N(\Lambda_\alpha \hat{x})^\tau}_{=\alpha^\tau \nu^\tau} \underbrace{V(\Lambda_\alpha \hat{x})}_{=\alpha^p V(\hat{x})} + \underbrace{\alpha^{p + \tau + \eta} M_R}_{\leq (1 - \delta) \sigma \alpha^{p + \tau} V(\hat{x}) \nu^\tau} \\ &\leq -\delta \sigma \alpha^{p + \tau} \nu^\tau V(\hat{x}) = -\delta \sigma N(\Lambda_\alpha \hat{x})^\tau V(\Lambda_\alpha \hat{x}) = -\delta \sigma N(x)^\tau V(x). \end{aligned}$$

This shows the claim. \square

4.2 Cost controllability via homogeneous control Lyapunov functions

We now show that cost controllability can be concluded from the existence of a homogeneous control Lyapunov function for the homogeneous approximation. To this end, we

use the observation made in the proof of [30, Theorem 1] (based on a similar statement for differential inclusions from [3]), that Inequality eq. (24) implies the existence of a measurable control function $u : [0, t] \rightarrow \mathbb{R}^m$ for which the integral inequality

$$V(x(t, x^0, u)) - V(x^0) \leq \int_0^t -q\delta\sigma N(x(s; x^0, u))^\tau V(x(s, x^0, u)) ds \quad (26)$$

and the inclusion $u(s) \in W_{x(s, x^0, u)}$ hold for all almost all $s \geq 0$. While the construction in [30] is for global control Lyapunov functions, standard Lyapunov function arguments show that it remains valid if the control Lyapunov function is only defined in a neighbourhood of the origin. Of course, in this case eq. (26) will only hold for x^0 from a (possibly different) neighbourhood of the origin.

Theorem 3. *Consider a vector field f with homogeneous approximation h of degree $\tau \leq 0$ according to definition 3. Assume that the homogeneous system with vector field h is globally asymptotically null controllable to the origin. Then there exists a neighborhood \mathcal{N} of the origin such that eq. (3) is satisfied with $K = \mathcal{N}$, i.e. the cost controllability is satisfied for the vector field f and $B_K(t)$ is uniformly bounded. In particular, given $\delta > 0$, for any sufficiently large $T > 0$ the origin is locally asymptotically stable for the MPC closed loop.*

Proof. By [14, Theorem 3.5] and proposition 4 there exists a control Lyapunov function V satisfying Inequality eq. (24) and, by suitably scaling, we have

$$V(\Lambda_\alpha x) = \alpha^d V(x) \quad (27)$$

and the inequality

$$\min_{v \in \text{co}f(x, W_x)} DV(x; v) \leq -\gamma N(x)^\tau V(x) \quad (28)$$

for $\gamma > 0$ in a neighbourhood \mathcal{N} of the origin. From Equation eq. (27) it follows that there exists $C_1 > 0$ with

$$V(x) \leq C_1 \inf_{u \in \mathbb{R}^m} \ell(x, u) \quad (29)$$

for all $x \in \mathbb{R}^n$. Now pick an initial condition x^0 from the neighbourhood \mathcal{N} of the origin on which eq. (26) with $q = d/p$ holds. Then V decreases along the solution which implies that $V(x(t, x^0, u))$ and thus $N(x(t, x^0, u))$ are bounded uniformly in t and x^0 . As a consequence, there exists $C_2 > 0$, independent of x^0 , such that $N(x(t, x^0, u))^\tau \geq C_2$ for all $t \geq 0$ using $\tau \leq 0$.

Then, along the solution in eq. (26), again eq. (27) and the inclusion $u(s) \in W_{x(s, x^0, u)}$ (which implies $N(u(s)) \leq DN(x(s, x^0, u))$ for an appropriate constant $D > 0$) yield the existence of $C_3 > 0$ with

$$\ell(x(s, x^0, u), u(s)) \leq C_3 V(x(s, x^0, u), u(s))$$

for almost all $s \in (0, t)$. Thus, using eq. (26) we obtain

$$\begin{aligned} V_i(x^0) &\leq \int_0^t \ell(x(s, x^0, u), u(s)) ds \leq \int_0^t C_3 V(x(s, x^0, u), u(s)) ds \\ &\leq \frac{C_3}{C_2 \gamma} \left(V(x^0) - V(x(t, x^0, u)) \right) \leq \frac{C_3}{C_2 \gamma} V(x^0) \\ &\stackrel{\text{eq. (29)}}{\leq} \frac{C_1 C_3}{C_2 \gamma} \inf_{u \in \mathbb{R}^m} \ell(x^0, u). \end{aligned}$$

This shows the first claim for $C = C_1 C_3 / (C_2 \gamma)$. Asymptotic stability of the origin for the MPC closed loop then follows from theorem 1. \square

4.3 Example: Nonholonomic Robot

We illustrate our findings by means of the following example.

Example 3. We consider the nonholonomic mobile robot given by

$$\dot{x}(t) = \tilde{f}(x(t), u(t)) := \begin{pmatrix} \cos(x_3(t)) \\ \sin(x_3(t)) \\ 0 \end{pmatrix} u_1(t) + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} u_2(t), \quad (30)$$

which is a control affine system, i.e. $\tilde{f}(x, u) = \tilde{f}_0(x) + \sum_{i=1}^m u_i \tilde{f}_i(x)$ with mappings $\tilde{f}_0, \tilde{f}_1, \dots, \tilde{f}_m : \mathbb{R}^n \rightarrow \mathbb{R}^n$. The system dynamics eq. (30) describe a driftless system since $\tilde{f}_0(x) \equiv 0$.

Let us recall that, as shown by Brockett in [7] (see also [9] and [10, Theorem 11.1]), the control system eq. (30) cannot be locally asymptotically stabilizable by means of continuous feedback laws. We are going to show that the MPC method allows to locally asymptotically stabilize the control system eq. (30).

Firstly, we state the homogeneous approximation and show that the approximation property, i.e. Inequality eq. (20), is satisfied. Note that the parameters r_i and s_j are not unique because the homogeneity condition eq. (4) remains satisfied if they are multiplied by an arbitrary positive scalar.

Proposition 5. Inequality eq. (20) holds for the homogeneous approximation of example 3 at the origin, which is given by

$$\dot{x}(t) = f(x(t), u(t)) := \begin{pmatrix} 1 \\ x_3(t) \\ 0 \end{pmatrix} u_1(t) + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} u_2(t) \quad (31)$$

with $\tau = 0$, $r_1 = r_3 = s_1 = s_2 = 1$, and $r_2 = 2$.

Proof. For $i = 3$, the approximation error is zero. For $i = 1$, we have to show the inequality $|(\cos(\alpha^{r_3} x_3) - 1) \alpha^{r_1} u_1| \leq M_1 \alpha^{r_1 + \tau + m_1}$. Using $r_1 = r_3 = 1$, we have $|\cos(\alpha x_3) -$

$1| \leq \alpha^2 x_3^2/2$. Hence, $|(\cos(\alpha^{r_3} x_3) - 1)\alpha^{r_1} u_1|$ is smaller than $\alpha^3 \rho^3/2$, which shows the assertion for $M_1 := \rho^3/2$ and $\eta_1 \in [0, 2]$. For $i = 2$, we analogously derive the estimate

$$|\sin(\alpha^{r_3} x_3) - \alpha^{r_3} x_3| \alpha^{r_1} |u_1| \leq \alpha^4 |x_3|^3 |u_1|/6 \leq \alpha^4 \rho^4/6$$

and, thus, get the estimate for $M_2 := \rho^4/6$ and $\eta_2 \in [0, 2]$. Overall, the desired property holds with $M := \max\{\rho^3/2, \rho^4/6\}$ and $\eta \in (0, 2]$. \square

For the nonholonomic mobile robot, i.e. example 3, MPC does not work if (purely) quadratic stage cost are used, i.e. $\ell(x, u) := x^\top Qx + u^\top Ru$ with matrices $Q \in \mathbb{R}^{n \times n}$ and $R \in \mathbb{R}^{m \times m}$ as rigorously shown in [26]. Using the proposed stage cost eq. (9) reads for the nonholonomic robot (and its homogeneous approximation at the origin) as

$$\ell(x, u) = x_1^4 + x_2^2 + x_3^4 + u_1^4 + u_2^4, \quad (32)$$

i.e. precisely the stage costs used in [33], for which asymptotic stability of the origin w.r.t. the MPC closed loop holds if a sufficiently large prediction horizon T is employed, cf. [34]. This assertion is, using the results derived in this paper, a direct corollary of the simple calculations presented in proposition 5. Moreover, we conjecture that the observations w.r.t. *essentially using the homogeneous approximation* may significantly facilitate (and improve) the quantitative estimates on the length of the prediction horizon.

Remark 2. *Note that the particular definition of d has not been used so far. This explains, e.g., why the stage cost eq. (32) for the nonholonomic robot may be replaced by*

$$\ell(x, u) = \sum_{i=1}^n q_{x_i} |x_i|^{d/r_i} + \sum_{j=1}^m q_{u_j} |u_j|^{d/s_j}$$

with arbitrary $d \in (0, \infty)$ where q_{x_i} , $i \in \{1, 2, \dots, n\}$, and q_{u_j} , $j \in \{1, 2, \dots, m\}$ are positive weighting factors, see [35, Subsection 2.2] for an example.³

Remark 3. *Other homogeneous approximations can also be used to locally stabilize in small-time the control system eq. (30) by means of periodic time-varying feedback laws as shown in [1].*

5 Conclusions

For nonlinear systems with stabilizable linearization at the origin, MPC *works* (with quadratic stage cost) for sufficiently large prediction horizon (without adding stabilizing terminal costs and/or constraints). We have generalized this methodology to nonlinear systems with globally asymptotically null controllable homogeneous approximation at the desired set point. To this end, we have set up a general framework for rigorously showing cost controllability and, thus, local asymptotic stability of the origin w.r.t. the

³In this reference, Brockett's nonholonomic integrator example was considered. However, this example is (locally) equivalent to the nonholonomic robot (unicycle), see [23, pp. 83-89].

MPC closed loop for systems by designing the stage cost based on the homogeneous approximation.

A major advantage of the presented approach is its simplicity from the user's perspective: calculating the homogeneous approximation, checking (local) asymptotic stabilizability using the already existing results from the literature, and finally running MPC with the stage costs based on the coefficients from the homogeneity definition.

References

- [1] d'Andréa-Novel, B., Jean-Michel Coron, and Wilfrid Perruquetti. Small-time stabilization of nonholonomic or underactuated mechanical systems: the unicycle and the slider examples. *Preprint*, 2019. <https://hal.archives-ouvertes.fr/hal-02140549>.
- [2] Vincent Andrieu, Laurent Praly, and Alessandro Astolfi. Homogeneous approximation, recursive observer design, and output feedback. *SIAM Journal on Control and Optimization*, 47(4):1814–1850, 2008.
- [3] J.-P. Aubin and A. Cellina. *Differential Inclusions*. Springer, 1984.
- [4] Andrea Bacciotti and Lionel Rosier. *Liapunov functions and stability in control theory*. Communications and Control Engineering Series. Springer-Verlag, Berlin, second edition, 2005.
- [5] Sanjay P Bhat and Dennis S Bernstein. Geometric homogeneity with applications to finite-time stability. *Mathematics of Control, Signals and Systems*, 17(2):101–127, 2005.
- [6] Andrea Boccia, Lars Grüne, and Karl Worthmann. Stability and feasibility of state constrained mpc without stabilizing terminal constraints. *Systems & control letters*, 72:14–21, 2014.
- [7] Roger W. Brockett. Asymptotic stability and feedback stabilization. In *Differential geometric control theory (Houghton, Mich., 1982)*, volume 27 of *Progr. Math.*, pages 181–191. Birkhäuser Boston, Boston, MA, 1983.
- [8] Hong Chen and Frank Allgöwer. A quasi-infinite horizon nonlinear model predictive control scheme with guaranteed stability. *Automatica*, 34(10):1205–1217, 1998.
- [9] Jean-Michel Coron. A necessary condition for feedback stabilization. *Systems Control Lett.*, 14(3):227–232, 1990.
- [10] Jean-Michel Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [11] Jean-Michel Coron and Laurent Praly. Adding an integrator for the stabilization problem. *Systems & Control Letters*, 17(2):89–104, 1991.

- [12] Michael G Forbes, Rohit S Patwardhan, Hamza Hamadah, and R Bhushan Gopaluni. Model predictive control in industry: Challenges and opportunities. *IFAC-PapersOnLine*, 48(8):531–538, 2015.
- [13] L. Grüne. Analysis and design of unconstrained nonlinear MPC schemes for finite and infinite dimensional systems. *SIAM Journal on Control and Optimization*, 48:1206–1228, 2009.
- [14] Lars Grüne. Homogeneous state feedback stabilization of homogenous systems. *SIAM Journal on Control and Optimization*, 38(4):1288–1308, 2000.
- [15] Lars Grüne and Jürgen Pannek. Nonlinear model predictive control. In *Nonlinear Model Predictive Control*. Springer, 2017.
- [16] Lars Grüne, Jürgen Pannek, Martin Seehafer, and Karl Worthmann. Analysis of unconstrained nonlinear mpc schemes with time varying control horizon. *SIAM Journal on Control and Optimization*, 48(8):4938–4962, 2010.
- [17] Henry Hermes. Homogeneous feedback controls for homogeneous systems. *Systems & Control Letters*, 24(1):7–11, 1995.
- [18] Matthias Kawski. Stability and nilpotent approximations. In *Proc. 27th Conf. on Decision and Control*, pages 1244–1248. IEEE, 1988.
- [19] Matthias Kawski. Homogeneous stabilizing feedback laws. *Control Theory and advanced technology*, 6(4):497–516, 1990.
- [20] Christopher M Kellett. A compendium of comparison function results. *Mathematics of Control, Signals, and Systems*, 26(3):339–374, 2014.
- [21] B. Kouvaritakis and M. Cannon. *Model Predictive Control*. Springer, Basel, 2016.
- [22] Jay H Lee. Model predictive control: Review of the three decades of development. *International Journal of Control, Automation and Systems*, 9(3):415, 2011.
- [23] Daniel Liberzon. Switching in systems and control, ser. systems & control: Foundations & applications. *Birkhauser*, 2003.
- [24] P Morin and C Samson. Time-varying exponential stabilization of chained form systems based on a backstepping technique. In *Proc. 35th IEEE Conf. Decision and Control*, volume 2, pages 1449–1454, 1996.
- [25] Pascal Morin, Jean-Baptiste Pomet, and Claude Samson. Design of homogeneous time-varying stabilizing control laws for driftless controllable systems via oscillatory approximation of lie brackets in closed loop. *SIAM Journal on Control and Optimization*, 38(1):22–49, 1999.
- [26] Matthias A. Müller and Karl Worthmann. Quadratic costs do not always work in MPC. *Automatica*, 82:269–277, 2017.

- [27] J.B. Rawlings, D.Q. Mayne, and M.M. Diehl. *Model Predictive Control: Theory, Computation, and Design*. Nob Hill Publishing, 2017.
- [28] Marcus Reble and Frank Allgöwer. Unconstrained model predictive control and suboptimality estimates for nonlinear continuous-time systems. *Automatica*, 48(8):1812–1817, 2012.
- [29] Lionel Rosier. Homogeneous lyapunov function for homogeneous continuous vector field. *Systems & Control Letters*, 19(6):467–473, 1992.
- [30] E.D. Sontag and H.J. Sussmann. Nonsmooth control-Lyapunov functions. In *Proc. 34th IEEE Conf. Decision and Control*, New Orleans, LA, USA, 1995.
- [31] S.E. Tuna, M.J. Messina, and A.R. Teel. Shorter horizons for model predictive control. In *Proc. Amer. Control Conf.*, pages 863–868, Minneapolis, MN, USA, 2006.
- [32] K. Worthmann. *Stability Analysis of Unconstrained Receding Horizon Control Schemes*. PhD thesis, University of Bayreuth, 2011.
- [33] K. Worthmann, M.W. Mehrez, M. Zanon, R.G. Gosine, G.K.I. Mann, and M. Diehl. Regulation of differential drive robots using continuous time mpc without stabilizing constraints or costs. *IFAC-PapersOnLine*, 48(23):129–135, 2015.
- [34] Karl Worthmann, Mohamed W Mehrez, Mario Zanon, George KI Mann, Raymond G Gosine, and Moritz Diehl. Model predictive control of nonholonomic mobile robots without stabilizing constraints and costs. *IEEE Transactions on Control Systems Technology*, 24(4):1394–1406, 2016.
- [35] Karl Worthmann, Marcus Reble, Lars Grüne, and Frank Allgöwer. The role of sampling for stability and performance in unconstrained nonlinear model predictive control. *SIAM Journal on Control and Optimization*, 52(1):581–605, 2014.