



HAL
open science

Chitin increases *Mycobacterium ulcerans* growth in acidic environments

Daniel Sanhueza, Christine Chevillon, Nicolas Bouzinbi, Sylvain Godreuil,
Jean-François Guégan

► **To cite this version:**

Daniel Sanhueza, Christine Chevillon, Nicolas Bouzinbi, Sylvain Godreuil, Jean-François Guégan. Chitin increases *Mycobacterium ulcerans* growth in acidic environments. *Microbes and environments / JSME*, 2018, 33 (2), pp.234-237. 10.1264/jsme2.ME17160 . hal-02152436

HAL Id: hal-02152436

<https://hal.science/hal-02152436>

Submitted on 11 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short Communication

Chitin Increases *Mycobacterium ulcerans* Growth in Acidic Environments

DANIEL SANHUEZA^{1*}, CHRISTINE CHEVILLON¹, NICOLAS BOUZINBI^{1,2}, SYLVAIN GODREUIL^{1,2}, and JEAN-FRANÇOIS GUÉGAN^{1,3}

¹MIVEGEC, IRD, CNRS, Université de Montpellier, Montpellier, France; ²Laboratoire de bactériologie, CHU de Montpellier, Université de Montpellier, France; and ³International United Nations Program FutureEarth, OneHealth Core Research Program, Montreal, Canada

(Received October 14, 2017—Accepted April 11, 2018—Published online June 16, 2018)

Species with a chitinous exoskeleton are overrepresented among the aquatic organisms carrying *Mycobacterium ulcerans* (MU) in nature and laboratory experiments have demonstrated the enhancing effects of chitin on the growth of MU. Field surveys identified pH as one of the key parameters delineating the distribution of MU in tropical regions. The present study investigated the relationship between chitin and pH in MU growth. By focusing on pH variations in the field, our results revealed that chitin enhanced MU growth in acidic environments. The present study provides new information on the ecological conditions favoring the development of this mycobacterium in nature.

Key words: *Mycobacterium ulcerans*, environmental mycobacteria, experimental test, ecological niche, neglected tropical disease

Mycobacterium ulcerans (MU) is the etiological agent of Buruli ulcer (BU), a neglected human infectious disease reported in more than 30 tropical and subtropical countries that is responsible for skin ulcers and/or disabilities in the absence of treatment (1). This mycobacterium has been associated with a large diversity of aquatic host taxa (2, 7, 8, 13, 16, 23–25), and recent field observations identified several environmental conditions that appear to be associated with a shift in the distribution of MU in tropical freshwater ecosystems. These key conditions involve not only the composition and diversity of the assemblages of aquatic species that are locally present (9, 10, 12, 21), but also physicochemical properties, such as the water flow speed, quantity of dissolved oxygen, and pH (8, 9, 11, 18). If field data remain scarce, it is important to note that the overlaps in pH values recorded across MU-positive sites from different tropical regions (8, 15) range between 4.5 and 7.5 (Supplementary file 1). This finding is consistent with laboratory research showing MU optimal growth *in vitro* for pH values ranging between 5.4 and 7.2 (19).

Since chitin is a major component of the cuticle of arthropods and arthropods are overrepresented in MU-associated taxa in BU endemic areas (9, 14, 16), the hypothesis that chitin may directly play a role in the development of MU was proposed. An experimental test recently confirmed this hypothesis and the availability of chitin was found to promote MU growth *in vitro* more strongly than other modifications in the concentrations of inorganic or organic elements commonly observed in the field (22).

The present study attempted to provide a clearer understanding of the biology of MU by (i) using pH variations recorded in BU endemic areas to investigate the impact of pH modifications on MU growth under controlled laboratory conditions (Supplementary file 1), and (ii) testing whether the impact of pH on MU growth is associated with the presence/absence of chitin in culture medium.

The present study did not involve any human or animal participants, only an *in vitro* culture of the 1G897 laboratory strain isolated from human biopsies in 1991 (6). We cultured the 1G897 strain at 30°C in tubes containing 27 mL of 7H9 medium and 3 mL of the growth supplement OADC (Oleic, Albumin, Dextrose, and Catalase) (Sigma-Aldrich, St Louis, MO, USA). This standard medium culture defines 6.7 as the control pH value. We also used seven experimental pH values (4.5, 5.0, 5.5, 6.0, 6.5, 7.0, and 7.5) in order to cover the field variations observed in BU endemic areas (9, 16) (Supplementary file 1). These experimental pH values were obtained by adding a 1:10 v/v dilution of acid phosphoric solution or sodium hydroxide to 7H9 growth medium where appropriate. Under each pH condition, we tested the effects of the presence of chitin by starting experiments with 10 tubes per pH condition: 5 replicates without chitin and 5 replicates with chitin-supplemented culture medium at a final chitin concentration of 0.5 g 100 mL⁻¹. Commercial alpha chitin extracted from shrimp shells (Sigma-Aldrich) was used in the present study.

We monitored pH and the abundance of MU at the launching date ($t=0$) and at $t=25$, 35, and 50 d post-inoculation. In order to modify the experimental volume as little as possible and avoid any pH changes caused by measurements, pH values were recorded with a micro-electrode (Thermo Scientific) adapted to small volumes and by only collecting 500 µL at each measurement. No changes in pH values were observed in any tubes throughout the experiment. DNA was extracted using 100 µL of each bacterium culture by employing a DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) according to the “bacteria protocol” recommended by the manufacturer. MU quantification was based on the Real-Time PCR protocols described previously (22). The accuracy of these protocols was tested with external standard curves obtained from serial dilutions of MU (strain 1G897) DNA over five logs (from 10⁶ to 10² units mL⁻¹). This allowed the mean inoculums added to each experimental tube at the launching date of the experiment ($t=0$) to be estimated as ~3,000 MU cells per tube.

* Corresponding author. E-mail: daniel.sanhueza@ird.fr;
Tel: +33-467-41-62-05; Fax: +33-467-41-63-30.

We defined the estimated number of MU cells per tube as the response variable and analyzed its variation using general linear mixed model (GLMM) procedures (5) in which replicates were fit as random factors, while pH values, chitin concentrations, and time were fixed factors. The maximal model, including all interactions among fixed factors, was simplified by sequentially eliminating non-significant terms, starting by the highest order of interactions among factors, to finally establish a minimal model (5). The significance of each term was established using either a likelihood ratio (LLR) test, which is approximately distributed as a χ^2 distribution (4), or an *F* test (5). Analyses and graphics were performed with the *R* statistical software, version 3.3.0 (20) and the *R* packages *MASS*, *nlme*, *lme4*, *beeswarm*, *RColorBrewer*, and *ggplot2*, or with the software GraphPad prism 5, version 5.02.

The experimental set-up confirmed that MU exhibited the ability to survive and grow over all the pH ranges observed in BU endemic regions, with the extremes (pH=4.5 or 7.5) being less suitable for its development *in vitro* (Fig. 1). The minimal GLMM evidenced independently significant effects of the factors ‘Time’ (*F* value=13.261, *P* value<0.0001) and ‘Chitin’ concentrations (*F* value=4.679, *P* value=0.0379) as well as of the interaction between ‘pH’ and ‘Time’ (*F* value=1.845, *P* value=0.0211) or ‘Chitin’ and ‘Time’ (*F* value=9.861, *P* value=<0.0001; Supplementary file 2). In contrast, the interaction between ‘pH’ and ‘Chitin’ concentrations was not significant (*F* value=1.980, *P* value=0.088; Supplementary file 2). Nevertheless, MU grew faster in chitin-supplemented media with pH≤6.5 than in chitin-free media with 6.7≤pH≤7.5; if the maximal abundance of MU was similar across

these experimental conditions (*F* value=3.587, *P* value=0.22), the maximum was reached within 25 to 35 d in chitin-supplemented media with pH≤6.5, but only in 50 d in chitin-free media with pH≥6.7 (Fig. 1).

In order to clarify the relationship between chitin concentrations and pH conditions, we pooled pH experimental conditions into three classes: ‘low’ regroups the three lowest experimental values (4.5, 5.0, and 5.5), ‘middle’ regroups the two intermediate values (6.0 and 6.5), and ‘high’ joins the two remaining experimental values (7.0 and 7.5). This led to the same results as those obtained when considering experimental pH values individually (LLR=33.202, *P* value=0.13). The effects of ‘Chitin’ concentrations on variations in the estimates of bacterial cells across replicates and experimental conditions remained significant in the low and middle groups (*i.e.*, for 4.5≤pH≤6.5; *F* value=4.394, *P* value=0.0055; Supplementary Material 3), but not in the high group (*i.e.*, for pH>6.5; *F* value=2.758, *P* value=0.0747). Interestingly, this may be correlated with other differences among experimental conditions: MU kinetic growth stopped after 25 to 35 d in chitin-supplemented media for pH values in the low or middle groups, but otherwise remained constant throughout the 50 d of monitoring (Fig. 2).

Several laboratory experiments and/or field surveys have attempted to elucidate the impact of environmental variables on MU growth *in vitro* or its presence in natural ecosystems. However, the present study was the first to experimentally investigate the impact of pH field variations reported in BU endemic areas on MU growth *in vitro*. Since the addition of chitin to standard culture medium was previously reported to stimulate MU growth *in vitro* at pH=6.7 (22), the present study attempted to elucidate the mechanisms by which the growth stimulating effects of chitin interplay with the effects on MU growth caused by variations in pH.

The present results on the individual effects of pH values are consistent with previous findings on the optimal conditions for *in vitro* MU cultivation (19). The optimal conditions for *in vitro* MU cultivation correspond to the present ‘control’ conditions under which the MU doubling time was 3.5 d (15). The difference in MU growth between optimal pH (6.7 in the control) and the experimental value of 6.5 may be attributed to H₃PO₄ being added to standard culture medium in order to reach more acidic experimental values. Some bacteria are known to use PO₄³⁻ as a phosphate source, which is essential for the biosynthesis of nucleic acids (12); however, this has not yet been examined in MU. It is important to note that we previously demonstrated that the concentration of PO₄³⁻ did not significantly affect MU growth (22).

The present results revealed the pH-mediated effects of chitin on MU growth, namely, the enhanced development of MU cells in acidic environments in the presence of chitin. Based on the presence of three genes coding for chitinases (MUL_2681, MUL_2210, MUL_0371) in the MU genome, one likely explanation for these complex MU kinetics in chitin-supplemented culture media is a decrease over time in the availability of chitin. Alternatively, since growth media was not renewed throughout the experiment, MU growth may have stopped after ~25 to 35 d in chitin-supplemented media because of the accumulation of sub-products resulting from the MU metabolic use of chitin in the early stages of experi-




Fig. 1. Mean estimates in numbers of bacterial cells across pH conditions in the presence (panel A) or absence (panel B) of chitin. The experiment started on day 0; pH=6.7 in the control.




Fig. 2. Interactions between pH conditions and the presence/absence of chitin on MU kinetic growth. pH values were pooled into three groups for clarity: (A) low ($4.5 \leq \text{pH} \leq 5.5$), (B) medium ($6.0 \leq \text{pH} \leq 6.5$), and (C) high ($7.0 \leq \text{pH} \leq 7.5$).

mentation. Further research, including the monitoring of chitin availability during MU growth, appears to be necessary in order to clarify the exact mechanisms underlying the relationship among chitin, pH, and MU, as revealed in the present study.

Because we have found no statistical effect of this solution on MU growth, it seems plausible that acid phosphoric is not used by MU as a phosphate source, thus it may affect MU growth indirectly e.g. by changing the available concentrations and compositions of chitin-derived nutrients in culture

medium. Although this study did not confirm this hypothesis in a culture media with and without H_3PO_4 , further investigations will clarify the effects of PO_4^{-3} on the physical and chemical states of chitin in media.

If the initial effects of chitin supplementation on MU growth is clarified (e.g., fixation support, nutrients used for MU metabolism, or both) (22), it is noteworthy that MU is not the only environmental bacterium sensitive to the presence of chitin. The presence of this polysaccharide accounts for approximately 40% of the variations observed in the abundance of *Vibrio cholerae* in natural environments (17). Moreover, longitudinal field surveys performed in BU endemic areas already demonstrated the co-occurrence of temporal seasonality (between dry and rainy seasons) and spatial variations in pH and the abundance of MU with fluctuations in the abundance of aquatic arthropods, and, thus, in chitin availability (8, 9, 14).

The present results, which were obtained from a rather simple experimental design, provide some explanations from these complex field observations: seasonal variations in pH and species assemblages within aquatic ecosystems may account for most of the seasonal variations in MU abundance, as exemplified by the relationship between low pH values and chitin availability. Furthermore, chitin is known for its antimicrobial effects against many MU competitors, including bacteria, yeast, and fungi (3). Since many of these competitors grow slightly faster than MU, the stimulating effects of chitin on the growth of MU at low pH (i.e., under conditions that are not suitable for the growth of MU in the absence of chitin, Fig. 1) combined with the adverse impact of chitin on many MU competitors suggest that the presence of chitin in nature may increase its growth rate, thereby indirectly increasing the competitive ability of MU towards co-existing microorganisms under these environmental conditions.

To conclude, the present study elucidated the roles of pH and chitin in MU cell growth *in vitro* and also demonstrated for the first time a strong relationship between chitin and pH, indicating that the enhancing effects of chitin on the growth of MU are the strongest in more acidic environments. This study provides new information on the multifactorial process playing a role in the development of MU, and gives a novel insight into the ecological niche of this environmentally persistent mycobacterium.

Acknowledgements

We are grateful to Pr. Françoise Portaels for her insightful comments on earlier drafts. Financial support was provided by funding from Centre National de la Recherche Scientifique (CNRS) and Institut de Recherche pour le Développement (IRD) as well as from 'an Investissement d'Avenir' grant from the Agence Nationale de la Recherche (CEBA ANR-10-LABX-2501) and from a PhD thesis and postdoctoral fellowship from BECAS-Chile (DS) (CONICYT PAI/INDUSTRIA 79090016).

References

- Asiedu, K., R. Scherpier, M. Raviglione (ed.). 2000. Buruli Ulcer: *Mycobacterium Ulcerans Infection*. World Health Organization, Geneva, Switzerland.

2. Benbow, M.E., H. Williamson, R. Kimbirauskas, M.D. McIntosh, R. Kolar, C. Quaye, F. Akpabey, D. Boakye, P. Small, and R. Merrit. 2008. Aquatic invertebrates as unlikely vectors of Buruli ulcer disease. *Emerg. Infect. Dis.* 14:1247–1254.
3. Benhabiles, M.S., R. Salah, H. Lounici, N. Drouiche, M.F.A. Goosen, and N. Mameri. 2012. Antibacterial activity of chitin, chitosan and its oligomers prepared from shrimp shell waste. *Food Hydrocolloids* 29:48–56.
4. Bolker, B.M. 2008. *Ecological models and data* in R. Princeton University Press, Princeton.
5. Crawley, M.J. 2007. *The R Book*, 15th ed. Wiley, London.
6. De Gentile, P.L., C. Mahaza, F. Rolland, B. Carbone, J.L. Verret, and D. Chabasse. 1992. Cutaneous ulcer from *Mycobacterium ulcerans*. A propos of 1 case in French Guyana. *Bull. Soc. Pathol. Exot. Ses Fil.* 85:212–214.
7. Fyfe, J.M., C.J. Lavender, K. Handasyde, *et al.* 2010. A major role for mammals in the ecology of *Mycobacterium ulcerans*. *PLoS Neglected Trop. Dis.* 4:e791.
8. Garchitorena, A., B. Roche, R. Kamgang, *et al.* 2014. *Mycobacterium ulcerans* ecological dynamics and its association with freshwater ecosystems and aquatic communities: Results from a 12-month environmental survey in Cameroon. *PLoS Neglected Trop. Dis.* 8:e2879.
9. Garchitorena, A., J.F. Guégan, L. Léger, S. Eyangoh, L. Marsollier, and B. Roche. 2015. *Mycobacterium ulcerans* dynamics in aquatic ecosystems are driven by a complex interplay of abiotic and biotic factors. *eLife* 4:e07616.
10. Garchitorena, A., C.N. Ngonghala, G. Texier, J. Landier, S. Eyangoh, M.H. Bonds, J.F. Guégan, and B. Roche. 2015. Environmental transmission of *Mycobacterium ulcerans* drives dynamics of Buruli ulcer in endemic regions of Cameroon. *Sci. Rep.* 5:18055.
11. Garcia-Peña, G.E., A. Garchitorena, K. Carolan, E. Canard, A.H. Prieur-Richard, G. Suzan, J. Mills, B. Roche, and J.F. Guégan. 2016. Niche-based host extinction increases prevalence of an environmentally acquired pathogen. *Oikos* 125:1508–1515.
12. Kulakovskaya, T.V., L.P. Lichtko, and L.P. Ryazanova. 2014. Diversity of phosphorus reserves in microorganisms. *Biochemistry (Moscow)* 54:385–412.
13. Lavender, C.J., J.A. Fyfe, J. Azuolas, K. Brown, R.N. Evans, L.R. Ray, and P.D. Johnson. 2011. Risk of Buruli ulcer and detection of *Mycobacterium ulcerans* in mosquitoes in southeastern Australia. *PLoS Neglected Trop. Dis.* 5:e1305.
14. Marion, E., S. Eyangoh, E. Yeramian, *et al.* 2010. Seasonal and regional dynamics of *M. ulcerans* transmission in environmental context: deciphering the role of water bugs as hosts and vectors. *PLoS Neglected Trop. Dis.* 4:e731.
15. Marsollier, L., N. Honoré, P. Legras, A.L. Manceau, H. Kouakou, B. Carbone, and S.T. Cole. 2003. Isolation of three *Mycobacterium ulcerans* strains resistant to rifampin after experimental chemotherapy of mice. *Antimicrob. Agents Chemother.* 47:1228–1232.
16. Morris, A., R. Gozlan, E. Marion, L. Marsollier, D. Andreou, D. Sanhueza, R. Ruffine, P. Couppié, and J.F. Guégan. 2014. First detection of *Mycobacterium ulcerans* DNA in environmental samples from South America. *PLoS Neglected Trop. Dis.* 8:10.
17. Neogi, S.B., M.S. Islam, G.B. Nair, S. Yamasaki, and R.J. Lara. 2012. Occurrence and distribution of plankton-associated and free-living toxigenic *Vibrio cholerae* in a tropical estuary of a cholera endemic zone. *Wetlands Ecol. Manage.* 20:271–285.
18. Pileggi, S.M., H. Jordan, J.A. Clennon, *et al.* 2017. Landscape and environmental influences on *Mycobacterium ulcerans* distribution among aquatic sites in Ghana. *PLoS One* 12:e0176375.
19. Portaels, F., and S.R. Pattyn. 1982. Growth of mycobacteria in relation to the pH of the medium. *Ann. Microbiol. (Paris)* 133:213–221.
20. R Core Team. 2013. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
21. Roche, B., M.E. Benbow, R. Merritt, R. Kimbirauskas, M. McIntosh, P.L.C. Small, H. Williamson, and J.F. Guégan. 2013. Identifying the Achilles' heel of multi-host pathogens: The concept of keystone "host" species illustrated by *Mycobacterium ulcerans* transmission. *Environ. Res. Lett.* 8:45009.
22. Sanhueza, D., C. Chevillon, R. Colwell, J. Babonneau, E. Marion, L. Marsollier, and J.F. Guégan. 2016. Chitin promotes *Mycobacterium ulcerans* growth. *FEMS Microbiol. Ecol.* 92:1–6.
23. Van Ravensway, J., M.E. Benbow, A. Tsonis, S.J. Pierce, L.P. Campbell, J.A.M. Fyfe, J.A. Hayman, P.D.R. Johnson, J.R. Wallace, and J. Qi. 2012. Climate and landscape factors associated with Buruli ulcer incidence in Victoria, Australia. *PLoS One* 7:1–11.
24. Wallace, J.R., M.C. Gordon, L. Hartsell, L. Mosi, E. Benbow, R.W. Merrit, and P.L.C. Small. 2010. Interaction of *Mycobacterium ulcerans* with mosquito species: Implications for transmission and trophic relationships. *Appl. Environ. Microbiol.* 76:6215–6222.
25. Williamson, H.R., M.E. Benbow, K.D. Nguyen, *et al.* 2008. Distribution of *Mycobacterium ulcerans* in Buruli ulcer endemic and non-endemic aquatic sites in Ghana. *PLoS Neglected Trop. Dis.* 2:e205.