

Synthesis of Ruthenium Tris-Diimine Photosensitizers Substituted by Four Methylphosphonate Anchoring Groups for Dye-Sensitized Photoelectrochemical Cell Applications

Nicolas Queyriaux, Emmanouil Giannoudis, Jean-François Lefebvre, Vincent Artero, Murielle Chavarot-Kerlidou

▶ To cite this version:

Nicolas Queyriaux, Emmanouil Giannoudis, Jean-François Lefebvre, Vincent Artero, Murielle Chavarot-Kerlidou. Synthesis of Ruthenium Tris-Diimine Photosensitizers Substituted by Four Methylphosphonate Anchoring Groups for Dye-Sensitized Photoelectrochemical Cell Applications. European Journal of Inorganic Chemistry, 2019, 2019 (15 Spécial Issue), pp.2154-2161. 10.1002/ejic.201900151. hal-02152654

HAL Id: hal-02152654 https://hal.science/hal-02152654

Submitted on 18 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis of ruthenium tris-diimine photosensitizers substituted by four methylphosphonate anchoring groups for dye-sensitized photoelectrochemical cell applications

Nicolas Queyriaux,^[a] Emmanouil Giannoudis,^[a] Jean-François Lefebvre,^[a,b] Vincent Artero,^[a] Murielle Chavarot-Kerlidou^{*[a]} ^[a] Univ. Grenoble Alpes, CNRS, CEA, LCBM, 38000 Grenoble, France. ^[b] Univ. Grenoble Alpes, CNRS, DPM, 38000 Grenoble, France. e-mail: murielle.chavarot-kerlidou@cea.fr

Keywords: Ruthenium, photosensitizers, anchoring groups, photoelectrochemistry

Abstract

The design and synthesis of ruthenium tris-diimine photosensitizers appropriately functionalized to be (*i*) anchored onto transparent conductive oxides (TCO) and (*ii*) covalently coupled with a water-splitting catalyst represents an important target for solar fuel production in dye-sensitized photoelectrochemical cells (DS-PECs). In this study, two different synthetic routes to prepare heteroleptic [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(N^N)](PF₆)₂ complexes are evaluated, the scope and limitations of the organometallic pathway involving half-sandwich η^{6} -arene ruthenium complexes as synthetic intermediates being especially studied. The spectroscopic and electrochemical characterizations of a series of novel structures varying by the nature of the third diimine N^N ligand are reported.

Introduction

One promising chemical approach to artificial photosynthesis relies on the development of tandem dye-sensitized photoelectrochemical cells (DS-PECs) for solar fuel production.^[1-4] Tandem DS-PECs combine two photoelectrodes, (*i*) the photoanode where the light-driven oxidation of water occurs to supply (*ii*) the photocathode with the required amount of electrons and protons to drive the reduction of interest. A major requirement for this approach to be effective is a stable grafting of the molecular photosensitizer/catalyst assemblies onto suitable transparent conductive oxides (TCO). The most commonly employed TCOs are *n*-type TiO₂ at the photoanode and *p*-type NiO at the photocathode; similar grafting strategies

have nevertheless been applied at both electrodes, drawn on the experience gained from the dye-sensitized solar cell (DSSC) technology.^[5] In that respect, the covalent grafting of carboxylic acid-substituted photosensitizers has been widely employed. Considering the specific aqueous conditions required for the DS-PECs, it is however relevant to focus on more stable anchoring systems, particularly with respect to hydrolysis. An improved stability of the grafting was observed for phosphonic acids^[6-8] covalently anchored onto oxide films compared to carboxylic acids, especially at low pH (\leq 5), making them anchoring groups of choice for DS-PEC applications.^[9, 10] In a previous study,^[11] we established that the number of phosphonate anchoring groups present on the Ru photosensitizer significantly improves the grafting efficiency on nickel oxide films, the surface concentration with four phosphonate anchoring groups being roughly twice higher than the one determined with only two anchoring groups. In addition, insertion of a methylene spacer between the bipyridine (bpy) ligand and the phosphonate group electronically insulates the anchors from the complex, which could help stabilizing the excess electron density on the third diimine ligand present in the Ru coordination sphere, thus away from the surface.^[11, 12] This warrants a better hole injection efficiency in the context of photocathode applications. Thus, bis-heteroleptic ruthenium tris-diimine photosensitizers bearing four methylphosphonate anchoring groups represent appealing targets for DS-PEC applications. Although synthetic routes were first reported in the 90's, an easy access to these complexes remains challenging due to low-yield procedures and difficult purification steps. The most straightforward route relies on the preparation of the *cis*-Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl₂ intermediate,^[13-17] whereas a second approach exploits the properties of organometallic half-sandwich η^6 -arene ruthenium (II) complexes.^[18-20] The former procedure, called "traditional approach" below, is similar to the one previously developed for the synthesis of *cis*-Ru(bpy)₂Cl₂, however with a yield not exceeding 32% for isolated *cis*-Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl₂.^[14] The organometallic route was first employed in 2010 by Bignozzi and coworkers,^[18] and later reexamined by the group of T.J. Meyer,^[19, 20] in particular to study a novel photosensitizer-catalyst assembly anchored to TiO₂.^[19]

In this study, we evaluate these two procedures for the synthesis of $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(N^N)](PF_6)_2$ complexes (**RuP**₄^{OEt}-**N**^N), N^N being a third diimine ligand different from bpy, if possible functionalized for an ultimate coupling with a catalytic center. The scope and limitations of the organometallic pathway involving half-sandwich η^6 -arene ruthenium complexes as synthetic intermediates is especially studied. The spectroscopic and

electrochemical characterizations of all novel structures are also reported and compared to the reference $[Ru(bpy)_2(N^N)](PF_6)_2$ complexes.

Results and discussion

Previous studies from the literature focused on the preparation of complexes bearing four methylphosphonic acid anchoring groups, the esters being most of the time hydrolyzed before the last purification step^[13, 14, 18, 20] to make it easier or the phosphonic acid ligand being employed at the beginning of the procedure.^[19] In this study, we decided to isolate, purify and characterize the **RuP4^{OEt-N^N}** complexes under their phosphonate ester form, in order to warrant good solubility in organic media allowing further synthetic steps on the N^N ligand. In addition to the reference 2,2'-bipyridine (bpy) ligand, we selected 1,10-phenantroline (phen) and phen-substituted derivatives, namely 5-amino-1,10-phenantroline (phenamine), dipyrido[3,2-a:2',3'-c]phenazine (dppz), 1,10-phenantroline-5,6-dione (phendione) and 2-(4-ethynylphenyl)-1H-imidazo[4,5-f][1,10]phenanthroline (EPIP) (Figure 1). The latter allows further functionalization of the **RuP4^{OEt-N^N}** complex by copper-catalyzed azide-alkyne cycloaddition (CuAAC).^[21]



Figure 1. Schematic representation of the two synthetic routes to access the **RuP**₄^{OEt}-**N**^**N** complexes and structure of the N^N ligands employed in this study.

The major difference between the two reported procedures lies in the sequential chelation order of the two 4,4'-(CH₂PO₃Et₂)₂-bpy and N^N to the Ru core, occurring at the last step for N^N in the traditional approach against the first step in the organometallic route (Figure 1). We initially attempted to prepare our functionalized complexes by the former procedure in order to take advantage of the introduction of the functionalized ligand at the last step of the process. However, in our hands, the required *cis*-Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl₂ intermediate was isolated pure in very low yields (10-20 %, using either RuCl₃ or Ru(DMSO)₄Cl₂ as

precursors), which was a strong limitation, as previously mentioned by T.J. Meyer.^[20] We thus decided to evaluate the scope of the organometallic route in order to extend it to substituted N^N ligands.

Preparation of the different organometallic $[(\eta^6-arene)Ru(N^{N})Cl](PF_6)$ and $[(\eta^6-arene)Ru(N^{N})(OTf)](OTf)$ precursors (Ar = benzene or p-cymene). The organometallic (η^6 -arene) ruthenium complexes $[(\eta^6\text{-arene})Ru(N^{N})Cl](PF_6)$ were readily prepared by reacting the dinuclear $[(\eta^6\text{-arene})RuCl_2]_2$ precursor (benzene or p-cymene) with two equivalents of substituted diimine N^N ligands under classical thermal conditions. Protection of the EPIP ligand by a trimethylsilyl group (TMS) proved to be mandatory to avoid alkyne coordination to the Ru center or formation of a Ru-alkynyl^[22] complexes. Overall, the $[(\eta^6\text{-arene})Ru(N^{N})Cl](PF_6)$ complexes were isolated as PF₆ salts after anion metathesis, with \geq 90 % yields.

These complexes being relatively kinetically inert, substitution of the chloride ligand for a triflate (OTf) one was reported to facilitate thermal removal of the arene ligand and the coordination of the two 4,4'-(CH₂PO₃Et₂)₂-bpy.^[19, 20] The [(η^6 -arene)Ru(N^N)(OTf)](OTf) derivatives were prepared by reacting [(η^6 -arene)Ru(N^N)Cl](PF₆) with triflic acid overnight at room temperature; complexes with bpy, phen or phenamine ligands were isolated with yields > 90 %. We however observed that alkyne hydration occurred for [(η^6 -benzene)Ru(TMS-EPIP)Cl](PF₆) and an alternative procedure relying on the use of AgOTf to displace the chloride ligand was unsuccessful. Isolation of the triflate complex also failed for [(η^6 -arene)Ru(phendione)Cl](PF₆) (either benzene or *p*-cymene). In the following, the TMS-EPIP and the phendione complexes were tested under their chloride form for the preparation of **RuP4^{OEt}-N^N** by the organometallic route. The [(η^6 -arene)Ru(dppz)Cl](PF₆) complex proved to be insoluble in most organic solvents (except for DMSO), preventing the preparation of [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(dppz)](PF₆)₂ by this approach.

Comparison of the thermal and microwave conditions on the RuP_4^{OEt} -bpy model complex. In order to better understand and to optimize the synthesis of the different diethylmethylphosphonate RuP_4^{OEt} -N^N complexes, we first studied various experimental conditions on the simplest case of the series, RuP_4^{OEt} -bpy. The reaction in refluxing ethanol between two equivalents of 4,4'-(CH₂PO₃Et₂)₂-bpy and four different organometallic precursors varying by the nature of either the arene (benzene or *p*-cymene) or the anionic ligand (OTf or Cl) was monitored by UV-vis absorption spectroscopy (Figure 2). The wavelength of 459 nm was selected to detect the formation of $\mathbf{RuP4^{OEt}}$ -bpy (MLCT transition)^[11] and to determine its conversion yield. As expected, the triflate [(η^{6} - arene)Ru(bpy)(OTf)](OTf) complexes are more reactive than their chloride counterparts with 30-35 % conversion for the former versus < 10 % for the latter. However, the conversion started to plateau after roughly 6 hours of reflux; this observation was confirmed by reacting the triflate precursors overnight at 90°C in a H₂O/EtOH mixture: isolated yields in $\mathbf{RuP4^{OEt}}$ -bpy didn't exceed 35 % (Table 1).



Figure 2. UV-vis monitoring (Abs @ $\lambda = 459$ nm) of the reaction between [(η^{6} -arene)Ru(bpy)L](PF₆) (arene = benzene or *p*-cymene; L = Cl or OTf) and 2 equivalents of 4,4'-(CH₂PO₃Et₂)₂-bpy ligand in refluxing ethanol. Right: Conversions in **RuP**4^{OEt}-bpy calculated from the absorbance.

Table 1.	Thermal	activation:	Isolated	yields	(after	purification	by	chromatography	on	silica
gel – see	experime	ntal section)) in RuP	4 ^{0Et} -bp	y for a	lifferent read	ctio	n conditions.		

Arene	L	Solvent, T, overnight	4,4'-(CH2PO3Et2)2-bpy	RuP4 ^{OEt} -bpy
	Cl	2-MeOEtOH/H ₂ O	2 eq	40 %
benzene	CI	(4 :1), 120°C	2.3 eq	58 %
	OTf	EtOH/H ₂ O (4 :1), 90°C	2 eq	35 %
n cymono	Cl	2-MeOEtOH/H ₂ O (4 :1), 120°C	2 eq	< 5 %
<i>p</i> -cymene	OTf	EtOH/H ₂ O (4 :1), 90°C	2 eq	33 %

The complex $[(\eta^6\text{-benzene})\text{Ru}(\text{bpy})\text{Cl}](\text{PF}_6)$ displays a very modest (6% conversion) but still slightly higher reactivity than $[(\eta^6\text{-}p\text{-}\text{cymene})\text{Ru}(\text{bpy})\text{Cl}](\text{PF}_6)$ (< 2% conversion) (Figure 2). Taking into consideration that triflate complexes couldn't be prepared for two ligands of the series (TMS-EPIP and phendione), we decided to explore other reaction conditions in order to increase the reactivity of $[(\eta^6\text{-benzene})\text{Ru}(\text{bpy})\text{Cl}](\text{PF}_6)$. This complex being kinetically inert, higher temperatures are required for its thermal activation compared to its triflate counterpart.

We thus switched to aqueous/organic solvent mixtures, and replaced EtOH by 2methoxyethanol (2-MeOEtOH). Results are summarized in Table 1. **RuP4**^{OEt}-**bpy** was isolated in 40 % and < 5 % yield, respectively, by refluxing overnight 2 equivalents of 4,4'-(CH₂PO₃Et₂)₂-bpy with either [(η^6 -benzene)Ru(bpy)Cl](PF₆) or [(η^6 -*p*cymene)Ru(bpy)Cl](PF₆) in a 2-MeOEtOH/H₂O 4:1 (v:v) solvent mixture; this result is in agreement with the difference of reactivity observed for these two precursors during the course of the UV-vis monitored experiment in EtOH. Increasing the amount of 4,4'-(CH₂PO₃Et₂)₂-bpy from 2 to 2.3 equivalents allowed to prepare **RuP4**^{OEt}-**bpy** from [(η^6 benzene)Ru(bpy)Cl](PF₆) with a satisfactory yield of 58 % after purification.

Microwave (MW) activation also represents an interesting alternative to thermal activation, as previously described by group of T. J. Meyer for the synthesis of **RuP4^{OH}-bpy** from $[(\eta^{6}-benzene)Ru(bpy)(OTf)](OTf).^{[20]}$ This procedure is appealing as it proceeds faster and with a lower energy consumption, comparatively to the thermal activation.^[23, 24] Using the previously reported experimental conditions,^[20] **RuP4^{OEt}-bpy** was isolated in 58 and 60 % yield, using $[(\eta^{6}-benzene)Ru(bpy)(OTf)](OTf)$ and $[(\eta^{6}-p-cymene)Ru(bpy)(OTf)](OTf)$ respectively, which is very close to the 62 % reported yield.^[20] We then examined whether such conditions could increase the reactivity of $[(\eta^{6}-aene)Ru(N^{N})Cl](PF_{6})$ precursors (Table 2). Under the conditions applied for the triflate precursors (EtOH, 150°C, 20 min), **RuP4^{OEt}-bpy** was isolated in 42 % yield from $[(\eta^{6}-benzene)Ru(bpy)Cl](PF_{6})$; this yield couldn't be improved by a longer reaction time (two 20 min cycles) or by adding an excess of 4,4'-(CH₂PO₃Et₂)₂-bpy ligand (2.3 instead of 2 eq). As previously observed, $[(\eta^{6}-p-cymene)Ru(bpy)Cl](PF_{6})$, with isolated yields in **RuP4^{OEt}-bpy** not exceeding 25%.

Arene	L	Reaction time	4,4'-(CH ₂ PO ₃ Et ₂) ₂ -bpy	RuP4 ^{OEt} -bpy	Monohydrolyzed RuP4 ^{OEt} -bpy	
	OTf	20 min	2 eq	58%	n.d.	
benzene	Cl	20 min	2 eq	42%	14%	
		20 min	2.3 eq	42%	26%	
		2 x 20 min	2 eq	30%	n.d.	
		2 x 20 min ^a	2.3 eq	30%	34%	
<i>p</i> - cymene	OTf	20 min	2 eq	60%	n.d.	
	Cl	20 min	2 eq	15%	13%	

Table 2. *Microwave activation*: Isolated yields in **RuP**₄^{OEt}-bpy for different reaction conditions (EtOH, 150°C).

2 x 20 min	2 eq	20%	14%
2 x 20 min	2.3 eq	25%	22%

^a2-methoxyethanol instead of ethanol.

It is important to underline the formation of side-products bearing partially hydrolyzed phosphonate groups. The mono-hydrolyzed derivative could be isolated with yields up to 34 % (Table 2), by chromatography on silica gel (eluted after **RuP4^{OEt}-bpy** with 30% aqueous KNO₃ (0.4 M) in CH₃CN), whereas complexes with a higher degree of hydrolysis were irreversibly adsorbed on silica. Formation of these hydrolyzed derivatives was previously mentioned in the literature^[25] and can account for the reported hydrolysis of the phosphonate ester groups before the purification.^[13, 14, 18, 20] It is however detrimental when preparation of the complexes under their phosphonate ester form is required.

Overall, from this set of data, preparation of **RuP**⁴^{OEt}-**bpy** can be smoothly achieved from $[(\eta^6\text{-}arene)Ru(bpy)(OTf)](OTf)$ under microwave activation (Method B: EtOH, 150°C, 20 min; 60 % isolated yield); in addition, if the triflate precursor is not accessible, thermal conditions can be applied to [(benzene)Ru(N^N)Cl](PF₆) (Method A: MeOEtOH/H₂O, 120°C, overnight; 58 % isolated yield) using a 2.3 eq excess of 4,4⁻-(CH₂PO₃Et₂)₂-bpy.

Extension to various N^N ligands. The conditions defined above were tested for the synthesis of novel complexes, with different N^N ligands. Method B was employed each time the triflate precursor complex could be prepared. RuP4^{OEt}-phen and RuP4^{OEt}-phenamine were in from $[(\eta^{6}-arene)Ru(phen)(OTf)](OTf)$ this way obtained and $[(n^{6}$ arene)Ru(phenamine)(OTf)](OTf), in 52 % and 55 % yield, respectively (Table 3). As already mentioned above, the lack of solubility of $[(\eta^6-p-cymene)Ru(dppz)Cl](PF_6)$ prevented us to test the preparation of RuP4^{OEt}-dppz by the organometallic route. Synthesis of RuP4^{OEt}phendione was attempted via method A, but proved to be unsuccessful; formation of the trishomoleptic [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₃](PF₆)₂ complex was instead observed under these conditions, the electron-deficient phendione probably being displaced by the more electrondonating 4,4'-(CH₂PO₃Et₂)₂-bpy. Finally, method A allowed isolating **RuP**₄^{OEt}-**TMS-EPIP** in 20 % yield.

Table 3. Synthesis of various **RuP**₄^{OEt}-**N**^**N** complexes.

	RuP4 ^{OEt} -phen	RuP4 ^{OEt} -phenamine	RuP4 ^{OEt} -phendione	RuP4 ^{OEt} -dppz	RuP4 ^{OEt} -TMS-EPIP	RuP4 ^{OEt} -EPIP
Method A	_	—	0 % ^a	_	20 % (13 % ^b)	—

Method B	52 %	55 %	_		_	—
Traditional approach ^[16]		_	78 %°	55 %°	20 %°	30 %°

a- $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_3](PF_6)_2$ was solely formed. b- Overall yield from $RuCl_3.3H_2O$. b- Overall yield from $Ru(DMSO)_4Cl_2$.

The synthetic limitations (poor orthogonality due to reactive organometallic intermediates, notably) observed at different stages of the organometallic route prompted us to reexamine the traditional approach. The group of S. Rau optimized it in order to prepare the phendione complex **RuP**₄^{OEt}-phendione, key to access an immobilizable RuPd dyad.^[16] Starting from Ru(DMSO)₄Cl₂ instead of RuCl₃, the *cis*-Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl₂ intermediate – better described as a mixture of $[cis-Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2Cl_{2-x}(DMSO)_x]Cl_x - was$ isolated without any purification and directly used for the last step of the process, *i.e.* coordination of the functionalized N^N ligand. The use of Ru(DMSO)₄Cl₂ has been previously reported by F. Odobel and coworkers to dramatically improved the first step of the procedure for related Ru complexes.^[25] The synthesis of **RuP**₄^{OEt}-phendione, reported with a 86% crude yield,^[16] was reproduced with an overall 78% yield in our hands. Moreover, we succeeded to prepare RuP4^{OEt}-dppz in 55%. By contrast, RuP4^{OEt}-TMS-EPIP was isolated with a disappointing 10 % yield, in a first attempt. It could be improved to 20 % by increasing the amount of TMS-EPIP to 1.5 equivalent. Interestingly, protection of the terminal alkyne is not required anymore in this procedure: we succeeded to prepare RuP4^{OEt}- EPIP in 30 % yield by directly reacting 1.5 equivalent of EPIP ligand with [cis-Ru(4,4'-(CH₂PO₃Et₂)₂ $bpy_2Cl_{2-x}(DMSO)_x]Cl_x$. This procedure thus proved to be more efficient and straightforward - only two steps only from $Ru(DMSO)_4Cl_2$ - compared to the organometallic route and its overall 13 % yield (from RuCl₃.3H₂O) for RuP₄OEt-TMS-EPIP.

Spectroscopic and redox properties of the new RuP_4^{OEt} -N^N complexes. The UV/vis absorption spectra of RuP_4^{OEt} -bpy, RuP_4^{OEt} -phen, RuP_4^{OEt} -phenNH₂, RuP_4^{OEt} -dppz and RuP_4^{OEt} -EPIP were recorded in acetonitrile and compared to the parent $[Ru(bpy)_2(N^N)](PF_6)_2$ compounds (Figures S7-11 in Supporting Information). Spectroscopic data are summarized in Table 4. These spectra display intense ligand-centered π - π^* absorption bands between 260 and 300 nm; RuP_4^{OEt} -dppz displays an additional band at 358 nm attributed to a dppz-centered transition.^[26] The visible part of the spectra is dominated by the classical metal-to-ligand charge-transfer (MLCT) transitions centered around 450-460 nm with a shoulder at 430 nm. In comparison to the parent complexes, introduction of four weak electron-donating methyl phosphonate substituents induces a slight bathochromic shift (5 to 7 nm) of the MLCT bands, due to stabilization of bpy-based π^* orbital.

Table 4. Spectroscopica and redox propertiesb of complexes RuP_4^{OEt} -bpy, RuP_4^{OEt} -phen,
 RuP_4^{OEt} -phenNH2, RuP_4^{OEt} -dppz, and RuP_4^{OEt} -EPIP together with the parent
 $[Ru(bpy)_3](PF_6)_2$, $[Ru(bpy)_2(phen)](PF_6)_2$, $[Ru(bpy)_2(phenNH_2)](PF_6)_2$,
 $[Ru(bpy)_2(dppz)](PF_6)_2$ and $[Ru(bpy)_2(EPIP)](PF_6)_2$ as reference complexes.

Photosensitizer	Ref	$\lambda_{abs}\left(\epsilon ight)^{c}$	Eox	Ered1	E _{red2}	Ered3	Ered4
RuP4 ^{OEt} -bpy	[11]	288 (85370), 456 (14830)	+0.87	-1.71	-1.90	-2.15	_
$[Ru(bpy)_3](PF_6)_2$	[11]	286 (94480), 451 (14710)	+0.89	-1.73	-1.92	-2.17	_
RuP4 ^{OEt} -phen	this work	264 (57120), 289 (67990), 455 (16580)	+0.87	-1.72	-1.90	-2.16	_
[Ru(bpy) ₂ (phen)](PF ₆) ₂	this work	264 (56150), 286 (63875), 449 (15900)	+0.88	-1.75	-1.94	-2.20	_
RuP4 ^{OEt} -phenNH2	this work	289 (60145), 464 (14025)	+0.95	-1.74	-1.94	-2.20	_
[Ru(bpy) ₂ (phenNH ₂)](PF ₆) ₂	this work	285 (59635), 458 (13540)	+0.93	-1.78	-1.98	-2.23	_
RuP4 ^{OEt} -dppz	[11]	285 (97975), 368 (13990), 457 (14331)	+0.88	-1.38	-1.72	-1.93	-2.34
$[Ru(bpy)_2(dppz)](PF_6)_2$	[11]	283 (85090), 367 (14780), 449 (14385)	+0.88	-1.38	-1.84	-2.05	-2.31
RuP4 ^{OEt} -EPIP	this work	288 (131360), 457 (20090)	+0.85	-1.74	-1.96	-2.38	_
$[Ru(bpy)_2(EPIP)](PF_6)_2$	this work	290 (106085), 463 (19250)	+0.88	-1.78	-2.02	-2.39	_

a. Absorption spectra were recorded in acetonitrile.

b. in V vs Fc⁺/Fc. Cyclic voltammograms were recorded at a complex concentration of 1 mM in a 0.1 M solution of n-Bu₄NBF₄ in degassed acetonitrile and at a scan rate of 100 mV.s⁻¹.

c. λ_{abs} in nm; ϵ in L.mol⁻¹.cm⁻¹.

Cyclic and square wave voltammograms were recorded in degassed 0.1 M n-Bu₄NBF₄ solution in acetonitrile (Figures S12-16 in Supporting Information) and the redox properties of the different complexes are listed in Table 4. The quasi-reversible one-electron process

observed at a potential comprised between +0.85 and +0.95 V vs Fc⁺/Fc was unambiguously attributed to the metal-centered Ru^{III/II} oxidation, according to previous studies on related ruthenium tris-diimine complexes,^[9, 27] For all complexes, the three quasi-reversible reductions on the diimine ligands are recorded on a range of potentials varying from -1.7 to -2.4 V vs Fc^{+/0}. The cyclic voltammograms of **RuP4^{OEt}-dppz** and of the parent [Ru(bpy)₂(dppz)](PF₆)₂ are characterized by the presence of an additional process at -1.38 V vs Fc^{+/0}, assigned to the pyrazine-based reduction of the dppz ligand.^[28, 29]

Overall, the introduction of four methylphosphonate anchoring groups in the coordination sphere of $[Ru(bpy)_2(N^N)](PF_6)_2$ photosensitizers does not significantly modify their properties, consistent with the electronic decoupling provided by the methylene spacer.^[11, 12]

Conclusion

Novel ruthenium trisdiimine photosensitizers of general formula $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(N^N)](PF_6)_2$, bearing four methylphosphonate anchors, were prepared by two different synthetic procedures. The scope of the organometallic procedure was first studied; some synthetic limitations are highlighted, related to the introduction of the N^N ligand at the very beginning of the process. On the other hand, the traditional approach offers wider versatility, provided that the *cis*-Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2Cl_2 intermediate is used as prepared. Notably, synthesis of a novel alkyne-functionalized photosensitizer, $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(EPIP)](PF_6)_2$, is reported. Although increasing its isolated yield remains challenging, this complex offers interesting perspectives for the immobilization of dye-catalyst assemblies onto semi-conductor surfaces. Finally, the electronic properties of these light-harvesting units were studied and proved not to be significantly altered by the introduction of four phosphonate anchors thanks to the presence of a methylene spacer, which is an important parameter for future applications in the field of DS-PECs.

Experimental section

All reagents were purchased from Sigma Aldrich or Strem and used as obtained unless otherwise stated. Reagent-grade solvents were used without further purification. The 4,4'-bis(diethylphosphonomethyl)-2,2'-bipyridine ligand (4,4'-(CH₂PO₃Et₂)₂-bpy) was custom-synthesized by the company Oribase Pharma, according to a previously reported procedure.^[9] The dipyrido[3,2-a:2',3'-c]phenazine (dppz),^[29] phendione,^[30] 2-(4-trimethylsilylethynylphenyl)-1H-imidazo[4,5-f][1,10]phenanthroline (epip-TMS)^[31] and 2-(4-ethynylphenyl)-1H-imidazo[4,5-f][1,10]phenanthroline (epip)^[21] ligands were prepared

according to previously reported procedures. Spectroscopic and electrochemical characterizations of complexes **RuP**₄^{OEt}-**bpy** and **RuP**₄^{OEt}-**dppz** were previously reported. ¹H NMR experiments were carried out on a Bruker Avance 300 MHz and the resulting spectra are referenced to the residual solvent peak and reported in relative to tetramethylsilane reference ($\delta = 0$ ppm). UV-vis absorption spectra were recorded either on a Shimadzu UV-1800 spectrometer or on an Agilent Cary 60 UV-Vis spectrometer. ESI-MS measurements were carried out on a Thermoquest Finnigan LCQ spectrometer.

[**Ru**(η^{6} -benzene)₂Cl₂]₂.^[32] A solution of RuCl₃.3H₂O (3 g, 11.5 mmol) and *1*,4cyclohexadiene (11 mL, 115 mmol) in absolute ethanol (150 mL) was heated to reflux for 8 hours. The reaction mixture was then cooled down to -40°C overnight. The dark orange precipitate is finally filtrated, washed with diethyl ether and dried under vacuum to yield 2.02 g (71 %) of [Ru(η^{6} -benzene)₂Cl₂]₂. ¹H NMR (300 MHz, CDCl₃): δ (ppm) 5.71 (s, 6H).

[**Ru**(η^{6} -*p*-cymene)₂Cl₂]₂.^[32] A solution of RuCl₃.3H₂O (3 g, 11.5 mmol) and α -phellandrene (19 mL, 115 mmol) in absolute ethanol (150 mL) was heated to reflux for 8 hours. The reaction mixture was then cooled down to -40°C overnight. The dark red crystalline product is finally filtrated, washed with diethyl ether and dried under vacuum to yield 3.39 g (96 %) of [Ru(η^{6} -*p*-cymene)₂Cl₂]₂. ¹H NMR (300 MHz, CDCl₃): δ (ppm) 5.47 (d, 2H), 5.33 (d, 2H), 2.92 (hept., 1H), 2.15 (s, 3H), 1.27 (d, 6H).

General procedure for the synthesis of $[(\eta^6\text{-benzene})Ru(N^N)Cl](PF_6)(N^N = bpy, phen, phenamine, phendione, epip-TMS)$. A solution of $[(\eta^6\text{-benzene})RuCl_2]_2$ (150 mg, 0.3 mmol) and N^N diimine ligand (0.6 mmol) in methanol (50 mL) was heated to reflux overnight. Solvent was removed under reduced pressure and the resulting precipitate dissolved in water (5 mL). Dropwise addition of a saturated KPF_6 aqueous solution gave a dark to bright yellow solid, which was collected, washed thoroughly with water (5 x 10 mL), diethyl ether (3 x 10 mL) and finally dried under vacuum.

[(η⁶- benzene)Ru(bpy)Cl](PF₆). Yield: 91 %. ¹H NMR (300 MHz, MeOD): δ (ppm) 9.40 (d, J = 5.6 Hz, 2H), 8.31 (d, J = 8.1 Hz, 2H), 8.18 (t, J = 7.8 Hz, 2H), 7.68 (t, J = 6.6 Hz, 2H), 6.01 (s, 6H). ESI-MS: m/z 371.0 [M-PF₆]⁺.

[(η⁶-benzene)Ru(phen)Cl](PF₆). Yield: 93 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.74 (d, J = 5.2 Hz, 2H), 8.75 (d, J = 8.2 Hz, 2H), 8.14 (s, 2H), 8.02 (dd, J = 8.2, 5.3 Hz, 2H), 6.10 (s, 6H). ESI-MS: m/z 395.0 [M-PF₆]⁺.

[(η⁶-benzene)Ru(phenamine)Cl](PF₆). Yield: 94 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.71 (d, J = 5.3 Hz, 1H), 9.34 (d, J = 5.2 Hz, 1H), 8.73 (d, J = 8.5 Hz, 1H), 8.31 (d, J = 8.3 Hz, 1H), 7.97 (dd, J = 8.5, 5.3 Hz, 1H), 7.76 (dd, J = 8.3, 5.2 Hz, 1H), 7.05 (s, 1H), 6.05 (s, 6H), 5.48 (s, 2H). ESI-MS: m/z 409.6 [M-PF₆]⁺.

[(η⁶- benzene)Ru(phendione)Cl](PF₆). Yield: 93 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.81 (dd, J = 5.6, 1.2 Hz, 2H), 8.65 (dd, J = 7.9, 1.2 Hz, 2H), 7.99 (dd, J = 7.9, 5.6 Hz, 2H), 6.31 (s, 6H). ESI-MS: m/z 425.0 [M-PF₆]⁺.

[(η⁶- benzene)Ru(TMS-EPIP)Cl](PF₆). Yield: 95 %. ¹H NMR (300 MHz, DMSO d6): δ (ppm) 9.97 (d, J = 5.2 Hz, 2H), 9.21 (d, J = 9.0 Hz, 2H), 8.31 (d, J = 8.3 Hz, 2H), 8.28 – 8.15 (m, 2H), 7.76 (d, J = 8.5 Hz, 2H), 6.33 (s, 6H), 0.27 (s, 9H). ESI-MS: m/z 607.2 [M-PF₆]⁺.

General procedure for the synthesis of $[(\eta^6 - p - cymene)Ru(N^N)Cl](PF_6)(N^N = bpy, phen, phenamine, phendione, dppz):$

A solution of $[(\eta^6-p\text{-cymene})\text{RuCl}_2]_2$ (150 mg, 0.25 mmol) and N^N ligand (0.5 mmol) in methanol (50 mL) was heated to reflux overnight. Solvent was removed under reduced pressure and the resulting precipitate dissolved in water (5 mL). Dropwise addition of a saturated KPF₆ aqueous solution gave a dark to bright yellow solid, which was collected, washed thoroughly with water (5 x 10 mL), diethyl ether (3 x 10 mL) and finally dried under vacuum.

[(η⁶-*p*-cymene)**Ru(bpy)Cl**] (**PF**₆). Yield: 89 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.32 (d, J = 5.6 Hz, 2H), 8.31 (d, J = 8.1 Hz, 2H), 8.18 (t, J = 7.9 Hz, 2H), 7.69 (t, J = 6.6 Hz, 2H), 5.91 (d, J = 6.3 Hz, 2H), 5.71 (d, J = 6.3 Hz, 2H), 2.65 (hept., 1H), 2.20 (s, 3H), 1.02 (d, J = 6.9 Hz, 6H). ESI-MS: m/z 427.1 [M-PF₆]⁺.

[(η^{6} -*p*-cymene)**Ru**(**phen**)**Cl**] (**PF**₆). Yield : 90 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.66 (d, J = 5.3 Hz, 2H), 8.75 (d, J = 8.3 Hz, 2H), 8.15 (s, 2H), 8.03 (dd, J = 8.2, 5.3 Hz, 2H), 6.02 (d, J = 6.3 Hz, 2H), 5.83 (d, J = 6.3 Hz, 2H), 2.69 (hept., 1H), 2.17 (s, 3H), 1.00 (d, J = 6.9 Hz, 6H). ESI-MS: m/z 451.1 [M-PF₆]⁺.

[(η⁶-*p*-cymene)Ru(phenamine)Cl] (PF₆). Yield : 94 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.63 (d, J = 5.3 Hz, 1H), 9.26 (d, J = 5.2 Hz, 1H), 8.73 (d, J = 8.4 Hz, 1H), 8.30 (d, J = 8.4 Hz, 1H), 7.98 (dd, J = 8.5, 5.3 Hz, 1H), 7.77 (dd, J = 8.3, 5.2 Hz, 1H), 7.04 (s, 1H), 5.97 (s, 2H), 5.78 (s, 2H), 5.48 (s, 2H), 2.65 (hept., 1H), 2.05 (s, 3H), 0.97 (dd, J = 6.9, 3.1 Hz, 6H). ESI-MS: m/z 465.6 [M-PF₆]⁺.

[(η^{6} -*p*-cymene)**Ru**(**phendione**)**Cl**] (**PF**₆). Yield: 91 %. ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.69 (d, J = 5.3 Hz, 2H), 8.64 (d, J = 7.9 Hz, 2H), 7.98 (dd, J = 7.7, 5.7 Hz, 2H), 6.30 (d, J = 6.4 Hz, 2H), 6.06 (d, J = 6.0 Hz, 2H), 2.61 (hept. , 1H), 2.22 (s, 3H), 0.96 (d, J = 6.8 Hz, 6H). ESI-MS: m/z 481.2 [M-PF₆]⁺.

[(η⁶-*p*-cymene)Ru(dppz)Cl] (PF₆). Yield : 95 %. ¹H NMR (300 MHz, DMSO d6): δ (ppm) 10.01 (d, J = 5.2 Hz, 2H), 9.74 (d, J = 8.1 Hz, 2H), 8.49 (dd, J = 6.3, 3.4 Hz, 2H), 8.31 (dd, J = 8.1, 5.4 Hz, 2H), 8.17 (dd, J = 6.4, 3.4 Hz, 2H), 6.39 (d, J = 6.2 Hz, 2H), 6.16 (d, J = 6.2 Hz, 2H), 2.79 – 2.62 (hept., 1H), 2.21 (s, 3H), 1.00 (d, J = 6.8 Hz, 6H). ESI-MS: m/z 553.3 [M-PF₆]⁺.

General procedure for the synthesis of $[(\eta^6-arene)Ru(N^N)OTf](OTf)$ (arene = bpy or pcymene; $N^N = bpy$, phen, phenamine):

Trifluoromethanesulfonic acid (500 μ L, 5.65 mmol, 100 eq) was carefully added to a suspension of [Ru(η^6 -arene)(N^N)Cl](PF₆) (150 mg) in dichloromethane (50 mL). The reaction mixture was stirred at room temperature in the dark overnight. Addition of diethyl ether gave a yellowish precipitate, which was collected, washed thoroughly with diethyl ether and finally dried under vacuum.

[(η⁶- benzene)Ru(bpy)(OTf)](OTf). Yield: 91 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm) 9.81 (d, J = 5.5 Hz, 2H), 8.59 (d, J = 8.2 Hz, 2H), 8.39 (t, J = 7.9 Hz, 2H), 7.92 (t, J = 6.6 Hz, 2H), 6.33 (s, 6H). ESI-MS: m/z 485.1 [M-OTf]⁺, 168.0 [M-2OTf]²⁺.

[(η⁶-*p*-cymene)Ru(bpy)(OTf)](OTf). Yield: 94 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm) 9.82 (d, J = 5.7 Hz, 2H), 8.63 (d, J = 8.1 Hz, 2H), 8.42 (t, J = 8.0 Hz, 2H), 7.96 (t, J = 6.6 Hz, 2H), 6.45 (d, J = 6.3 Hz, 2H), 6.23 (d, J = 6.3 Hz, 2H), 2.57 (hept., J = 6.8 Hz, 1H), 2.30 (s, 3H), 1.03 (d, J = 6.9 Hz, 6H). ESI-MS: m/z 540.8 [M-OTf]⁺, 196.2 [M-2OTf]²⁺.

[(η⁶- benzene)Ru(phen)(OTf)](OTf). Yield: 93 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm) 10.15 (d, J = 5.3 Hz, 2H), 9.04 (d, J = 8.2 Hz, 2H), 8.46 (s, 2H), 8.34 (dd, J = 8.2, 5.1 Hz, 2H), 6.35 (s, 6H). ESI-MS: m/z 508.7 [M-OTf]⁺, 179.4 [M-2OTf]²⁺.

[(η⁶-*p*-cymene)Ru(phen)(OTf)](OTf). Yield 91 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm) 10.19 (d, J = 5.3 Hz, 2H, 9.02 (d, J = 8.3 Hz, 2H), 8.35 – 8.20 (m, 4H), 6.57 (d, J = 6.4 Hz, 2H), 6.37 (d, J = 6.4 Hz, 2H), 2.58 (hept., 1H), 2.29 (s, 3H), 0.97 (d, J = 6.9 Hz, 6H). ESI-MS: m/z 564.9 [M-OTf]⁺, 207.4 [M-2OTf]²⁺.

[(η⁶- benzene)Ru(phenamine)(OTf)](OTf). Yield: 92 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm)10.21 (d, J = 5.1 Hz, 1H), 9.45 (d, J = 5.3 Hz, 1H), 8.92 (d, J = 8.2 Hz, 1H), 8.53 (d, J = 8.5 Hz, 1H), 8.26 (dd, J = 8.2, 5.4 Hz, 1H), 8.02 (dd, J = 8.2, 5.4 Hz, 1H), 7.16 (s, 1H), 6.23 (s, 6H), 5.62 (s, 2H). ESI-MS: m/z 525.5 [M-OTf]⁺, 187.2 [M-2OTf]²⁺.

[(η⁶-*p*-cymene)Ru(phenamine)(OTf)](OTf). Yield 93 %. ¹H NMR (300 MHz, CD₃OD): δ (ppm)10.04 (d, J = 5.2 Hz, 1H), 9.47 (d, J = 5.3 Hz, 1H), 8.95 (d, J = 8.2 Hz, 1H), 8.52 (d, J = 8.1 Hz, 1H), 8.12 (dd, J = 8.4, 5.4 Hz, 1H), 7.95 (dd, J = 8.1, 5.2 Hz, 1H), 7.36 (s, 1H), 6.19 (s, 2H), 6.02 (s, 2H), 5.73 (s, 2H), 2.63 (s, 1H), 2.24 (s, 3H), 1.02 (dd, J = 6.9, 3.1 Hz, 6H). ESI-MS: m/z 579.6 [M-OTf]⁺, 215.3 [M-2OTf]²⁺.

General Procedure for the synthesis of $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(N^N)](PF_6)_2$ by the organometallic route:

Method A (thermal conditions): A solution of $[(\eta^{6}\text{-arene})\text{Ru}(N^{N})\text{Cl}](\text{PF}_{6})$ (150 mg, 1 equivalent) and 4,4'-(CH₂PO₃Et₂)₂-bpy (2 to 2.3 equivalents) in a mixture of water (9 mL) and 2-methoxyethanol (36 mL) was refluxed overnight. After cooling down to room temperature, 5 mL of a saturated KPF₆ aqueous solution was added to the reaction mixture. Extraction with dichloromethane and removal of the organic solvent yield a reddish solid which was then subjected to chromatography on silica gel (MeCN / aqueous 0.4 M KNO₃, 80:20). After removal of MeCN under reduced pressure and addition of a saturated KPF₆ aqueous solution (5 mL), the aqueous phase was further extracted (3 x 50 mL) with dichloromethane and the organic phases washed with water. Evaporation of dichloromethane under reduced pressure gave a red-orange solid, which was collected and vacuum-dried overnight.

Method B (microwave conditions): A solution of $[(\eta^6\text{-arene})\text{Ru}(N^N)(\text{OTf})](\text{OTf})$ (150 mg, 1 equivalent) and 4,4'-(CH₂PO₃Et₂)₂-bpy (2 equivalents) in ethanol (10 mL) was put in a sealed tube before being subjected to microwave irradiation (150°C) for 20 min. After cooling down to room temperature, 5 mL of a saturated KPF₆ aqueous solution was added to the reaction mixture. Extraction with dichloromethane and removal of the organic solvent yield a reddish solid which was then subjected to chromatography on silica gel (MeCN / aqueous 0.4M KNO₃, 80:20). After removal of MeCN under reduced pressure and addition of a saturated KPF₆ aqueous solution (5 mL), the aqueous phase was further extracted (3 x 50 mL) with dichloromethane and the organic phases washed with water. Evaporation of dichloromethane

under reduced pressure gave a red-orange solid, which was collected and vacuum-dried overnight.

 $[\mathbf{Ru}(4,4'-(\mathbf{CH}_2\mathbf{PO}_3\mathbf{Et}_2)_2-\mathbf{bpy})_2(\mathbf{bpy})](\mathbf{PF}_6)_2$ ($\mathbf{RuP}_4^{\mathbf{OEt}}-\mathbf{bpy}$).^[11] Yield: 58% (method A: benzene); 58% (method B: benzene); 60% (method B: *p*-cymene).

[**Ru**(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(phen)](PF₆)₂ (**Ru**P₄^{OEt}-phen). Yield: 52% (method B: *p*-cymene). ¹H NMR (300 MHz, CD₃CN): δ (ppm) 8.62 (d, J = 8.2 Hz, 2H), 8.44 (d, J = 15.1 Hz, 4H), 8.24 (s, 2H), 8.06 (d, J = 5.2 Hz, 2H), 7.81 – 7.69 (m, 4H), 7.43 (d, J = 5.8 Hz, 2H), 7.39 (d, J = 5.7 Hz, 2H), 7.14 (d, J = 5.7 Hz, 2H), 4.11 – 3.88 (m, 16H), 3.42 (d, J = 22.6 Hz, 4H), 3.33 (d, J = 22.5 Hz, 4H), 1.25 – 1.03 (m, 24H). ESI-MS: m/z 597.1 [M–2PF₆]²⁺.

[**Ru**(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(phenNH₂)](PF₆)₂ (**Ru**P₄^{OEt}-phenNH₂). Yield: 55% (method B: *p*-cymene). ¹H NMR (300 MHz, CD₃CN): δ (ppm) 8.61 (m, J = 8.5 Hz, 2H), 8.44 (s, 2H), 8.39 (s, 2H), 8.20 (d, J = 8.4 Hz, 1H), 8.02 (d, J = 5.1 Hz, 1H), 7.80 – 7.60 (m, 4H), 7.54 – 7.41 (m, 3H), 7.37 (s, 2H), 7.17 (s, 2H), 5.57 (s, 2H), 4.17 – 3.82 (m, 16H), 3.55 – 3.16 (m, 8H), 1.32 – 0.99 (m, 24H). ESI-MS: m/z 604.3 [M–2PF₆]²⁺.

[**Ru**(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(TMS-EPIP)](PF₆)₂ (**Ru**P₄^{OEt}-TMS-EPIP). Yield: 20% (method A: benzene; 2.3 eq). ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.04 – 8.91 (m, 2H), 8.43 (d, *J* = 14.7 Hz, 4H), 8.26 (d, *J* = 8.1 Hz, 2H), 7.99 (d, *J* = 4.2 Hz, 2H), 7.79 – 7.66 (m, 4H), 7.50 (d, *J* = 5.7 Hz, 2H), 7.40 (dd, *J* = 5.4 & 1.8 Hz, 4H), 7.15 (d, *J* = 5.7 Hz, 2H), 4.10 – 3.89 (m, 16H), 3.43 (d, J = 22.5 Hz, 4H), 3.33 (d, J = 22.5 Hz, 4H), 1.23 – 1.05 (m, 24H), 0.28 (s, 9H). ESI-MS: m/z 703.4 [M–2PF₆]²⁺.

General Procedure for the synthesis of $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(N^N)](PF_6)_2$ by the traditional approach^[16] (N^N: phendione, dppz, epip-TMS, epip): A solution of commercially available Ru(DMSO)_4Cl₂ (420 mg, 0.88 mmol) and 4,4'-(CH₂PO₃Et₂)₂-bpy (800 mg, 1.75 mmol) in methanol (100 mL) was refluxed overnight under argon. Removal of the solvent and drying under vacuum yielded the dark-red highly hygroscopic intermediate [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl_x(DMSO)_{2-x}]Cl_{2-x}. A solution of crude [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂Cl_x(DMSO)_{2-x}]Cl_{2-x} (1 equivalent, using an average molecular weight) and N^N (1 to 1.5 equivalent) in a water/ethanol (1:3) mixture was refluxed overnight. After cooling down to room temperature, 5 mL of a saturated KPF₆ aqueous solution was added to the reaction mixture. Extraction with dichloromethane and removal of the organic solvent yield a crude reddish solid. After purification by flash chromatography on silica gel (MeCN/aqueous KNO₃ (0.4 M), 80:20), MeCN was removed under reduced pressure and a saturated KPF₆ aqueous

solution (5 mL) added. The aqueous phase was further extracted (3 x 50 mL) with dichloromethane and the organic phases washed with water. Evaporation of dichloromethane under reduced pressure gave red-orange solids, which were collected and vacuum-dried overnight.

[Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(phendione)](PF₆)₂ (RuP₄^{OEt}-phendione).^[16, 17] Yield: 78%. [Ru(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(dppz)](PF₆)₂ (RuP₄^{OEt}-dppz).^[11] Yield: 55%.

 $[Ru(4,4'-(CH_2PO_3Et_2)_2-bpy)_2(TMS-EPIP)](PF_6)_2$ (RuP_4^{OEt} -TMS-EPIP). Yield: 20% (1.5 eq of TMS-EPIP ligand).

[**Ru**(4,4'-(CH₂PO₃Et₂)₂-bpy)₂(EPIP)](PF₆)₂ (**Ru**P₄^{OEt}-EPIP). Yield: 30% (1.5 eq of EPIP ligand). ¹H NMR (300 MHz, CD₃CN): δ (ppm) 9.09 (d, *J* = 8.5 Hz, 1H), 8.87 (d, *J* = 8.1 Hz, 1H), 8.42 (d, *J* = 15.4 Hz, 4H), 8.28 (d, *J* = 8.5, 2H), 8.00 (d, *J* = 5.1 Hz, 2H), 7.81 – 7.74 (m, 6H), 7.49 (t, *J* = 6.0 Hz, 2H), 7.40 (d, *J* = 5.4 Hz, 2H), 7.15 (s broad, 2H), 4.10 – 3.88 (m, 16H), 3.60 (s, 1H), 3.42 (d, *J* = 22.6, 4H), 3.32 (d, *J* = 22.6, 4H), 1.23 – 1.05 (m, 24H). ESI-MS: m/z 667.3 [M–2PF₆]²⁺.

Supporting Information. ¹H NMR spectra, UV-vis absorption spectra and cyclic voltammograms for all new complexes.

Acknowledgements

Colette Lebrun (CEA/DRF/INAC/SyMMES) is acknowledged for the ESI-MS measurements. This work was supported by the Labex program ARCANE (ANR-11-LABX-0003-01), the graduate school in Chemistry, Biology and Health of Univ. Grenoble Alpes (CBH-EUR-GS, ANR-17-EURE-0003), the European Research Council under the European Union's Seventh Framework Program FP/2007-2013 (ERC Grant Agreement n.306398).

Bibliography

- [1] P. Xu, N. S. McCool, T. E. Mallouk, *Nano Today* **2017**, *14*, 42-58.
- [2] M. K. Brennaman, R. J. Dillon, L. Alibabaei, M. K. Gish, C. J. Dares, D. L. Ashford, R. L. House, G. J. Meyer, J. M. Papanikolas, T. J. Meyer, J. Am. Chem. Soc. 2016, 138, 13085-13102.
- [3] Z. Yu, F. Li, L. Sun, *Energy Environ. Sci.* **2015**, *8*, 760-775.
- [4] S. Berardi, S. Drouet, L. Francas, C. Gimbert-Surinach, M. Guttentag, C. Richmond, T. Stoll, A. Llobet, *Chem. Soc. Rev.* **2014**, *43*, 7501-7519.
- [5] A. Hagfeldt, G. Boschloo, L. Sun, L. Kloo, H. Pettersson, *Chem. Rev.* 2010, 110, 6595-6663.
- [6] C. Queffelec, M. Petit, P. Janvier, D. A. Knight, B. Bujoli, *Chem. Rev.* 2012, 112, 3777-3807.
- [7] P. H. Mutin, G. Guerrero, A. Vioux, J. Mater. Chem. 2005, 15, 3761-3768.
- [8] L. Zhang, J. M. Cole, ACS Appl. Mater. Interfaces 2015, 7, 3427-3455.

- [9] I. Gillaizeau-Gauthier, F. Odobel, M. Alebbi, R. Argazzi, E. Costa, C. A. Bignozzi, P. Qu, G. J. Meyer, *Inorg. Chem.* **2001**, *40*, 6073-6079.
- [10] K. Hanson, M. K. Brennaman, H. Luo, C. R. K. Glasson, J. J. Concepcion, W. Song, T. J. Meyer, ACS Appl. Mater. Interfaces 2012, 4, 1462-1469.
- [11] N. Queyriaux, R. A. Wahyuono, J. Fize, C. Gablin, M. Wächtler, E. Martinez, D. Léonard, B. Dietzek, V. Artero, M. Chavarot-Kerlidou, *J. Phys. Chem. C* **2017**, *121*, 5891-5904.
- [12] Y. Pellegrin, L. Le Pleux, E. Blart, A. Renaud, B. Chavillon, N. Szuwarski, M. Boujtita, L. Cario, S. Jobic, D. Jacquemin, F. Odobel, J. Photochem. Photobiol. A: Chemistry 2011, 219, 235-242.
- [13] A. Zaban, S. Ferrere, B. A. Gregg, J. Phys. Chem. B 1998, 102, 452-460.
- [14] G. Will, G. Boschloo, S. N. Rao, D. Fitzmaurice, J. Phys. Chem. B 1999, 103, 8067-8079.
- [15] J. Sotomayor, G. Will, D. Fitzmaurice, J. Mater. Chem. 2000, 10, 685-692.
- [16] M. Braumuller, M. Schulz, D. Sorsche, M. Pfeffer, M. Schaub, J. Popp, B.-W. Park, A. Hagfeldt, B. Dietzek, S. Rau, *Dalton Trans.* **2015**, *44*, 5577-5586.
- [17] M. Braumuller, M. Schulz, M. Staniszewska, D. Sorsche, M. Wunderlin, J. Popp, J. Guthmuller, B. Dietzek, S. Rau, *Dalton Trans.* **2016**, *45*, 9216-9228.
- [18] S. Caramori, V. Cristino, R. Argazzi, L. Meda, C. A. Bignozzi, *Inorg. Chem.* 2010, 49, 3320-3328.
- [19] D. L. Ashford, W. Song, J. J. Concepcion, C. R. K. Glasson, M. K. Brennaman, M. R. Norris, Z. Fang, J. L. Templeton, T. J. Meyer, J. Am. Chem. Soc. 2012, 134, 19189-19198.
- [20] M. R. Norris, J. J. Concepcion, C. R. K. Glasson, Z. Fang, A. M. Lapides, D. L. Ashford, J. L. Templeton, T. J. Meyer, *Inorg. Chem.* 2013, 52, 12492-12501.
- [21] N. Queyriaux, E. S. Andreiadis, S. Torelli, J. Pecaut, B. S. Veldkamp, E. A. Margulies, M. R. Wasielewski, M. Chavarot-Kerlidou, V. Artero, *Faraday Discuss.* 2017, 198, 251-261.
- [22] C. Menéndez, D. Morales, J. Pérez, V. Riera, D. Miguel, *Organometallics* **2001**, *20*, 2775-2781.
- [23] C. O. Kappe, Angew. Chem. Int. Ed. 2004, 43, 6250-6284.
- [24] S. Rau, B. Schäfer, A. Grüßing, S. Schebesta, K. Lamm, J. Vieth, H. Görls, D. Walther, M. Rudolph, U. W. Grummt, E. Birkner, *Inorg. Chim. Acta* **2004**, *357*, 4496-4503.
- [25] H. Zabri, I. Gillaizeau, C. A. Bignozzi, S. Caramori, M.-F. Charlot, J. Cano-Boquera, F. Odobel, *Inorg. Chem.* **2003**, *42*, 6655-6666.
- [26] E. Sabatani, H. D. Nikol, H. B. Gray, F. C. Anson, J. Am. Chem. Soc. 1996, 118, 1158-1163.
- [27] M. Montalti, S. Wadhwa, W. Y. Kim, R. A. Kipp, R. H. Schmehl, *Inorg. Chem.* 2000, *39*, 76-84.
- [28] J. Fees, W. Kaim, M. Moscherosch, W. Matheis, J. Klima, M. Krejcik, S. Zalis, *Inorg. Chem.* 1993, 32, 166-174.
- [29] E. Amouyal, A. Homsi, J.-C. Chambron, J.-P. Sauvage, J. Chem. Soc., Dalton Trans. 1990, 1841-1845.
- [30] J. Ettedgui, R. Neumann, J. Am. Chem. Soc. 2009, 131, 4-5.
- [31] S. Ramachandra, K. C. Schuermann, F. Edafe, P. Belser, C. A. Nijhuis, W. F. Reus, G. M. Whitesides, L. De Cola, *Inorg. Chem.* **2011**, *50*, 1581-1591.
- [32] M. A. Bennett, A. K. Smith, J. Chem. Soc., Dalton Trans. 1974, 233-241.