



**HAL**  
open science

## The effect of ageing errors on Von Bertalanffy parameters estimation using a Bayesian sensitivity analysis approach

Lucie Aulus Giacosa, Philippe Gaudin, Jean-Christophe Aymes, Francois Gueraud, Matthias Vignon

### ► To cite this version:

Lucie Aulus Giacosa, Philippe Gaudin, Jean-Christophe Aymes, Francois Gueraud, Matthias Vignon. The effect of ageing errors on Von Bertalanffy parameters estimation using a Bayesian sensitivity analysis approach. 6. International Otolith Symposium IOS 2018, Apr 2018, Keelung, Taiwan. 295 p., 2018, Sixth International Otolith Symposium Handbook. hal-02154547

**HAL Id: hal-02154547**

**<https://hal.science/hal-02154547>**

Submitted on 12 Jun 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Distributed under a Creative Commons Attribution 4.0 International License


### Context

The analysis of age and growth information contained in the **scales of fish** has become a standard technique in fish population dynamics model. Commonly modeled using Von Bertalanffy growth function (VBGF), inferring growth for fish is dependent upon an accurate description of the age-length relationship, which may be undermined by ageing errors. Ageing errors frequently arise either from disagreements between scales, readers or inability of scales to reflect true age (i.e. erosion, false growth ring). Here we use a Bayesian approach to infer *Salmo trutta* (L.) VBGF's parameters from scale samples. Inferred from scale readings of 60 fish, age at capture (for all fish) and age at migration (for the anadromous fish) were extracted based on a hierarchical design with blind double reading using 4 scales per fish. We estimated VBGF's parameters under different parametrization in the number of scales and reader per fish.


### Hierarchical design (Nested sampling)



### Localization




### Variance analyses



$$AGE \sim rpt(age \sim (1|Fish)+(1|Reader)+(1|Scale), datatype=Poisson, \dots)^*$$

$$MEASURE \sim rpt(rt \sim (1|Fish)+(1|Reader)+(1|Scale), datatype=Gaussian, \dots)^*$$



### Models



$$L_{i,t} = L_{\infty i} + e^{-K_i(t_i - t_0)}$$

$$L_{\infty i} = a + \varepsilon_{1i}, \quad a \sim \text{dunif}(500, 1000)$$

$$K_i = b + \varepsilon_{2i}, \quad b \sim \text{dnorm}(0.6, 0.001)$$

$$t_0 \sim \text{dunif}(-10, 0.5)$$

### Results



Age accuracy can be proven or estimated with methods such as capture-mark-recapture (CMR). Agreement in the age determination between reader (consensual readings) is a selection of fish for which precision is 100% but it does not mean that the determination is accurate and represents the entire population. \*\*

Sources of variance in the age determination should be investigated before any results interpretation. By doing so, we can empirically determine the robustness of VBGF's parameters at a population scale in our study. \*\*\* Going further, variance on age should be integrated at the individual level.

\* Stoffel, M. A., Nakagawa, S. and Schielzeth, H. (2017). rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol Evol*, 8: 1639-1644.

\*\* Beamish, R. J., et G. A. McFarlane. « The Forgotten Requirement for Age Validation in Fisheries Biology ». *Transactions of the American Fisheries Society* 112

\*\*\* Cope, Jason M., et André E. Punt. « Admitting ageing error when fitting growth curves: an example using the von Bertalanffy growth function with random effects ». *Canadian Journal of Fisheries and Aquatic Sciences* 64, no 2 (1 février 2007): 205-18.