



Spatial complexity and temporal dynamics in viticulture: A review of climate-driven scales

Etienne Neethling, Gérard Barbeau, Cécile Coulon-Leroy, Hervé Quénol

► To cite this version:

Etienne Neethling, Gérard Barbeau, Cécile Coulon-Leroy, Hervé Quénol. Spatial complexity and temporal dynamics in viticulture: A review of climate-driven scales. *Agricultural and Forest Meteorology*, 2019, 276-277, pp.107618. <10.1016/j.agrformet.2019.107618>. <hal-02156533>

HAL Id: hal-02156533

<https://hal.science/hal-02156533v1>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Spatial complexity and temporal dynamics in viticulture: A review of climate-driven scales

Etienne Neethling^{1*}, Gérard Barbeau, Cécile Coulon-Leroy¹ and Hervé Quéno²

¹ESA, USC 1422 INRA-GRAPPE, Ecole Supérieure d'Agricultures, 55 rue Rabelais, 49007 Angers, France

²CNRS, UMR 6554 LETG, Université Rennes 2, Place du Recteur Henri Le Moal, 35043 Rennes, France

*Corresponding author: E. Neethling, Email: e.neethling@groupe-esa.com

Abstract

Viticulture is a complex and dynamic system, where climate is a key environmental component of plant suitability and productivity. From field to farm level, climate also plays a prominent role in ongoing practices, shaping winegrowers' decision making and adaptive management. With a changing climate, the wine sector faces many environmental and socio-economic challenges, to which winegrowers are required to adapt. Given the perennial nature of grape growing, there is a need to develop strategies that deal with both short- and long-term climate changes, while likewise accounting for contextual vulnerability. The content of this article aims to provide an overview of climate-driven scales, outlining the spatial complexity and temporal dynamics in viticulture. In addressing these aspects, this literature review offers a framework of scale and cross-scale interactions for policymakers and stakeholders to use when considering responses to attenuate climate change and to reduce its impacts on grape and wine production. The article concludes by discussing the local and context-dependent nature of viticulture in a changing global climate, by emphasizing that the question of scale is fundamental to assessing expected impacts, understanding uncertainty and framing sustainable policies and responses over space and in time.

Key words: Climate, global change, spatial complexity, temporal dynamics, viticulture

1. Introduction

Cultivated and shaped by human uses for many centuries, the grapevine (*Vitis vinifera* L.) has a rich geographical spread, covering more than seventy countries today (OIV, 2018). Given the genetic diversity found in grapevine

varieties (This et al., 2006), it can be grown in diverse climates, ranging from Mediterranean (e.g. Mediterranean Basin) to continental (e.g. Hungary), oceanic (e.g. New Zealand), dry subtropical (e.g. Argentina) and humid subtropical conditions (e.g. Uruguay). Over these geographical areas, long-term climate structures influence regional grape growing and winemaking potential (Tonietto and Carbonneau, 2004), while seasonally, affects grape productivity and wine quality (Salinger et al., 2015; Real et al., 2017). Many studies have demonstrated this central role of climate on vine parameters such as perennial biomass, yield and berry composition (Buttrose et al., 1971; Coombe, 1987; Jackson and Lombard, 1993; Jones and Davis, 2000; Van Leeuwen et al., 2004). As wine structure and flavour are much dependent on grape attributes (Cadot et al., 2012; Bindon et al., 2013), any climate changes impact wine quality and volume produced, resulting in considerable effects on the economic viability of the wine sector (Haeger and Storchmann, 2006; Ashenfelter and Storchmann, 2016). For example, world wine production was 8.2% lower in 2017, than compared with 2016, particularly in the European Union following an extreme frost event in April (OIV, 2017). Viticulture is therefore highly sensitive to changing climate conditions, both temporally and spatially (Jones and Webb, 2010).

Although until the year 2000, few studies about climate change and its impacts on viticulture emerged (Dry, 1988; Smart, 1989; Kenny and Harrison, 1992; Bindi et al., 1996), this topic has received much attention in recent years. Across wine growing regions, climate change has essentially resulted in regional warming (Jones et al., 2005; Cook and Wolkovich, 2016), and to some extent, though less spatially coherent and statistically significant, in greater variability in seasonal rainfall (Laget et al., 2008; Ramos et al., 2015). In response to increasing temperatures and declining soil water contents (Webb et al., 2012), grapevine phenology has shifted earlier by several days (Jones and Davis, 2000; Tomasi et al., 2011), while a warmer ripening period has resulted in changes in berry composition, increased alcohol contents and altered wine sensory profiles (de Orduña, 2010; Neethling et al., 2012). With climate playing a key role in viticulture, a changing climate raises many questions ranging from regional wine quality and style to issues such as geographical shifts in suitable grapevine varieties and production areas over the long term (Schultz and Jones, 2010; Ollat et al., 2016). Vineyards planted over the next decade, and remaining economically productive for many years, are likely to be exposed to unprecedented climate conditions (Mora et al., 2013). Adjusting grape growing and winemaking practices and techniques are therefore an immediate priority to attenuate climate change and to reduce its impacts (Van Leeuwen and Darriet, 2016).

When addressing these issues, one of the key challenges is strategically planning mitigation efforts and adaptation measures across and within different sectors and scales (Tol, 2005). Firstly, they act at different time periods, where for instance the beneficial effects of mitigation efforts may take many years to emerge (Meehl et

al., 2012). Secondly, despite the global extent of climate change (IPCC, 2014), the relevant factors and processes that contribute to increasing greenhouse gases or the consequences of a changing climate are local and context-dependent in nature, varying from one location to another (Mimura et al., 2014). Hence, the question of scale is central to understanding the challenges, or even the opportunities, brought by a changing climate (O'Brien et al., 2004). For viticulture, the understanding of scale and cross-scale interactions is specifically important as produced wines are the result of complex interactions between physical, biological and human factors (Van Leeuwen and Seguin, 2006), taking place at different levels and times, known as the components of terroir (OIV, 2010). The content of this article aims to provide an overview of climate-driven scales, outlining the spatial complexity and temporal dynamics in viticulture. In the context of climate change mitigation and adaptation, the article concludes with a discussion of the implications of this literature review on assessing expected impacts, understanding uncertainty and framing sustainable policies and responses in viticulture.

2. Spatial complexity in viticulture

The concept of spatial scale is fundamental to viticulture (Vaudour, 2002). At large scales, it explains wine geography and varietal distribution, while at fine scales, it accentuates the variability that exist from one location to another, enhancing wine identity and diversity (White et al., 2009). Since vine behaviour is much dependent on climate, the spatial description of viticulture closely follows its organizational structure (Asselin et al., 2001; Vaudour et Shaw, 2005), where two aspects are vital. Firstly, climate has a multi-level structure of four spatial scales with distinct features (Geiger et al., 1995): the macroclimate applying to a large geographical area (e.g. horizontal range greater than 200km); the mesoclimate corresponding to the climate of a region of a variable size, ranging from 1km to 200km; the local climate which is the climate of a smaller area; and lastly, the microclimate referring to a very small geographic unit (Figure 1). The horizontal or vertical boundaries of the four levels vary with the nature of the underlying surface (Geiger et al., 1995). For instance, the local climate has a horizontal distance of 10km in a flat area, while a characteristic distance of 1 km in a mountain area (Guyot, 1999).

Secondly, the hierarchical organization of climate must be considered, as lower spatial levels (i.e. with a finer spatial resolution) are embedded or nested within higher levels (Ackerly et al., 2010; Quéno and Bonnardot, 2014). The reason for this organization is related to the functioning of the atmosphere, knowing that the same air mass will generate similar weather patterns for an extended territory, and it is certain that at a smaller spatial entity within this territory, common climate features will be present (Geiger et al., 1995; Guyot, 1999). The higher level

therefore provides a climatic context that imposes conditions and constraints in a top-down approach towards lower levels (Wu and David, 2002). Still, local terrain effects (e.g. altitude and aspect) lead to large climate variations at smaller scales (Geiger et al., 1995; Quéno1, 2014), which means that lower levels also impose climate feedbacks that modify large-scale patterns, by following an upward or bottom-up approach. The individual characteristics of the grapevine are therefore dependent on these environmental processes, taking place at different spatial levels. As each level has its own attributes, it is important to understand what are the main influencing factors that emerge (Willis and Whittaker, 2002), while also knowing that spatial heterogeneity becomes increasingly important at finer scales (Quéno1, 2014).




Figure 1: Example of the schematic representation of the different spatial levels of viticulture for cv. Chenin blanc in the Loire Valley, France, from national to field level.

2.1. Global level

Apart from a few exceptions (e.g. Brazil, India), the main wine producing regions are roughly located between latitudes 30 to 50 degrees, in both hemispheres. This geographical distribution of wine grape varieties is shaped

by the spatial variability of large-scale macroclimatic conditions, which are strongly determined by the latitudinal differences in solar energy intercepted at the surface of the earth (Geiger et al., 1995; Ricklefs and Miller, 2005). As the equator receives more incoming radiation during the year, the average annual temperature decreases with latitude, from the equator to the poles (Guyot, 1999). Consequently, it appears that favourable macroclimate conditions for cultivating grapevines are found between annual isotherms of 10°C to 20°C (De Blij, 1983; Spellman, 1999). Schultz and Jones (2010) indicated that the geographical extension of viticulture is best represented by growing season isotherms of 12°C to 22°C. Regions with seasonal average temperatures below 12°C are unfavourable due to short growing seasons and low amounts of sunshine hours and accumulated heat (Jones et al., 2012). In these areas, the high frequency and intensity of extreme cold temperatures (e.g. winter or spring frosts) also makes grapevine growing very difficult. Conversely, in regions with mean seasonal temperatures above 22°C, grapevines are regularly exposed to very warm summers and extreme heat events. And depending on the seasonal rainfall rhythm, they may also be subject to either very arid conditions (e.g. North Africa), or very humid conditions where pest and disease pressures are particularly high (e.g. Thailand). Regarding the distribution of the most recognized cultivars (e.g. Cabernet Sauvignon), Jones (2006) also showed that the production of high quality wines from these varieties is located between isotherms 13°C and 21°C. While there are more than 6 000 wine grape varieties known (This et al., 2006; Myles et al., 2011), 13 cover more than one-third of the world's vineyard area and 33 varieties cover 50% (OIV, 2017).

2.2. Regional level

The wine sector is growing every day with emerging wine regions and markets. In response to an increasingly competitive global industry, each region seeks to differentiate itself by promoting wines with a strong identity (Hayward and Lewis, 2008; Easingwood et al., 2011). In order to maintain a premium in the market by ensuring high quality products, the cultivated grape variety and in particular its ripening period must be in relation with the surrounding environment (Jackson and Lombard, 1993). When berry maturation is reached too late, grapes remain unripe, with high acidity, low sugar contents and herbaceous flavours, whereas reached too soon, grapes are rich in sugars, low in acidity, giving place to unbalanced wines that lack in aromatic complexity (Van Leeuwen and Seguin, 2006). Since vine phenology and ripening earliness are measurable (Parker et al., 2013), berry maturation under favourable conditions is usually achieved when varieties are grown at their northern limits in the northern hemisphere, or southern limits in the southern hemisphere. In European wine regions, extended grape growing

through several centuries has allowed the emergence of the most adapted varieties in regional climates that are representative of their northern limits (Barbeau et al., 2015). For example, Pinot Noir and Chardonnay are the emblematic varieties of Burgundy wines, Cabernet Sauvignon and Merlot of Bordeaux red wines (France), Sangiovese of Tuscany red wines (Italy) and Riesling of Rheingau white wines (Germany). Conversely, in new world wine regions, one of the main challenges in defining their success and competitiveness has been to identify the most fitted grapevine varieties (Van Leeuwen and Seguin 2006). Varietal suitability is strongly related to the regional attributes of growing season accumulated heat (Tonietto and Carbonneau, 2004; Jones et al., 2012), i.e. sum of daily temperatures above a threshold where vine development is active. Still, for a given variety, cultivated in different regions but under similar heat sums, climatic properties, such as rainfall regime and thermal conditions during ripening, contribute to the unique traits of produced wines (Shaw, 2012). With this context, several studies have led to the conception of bioclimatic indices, specific to viticulture. By integrating, in one way or another, the temperature factor, there are simple or complex indices (i.e. mono- or multi-factorial), which enables to define a region's ability to produce wine, varietal suitability and vine phenology, possible environmental risks and to some extent, potential wine styles. Today, the application of these indices allows an understanding of the spatial variation in viticultural potentialities and limitations, as e.g., in Australia (Hall and Jones, 2010), Chile (Montes et al., 2012), United States (Jones et al., 2010), New Zealand (Anderson et al., 2012). Figure 2 illustrate the regional climate differences and their viticultural potentialities encountered over Europe, which are essentially due to latitudinal position and degree of continentality (e.g. rainfall patterns).




Figure 2: The average values of a) Winkler Index (WI), b) Huglin Index (HI), c) Cool Night Index (CI) and d) Dryness Index (DI), calculated at the European scale for the period from 1950-2009 (adapted from Santos et al., 2012).

2.3. Local level

Within a wine growing region, viticultural potentialities and limitations are described by local terrain features. Indeed, atmospheric boundary layer characteristics are dependent on surface conditions that modify fine-scale environmental conditions (Geiger et al., 1995). Accordingly, each wine region is composed of various local climates depending on its terrain complexity (Figure 3). Through interactions with the atmosphere and different weather patterns, topographical factors (e.g. elevation, aspect) influence heat and moisture exchange at the earth's surface, shaping topoclimates (Jacquet and Morlat, 1997; Orlandini et al., 2006; Bonnefoy et al., 2012). Besides the topographic context, soil parameters (e.g. texture, depth) and other proximity features (e.g. water mass, forest) also have significant effects on local climate or pedoclimate conditions (Guyot, 1999). The latter refers to soil moisture and temperature, which is a key factor of the earliness of grapevine growth in spring (Morlat and Jacquet, 1993). All the elements of the surrounding landscape generate a strong spatial climate variability, as for example

described in the Coteaux du Layon France (Neethling et al., 2014), Saint Emilion France (De Resséquier et al., 2016), Canelones Uruguay (Fourment et al., 2013), Uco Valley Argentina (Grassin et al., 2014), Douro Valley Portugal (Jones and Alves 2012) and Stellenbosch South Africa (Bonnardot et al., 2012). This spatial heterogeneity in local climates is a critical element in viticulture, making it possible for winegrowers to mitigate the conditions and constraints imposed by the zonal climate (Quénol, 2014). For example, air temperature generally decreases by 1°C per 150m increasing elevation, except during thermal inversions (Guyot, 1999). In areas with very warm regional climates, the effects brought by ascending in elevation are beneficial for growing wine grapes, as in the Uco Valley (1000m-1200m, Argentina), or the Central Valley of Tarija (1600m-2150m, Bolivia). At higher altitudes, vines profit from lower temperatures, allowing slower berry maturation and higher concentrations in organic acidity (Miguel-Tabares et al., 2002). The light intensity is also higher, favouring phenolic compound synthesis in red varieties. Conversely, in cool climate regions, it is worth mentioning the importance of slope orientation and inclination, the latter describing the angle between orientation and incoming sunlight. In high latitude regions of the northern hemisphere (e.g. Burgundy), south facing slopes allow for a higher amount of solar radiation reaching the surface, whereas the steeper slopes of esteemed plots (e.g. Burgundy crus) increases light and heat energy interception (Huggett, 2006; Jackson, 2008). Although steep slopes enlarge erosion risks, they enable cold air to drain away during radiative frost events, a frequent event in high latitude vineyards. Similarly, in warm and dry Mediterranean climates (e.g. Western Cape South Africa), vineyards may benefit significantly from the cooling effect brought by sea breezes (Bonnardot et al., 2005), whereas in Oceanic climates with regular rainfall patterns (e.g. Loire Valley France), soils with low water holding capacities favour grapevine earliness and wine production (Barbeau et al., 1998). Each wine growing region therefore consists of unique environmental contexts, providing winegrowers with a range of local climates and different viticultural potentialities.




Figure 3: Nested within a specific zonal climate, local climate conditions in viticultural areas are shaped by different landscape elements, which varies strongly over space (adapted from Quénol 2011).

2.4. Field level

The grapevine is a perennial crop and requires a few years to reach reproductive maturity, remaining then economically productive for many years (Orlandini et al., 2009). As the lifespan of a plot may correspond to two generations of winegrowers, management decisions prior to planting are very important. At this level, decision making is closely related to winegrowers' production objectives and the physical attributes and constraints of planting sites (Coulon-Leroy et al., 2012). To ensure sustainable productivity, winegrowers strive for appropriate measures in terms of perennial practices in order to optimise seasonal grapevine growth and fruit development. These practices vary considerably, from the choice of plant material to decisions involved in vineyard design and layout, each practice affecting microclimate conditions at field levels (Smart, 1988; Matese et al., 2014). For example, row orientation greatly affects light interception and wind velocity and as a result, canopy and fruit

temperature (Giacosa et al., 2015; Hunter et al., 2016). For vertically positioned shoots with East-West orientated rows, less direct sunlight is reach in fruits than compared to North-South orientated rows (Hunter et al., 2016). East-West orientated rows also expresses greater thermal variations between exposed and shaded leaves. These alterations in light exposure explains the greater suitability of North-South orientated rows in cool growing areas. Other notable perennial practices are the choice in planting density (Archer and Strauss, 1989, 1990), training system (Reynolds and Vanden Heuvel, 2009) and trunk height, where these practices are to some extent interconnected. For example, planting density is likely to determine the fitted training system, whereas the latter may condition trunk height. Vineyards with higher planting densities generally display improved crop yield and quality (i.e. per ha), which are mainly due to greater root competition (Jackson, 2008). Increased root competition leads to lower vegetative growth and an enhanced microclimate, while also allowing for deeper root penetration and a greater soil volume to be explored. Still, in shallow soils with low water availability, high planting densities are likely to cause severe vine water deficiencies, or conversely, excessive vegetative growth in fertile soils with high water and nutrient availability. In this context, the microclimate is strongly dependent on human decisions, which makes the spatial variability in viticulture even more complex. Winegrowers are therefore able to influence plant exposure to different climate conditions (e.g. sunlight and air temperature) and resources (e.g. soil water and nutrition). They accordingly intervene at field levels and further modify the viticultural potential of their vineyard sites, by implementing practices that attenuate the undesirable impacts of the natural environment.

3. Temporal dynamics in viticulture

The notion of temporality is also central in viticulture. In the long term, natural attributes of a vineyard site (e.g. zonal climate, topographic context) largely determines its viticultural potential, which can be expressed according to two ecophysiological variables, namely vine earliness and vine vigour (Bodin and Morlat, 2006; Coulon-Leroy et al., 2012). Perennial practices are then fixed and superimposed on natural site features and act as a functioning parameter over many years (Figure 4). In this context, these long-term practices will either enhance or modify vine earliness and vigour as conferred by the planting site. Indeed, grapevine performance is predictable depending on the timing and duration of its phenological stages (i.e. level of vine earliness) and the rhythm and amount of vegetative growth (i.e. level of vine vigour). These variables are strongly linked with vine water supply (Morlat and Jacquet, 1997, Barbeau et al., 1998), which is a key determining factor of crop productivity and fruit quality at harvest (Matthews et al., 1990; Kennedy et al., 2002; Van Leeuwen et al., 2009). As they are measurable in

time, these two ecophysiological variables are used by winegrowers to monitor achieved harvest and wine quality (Coulon-Leroy et al., 2012), depending on annual functioning parameters, namely climate variability and management practices (Neethling et al., 2017).




Figure 4: Temporal representation of long- and short-term factors affecting vine ecophysiological variables and subsequently, harvest and wine quality.

3.1. Annual climate variability

Annual climate variability, known as the vintage effect, refers to the natural variation of climate conditions between adjacent years. Given its strong temporality, i.e. between and within growing seasons, short-term climate variations are key influencing factors of seasonal fruit yield and quality (Jackson and Lombard, 1993; Van Leeuwen et al., 2004; Ubalde et al., 2007). Climate conditions (e.g. air temperature, rainfall) and associated factors (e.g. water supply) influence grapevine behaviour and characterize vine earliness and vine vigour according to their means or sums, critical thresholds or distribution during vine growing cycle (Jones et al., 2012). Warmer than normal growing season conditions results in higher heat accumulation to which generally vines respond by an earlier onset of phenological stages. Depending on the cultivated variety and its geographical position, improved vine earliness is likely to favour grape quality, as for example in high latitude vineyards, where vine earliness shifts berry ripening to a warmer period of the year (Barbeau et al., 1998). Conversely, cooler than normal growing seasons are challenging as accumulated heat is lower than culture requirements, causing delayed vine phenology and unripe grapes (e.g. low sugar contents, high acidity levels). For red varieties, insufficient amounts of heat also affect fruit phenolic and flavour ripeness (Barbeau et al., 2004). Climate thresholds, especially critical air

temperatures (e.g. during spring frosts), may also greatly affect the intended outcome of yield (Neethling et al., 2017). Other examples are episodes of extreme weather events (e.g. hail, heatwaves) or the seasonal distribution and timing of rainfall. Seasonal rainfall distribution plays a central role in vine water supply and fungal pathogen outbreaks, whereas rainfall timing is associated with vine phenology when dry conditions are crucial (e.g. during bloom). Annual climate variability is therefore a decisive factor of vine productivity (e.g. plant biomass, yield) and fruit composition. Strongly linked to berry biochemical characteristics (Cadot et al., 2012; Bindon et al., 2013), the qualitative attributes of wines (e.g. its sensory profile, aging potential) are also much climate dependent (Grifoni et al., 2006; Sadras et al., 2007; Baciocco et al., 2014). In this perspective, and in relation to its impact on the amount volume produced, annual climate variability has considerable effects on the economic viability of the wine sector (Haeger and Storchmann, 2006; Ashenfelter and Storchmann, 2016).

3.2. Annual management practices

In time, winegrowers continually deal with inter- and intra-seasonal climate variability (Neethling et al., 2017). As climate influence vine growth and development, winegrowers seek to optimize its behaviour by selecting a fitting technical itinerary for each plot (Jackson and Lombard, 1993; Hunter et al., 2010). This itinerary involves all annual practices, from winter pruning to harvest date, including the different techniques of soil, vigour and disease management. Depending on seasonal climate conditions, a vineyard plot may follow various technical itineraries, which will consequently cause different grape qualities at harvest (Coulon-Leroy et al., 2012). Likewise, the same attributes in berry qualities can be achieved through different combinations of practices and techniques. In this context, the long-term viticultural potential of a plot can be mitigated by farming strategies. For example, winegrowers are able to manage temporal variations in vine water supply through different vine inter-row soil management techniques (Wheeler and Pickering, 2003). Within a growing season, most adaptive responses generally occur at harvest or in the cellar (Neethling et al., 2017). During fruit development, winegrowers will closely follow the berry ripening process in order to adjust their harvest management decisions and pick grapes at the most adequate period (Coulon-Leroy et al., 2012). From here, winegrowers will use an array of winemaking techniques to produce a wine with a distinct structure, style, and sensory profile (Cadot et al., 2012). Still, the impacts of climate variations are not new and adapting to those conditions has always been a constant challenge faced by winegrowers. Through constant learning experiences, decision making in viticulture is an ongoing process, where winegrowers perform various forms and types of adjustments to their existing

management practices and techniques (Nicholas and Durham, 2012; Lereboullet et al., 2013; Neethling et al., 2017). To that end, the term practice describes the action exercised, while technique refers to the knowledge of exercising the activity (Landais et al., 1988). Adaptation in viticulture is an iterative process, involving many factors that assist or constrain the process of adjusting practices and techniques, and understanding this dynamic nature of viticultural and winemaking activities is crucial (Neethling et al., 2017).

4. Discussing viticulture in a changing climate

In viticulture, contemporary climate change is positioning itself as a major environmental challenge. Regional climate changes and its impacts are taking place at a substantial rate, affecting grapevine productivity and fruit quality. As temperatures are projected to continue warming, with likely changes in rainfall patterns and extreme weather events (IPCC, 2014), the findings reported in this review disclose that the territorial identity of wine growing regions may significantly evolve over time. This identity being shaped by a distinct product quality and style, which are strongly related to its geographical origin. In the short-term, as vine earliness is expected to keep improving under higher seasonal temperatures, resulting in warmer ripening periods, berry metabolism and hence wine composition and sensory properties will strongly be altered. In the long-term, varieties currently cultivated at their upper suitability threshold may accordingly become less fitted with future climate changes, where a varietal change will substantially modify the territorial product identity. While the latter has a temporal characteristic, evolving according to context, the evidence of rapid climate changes brings strong attention to the socio-economic impacts on the wine sector. With a changing climate, review findings show likewise that important geographical shifts in viticultural potentialities are likely to occur. At field to local levels, land considered limiting for viticulture in the past could develop into favourable planting sites (e.g. sites with cooler facing slopes or with larger water holding capacities). At the same time, traditional planting sites may become less adapted to continued climate trends, creating an important relocation of growing areas within wine regions, placing policy pressure on existing boundaries of geographical indications. At regional to global levels, the northern and southern limits of vine distribution are also likely to shift, an event already occurring with developing regions in Canada, Denmark and even Sweden. Conversely, with increasing vine water demands in warm and dry regions, such as in the Mediterranean Basin, the substantial loss in crop productivity may hinder the long-term economic viability of those traditional wine regions (Costa et al., 2016).

Global climate change is affecting more than just the local growing environment, as it also denotes an increase in uncertainty (Dessai et al., 2007). Indeed, the extent to which climate will evolve as a result of natural processes and human activities is unclear, and likely to remain unknown. Besides temporal uncertainties in climate estimates, the review has demonstrated that each vineyard site is unique, defined by specific natural, biological and human properties and interactions. These results support the growing and urgent need for addressing spatial uncertainties in expected vulnerability, which will vary considerably with location, relative to its exposure to extrinsic environmental changes and to the intrinsic features defining its sensitivity (e.g. site and plant characteristics) and adaptive capacity (e.g. resources, technology, skills) to such changes and impacts. Although this strong heterogeneity in natural site attributes lowers the transparency of adaptation and mitigation actions, review findings suggest that it should also constitute an important buffer in response to future climate changes, allowing producers to spatially manage expected climate changes. The review has highlighted the influence of a strong seasonal climate variability on ongoing practices and techniques. Temporal and spatial uncertainties are therefore likely to be further aggravated by the difficulty of separating the effects of long-term climate changes from natural and short-term climate variations. This may hinder the perception of climate changes and the recognition of its impacts, resulting in delayed implementations of mitigation and adaptation priorities (Neethling et al., 2017).

In response to global climate change, adjustments made in practices and techniques will need to be place-based and context-specific, as the relevant factors and processes contributing to increasing greenhouse gases or the consequences of a changing climate are local in nature (Mimura et al., 2014). To that end, many different strategies are being discussed with ongoing research projects, for example, changes in the areas of cultural practices related to vigour or soil management. Still, based on the results of this literature review, flexibility and robustness become two key elements in framing sustainable policies and responses over space and in time (Hallegatte, 2009; White et al., 2009). Flexibility refers here to the ability of a viticultural system or itinerary to adjust to changing circumstances, strengthening its resilience and lessening its vulnerability to climate change. Whereas robust strategies should allow winegrowers to manage uncertain climate impacts, by developing practices and techniques that are able to deal with various climate outcomes. Taking the example found in regions like the Douro Valley in Portugal, robust strategies can imply planting various varieties, and clones of the same variety, each developing and ripening at different stages, allowing producers to be more responsive to climate impacts. Winegrowers can also look at implementing no-regret strategies (Hallegatte, 2009), which can benefit grape and wine quality in the absence of a changing climate. For instance, adjusting rootstock varieties to actual environmental conditions and

site-specific soil properties. Goulet and Morlat (2010) illustrated that very few vineyards in the middle Loire Valley (France) are planted with the most suitable rootstock variety. As in the past with other global drivers of change (e.g. ramifications caused by Phylloxera), there is a therefore great need for policy and research to assist winegrowers. They may range from informing and educating stakeholders to providing institutional and technological support (e.g. varietal creation, flexible regulations). While winegrowers will seek to manage and adapt to the expected changes in grapevine performance and wine quality, consumers will also have to adjust. The latter will need to be educated about the issues brought about a changing climate and its impacts on the wine sector.

5. Conclusion

Through time and space, climate has always played a central role in viticulture. The literature review provided in this article has identified the role of climate, outlining the spatial complexity and temporal dynamics in grape growing and wine production. By exploring climate-driven scales, the findings are of relevance in the context of adapting viticulture to climate change, offering support to arising questions that increasingly require attention. Apart from discussing climate change impacts and uncertainties, the perspectives brought by this article argue that flexibility and robustness will be two key areas to focus on when framing sustainable policies and responses. While there are many adjustments possible, there is a need and urgency in developing strategies that are adaptable and flexible in time and over space (e.g. accounting for short-term climate variability and long-term changes), while also being planned and implemented to be more resilient to endure various climate outcomes of an inherently unpredictable future. Therefore, from climate change as a global driver to applying context-specific actions, this understanding of the spatial complexity and temporal dynamics in viticulture should provide a good conceptual and theoretical framework that can be used to address adapting viticulture to climate change.

Acknowledgements

This study was carried out with the contribution of the LIFE financial instrument of the European Union, as part of the LIFE-ADVICLIM project (LIFE13 ENV/FR/001512).

References

384 Ackerly, D.D., Loarie, S.R., Cornwell, W.K., Weiss, S.B., Hamilton, H., et al., 2010. The geography of climate
385 change: implications for conservation biogeography. *Diversity and Distributions*, 16, 476-487

386 Anderson, J.D., Jones, G.V., Tait, A., Hall, A., Trought, C.T., 2012. Analysis of viticulture region climate structure
387 and suitability in New Zealand. *J. Int. Sci. Vigne Vin*, 46(3), 149-165

388 Archer, E., Strauss, H.C., 1989. The effect of plant spacing on the water status of soil and grapevines. *S. Afr. J.*
389 *Enol. Vitic.*, 10, 49-58

390 Archer, E., Strauss, H.C., 1990. The effect of vine spacing on some physiological aspects of *Vitis vinifera* L. (cv.
391 Pinot noir). *S. Afr. J. Enol. Vitic.*, 11, 76-87

392 Ashenfelter, O., Storchmann, K., 2016. The economics of wine, weather, and climate change. *Review of*
393 *Environmental Economics and Policy*, 10, 25-46

394 Asselin, C., Barbeau, G., Morlat, R., 2001. Approche de la composante climatique à diverses échelles dans le
395 zonage viticole. *Bulletin de l'OIV.*, 883-844

396 Baciocco, K.A., Davis, R.E., Jones, G.V. 2014. Climate and Bordeaux wine quality: identifying the key factors
397 that differentiate vintages based on consensus rankings. *J. Wine Res.*, 25, 75-90

398 Barbeau, G., Morlat, R., Asselin, C., Jacquet, A., Pinard, C., 1998. Comportement du cépage Cabernet Franc dans
399 différentes terroirs du Val de Loire. Incidence de la précocité sur la composition de la vendange en année
400 climatique normale (exemple de 1988). *J. Int. Sci. Vigne. Vin*. 32(2), 69-81

401 Barbeau, G., Bournand, S., Champenois, R., Bouvet, M.H., Blin, A., Cosneau, A. 2004. Comportement de 4
402 cépages rouges du val de Loire. Relations entre les variables climatiques et la composition des baies. *J. Int.*
403 *Sci. Vigne, Vin* 38(1), 35-40

404 Barbeau, G., Neethling, E., Ollat, N., Quénot, H., Touzard, J.M., 2015. Adaptation au changement climatique en
405 agronomie viticole. *L'agronomie environnement et sociétés* 5(1), 67-75

406 Bindi, M., Fibbi, L., Gozzini, B., Orlandini, S., Miglietta, F., 1996. Modelling the impact of future climate scenarios
407 on yield and yield variability of grapevine. *Clim. Res.*, 7, 213-224

408 Bindon, K., Varela, C., Kennedy, J., Holt, H., Herderich, M., 2013. Relationships between harvest time and wine
409 composition in *Vitis vinifera* L. cv. Cabernet Sauvignon 1. *Grape and wine chemistry. Food Chem.*, 138, 1696-
410 1705

411 Bodin, F., Morlat, R., 2006. Characterization of viticultural terroirs using a simple field model based on soil depth
412 I. Validation of the water supply regime, phenology and vine vigour, in the Anjou vineyard (France). *Plant*
413 *Soil*, 281, 37-54

414 Bonnardot, V., Planchon, O., Cautenet, S., 2005. Sea breeze development under an offshore synoptic wind in the
415 South-Western Cape and implications for the Stellenbosch wine-producing area. *Theor. Appl. Climatol.* doi
416 10.1007/s00704-004-0087-y

417 Bonnardot, V., Carey, V., Madelin, M., Cautenet, S., Coetzee, Z., Quénot, H., 2012. Spatial variability of night
418 temperatures at a fine scale over the Stellenbosch wine district, South Africa. *J. Int. Sci. Vigne Vin*, 46(1), 1-
419 13

420 Bonnefoy, C., Quénot, H., Bonnardot, V., Barbeau, G., Madelin, M., et al., 2012. Temporal and spatial analyses
421 of temperature in a French wine-producing area: The Loire Valley. *Int. J. Climatol.*, 33, 1849-1862

422 Buttrose, M.S., Hale, C.R., Kleiwer, W.M., 1971. Effect of temperature on composition of Cabernet Sauvignon
423 berries. *Am. J. Enol. Vitic.*, 22, 71-75

424 Cadot, Y., Caillé, S., Thiollet-Scholtus, M., Samson, A., Barbeau, G., Cheynier, V., 2012. Characterisation of
425 typicality for wines related to terroir by conceptual and by perceptual representations. An application to red
426 wines from the Loire Valley. *Food Qual. Prefer.*, 24(1), 48-58

427 Cook, B.I., Wolkovich, E.M., 2016. Climate change decouples drought from early wine grape harvests in France.
428 *Nat. Clim. Change*, 2960. doi: 10.1038

429 Coombe, B.G., 1987. Influence of temperature on composition and quality of grapes. *Acta Hortic.*, 206, 23-35

430 Costa, J.M., Vaz, M., Escalona, J., Egipto, R., Lopes, C., Medrano, H., Chaves, M.M., 2016. Modern viticulture
431 in southern Europe: vulnerabilities and strategies for adaptation to water scarcity, *Agric. Water Manage.* 164,
432 5-18.

433 Coulon-Leroy, C., Morlat, R., Barbeau, G., Gary, C., Thiollet-Scholtus, M., 2012. The vine functioning pathway,
434 a new conceptual representation. *Sustain. Agric. Rev.*, 11, 241-264

435 De Blij, H.J. 1983. Wine: a geographic appreciation. *Progress in Physical Geography*, 38(5), 674-684

436 De Resséquier, L., Le Roux, R., Quénot, H., Van Leeuwen, C., 2016. Spatial temperature variability and
437 distribution at local scale in Saint-Emilion and Pomerol. In 'Climwine Sustainable grape and wine production
438 in the context of climate change', Bordeaux, France, 10-13 April, 2016

439 Dessai, S., O'Brien, K., Hulme, M., 2007. Editorial: on uncertainty and climate change *Global Environmental*
440 *Change*, 17(1), 1-3

441 Dry, P.R., 1988. Climate change and the Australian grape and wine industry. *Aust. Grapegrower Winemaker*, 300,
442 14-15

443 Easingwood, C., Lockshin, L., Spawton, A., 2010. The drivers of wine regionality. *J. Wine Res.*, 22(1), 19-33

444 Fourment, M., Ferrer, M., González-Neves, G., Barbeau, G., Bonnardot, V., Quénol, H., 2013. Spatial variability
 445 of temperature and grape berry composition at terroir scale in Uruguay. *Ciência y Técnica Vitícola*, 28(1), 329-
 446 334.

447 Gale, G., 2011. *Dying on the Vine: How Phylloxera Transformed Wine*. University of California Press.

448 Geiger, R., Aron, R.H., Todhunter, P., 1995. *The climate near the ground*. 5th ed., Braunschweig, Germany.

449 Giacosa, S., Marengo, F., Guidoni, S., Rolle, L., Hunter, J.J., 2015. Anthocyanin yield and skin softening during
 450 maceration, as affected by vineyard row orientation and grape ripeness of *Vitis vinifera* L. cv. Shiraz Food
 451 Chemistry, 174, 8-15

452 Goulet, E., Morlat, R., 2010. The use of surveys among wine growers in vineyards of the middle-Loire Valley
 453 (France), in relation to terroir studies. *Land Use Policy* 28, 770-782

454 Grassin, M., Quénol H., Trapeteau L., Pinson, L., Barbeau, G., et al., 2014. Variabilité spatiale du climat dans les
 455 vignobles de la bodega Alta Vista. Proceedings from the Xth International Terroir Congress, Tokaj and Eger,
 456 Hungary, 7-10 July, 2014.

457 Grifoni, D., Mancini, M., Maracchi, G., Orlandini, S., Zipoli, G., 2006. Analysis of Italian wine quality using
 458 freely available meteorological information. *Am. J. Enol. Vitic.*, 57, 339–346

459 Guyot, G., 1999. *Climatologie de l'Environnement*. Edition Dunod, Paris, France

460 Haeger, J.W., Storchmann, K., 2006. Prices of North American Pinot Noir wines: climate, craftsmanship, critics.
 461 *Agricultural Economics*, 35, 67-78

462 Hall, A., Jones, G.V., 2010. Effect of potential atmospheric warming on temperature-based indices describing
 463 Australian winegrape growing conditions. *Aust. J. Grape Wine Res.*, 15, 97-119

464 Hallegatte, S., 2009. Strategies to adapt to an uncertain climate change. *Global Environmental Change*, 19, 240-
 465 247

466 Hayward, D., Lewis, N., 2008. Regional dynamics in the globalising wine industry, the case of Marlborough, New
 467 Zealand. *Geogr. J.*, 174(2), 124-137

468 Holland, T., Smit, B., 2010. Climate change and the wine industry, current research themes and new directions. *J.*
 469 *Wine Res.*, 21, 125-136

470 Huggett, J.M., 2006. Geology and wine, a review. *Proc. Geologists' Assoc.*, 117, 239-247

471 Huglin, P., Schneider, C., 1998. *Biologie et écologie de la vigne*. Ed. Lavoisier Tec et Doc, Paris

472 Hunter, J.J., Archer, E., Volschenk, C.G., 2010. Vineyard management for environment valorisation. Proceedings
 473 from the VIIIth International Terroir Congress, Soave, Italy, 14-18 June 2010

474 Hunter, J.J., Volschenk, C.G., Zorer, R., 2016. Vineyard row orientation of *Vitis vinifera* L. cv. Shiraz/101-14
475 Mgt, climatic profiles and vine physiological status. *Agricultural and Forest Meteorology*, 228, 104-119

476 IPCC, 2014. *Climate Change 2014, synthesis report. Contribution of Working Groups I, II and III to the Fifth*
477 *Assessment Report of the Intergovernmental Panel on Climate Change. Core Writing Team, Pachauri RK,*
478 *Meyer LA eds., IPCC, Geneva, Switzerland*

479 Jackson, D.I., Lombard, P.B., 1993. Environmental and management practices affecting grape composition and
480 wine quality, a review. *Am. J. Enol. Vitic.*, 44, 409-430

481 Jackson, R.S., 2008. *Wine science, principles and applications. Ed. Academic Press, New York*

482 Jacquet, A., Morlat, R., 1997. Caractérisation de la variabilité climatique des terroirs viticoles en val de Loire.
483 Influence du paysage et des facteurs physiques du milieu. *Agronomie*, 17, 465-480

484 Jones, G.V., Davis, R.E., 2000. Climate influences on grapevine phenology, grape composition, and wine
485 production and quality for Bordeaux, France. *Am. J. Enol. Vitic.*, 51, 249-261

486 Jones, G.V., White, M.A., Cooper, O.R., Storchmann, K., 2005. Climate change and global wine quality. *Clim*
487 *Change* 73, 319-343

488 Jones, G.V., 2006. Climate and terroir: Impacts of climate variability and change on wine. In *Fine Wine and*
489 *Terroir: The Geoscience Perspective. R.W. Macqueen and L.D. Meinert (eds.), pp. 203-216. Geoscience*
490 *Canada Reprint Series No. 9. Geological Association of Canada, St. John's, Newfoundland.*

491 Jones, G.V., Duff A., Hall, A., Myers, J., 2010. Spatial analysis of climate in winegrape growing regions in the
492 western United States. *Am. J. Enol. Vitic.*, 61, 313-326

493 Jones, G.V., Webb, L.B., 2010. Climate change, viticulture, and wine, challenges and opportunities. *J. Wine Res.*,
494 21, 103-106

495 Jones, G.V., Reid, R., Vilks, A., 2012. Climate, Grapes and Wine, Structure and Suitability in a Variable and
496 Changing Climate' pp. 109-133. In *The Geography of Wine, Regions, Terroir and Techniques, Ed. Dougherty,*
497 *P.H., Springer Press, Berlin.*

498 Jones, G.V., Alves, F., 2012. Impact of climate change on wine production: a global overview and regional
499 assessment in the Douro Valley of Portugal. *Int J Glob Warming* 4:383-406

500 Kennedy, J.A., Matthews, M.A., Waterhouse, A., 2002. Effect of maturity and vine water status on grape skin and
501 wine flavonoids. *Am. J. Enol. Vitic.*, 53, 268-274

502 Kenny, G.J., Harrison, P.A., 1992. The effects of climate variability and change on grape suitability in Europe. *J.*
503 *Wine Res.*, 3, 163-183

504 Laget, F., Tondut, J., Deloire, A., Kelly, M., 2008. Climate trends in a specific Mediterranean viticultural area
505 between 1950 and 2006. *J. Int. Sci. Vigne*, 42, 113-123.

506 Landais, E., Deffontaines, J.P., Benoit, M., 1988. Les pratiques des agriculteurs. Point de vue sur un nouveau
507 courant de la recherche agronomique. In *Études Rurales*, 109, 125-158

508 Lereboullet, A.L., Beltrando, G., Bardsley, D.K., 2013. Socio-ecological adaptation to climate change, a
509 comparative case study from the Mediterranean wine industry in France and Australia. *Agric. Ecosyst.*
510 *Environ.*, 164, 273-285

511 Matese, A., Crisci, A., Di Gennaro, S.F., Primicerio, J., Tomasi, D., Marcuzzo, P., Guidoni, S., 2014. Spatial
512 variability of meteorological conditions at different scales in viticulture *Agric. For. Meteorol.*, 159-167

513 Matthews, M.A., Ishii, R., Anderson, M.M., O'Mahony, M., 1990. Dependence of wine sensory attributes on vine
514 water status. *J. Sci. Food. Agric.*, 51(3), 321–335

515 Meehl, G.A., Hu, A., Tebaldi, C., Arblaster, J.M., Washington, W.M. et al., 2012. Relative outcomes of climate
516 change mitigation related to global temperature versus sealevel rise. *Nat. Clim. Change*, 2, 576–580

517 Meloni, G., Swinnen, J., 2014. The rise and fall of the world's largest wine exporter - and its institutional legacy
518 *J. Wine Econ.*, 9(1), 3-33

519 Miguel-Tabares, J.A., Martin-Luis, B., Carrillo-Lopez, M., Diaz-Diaz, E., Darias-Martin, J., 2002. Effect of
520 altitude on the wine-making potential of listan negro and ruby carbernet cultivars in the south of Tenerife
521 Island. *J. Int. Sci. Vigne Vin*, 36(4), 185-194

522 Mimura, N., Pulwarty, R., Duc, D., Elshinnawy, I., Redsteer, M.H., et al., 2014. Adaptation planning and
523 implementation. In, IPCC, Climate change 2014, Impacts, Adaptation, and Vulnerability. Part A, Global and
524 Sectoral Aspects. Working Group II Contribution to the Fifth Assessment Report of the Intergovernmental
525 Panel on Climate Change. Cambridge University Press, New York, NY, USA, pp. 869-898.

526 Montes, C., Perez-Quezada, J.F., Peña-Neira, A., Tonietto, J., 2012. Climatic potential for viticulture in Central
527 Chile. *Aust. J. Grape Wine Res.*, 18, 20-28

528 Mora, C., Fraizier, AG., Longman, RJ., Dacks, RS., Walton, M.M., et al., 2013. The projected timing of climate
529 departure from recent variability. *Nature*, 502:183-188

530 Morlat, R., Jacquet, A., 1993. The soil effects on the grapevine root system in several vineyards of the Loire valley
531 France. *Vitis*, 32, 35-42

532 Morlat R., Jacquet A., Asselin, C., 1997. Variabilité de la précocité de la vigne en Val de Loire : Rôle du terroir et
533 du millésime., conséquences sur la composition de la baie. *Revue française d'œnologie*, juillet, 165, 11-22

534 Myles, S., Boyko, AR., Owens, CL., Brown, PJ., Grassi, F., et al., 2011. Genetic structure and domestication
535 history of the grape. *PNAS*, 108, 3530-3535

536 Neethling, E., Barbeau, G., Bonnefoy, C., Quénel, H 2012. Change in climate and berry composition for grapevine
537 varieties cultivated in the Loire Valley. *Clim Res* 53: 89-101

538 Neethling, E., Petitjean, T., Barbeau, G., Quénel, H., 2014. Impacts of environmental variability and viticultural
539 practices on grapevine behaviour at terroir scales. *Proceedings from the Xth International Terroir Congress*,
540 Tokaj and Eger, Hungary, 7-10 July, 2014.

541 Neethling, E., Petitjean, T., Quénel, H., Barbeau, G., 2017. Assessing local climate vulnerability and winegrowers'
542 adaptive processes in the context of climate change. *Mitig. Adapt. Strateg. Glob. Change.*, 22, 777-803

543 Nicholas, KA., Durham, W.H., 2012. Farm-scale adaptation and vulnerability to environmental stresses: insights
544 from winegrowing in Northern California. *Glob. Environ. Ch.*, 22, 483-494

545 O'Brien, K., Sygna, L., Haugen, J.E., 2004. Vulnerable or resilient? A multi-scale assessment of climate impacts
546 and vulnerability in Norway. *Clim. Change*, 64,193-225

547 OIV, 2010. Definition of vitivinicultural terroir. Resolution OIV/VITI 333/2010. Retrieved from:
548 <http://www.oiv.int/public/medias/379/viti-2010-1-en.pdf>

549 OIV, 2018. OIV Statistical Report on World Vitiviniculture. Retrieved from: [http://www.oiv.int/en/oiv-life/oiv-](http://www.oiv.int/en/oiv-life/oiv-2018-report-on-the-world-vitivinicultural-situation)
550 [2018-report-on-the-world-vitivinicultural-situation](http://www.oiv.int/en/oiv-life/oiv-2018-report-on-the-world-vitivinicultural-situation)

551 OIV, 2017. Global Economic Vitiviniculture data. Retrieved from: [http://www.oiv.int/public/medias/5686/ptconj-](http://www.oiv.int/public/medias/5686/ptconj-octobre2017-en.pdf)
552 [octobre2017-en.pdf](http://www.oiv.int/public/medias/5686/ptconj-octobre2017-en.pdf)

553 Ollat, N., Touzard J.M., Van Leeuwen, C., 2016. Climate Change Impacts and Adaptations: New Challenges for
554 the Wine Industry. *Journal of Wine Economics*, 11, 139-149

555 Orduna, R.M., 2010. Climate change associated effects on grape and wine quality and production. *Food Res. Int.*
556 43:1844–1855

557 Orlandini, S., Bindi, M., Howden, M., 2009. Plant biometeorology and adaptation. In: Ebi KL., Burton I.,
558 McGregor GR eds. *Biometeorology for adaptation to climate variability and change*. Ed. Springer, Dordrecht,
559 107–129

560 Orlandini, S., Marta, AD., Mancini, M., 2006. The agroclimatic analysis at farm scale. *Meteorol. Appl.*, 87-93

561 Parker, A., Garcia de Cortázar, I., Chuine I., Barbeau, G., Bois, B., et al., 2013. Classification of varieties for their
562 timing of flowering and véraison using a modelling approach: a case study for the grapevine species *Vitis*
563 *vinifera* L. *Agric. For. Meteorol.* 180,249–264

564 QuénoI, H., 2011. Observation et modélisation spatiale du climat aux échelles fines dans un contexte de
 565 changement climatique. Habilitation à Dirigée des Recherches. Université Rennes 2
 566 QuénoI, H., 2014. Changement climatique et terroirs viticoles. Lavoisier Tec et Doc, Paris
 567 QuénoI, H., Bonnardot, V., 2014. A multi-scale climatic analysis of viticultural terroirs in the context of climate
 568 change: the « TERADCLIM » project. J. Sci. Vigne Vin., Special Issue Laccave, 25-34
 569 Ramos, M.C., Jones, G.V., Yuste, J., 2015. Spatial and temporal variability of cv. Tempranillo phenology and
 570 grape quality within the Ribera del Duero DO Spain. and relationships with climate. Int. J. Biometeorol. DOI:
 571 10.1007/s00484-015-0992-z
 572 Real, A.C., Borges, J., Cabral, J.S., Jones, G.V., 2017. A climatology of vintage Port quality. Int. J. Climatol., 37,
 573 3798–3809.
 574 Reynolds, A.G., Vanden Heuvel, J.E., 2006. Influence of grapevine training systems on vine growth and fruit
 575 composition: a review. Am. J. Enol. Vitic., 60, 251-268
 576 Ricklefs, R., Miller, G., 2005. Ecologie. De Boeck., Bruxelles
 577 Sadras, V.O., Soar, C.J., Petrie, P.R., 2007. Quantification of time trends in vintage scores and their variability for
 578 major wine regions of Australia. Aust. J. Grape Wine Res., 13, 117-123
 579 Salinger, M.J., Baldi, M., Grifoni, D., Jones, G., Bartolini, G., et al., 2015. Seasonal differences in climate in the
 580 Chianti region of Tuscany and the relationship to vintage wine quality. Int. J. Biometeorol., 59, 1799-1811
 581 Santos, J.A., Malheiro, A.C., Pinto, J.G., Jones, G.V., 2012. Macroclimate and viticultural zoning in Europe:
 582 observed trends and atmospheric forcing. Clim. Res., 51, 89-103
 583 Schultz, H.R., Jones, G.V., 2010. Climate induced historic and future changes in viticulture. J. Wine Res., 21, 137-
 584 145
 585 Shaw, T.B., 2012. A climatic analysis of wine regions growing pinot noir. J. Wine Res., 23(3), 203-228
 586 Smart, R.E., 1988. Shoot spacing and canopy light microclimate. Am. J. Enol. Vitic., 39, 325-333
 587 Smart, R.E., 1989. Climate change and the New Zealand wine industry. Prospects for the third millennium. Wine
 588 Ind. J., February 8-11
 589 Spellman, G., 1999. Wine, weather and climate. Weather, 54(8), 230-239
 590 This, P., Lacombe, T., Thomas, M.R., 2006. Historical origins and genetic diversity of wine grapes. Trends Genet,
 591 22, 511-519
 592 Tol, R.S.J., 2005. Adaptation and mitigation: trade-offs in substance and methods. Environ. Sci. Policy, 8, 572-
 593 578

594 Tomasi, D., Jones, G.V., Giust, M., Lovat, L., Gaiotti, F., 2011. Grapevine phenology and climate change:
595 relationships and trends in the Veneto region of Italy for 1964–2009. *Am. J. Enol. Vitic.*, 62, 329-339

596 Tonietto, J., Carbonneau, A., 2004. A multicriteria climatic classification system for grapegrowing regions
597 worldwide. *Agr. Forest. Meteorol.*, 124, 81-97

598 Ubalde, J.M., Sort, X., Zayas, A., Poch, R.M., 2010. Effects of Soil and Climatic Conditions on Grape Ripening
599 and Wine Quality of Cabernet Sauvignon, *Journal of Wine Research*, 21(1), 1-17

600 Van Leeuwen, C., Friant, P., Xavier, C., Tregoat, O., Koundouras, S., Dubourdieu, D., 2004. Influence of climate,
601 soil, and cultivar on terroir. *Am. J. Enol. Vitic.*, 55, 207-217

602 Van Leeuwen, C., Seguin, G., 2006. The concept of terroir in viticulture. *J. Wine Res.*, 17:1-10

603 Van Leeuwen, C., Trégoat, O., Choné, X., Bois, B., Pernet, D., Gaudillère, J.P., 2009. Vine water status is a key
604 factor in grape ripening and vintage quality for red Bordeaux wine. How can it be accessed for vineyard
605 management purposes. *J. Int. Sci. Vigne Vin*, 43(3), 121-134

606 Van Leeuwen, C., Darriet P., 2016. The impact of climate change on viticulture and wine quality. *Journal of Wine*
607 *Economics*, 11(1), 150-167

608 Vaudour, E., 2002. The quality of grapes and wine in relation to geography: Notions of terroir at various scales. *J.*
609 *Wine Res.*, 13(2), 117-141

610 Vaudour, E., Shaw, A.B., 2005. A worldwide perspective on viticultural zoning. *S. Afr. J. Enol. Vitic.*, 26(2), 106-
611 115

612 Webb, L.B., Whetton, P.H., Bhend, J., Darbyshire, R., Briggs, P.R., et al., 2012. Earlier wine-grape ripening driven
613 by climatic warming and drying and management practices. *Nat. Clim. Change*, 2, 259-264

614 Wheeler, S.J., Pickering, G.J., 2003. Optimizing grape quality through soil management practices. *Food Agric.*
615 *Environ.*, 1(2), 190-197

616 White, M.A., Whalen, P., Jones, G.V., 2009. Land and wine. *Nat. Geosci.*, 2, 82–84

617 Willis, K.J., Whittaker, R.J., 2002. Species diversity-scale matters. *Science*, 295, 1245-1248

618 Wu, J., David, J., 2002. A spatially explicit hierarchical approach to modeling complex ecological systems: theory
619 and applications. *Ecological Modelling*, 153, 7-26