



HAL
open science

Formation of methyl radicals derived from cumene hydroperoxide in reconstructed human epidermis: an EPR spin trapping confirmation by using 13 C-substitution

Fatma Sahli, Amélie Godard, Bertrand Vilen, Jean-Pierre Lepoittevin, Elena Giménez-Arnau

► To cite this version:

Fatma Sahli, Amélie Godard, Bertrand Vilen, Jean-Pierre Lepoittevin, Elena Giménez-Arnau. Formation of methyl radicals derived from cumene hydroperoxide in reconstructed human epidermis: an EPR spin trapping confirmation by using 13 C-substitution. *Free Radical Research*, 2019, pp.1-11. 10.1080/10715762.2019.1624741 . hal-02157705

HAL Id: hal-02157705

<https://hal.science/hal-02157705v1>

Submitted on 23 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Formation of methyl radicals derived from cumene hydroperoxide in reconstructed human epidermis: an EPR spin trapping confirmation by using ¹³C-substitution

Journal:	<i>Free Radical Research</i>
Manuscript ID	GFRR-OM-2019-0062.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Sahli, Fatma; University of Strasbourg, Dermatochemistry, CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal Godard, Amélie; University of Strasbourg, Dermatochemistry, CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal Vileno, Bertrand; University of Strasbourg, POMAM, CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal Lepoittevin, Jean-Pierre; University of Strasbourg, Dermatochemistry, CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal Gimenez-Arnau, Elena; University of Strasbourg, Dermatochemistry, CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal
Keywords:	cumene hydroperoxide, skin sensitizers, ¹³ C-substitution, EPR spin-trapping spectroscopy, reconstructed human epidermis

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Formation of methyl radicals derived from cumene hydroperoxide in reconstructed**
6
7 **human epidermis: an EPR spin trapping confirmation by using ^{13}C -substitution**
8
9

10
11
12 Fatma Sahli,^a Amélie Godard,^a Bertrand Vileno,^b Jean-Pierre Lepoittevin,^a Elena
13
14 Giménez-Arnau^{a,*}
15
16
17
18
19
20

21 ^a Dermatochemistry Laboratory, University of Strasbourg-CNRS UMR 7177, Institut le Bel, 4 Rue
22
23 Blaise Pascal, 67081 Strasbourg, France
24
25

26 ^b POMAM Laboratory, University of Strasbourg-CNRS UMR 7177, Institut le Bel, 4 Rue Blaise
27
28 Pascal, 67081 Strasbourg, France, and French EPR Federation of Research (REseau NAational de Rpe
29
30 interDisciplinaire (RENARD, Fédération IR-RPE CNRS #3443)
31
32
33
34

35 **Conflicts of interest:** none
36
37
38
39

40 **Corresponding author:** Elena Giménez-Arnau, Dermatochemistry Laboratory, University of
41
42 Strasbourg-CNRS UMR 7177, Institut le Bel, 4 Rue Blaise Pascal, 67081 Strasbourg, France. Phone:
43
44 +33 3 68 85 15 25, E-mail: egimenez@unistra.fr
45

46 **ORCID:**

47
48 Bertrand Vileno: 0000-0001-7527-2068

49
50 Elena Giménez-Arnau: 0000-0001-8275-5447
51
52
53
54
55
56
57
58
59
60

Abstract

1
2
3
4 Dermal exposure to cumene hydroperoxide (CumOOH) during manufacturing processes is a
5 toxicological issue for the industry. Its genotoxicity, mutagenic action, ability to promote skin tumor,
6 capacity to induce epidermal hyperplasia and aptitude to induce allergic and irritant skin contact
7 dermatitis are well known. These toxic effects appear to be mediated through the activation to free
8 radical species such as hydroxyl, alkoxy and alkyl radicals characterized basically by electron
9 paramagnetic resonance (EPR) and spin-trapping (ST) techniques. To be a skin sensitizer CumOOH
10 needs to covalently bind to skin proteins in the epidermis to form the antigenic entity triggering the
11 immunotoxic reaction. Cleavage of the O-O bond allows formation of unstable CumO[•]/CumOO[•]
12 radicals rearranging to longer half-life specific carbon-centered radicals R[•] proposed to be at the origin
13 of the antigen formation. Nevertheless, it is not still clear which R[•] are precisely formed in the
14 epidermis and thus involved in the sensitization process. The aim of this work was to elucidate in
15 conditions closer to real-life sensitization which specific R[•] are formed in a 3D reconstructed human
16 epidermis (RHE) model by using ¹³C-substituted CumOOH at carbon positions precursors of
17 potentially reactive radicals and EPR-ST. We demonstrated that **most probably** methyl radicals derived
18 from β-scission of CumO[•] radicals occur in RHE through a one-electron reductive pathway **suggesting**
19 that these **could** be involved in the antigen formation inducing skin sensitization. We also describe a
20 coupling between nitroxide radicals and β position ¹³C atoms that could be of an added value to the
21 very few examples existing for the coupling of radicals with ¹³C atoms.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Keywords:** cumene hydroperoxide, skin sensitizers, ¹³C-substitution, EPR spin-trapping spectroscopy,
44 reconstructed human epidermis
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Cumene hydroperoxide (CumOOH) is produced mainly through the oxidation of cumene in the so-called “cumene process”, an industrial procedure for synthesizing phenol and acetone from benzene and propylene based on the acid-catalyzed Hock rearrangement [1]. CumOOH is used as catalyst, curing agent, and initiator in the manufacture of polymers and fiber-reinforced plastics. Plus, it is a strong oxidizing agent making it an important intermediate in the fabrication of epoxy and polyester resin coatings. However, dermal exposure to CumOOH during manufacturing processes is an important toxicological issue for the industry, currently lacking of regulatory guidelines for occupational exposure limits [2]. CumOOH genotoxicity, mutagenic action and ability to promote skin tumor and induce epidermal hyperplasia have been described [3-5]. CumOOH is also known to cause allergic contact dermatitis (ACD) and has been reported as a significant sensitizer in guinea pigs, with positive specific responses reported in allergy cross-reactivity studies with allergenic cyclohexene hydroperoxides [6]. These toxic effects are believed to be mediated through the activation of the hydroperoxide to radical species such as hydroxyl (HO[•]), alkoxy (CumO[•]) and alkyl (R[•]) radicals.

Electron paramagnetic resonance (EPR) together with spin-trapping (ST) is a powerful combination for the characterization of transient radicals in chemical and biological systems [7]. Basically, the formed radical (either too short-lived or of too low concentration to be directly detected) reacts with a diamagnetic reagent (spin-trap) to form a more persistent radical (spin-adduct), whose EPR signature can help the characterization of the trapped transient reactive species. EPR-ST has been used for dermatological purposes mainly for the detection of free radicals in UV-irradiated skin, melanoma investigation and *in vivo* EPR imaging directly in skin of human volunteers [8-10]. EPR-ST has also been used to characterize radicals in isolated mouse keratinocytes and in murine skin treated with CumOOH [11-13]. In addition, EPR-ST proved the formation of *in vivo* lipid-derived free radicals triggered by CumOOH when mouse skin was exposed causing severe oxidative stress [14,15]. To understand ACD mechanisms for risk assessment purposes, we have proved that EPR-ST is also a

1
2 powerful technique for the detection and identification of radical species derived from sensitizing
3 hydroperoxides (ROOHs) present in consumer products [16-18]. To be skin sensitizers, ROOHs need
4 to covalently bind *via* radical processes to skin proteins after having penetrated the epidermis, to form
5
6 this way the antigenic entity trigger of the immunotoxic process. As for the CumOOH tumor promoting
7
8 action, initial cleavage of the O-O bond forms unstable RO•/ROO• radicals that further rearrange to
9
10 longer half-life carbon centered radicals R• proposed to be at the origin of the antigen formation
11
12 [15,19,20]. Still, it is not still clear which R• are precisely formed in the epidermis and thus involved in
13
14 the sensitization processes.
15
16
17
18
19

20 We have recently reported an EPR-ST methodology allowing to investigate the formation of free
21 radicals issued from skin allergens in a reconstructed human epidermis (RHE) 3D model, thus much
22 closer to what may happen *in vivo*, by using CumOOH as proof of concept [21]. RHE are multi-layered
23 keratinocyte cultures, major cell type in the epidermis playing a key role in skin inflammatory
24 reactions. We disclosed that in the EpiSkin™ RHE model carbon radicals are mainly formed, both by
25 incubating the assay medium and by topical application with CumOOH. Previous *in vitro* EPR-ST
26 studies reported in keratinocytes cultures suggested these carbon radicals as being methyl radicals
27 issued from β -scission fragmentation of tertiary alkoxy radicals (CumO•) formed through cleavage of
28 the O-O bond (Scheme 1) [10]. Unlike keratinocytes cultures isolated from skin biopsies, EpiSkin™
29 RHE is an *in vitro* reconstructed human epidermis histologically similar to *in vivo* human epidermis. It
30 is a complex 3D structure including a *stratum corneum* that allows assessing skin penetration concerns
31 with phospholipids and ceramides content influencing the RHE barrier properties [22]. The aim of the
32 present study was to elucidate, in the skin allergy context and in conditions closer to real-life
33 sensitization, if methyl radicals are also formed in EpiSkin™ RHE by using ¹³C-substituted CumOOH
34 derivatives **1** (¹³C-MeCumOOH) and **2** ((¹³C-Me)₂CumOOH) (Figure 1), and thus be able to suggest
35 their participation in the formation of the immunogenic complex inducing ACD.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and methods

Synthesis of ^{13}C -substituted CumOOHs

^{13}C -MeCumOOH **1** and (^{13}C -Me) $_2$ CumOOH **2** were synthesized ^{13}C -substituted at the methyl positions following the synthetic pathway shown in Scheme 2.

Chemicals and instrumentation

Starting materials were obtained from Sigma-Aldrich (Saint Quentin Fallavier, France) and used as received. ^{13}C -Labeled reagents and deuterated solvents were purchased from Euriso-Top (Saint Aubin, France). Air-/moisture-sensitive reactions were conducted in flame-dried glassware under an atmosphere of dry argon. Solvents were dried on alumina, under argon, using a Glass Technology GTS 100 device. Reactions were followed by thin layer chromatography (TLC) on 0.25 mm silica gel plates (60F $_{254}$; Merck, Darmstadt, Germany). After migration, the plates were inspected under UV light (254 nm) or sprayed with a solution containing phosphomolybdic acid (5 g), cerium (IV) sulfate (2 g) and sulfuric acid (12 mL) in water (188 mL), followed by heating. Column chromatography purifications were performed on silica gel 60 (Merck, Geduran $^{\text{®}}$, 40–63 μm). Neutralized silica was prepared by adding to a homogenous water solution of silica gel 60 a saturated solution of NaHCO_3 until pH about 8. After decantation the silica precipitate was washed with water to reach a pH of 7, filtered then dried in a oven for at least 24 h. ^1H and ^{13}C NMR spectra were recorded on a Bruker Avance 500 spectrometer at 500 MHz and 125 MHz, respectively. Chemical shifts (δ) are reported in ppm and are indirectly referenced to TMS via the solvent signal (CDCl_3 : $\delta^1\text{H} = 7.26$, $\delta^{13}\text{C} = 77.16$). Spectra multiplicities are denoted as d (doublet), dd (doublet of doublets), t (triplet) and m (multiplet).

Synthesis of 1-(^{13}C)acetophenone (4)

To a suspension of aluminum chloride (2.07 g, 15.5 mmol, 1.4 eq) in anhydrous benzene (6 mL) at 0°C , 2-(^{13}C)acetyl chloride was added drop wise (0.8 mL, 11.07 mmol, 1 eq) under argon atmosphere.

Once the addition completed, the mixture was stirred for 17 h at room temperature. After that time, the mixture was placed into an ice bath and distilled water (6 mL) was added drop wise until it became white, followed by addition of HCl 2 M (12 mL). The two phases were separated, and the aqueous phase was extracted with diethyl ether (4 × 40 mL). The organic phases were combined, washed with brine, dried over MgSO₄, filtered and concentrated to obtain 1-(¹³C)acetophenone (**4**) as a yellow oil (1.04 g, 8.58 mmol, 77 % yield). CAS Registry Number: [71777-36-1]. ¹H NMR (CDCl₃) δ 2.61 (d, 3H, ¹J_{H1-C1} = 127.6 Hz, H1), 7.44-7.50 (m, 2H, H5 and H7), 7.54-7.60 (m, 1H, H6), 7.95-7.99 (m, 2H, H4 and H8); ¹³C NMR (CDCl₃) δ 26.8 (C1), 128.4 (2C, C5 and C7), 128.7 (2C, C4 and C8), 133.2 (C6), 137.1 (d, ²J_{C3-C1} = 13.5 Hz, C3), 198.4 (d, ¹J_{C2-C1} = 42.5 Hz, C2).

Synthesis of 1-(¹³C)2-phenylpropan-2-ol (5)

In a two-necked flask under argon 1-(¹³C)acetophenone (**4**) (450 mg, 3.71 mmol, 1 eq) was dissolved in 6 mL of anhydrous tetrahydrofuran at -78 °C. Methyllithium (2.8 mL, 4.46 mmol, 1.6 M in Et₂O, 1.2 eq) was added drop by drop and the mixture was stirred at -78 °C for 21 h then at room temperature for 5 h. Ammonium chloride was added (7 mL) and the solution was extracted with diethyl ether (4 × 20 mL). Organic phases were combined, dried over MgSO₄, filtered and concentrated under reduce pressure to give an orange oil. The crude product was purified by chromatography on silica gel (pentane/EtOAc 9/1 then 5/5) to afford 1-(¹³C)2-phenylpropan-2-ol (**5**) as a yellow oil (269.4 mg, 1.96 mmol, 53 % yield). ¹H NMR (CDCl₃) δ 1.59 (d, 3H, ¹J_{H1-C1} = 126.5 Hz, H1), 1.59 (d, 3H, ³J_{H3-C1} = 4.1 Hz, H3), 7.24-7.27 (m, 1H, H7), 7.34-7.37 (m, 2H, H6 and H8), 7.49-7.51 (m, 2H, H5 and H9); ¹³C NMR (CDCl₃) δ 31.9 (2C, C1 and C3), 72.7 (d, ¹J_{C2-C1} = 39.3 Hz, C2), 124.5 (2C, C5 and C9), 126.8 (C7), 128.4 (2C, C6 and C8), 149.2 (C-4).

Synthesis of 1-(¹³C)cumene hydroperoxide (1)

To an aqueous solution of hydrogen peroxide (30 %, 12.7 mL) at 0°C, one drop of concentrated sulfuric acid was added. After 30 min stirring, 1-(¹³C)2-phenylpropan-2-ol (**5**) (268.1 mg, 1.95 mmol, 1

1
2 eq) was added and the reaction mixture was vigorously stirred at 0°C during 4 h, followed by
3
4 extraction with ethyl acetate (3 × 10 mL). The organic layers were combined, dried over MgSO₄,
5
6 filtered, and concentrated under reduce pressure. The crude product was purified by flash
7
8 chromatography on neutralized silica gel (pentane/EtOAc 85/15 then 50/50) to obtain 1-(¹³C)cumene
9
10 hydroperoxide (**1**) as an orange oil (136.0 mg, 0.88 mmol, 32 % yield). ¹H NMR (CDCl₃) δ 1.62 (d,
11
12 3H, ³J_{H3-C1} = 4.4 Hz, H3), 1.62 (d, 3H, ¹J_{H1-C1} = 127.7 Hz, H1), 7.29-7.34 (m, 2 × 1H, H6 and H8),
13
14 7.38-7.41 (m, 1H, H7), 7.47-7.49 (m, 2 × 1H, H5 and H9); ¹³C NMR (CDCl₃) δ 26.2 (2C, C1 and C3),
15
16 84.1 (d, ¹J_{C2-C1} = 39.9 Hz, C2), 125.5 (2C, C5 and C9), 127.6 (C7), 128.7 (2C, C6 and C8), 144.7 (C4).
17
18
19
20

21 *Synthesis of 1,3-(¹³C)2-phenylpropan-2-ol (6)*

22
23 Previous synthesis of (¹³C)methylmagnesium iodide: in a 50 mL two-necked flask under argon,
24
25 magnesium (305 mg, 12.53 mmol, 1.2 eq) was flame-dried before to add a spatula tip of iodine and
26
27 anhydrous diethyl ether (10 mL). (¹³C)Methyl iodide (0.65 mL, 10.44 mmol, 1 eq) was then added to
28
29 the reaction mixture heated slightly with a hairdryer to initiate the reaction. Diethyl ether was added
30
31 (10 mL) and the reaction heated again slightly. The reaction mixture was stirred at room temperature
32
33 during 24 h. A solution of (¹³C)methylmagnesium iodide 0.5 M in diethyl ether was in this way
34
35 obtained.
36
37
38

39
40 In a two-necked flask under argon, zinc chloride (79.1 mg, 0.58 mmol, 0.2 eq) was placed and
41
42 dried under reduced pressure. Lithium chloride (295.0 mg, 6.94 mmol, 2.4 eq) was added and the flask
43
44 was again dried under reduced pressure. To the mixture, (trimethylsilyl)methylmagnesium chloride
45
46 (2.30 mL, 1.2 mmol, 0.4 eq) was added drop wise at room temperature and the reaction was stirred for
47
48 15 min. (¹³C)Methylmagnesium iodide (10 mL, 4.9 mmol, 0.5 M in diethyl ether, 1.7 eq) was then
49
50 added drop wise and the reaction stirred for 45 min and then cooled to 0°C. 1-(¹³C)Acetophenone (**4**)
51
52 (350 mg, 2.89 mmol, 1 eq) was added during 1 h with the use of a syringe pump and the reaction was
53
54 stirred for 3 h at 0°C. After this time, the reaction mixture was quenched with ammonium chloride (10
55
56
57
58
59
60

mL), extracted with ethyl acetate (3×10 mL) and washed with brine (10 mL). The organic phases were combined and dried over MgSO_4 , filtered and concentrated under reduced pressure to obtain an orange oil. The crude product was purified by chromatography on silica gel (pentane/EtOAc 92/8) and 1,3- ^{13}C 2-phenylpropan-2-ol (**6**) was obtained as a yellow oil (303.7 mg, 2.19 mmol, 76% yield). ^1H NMR (CDCl_3) δ 1.59 (dd, 3H, $^1J_{\text{H1/3-C1/3}} = 121.3$ Hz, $^3J_{\text{H1/3-C1/3}} = 1.7$ Hz, H1 or H3), 1.59 (d, 3H, $^1J_{\text{H1/3-C1/3}} = 130.7$ Hz, H-1 or H-3), 7.25-7.28 (m, 1H, H7), 7.34-7.37 (m, 2H, H6 and H8), 7.50-7.51 (m, 2H, H5 and H9); ^{13}C NMR (CDCl_3) δ 31.8 (2C, C1 and C3), 74.0 (t, $^1J_{\text{C2-C1/3}} = 38.9$ Hz, C2), 124.5 (2C, C5 and C9), 126.8 (C7), 128.3 (2C, C6 and C8), 149.2 (C4).

Synthesis of 1,3- ^{13}C cumene hydroperoxide (2)

To an aqueous solution of hydrogen peroxide (30 %, 7.5 mL) was added at 0°C one drop of concentrated sulfuric acid. After 30 min, 1,3- ^{13}C 2-phenylpropan-2-ol (**6**) (155.9 mg, 1.13 mmol, 1 eq) was added and the reaction mixture was vigorously stirred at 0°C during 4 h, followed by extraction with ethyl acetate (3×10 mL). The organic layers were combined, dried over MgSO_4 , filtered, and concentrated under reduce pressure. The crude product was purified by flash chromatography on neutralized silica gel (pentane/EtOAc 5/1 and 1/1) to obtain 1,3- ^{13}C cumene hydroperoxide (**2**) as an orange oil (36.7 mg, 0.24 mmol, 21 % yield). ^1H NMR (CDCl_3) δ 1.62 (d, 3H, $^1J_{\text{H1/3-C1/3}} = 123.8$ Hz, H1 or H3), 1.62 (d, 3H, $^1J_{\text{H1/3-C1/3}} = 132.1$ Hz, H1 or H3), 7.29-7.32 (m, 1H, H7), 7.37-7.41 (m, 2H, H6 and H8), 7.47-7.49 (m, 2H, H5 and H9); ^{13}C NMR (CDCl_3) δ 26.2 (2C, C1 and C3), 84.1 (t, $^1J_{\text{C2-C1/3}} = 39.9$ Hz, C2), 125.5 (2C, C5 and C9), 127.6 (C7), 128.7 (2C, C6 and C8), 144.7 (C4).

EPR studies in solution

CumOOH was purchased from Sigma-Aldrich (Saint-Quentin Fallavier, France) and used as received. DEPMPO was synthesized as reported in the literature [23]. HEPES (≥ 99.5 %), ferrous sulfate heptahydrate $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ and acetonitrile CH_3CN (99.8%) were acquired from Sigma-Aldrich (Saint-Quentin Fallavier, France). Aqueous solutions were prepared with deionized water. HEPES buffer

1
2 solution (10 mM, pH 6.8) was prepared by dissolving 1.19 g HEPES in 400 mL deionized water, with
3
4 additional 4 g NaCl and 0.1 g KCl. To attain pH 6.8 NaOH pellets were added. If the pH went too high,
5
6 it was lowered back by carefully adding HCl (2 M) until pH remained stable to 6.8. Deionized water
7
8 was added for a final volume of 500 mL. Stock solutions were prepared for CumOOH, ^{13}C -
9
10 MeCumOOH and $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ (10 mM, HEPES/ CH_3CN 9/1), DEPMPO (100 mM in HEPES)
11
12 and $\text{FeSO}_4\cdot 7\text{H}_2\text{O}$ (10 mM in HEPES). 12.5 μL of DEPMPO solution were mixed with $\text{FeSO}_4\cdot 7\text{H}_2\text{O}$
13
14 (0.5 μL for a final 0.1 mM concentration in the reaction mixture), 5 μL of the hydroperoxide solution
15
16 added and final volume completed to 50 μL . This way, final concentrations in the reaction mixture
17
18 were 25 mM spin-trap and 0.54 mM hydroperoxide. The reaction mixture was subjected to stirring and
19
20 further introduced into the EPR quartz capillary tube.
21
22
23

24 25 ***EPR studies in reconstructed human epidermis***

26
27 EpiSkinTM (Lyon, France; 0.38 cm²) is a RHE from normal human keratinocytes cultured for 13 days
28
29 on a collagen matrix at the air-liquid interface. Immediately after arrival in the laboratory, the RHE
30
31 were removed from the agarose-nutrient solution in the shipping multiwell plate under a sterile airflow.
32
33 They were placed in a plate in which each well was previously filled with 2 mL EpiSkinTM
34
35 maintenance or growth medium at room temperature. Samples were placed in the incubator at 37 °C,
36
37 5% CO₂ and saturated humidity, at least 24 h before incubation. EpiSkinTM (Lyon, France) furnished
38
39 the assay medium used for incubations. RHE were topically treated first with DEPMPO (20 μL , 250
40
41 mM in HEPES) and post incubated (37 °C, 5% CO₂) during 15 min. After the incubation time, the
42
43 RHE were placed in the EPR tissue cell and the hydroperoxide (20 μL , 50 mM in acetone) was applied
44
45 to the epidermis taking care to ensure that the solution was only applied to it. EPR spectra were then
46
47 recorded.
48
49
50
51
52

53 54 ***EPR instrumentation***

1
2 EPR spectra were recorded on an EPR X-band spectrometer (ESP300E, Bruker Biospin GmbH,
3 Germany), equipped with a high sensitivity resonator (4119HS-W1, Bruker Biospin GmbH, Germany).
4
5 The g calibration was performed using Bruker standard (strong pitch) with known isotropic g factor of
6
7 2.0028. The principal experimental parameters values were: microwave power 5 mW, modulation
8
9 amplitude 0.5-1 G, and conversion time of ca. 160 ms resulting sweep time of ca. 3 min s for a single
10
11 scan. Spectra were recorded at room temperature ($295\text{K}\pm 1\text{K}$). Regarding the investigations in solution,
12
13 samples were introduced into glass capillaries (Hirschmann, 25 μL), sealed at both ends and transferred
14
15 into the EPR cavity for measurement as fast as possible after reagents mixing. RHE were placed in an
16
17 EPR tissue cell equipped with a silica window (Willmad, #ER162TC-Q) and resulting EPR spectra
18
19 recorded *in situ* right afterwards. Up to 10 scans were accumulated to improve the signal-to-noise ratio
20
21 (S/N).
22
23
24
25
26

27 ***EPR simulations***

28
29 All experimental spectra were analyzed by means of computer simulation using labmade scripts based
30
31 on Easyspin toolbox under Matlab (Mathworks) environment [24]. Relevant ^{14}N , ^1H , ^{31}P , ^{13}C hyperfine
32
33 coupling constants (*hfccs*) of detected spin-adducts are listed in Table 1. Coupling constants in RHE
34
35 were calculated following the best simulation attempts for such S/N.
36
37
38
39
40
41

42 **Results**

43 ***EPR studies in solution***

44
45 Analyses in solution were initially carried out to evaluate radicals formed from ^{13}C -substituted
46
47 CumOOHs in comparison with previous studies we achieved with non-substituted CumOOH [21].
48
49 DEPMPO was chosen as the spin-trap since we observed previously that it allowed a broad view of all
50
51 sort of radicals deriving from CumOOH [21]. Radical initiation was triggered by a catalytic amount of
52
53 Fe(II) (0.1 mM) inducing a Fenton-like reaction. DEPMPO was employed in an excess concentration
54
55
56
57
58
59

(25 mM) with respect to ^{13}C -substituted hydroperoxides (0.54 mM). First studies were completed with mono-substituted ^{13}C -MeCumOOH **1**. Figure 2(c) shows the experimental EPR spectrum obtained. Computer simulation of the signal displayed a mixture of different spin-adducts formed by addition of transient short-lived radicals derived from **1** to the α -carbon of DEPMPO nitronyl group. In agreement with our previous investigations on CumOOH [21], spin-adducts were identified through their hyperfine coupling constants (*hfccs*) [25, 26] as originating from HO^\bullet (Figure 2(e)), carbon centred radicals, probably methyl (Figure 2(f)), methoxy and/or possible peroxy radicals (Figure 2(h)). But here, when compared to the non-substituted sample, an additional hyperfine splitting appeared (Figure 2(g)), attributed to the presence of the paramagnetic probe of a ^{13}C nucleus with nuclear spin $I = \frac{1}{2}$. Analysis of the spectrum led to *hfccs* values of a_{N} (G) = 15.5, a_{H} (G) = 22.6, a_{P} (G) = 48.3 and a new value $a_{^{13}\text{C}}$ of 6.3 G (Table 1). As only half of methyl radicals were ^{13}C -substituted, it was possible to observe two spin-adducts corresponding to two types of carbon radicals trapped with similar a_{N} , a_{H} and a_{P} *hfccs*, being either non-substituted (Figure 2(f)) or ^{13}C -substituted (Figure 2(g)). There are only few examples in the literature describing *hfccs* for ^{13}C atoms, for example the description of *p*-benzyne radicals [27], β -substituted ethyl radicals [28] or some vinyl and allyl radicals [29]. However, no reference was found on *hfccs* values for the coupling with ^{13}C atoms on β position of a nitroxide radical. Thus, to corroborate that ^{13}C -methyl radicals had been trapped by DEPMPO, studies were conducted with di-substituted $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2** having ^{13}C -substituted methyl groups exclusively. Figure 3(c) shows the experimental EPR spectrum obtained. Here, computer simulation displayed a mixture of different spin-adducts formed again by trapping of HO^\bullet (Figure 3(e)), carbon centred (Figure 3(f)), methoxy and/or possible peroxy radicals (Figure 3(g)). Yet, previously assigned DEPMPO- $^{13}\text{CH}_3$ spin-adduct was again detected in a relative higher proportion when compared to the studies performed with ^{13}C -MeCumOOH **1** and having very comparable *hfccs* values of a_{N} (G) = 15.4, a_{H} (G) = 22.6, a_{P} (G) = 48.3 and $a_{^{13}\text{C}}$ (G) = 6.4. Thus, the relative intensity of the DEPMPO-carbon

1 radical spin-adduct derived from CumOOH (Figure 4(a), star) was significantly reduced while
2 introducing a single ^{13}C -methyl group in the molecule of CumOOH (Figure 4(b)) and was suppressed
3
4 in the case of doubly ^{13}C substitution (Figure 4(c)), to the benefit of the spin-adduct corresponding to
5
6 the ^{13}C -methyl trapped (Figures 4(b)(c), black circle). These results supported that CumOOH derived
7
8 carbon centred radicals were, in these experimental conditions, methyl radicals.
9
10
11
12

13 *EPR studies in reconstructed human epidermis*

14
15
16 In order to carry out the experiments, the RHE tissue was loaded beforehand with the spin-trap. As
17
18 described previously in studies with tumour promoting peroxides [12], radicals formed from CumOOH
19
20 in the skin model being both short lived and potentially generated at low concentrations, it was
21
22 desirable for the spin-trap to be as close as possible to the site of radical generation prior to the topical
23
24 application of the hydroperoxides. If similar EPR fingerprints to those of the experiments in solution
25
26 were obtained, it was noticeable that other features did not fully overlay with any of the solution
27
28 studies. Such differences can be explained by the much more complex system formed by the RHE
29
30 when compared to simplified in solution investigations. Figure 4 shows the spectra obtained in solution
31
32 along and in comparison with the spectra obtained in RHE for the three hydroperoxides tested. If the
33
34 appearance of the spin-adduct resulting from trapping the radical ^{13}C -substituted in RHE was barely
35
36 visible when using ^{13}C -MeCumOOH **1** (Figure 4(f)), it became more evident with $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$
37
38 **2** (Figure 4(g)). Major peaks could be assigned to DEPMPO-OH, DEPMPO- CH_3 and DEPMPO- $^{13}\text{CH}_3$
39
40 spin-adducts by comparison with previous data in solution (Table 1) [21]. Again, the difference
41
42 observed when comparing the *hfccs* values with those obtained in the EPR-ST solution studies could be
43
44 explained by the distinct environment of the spin-adducts in solution compared to that of a complex
45
46 heterogeneous tissue such as the RHE. Indeed, magnitudes of the *hfccs* pointing to the extent of
47
48 delocalization of the unpaired electron over the molecule may be affected by the local spin
49
50 environment and polarity. Moreover, control experiments were carried out for each compound by
51
52
53
54
55
56
57
58
59
60

1 simply incubating DEPMPO in the RHE and showed no EPR signal, corroborating that spin-adducts
2 resulted from trapping radicals issued from the hydroperoxide. Also, after penetrating the RHE,
3
4
5
6 CumOOH radical initiation has been induced by RHE matrix itself as no Fe(II) was supplemented, thus
7
8
9 better mimicking real life exposure scenarios. These results suggested methyl radicals being formed in
10
11 RHE after CumOOH exposure.
12
13
14
15

16 Discussion

17
18 In a recent work, we employed EPR-ST for the first time in the skin allergy field to study the potential
19 of this methodology to investigate *in situ* the formation of free radicals from skin allergens in 3D RHE
20 models [21]. CumOOH was used as proof of concept and we demonstrated the formation of carbon
21 radicals in RHE. However, no evidence could be obtained concerning their specific nature. In the
22 present work, this was investigated by using ^{13}C -substitution as a tool to unravel which carbon radicals
23 could be formed within the epidermis. As the major mechanism described for the formation of carbon
24 radicals from CumOOH is β -scission of cumyl oxyl radicals affording methyl radicals, we developed
25 the synthesis of CumOOH containing either one (^{13}C -MeCumOOH **1**) or both ($(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2**)
26 methyl groups ^{13}C -substituted and we further studied the formation of free radicals in solution and in
27 RHE.
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Preliminary studies in solution allowed setting up the optimal experimental conditions providing a
42 decent signal-to-noise ratio while keeping low CumOOHs concentrations, *i.e.* suitable with
43 physiological conditions. DEPMPO was chosen as an accurate spin-trap based upon (i) the high
44 persistency of its superoxide and alkylperoxyl spin-adducts when compared to other spin-traps (e.g.
45 DMPO); (ii) the clear distinction between oxygen- and carbon-derived spin-adducts [25, 30]. Radical
46 initiation was performed *via* a Fe(II) Fenton-like reaction. As for CumOOH, carbon centered, CumO \cdot
47 and HO \cdot radicals were identified for ^{13}C -MeCumOOH **1** and $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2**. A new signal
48
49
50
51
52
53
54
55
56
57
58
59

1
2 appeared in the case of mono-substituted **1** with $hfccs$ a_N (G) = 15.5, a_H (G) = 22.6, a_P (G) = 48.3 and
3
4 a_{13C} (G) = 6.3, whose fingerprint significantly increased in the case of the double-substituted
5
6 compound **2**, fully replacing the signal of carbon centered radicals trapped from non-substituted
7
8 CumOOH. Thus, when all methyl groups were ^{13}C -substituted, only DEPMPO- $^{13}CH_3$ spin-adducts
9
10 were detected confirming that methyl radicals were the carbon-centered trapped radicals. This
11
12 information was of high relevance for further studies in RHE.
13
14

15 RHE EpiSkin™ shows reasonable similarities to native human tissue in terms of morphology,
16
17 *stratum corneum* presence, lipid composition and biochemical markers [22]. It is today a useful tool for
18
19 testing irritation and penetration features of chemicals in contact with the skin as replacement of *in vivo*
20
21 testing [31]. For our studies on skin allergy the availability of EpiSkin™ RHE is a very good
22
23 alternative to the use of human and animal tissues as the model amends important legal and ethical
24
25 issues. Although considerably more permeable than human skin on average, the model appears to be
26
27 more consistent in permeability and responsiveness than human skin which is highly variable. **Similar**
28
29 **EPR-ST results were observed in RHE when testing CumOOH, ^{13}C -MeCumOOH **1** and (^{13}C -**
30
31 **Me) $_2$ CumOOH **2**. Though, new features in the RHE samples did not fully overlay with the reference**
32
33 **spectra in solution and were ascribed to the complexity of the RHE system. Even if the S/N was**
34
35 **significantly lower when compared to solution investigations and thus did not allow quantification of**
36
37 **the spin-adducts, we can still suggest that methyl radicals issued from CumOOH decomposition can be**
38
39 **formed in RHE EpiSkin™. Indeed, spectrum of the RHE experiment with twice ^{13}C -labeled CumOOH**
40
41 **(Figure 4(g)) was rather convincing as an additional spin-adduct assigned to DEPMPO- $^{13}CH_3$ spin-**
42
43 **adduct (black circles) was detected. Moreover, control experiments with DEPMPO, in absence of ^{13}C -**
44
45 **MeCumOOH **1** or (^{13}C -Me) $_2$ CumOOH **2**, did not exhibit any EPR signal (Figure 4(d)), suggesting that**
46
47 **the trapped $\cdot CH_3$ radicals originate from the hydroperoxide. Initially formed alkoxy radicals derived**
48
49 **from the O-O bond cleavage and giving the $\cdot CH_3$ radicals by β -scission could also react with**
50
51 **biomolecules in the RHE before DEPMPO trapping occurred. However, these potential side-reactions**
52
53
54
55
56
57
58
59
60

1
2 were not the purpose of the present study and will be part of future investigations. Worthy of note, no
3
4 Fe(II) was used for radical initiation when probing RHE, to mimic real life skin allergy situation,
5
6 whereas we previously showed that Fe(II) was required to initiate the free radical generation when
7
8 mixing CumOOH and DEPMPO in solution [21]. As it is shown in Figure 5(d), no EPR signal was
9
10 observed in the absence of Fe(II), whereas 0.1 mM Fe(II) was enough to initiate the radical reaction
11
12 with a Fe(II) concentration dependent answer on the amount of spin-adducts formed. We can then
13
14 hypothesize that reaction of organic peroxides in the skin in the presence of one-electron donor agents
15
16 (*i.e.* amino and thiol groups present in amino acids, metal complexes, enzymes) may initiate the
17
18 electron transfer processes and further skin proteins haptentation [32-34]. Also, the potential presence in
19
20 the skin of Fe(III) heme-containing enzymes could be responsible for the peroxide O-O bond cleavage
21
22 and start radical processes. Reactions of model Fe(III) porphyrin complexes with oxidants, such as
23
24 peroxyacids and hydroperoxides, have been extensively studied as models. The cleavage has been
25
26 confirmed but a clear mechanistic consensus has not evolved on the homolytic or heterolytic nature of
27
28 the O-O bond cleavage by these species, both kinds of mechanisms being possible [35-37].
29
30
31
32
33

34 While skin allergens (haptens) are unable to stimulate an adaptive immune response after
35
36 penetration into the epidermis, the immunogenicity is reached by reaction with skin proteins, to form
37
38 stable antigenic conjugates recognized and processed for presentation to the immune system [38].
39
40 Therefore, the hapten-protein interaction is the primary key step to induce sensitization and was shown
41
42 to occur relatively quickly in RHE [39]. The resulting sequence of events leading epidermal cells to
43
44 present the antigen so formed to the immune system and its activation take 10 to 15 days in humans.
45
46 The fast trapping by DEPMPO allowed EPR-ST characterization of the generated methyl radicals but
47
48 the reaction with epidermal proteins was not evidenced. Thus, if we cannot testify so far that these
49
50 radicals are the reacting species for the antigen formation we can strongly suggest their involvement.
51
52 Additional tests to decipher their reactivity towards amino acids in the skin could be isolation from
53
54
55
56
57
58
59
60

1
2 RHE of modified proteins (reactivity in the absence of spin-trap) by sonication followed by mass
3
4 spectrometry analysis and will be addressed in a near future.
5
6
7

8 9 **Conclusion**

10
11 This work shows by using ^{13}C -substituted compounds and EPR-ST that the production of methyl
12
13 radicals from CumOOH previously reported in isolated keratinocytes and other model systems is also
14
15 possible in reconstructed skin tissue through a one-electron reductive pathway, and suggest that these
16
17 methyl radicals could be involved in the antigen formation responsible for skin sensitization to
18
19 CumOOH.
20
21
22
23
24

25 **Acknowledgements**

26
27 This work was part of the DEF-CHEMSKALL Franco-German Collaborative International Research
28
29 Project financially supported by the ANR (Agence Nationale de la Recherche; project no. ANR-15-
30
31 CE15-0023-01) and the DFG (Deutsche Forschungsgemeinschaft; project no. DFG, BL340/6-1). The
32
33 authors thank the University of Strasbourg, the CNRS and the REseau NATional de Rpe
34
35 interDisciplinaire (RENARD, Fédération IR-RPE CNRS #3443).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- [1] Yaremenko IA, Vil' VA, Demchuk DV, Terent'ev AO. Rearrangements of organic peroxides and related processes. *Beilstein J Org Chem* 2016; 12: 1647-1748.
- [2] Rider CV, Chan P, Herbert RA, et al. Dermal exposure to cumene hydroperoxide: assessing its toxic relevance and oxidant potential. *Toxicol Pathol* 2016; 44: 749-762.
- [3] Lai DY, Woo Y, Argus MF, Arcos JC. Carcinogenic potential of organic peroxides: prediction based on structure-activity relationships (SAR) and mechanism-based short-term tests. *J Environ Sci Health Part C Environ Carcinog Ecotoxicol Rev* 1996; 14: 63-80.
- [4] Hanausek M, Walaszek Z, Viaje A, et al. Exposure of mouse skin to organic peroxides: subchronic effects related to carcinogenic potential. *Carcinogenesis* 2004; 25: 431-437.
- [5] Brink A, Richter I, Lutz U, et al. Biological significance of DNA adducts: comparison of increments over background for various biomarkers of genotoxicity in L5178Y tk(+/-) mouse lymphoma cells treated with hydrogen peroxide and cumene hydroperoxide. *Mutat Res* 2009; 678: 123-128.
- [6] Bråred Christensson J, Matura M, Bäcktorp C, Börje A, Nilsson JL, Karlberg AT. Hydroperoxides form specific antigens in contact allergy. *Contact Dermatitis* 2006; 55: 230-237.
- [7] Lauricella R, Tuccio B. Détection et caractérisation de radicaux libres après piégeage de spins. In: *La Spectroscopie de Résonance Paramagnétique Électronique, 2014*, Bertrand P Ed. edp Sciences, Collection Grenoble Sciences, Les Ulis, France, pp. 49-78.
- [8] He G, Samouilov A, Kuppusamy P, Zweier JL. In vivo EPR imaging of the distribution and metabolism of nitroxide radicals in human skin. *J Magn Reson* 2001; 148: 155-164.
- [9] Jurkiewicz BA, Buettner GR. EPR detection of free radicals in UV-irradiated skin: mouse versus human. *Photochem Photobiol* 1996; 64: 918-922.
- [10] Plonka PM. Electron paramagnetic resonance as a unique tool for skin and hair research. *Exp Dermatol* 2009; 18: 472-484.
- [11] Taffe BG, Takahashi N, Kensler TW, Mason RP. Generation of free radicals from organic hydroperoxide tumor promoters in isolated mouse keratinocytes. *J Biol Chem* 1987; 262: 12143-12149.
- [12] Timmins GS, Davies MJ. Free radical formation in murine skin treated with tumour promoting organic peroxides. *Carcinogenesis* 1993; 14: 1499-1503.
- [13] Timmins GS, Davies MJ. Free radical formation in isolated murine keratinocytes treated with organic peroxides and its modulation by antioxidants. *Carcinogenesis* 1993; 14: 1615-1620.
- [14] Shvedova AA, Kisin ER, Murray AR, et al. Antioxidant balance and free radical generation in vitamin E-deficient mice after dermal exposure to cumene hydroperoxide. *Chem Res Toxicol* 2002; 15: 1451-1459.
- [15] Shvedova AA, Kisin ER, Murray AR, et al. Pro/antioxidant status in murine skin following topical exposure to cumene hydroperoxide throughout the ontogeny of skin cancer. *Biochemistry (Moscow)* 2004; 69: 23-31.
- [16] Sahli F, Silva e Sousa M, Vileno B, Lichter J, Lepoittevin JP, Blömeke B, Giménez-Arnau E. Understanding the skin sensitization capacity of ascaridole: a combined study of chemical reactivity and activation of the innate immune system (dendritic cells) on the epidermal environment. *Arch Toxicol* 2019, DOI: 10.1007/s00204-019-02444-3.

- 1
2
3 [17] Kao D, Chaintreau A, Lepoittevin JP, Giménez-Arnau E. Synthesis of allylic hydroperoxides and
4 EPR spin-trapping studies on the formation of radicals in iron systems as potential initiators of the
5 sensitizing pathway. *J Org Chem* 2011; 76: 6188-6200.
6
7 [18] Kao D, Chaintreau A, Lepoittevin JP, Giménez-Arnau E. Mechanistic studies on the reactivity of
8 sensitizing allylic hydroperoxides: investigation of the covalent modification of amino acids by
9 carbon-radical intermediates. *Toxicol Res* 2014; 3: 278-289.
10
11 [19] Bråred-Christensson J, Johansson S, Hagvall L, Jonsson C, Börje A, Karlberg AT. Limonene
12 hydroperoxide analogues differ in allergenic activity. *Contact Dermatitis* 2008; 59: 344-352.
13
14 [20] Deza G, García-Bravo B, Silvestre JF, Pastor MA, González-Pérez R, Heras-Mendoza F, Mercader
15 P, Fernández-Redondo V, Niklasson B, Giménez-Arnau A, GEIDAC. Contact sensitization to
16 limonene and linalool hydroperoxides in Spain: a GEIDAC prospective study. *Contact Dermatitis*
17 2017; 76: 74-80.
18
19 [21] Kuresepi S, Vileno B, Turek P, Lepoittevin JP, Giménez-Arnau E. Potential of EPR spin-trapping
20 to investigate in situ free radicals generation from skin allergens in reconstructed human
21 epidermis: cumene hydroperoxide as proof of concept. *Free Radic Res* 2018; 52: 171-179.
22
23 [22] Netzlaff F, Lehr CM, Wertz PW, Schaefer UF. The human epidermis models EpiSkin[®],
24 SkinEthic[®] and EpiDerm[®]: an evaluation of morphology and suitability for testing phototoxicity,
25 irritancy, corrosivity, and substance transport. *Eur J Pharm Biopharm* 2005; 60: 167-178.
26
27 [23] Barbati S, Clément JL, Olive G, Roubaud V, Tuccio B, Tordo P. In Minisci F. Ed. *Free Radicals in*
28 *Biology and Environment*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997, pp.
29 39-47.
30
31 [24] Stoll S, Schweiger A. EasySpin, a comprehensive software package for spectral simulation and
32 analysis in EPR. *J Magn Res* 2006; 178: 42-55.
33
34 [25] Frejaville C, Karoui H, Tuccio B, et al. 5-(Diethoxyphosphoryl)-5-methyl-1-pyrroline-*N*-oxide: A
35 new efficient phosphorylated nitron for the *in vitro* and *in vivo* spin trapping of oxygen-centered
36 radicals. *J Med Chem* 1995; 38: 258-265.
37
38 [26] Karoui H, Chalier F, Finet JP, Tordo P. DEPMPO: an efficient tool for the coupled ESR-spin trapping
39 of alkylperoxyl radicals in water. *Org Biomol Chem* 2011; 9: 2473-2480.
40
41 [27] Usuki T, Mita T, Lear MJ, et al. Spin trapping of ¹³C-labeled *p*-benzynes generated by Masamune-
42 Bergman cyclization of bicyclic nine-membered enediynes. *Angew Chem Int Ed* 2004; 43: 5249-
43 5253.
44
45 [28] Griller D, Ingold KU. On the conformation of β -substituted ethyl radicals. *J Am Chem Soc* 1974:
46 96; 6715-6720.
47
48 [29] Griller D, Cooper JW, Ingold KU. Kinetic applications of electron paramagnetic resonance
49 spectroscopy. XVIII. Persistent vinyl, alkyl, and allyl radicals. *J Am Chem Soc* 1975; 97: 4269-
50 4275.
51
52 [30] Stolze K, Udilova N, Nohl H. Spin trapping of lipid radicals with DEPMPO-derived spin traps:
53 detection of superoxide, alkyl and alkoxy radicals in aqueous and lipid phase. *Free Radic Biol*
54 *Med* 2000; 29: 1005-1014.
55
56 [31] OECD testing guideline 439, 2015. *In vitro* skin irritation: RHE test method. DOI:
57 10.1787/9789264242845-en.
58
59
60

- 1
2 [32] Moriarty-Craige SE, Jones DP. Extracellular thiols and thiol/disulphide redox in metabolism. *Annu*
3 *Rev Nutr* 2004; 24: 481-509.
4
5 [33] Fujimoto A, Masuda T. Chemical interaction between polyphenols and a cysteinyl thiol under
6 radical oxidation controls. *J Agric Food Chem* 2012; 60: 5142-5151.
7
8 [34] Grossi L. The mechanism of oxidation of aniline by peroxy-compounds: can EPR spectroscopy
9 help to discriminate between one-electron transfer and a nucleophilic substitution process? *Res*
10 *Chem Intermed* 1993; 19: 697-705.
11
12 [35] Traylor TJ, Kim C, Fann WP, Perrin CL. Reactions of hydroperoxides with iron (III) porphyrins:
13 heterolytic cleavage followed by hydroperoxide oxidation. *Tetrahedron* 1998; 54: 7977-7986.
14
15 [36] Almarsson O, Bruice TC. A Homolytic mechanism of O-O bond scission prevails in the reactions
16 of alkyl hydroperoxides with an octacationic tetraphenylporphinato-iron(III) complex in aqueous
17 solution. *J Am Chem Soc* 1995; 117: 4533-4544.
18
19 [37] Labeque R, Marnett LJ. Homolytic and heterolytic scission of organic hydroperoxides by (meso-
20 tetraphenylporphinato)iron(III) and its relation to olefin epoxidation. *J Am Chem Soc* 1989; 111:
21 6621-6627.
22
23 [38] Lepoittevin JP. Molecular aspects in allergic and irritant contact dermatitis. In Johansen JD, Frosch
24 PJ, Lepoittevin JP (Eds.) *Contact Dermatitis* 5th Ed. Springer-Verlag, Berlin, Heidelberg, 2011, pp.
25 91-110.
26
27 [39] Elbayed K, Berl V, Debeuckelaere C, Moussallieh FM, Piotto M, Namer IJ, Lepoittevin JP. HR-
28 MAS NMR spectroscopy of reconstructed human epidermis: potential for the in situ investigation
29 of the chemical interactions between skin allergens and nucleophilic amino acids. *Chem Res*
30 *Toxicol* 2013; 26: 136-145.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Table 1

Table 1. Hyperfine coupling constants (*hfccs*) of spin-adducts identified in solution (sol.) and in RHE*

Cpd.	Coupling constants					Assignment
	a_N (G)	a_H (G)	a_P (G)	$a_{^{13}C}$ (G)	g	
CumOOH (sol.) [21]	14.2	13.5	47.4		2.0055	DEPMPO-OH
	15.4	22.6	48.1		2.0053	DEPMPO-carbon R
	11.3	13.7	48.5		2.0055	DEPMPO-OCH ₃ and/or DEPMPO-OOCum
1 (sol.)	14.2	13.3	47.3		2.0056	DEPMPO-OH
	15.4	22.6	48.0		2.0053	DEPMPO-carbon R
	15.5	22.6	48.3	6.3	2.0053	DEPMPO- ¹³ CH ₃
	11.3	13.7	48.2		2.0055	DEPMPO-OCH ₃ and/or DEPMPO-OOCum
2 (sol.)	14.2	13.4	47.4		2.0053	DEPMPO-OH
	15.4	22.6	48.3	6.4	2.0051	DEPMPO- ¹³ CH ₃
	11.6	14.4	48.1		2.0055	DEPMPO-OCH ₃ and/or DEPMPO-OOCum
CumOOH (RHE)	12.5	13.6	46.0		2.0064	DEPMPO-OH
	14.8	21.4	48.0		2.0058	DEPMPO-carbon R
1 (RHE)	12.9	13.7	46.0		2.0064	DEPMPO-OH
	14.5	21.2	47.3		2.0056	DEPMPO-carbon R
	14.9	21.6	47.9	6.2	2.0056	DEPMPO- ¹³ CH ₃
2 (RHE)	13.1	14.0	46.4		2.0063	DEPMPO-OH
	15.1	21.4	48.2	6.4	2.0059	DEPMPO- ¹³ CH ₃

* *hfccs* in RHE were calculated following the best simulation attempts considering the S/N

Scheme 1



Scheme 2




Figure 1




Figure 2




Figure 3




Figure 4





Figure 5



Legends

Scheme 1. Radical process leading to the formation of methyl radicals from CumOOH.

Scheme 2. Synthetic pathway to obtain ^{13}C -MeCumOOH **1** and $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2**. 1- (^{13}C) Acetophenone **4** was the key starting material for both compounds.

Figure 1. Chemical structures of ^{13}C -MeCumOOH **1** and $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2**.

Figure 2. EPR spectra of ^{13}C -MeCumOOH **1**/DEPMPO/Fe(II) in solution. (a) Control with single DEPMPO (25 mM); (b) Control with DEPMPO (25 mM) and Fe(II) (1 mM); (c) Experimental spectrum obtained with ^{13}C -MeCumOOH **1** (0.54 mM), DEPMPO (25 mM) and Fe(II) (0.1 mM); (d) Computer simulation of spectrum (c); (e-h) deconvolution of (d) with: (e) DEPMPO-OH spin-adduct (45%); (f) DEPMPO-carbon R spin-adduct (28%); (g) DEPMPO- $^{13}\text{CH}_3$ spin-adduct (23%); (h) DEPMPO-OCH₃ and/or DEPMPO-OOCum spin-adduct (4%).

Figure 3. EPR spectra of $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2**/DEPMPO/Fe(II) in solution. (a) Control with single DEPMPO (25 mM); (b) Control with DEPMPO (25 mM) and Fe(II) (1 mM); (c) Experimental spectrum obtained with $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2** (0.54 mM), DEPMPO (25 mM) and Fe(II) (0.1 mM); (d) Computer simulation of spectrum (c); (e-g) deconvolution of (d) with: (e) DEPMPO-OH spin-adduct (55%); (f) DEPMPO- $^{13}\text{CH}_3$ spin-adduct (42%); (g) DEPMPO-OCH₃ and/or DEPMPO-OOCum spin-adduct (3%).

Figure 4. Comparative experimental EPR spectra in solution (a-c) and in RHE (d-g): (a) CumOOH (0.54 mM)/DEPMPO (25 mM)/Fe (II) (0.1 mM) [21]; (b) ^{13}C -MeCumOOH **1** (0.54 mM)/DEPMPO (25 mM)/Fe (II) (0.1 mM); (c) $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2** (0.54 mM)/DEPMPO (25 mM)/Fe (II) (0.1 mM); (d) DEPMPO (250 mM)/RHE; (e) CumOOH (10 mM)/DEPMPO (250 mM)/RHE; (f) ^{13}C -MeCumOOH **1** (50 mM)/DEPMPO (250 mM)/RHE; (g) $(^{13}\text{C}\text{-Me})_2\text{CumOOH}$ **2** (50 mM)/DEPMPO (250 mM)/RHE. Because of the low signal-to-noise ratio, DEPMPO-OCH₃ and/or DEPMPO-OOCum spin-adducts are not indicated for clarification purposes. Spin-adducts are indicated as follows: DEPMPO-OH white circle, DEPMPO-CH₃ star and DEPMPO- $^{13}\text{CH}_3$ black circle. Dotted arrows indicate the formation of signals related to DEPMPO- $^{13}\text{CH}_3$ spin-adduct in RHE.

Figure 5. EPR experimental spectra in solution of CumOOH (0.54 mM)/DEPMPO (25 mM) with varying concentrations of Fe (II): (a) Fe(II) 1.0 mM; (b) Fe(II) 0.5 mM; (c) Fe(II) 0.1 mM; (d) in the absence of Fe(II).