



Uniqueness of the entropy solution of a stochastic conservation law with a Q-Brownian motion

Tadahisa Funaki, Yueyuan Gao, Danielle Hilhorst

► To cite this version:

Tadahisa Funaki, Yueyuan Gao, Danielle Hilhorst. Uniqueness of the entropy solution of a stochastic conservation law with a Q-Brownian motion. 2019. hal-02159743v1

HAL Id: hal-02159743

<https://hal.science/hal-02159743v1>

Preprint submitted on 19 Jun 2019 (v1), last revised 19 Feb 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ARTICLE TYPE

Uniqueness of the entropy solution of a stochastic conservation law with a Q -Brownian motion

Tadahisa FUNAKI¹ | Yueyuan GAO² | Danielle HILHORST^{*3}

¹Department of Mathematics, School of Fundamental Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

²MathAM-OIL, AIST c/o AIMR, Tohoku University, 2-1-1, Katahira, Aoba-ku, Sendai, 980-8577, Japan

³CNRS and Laboratoire de Mathématiques d'Orsay, Université Paris-Sud, Orsay Cedex, 91405, France

Correspondence

*Danielle Hilhorst. Email:
danielle.hilhorst@math.u-psud.fr

Present Address

CNRS and Laboratoire de Mathématiques d'Orsay, Université Paris-Sud, Orsay Cedex, 91405, France

Summary

In this paper, we prove the uniqueness of the entropy solution for a first order stochastic conservation law with a multiplicative source term involving a Q -Brownian motion.

After having defined a measure-valued weak entropy solution of the stochastic conservation law, we present the Kato inequality and as a corollary we deduce the uniqueness of the measure-valued weak entropy solution which coincides with the unique weak entropy solution of the problem. The Kato inequality is proved by a doubling of variables method; to that purpose, we prove the existence and the uniqueness of the strong solution of an associated stochastic nonlinear parabolic problem by means of an implicit time discretization scheme; we also prove its convergence to a measure-valued entropy solution of the stochastic conservation law, which in turn coincides with its unique entropy solution.

KEYWORDS:

Stochastic first order conservation law, Q -Brownian motion, Kato inequality, Uniqueness of entropy solution, Associated parabolic problem

1 | INTRODUCTION

In this paper we prove the uniqueness of the entropy solution for the first order stochastic conservation law

$$\begin{cases} du + \operatorname{div}(\mathbf{v}f(u))dt = g(u)dW(x, t), & \text{in } \Omega \times \mathbb{T}^d \times (0, T), \\ u(\omega, x, 0) = u_0(x), & \omega \in \Omega, x \in \mathbb{T}^d. \end{cases} \quad (1)$$

where \mathbb{T}^d is the d -dimensional torus, $W(x, t)$ is a Q -Brownian motion, with the following hypotheses (H):

- $u_0 \in L^2(\mathbb{T}^d)$.
- $f : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz continuous function with Lipschitz constant C_f such that $f(0) = 0$.
- $g : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz continuous function with Lipschitz constant C_g ; moreover we suppose that g is bounded such that $|g| < M_g$ for a positive constant M_g .
- $\mathbf{v} \in C^1(\mathbb{T}^d \times [0, T])$ with $\operatorname{div} \mathbf{v} = 0$ for all $(x, t) \in \mathbb{T} \times [0, T]$, so that there exists $V < \infty$ such that $|\mathbf{v}(x, t)| \leq V$ for all (x, t) .

Several articles have been devoted to the study of stochastic perturbations of nonlinear first order hyperbolic problems. Let us mention the article of Bauzet-Vallet-Wittbold² who prove the existence and uniqueness of a stochastic entropy solution of the

Cauchy problem for a stochastic conservation law with a multiplicative one-dimensional noise in time. They apply an artificial viscosity method to prove the existence of a solution, and an adaptation of the Kruzhkov doubling of variables method to prove that any stochastic entropy solution coincides with the solution obtained by the artificial viscosity method. Then as an extension, Bauzet-Vallet-Wittbold³ prove an existence and uniqueness result for the stochastic entropy solution of a Dirichlet problem for the stochastic conservation law with multiplicative noise in time.

As for the uniqueness result, Debussche-Vovelle⁵ consider the d -dimensional problem with multiplicative noise. $du + f(u)_x dt = h(u)dW$. The noise term dW is a function of space and time. The authors prove the existence and uniqueness of the kinetic solution. We believe however that our present results are important since entropy solutions are much easier to approximate numerically than kinetic solutions.

Funaki-Gao-Hilhorst⁷ prove the convergence of a finite volume scheme for the stochastic conservation law (1). This convergence result implies the existence of a measure-valued weak entropy solution. In this paper, we extend the proof of the uniqueness of the weak entropy solution of Bauzet-Vallet-Wittbold^{2,3} to the case of equation (1) where the noise involves a Q -Brownian motion.

This paper is organized as follows: In section 2, we recall the definitions of a weak entropy solution and of a measure-valued entropy solution of Problem (1) and we present the associated parabolic problem and an inequality satisfied by its strong solution. We prove that the solution of the associated parabolic problem converges to a measure-valued entropy solution of the stochastic conservation law.

In section 3, we first present the Kato inequality and as a corollary, we deduce the uniqueness of the measure-valued weak entropy solution and then the uniqueness of the weak entropy solution. As a consequence of the proofs, the measure-valued weak entropy solution is unique and coincides with the unique weak entropy solution. The Kato inequality is proved by a doubling of variables method for which we need entropy type inequalities for measure-valued weak entropy solutions and for the strong solutions of the associated parabolic problem.

In section 4, we apply an implicit time discretization to obtain a semi-discrete solution of the associated stochastic nonlinear parabolic problem. By Itô's formula and a priori estimates, we prove that when the time step tends to zero, the limit of this discrete solution is a strong solution of the corresponding parabolic problem. We then prove the uniqueness of the strong solution.

2 | ENTROPY FORMULATION

We recall the definitions of a Q -Brownian motion and of an entropy solution of Problem (1). Let $Q \geq 0$ be a symmetric trace class operator on $L^2(\mathbb{T}^d)$, let $\{e_j\}_{j \geq 1}$ be an orthonormal basis in $L^2(\mathbb{T}^d)$ diagonalizing Q and $\{\lambda_j \geq 0\}_{j \geq 1}$ be the corresponding eigenvalues, such that

$$Qe_j = \lambda_j e_j$$

for all $j \geq 1$. We recall that Q is of trace class, namely that

$$\text{Tr } Q = \sum_{j=1}^{\infty} \langle Qe_j, e_j \rangle_{L^2(\mathbb{T}^d)} = \sum_{j=1}^{\infty} \lambda_j \leq \Lambda_0, \quad (2)$$

for some positive constant Λ_0 . Let (Ω, \mathcal{F}, P) be a probability space equipped with a filtration $(\mathcal{F}_t)^9$ and $\{\beta_j\}_{j \geq 1}$ be a sequence of independent (\mathcal{F}_t) -Brownian motions defined on (Ω, \mathcal{F}, P) . We recall that a Brownian motion $\beta(t)$ is called an (\mathcal{F}_t) -Brownian motion if it is (\mathcal{F}_t) -adapted and the increment $\beta(t) - \beta(s)$ is independent of \mathcal{F}_s for every $0 \leq s < t$. The process W defined by

$$W(x, t) = \sum_{j=1}^{\infty} \beta_j(t) Q^{\frac{1}{2}} e_j(x) = \sum_{j=1}^{\infty} \sqrt{\lambda_j} \beta_j(t) e_j(x) \quad (3)$$

is a Q -Brownian motion in $L^2(\mathbb{T}^d)$, and the series defined by (3) is convergent in $L^2(\Omega, C([0, T], L^2(\mathbb{T}^d)))$ [cf. ⁴]. We suppose furthermore that $e_j \in C(\mathbb{T}^d)$ for $j = 1, 2, \dots$ and that there exists a positive constant Λ_1 such that

$$\sum_{j=1}^{\infty} \lambda_j \|e_j\|_{L^{\infty}(\mathbb{T}^d)}^2 \leq \Lambda_1. \quad (4)$$

In the proof of the existence of a strong solution of the associated parabolic problem, we need the further assumption that there exists a positive constant Λ_2 , such that

$$\sum_{j=1}^{\infty} \lambda_j \|\nabla e_j\|_{L^\infty(\mathbb{T}^d)^d}^2 \leq \Lambda_2. \quad (5)$$

Definition 1 (Weak entropy solution of Problem (1)). A function $u \in \mathcal{N}_\omega^2(0, T; L^2(\mathbb{T}^d)) \cap L^\infty(0, T; L^2(\Omega \times \mathbb{T}^d))$ is a weak entropy solution of the stochastic scalar conservation law (1) with the initial condition $u_0 \in L^2(\mathbb{T}^d)$, if P-a.s. in Ω ,

$$\begin{aligned} & \int_{\mathbb{T}^d} \eta(u_0(x) - k)\varphi(x, 0)dx + \int_0^T \int_{\mathbb{T}^d} \eta(u - k)\partial_t \varphi(x, t)dxdt + \int_0^T \int_{\mathbb{T}^d} F^\eta(u, k)\mathbf{v} \cdot \nabla_x \varphi(x, t)dxdt \\ & + \int_{\mathbb{T}^d} \int_0^T \eta'(u - k)g(u)\varphi(x, t)dW(x, t)dx \\ & + \frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \eta''(u - k)g^2(u)\varphi(x, t)Q(x, x)dxdt \\ & \geq 0 \end{aligned} \quad (6)$$

where

$$F^\eta(a, b) = \int_b^a \eta'(\sigma - b)f'(\sigma)d\sigma \quad (7)$$

and

$$Q(x, y) = \sum_{j=1}^{\infty} \lambda_j e_j(x)e_j(y), \quad (8)$$

for all $\varphi \in \mathcal{C} := \{\varphi \in C_0^\infty(\mathbb{T}^d \times [0, T]), \varphi \geq 0, \varphi(\cdot, T) = 0\}$ and for all $\eta \in \mathcal{A}$ where \mathcal{A} is the set of nonnegative $C^2(\mathbb{R})$ convex functions η such that the support of η'' is compact and $\eta(0) = 0$. $\mathcal{N}_\omega^2(0, T; L^2(\mathbb{T}^d))$ is the subclass of $L^2(\Omega \times \mathbb{T}^d \times (0, T))$ consisting of predictable $L^2(\mathbb{T}^d)$ -valued processes [cf.⁴, page 98].

We remark that for all $x, y \in \mathbb{T}^d$

$$|Q(x, y)| \leq \sum_{j=1}^{\infty} \lambda_j \|e_j\|_{L^\infty(\mathbb{T}^d)}^2 \leq \Lambda_1. \quad (9)$$

Definition 2 (Measure-valued weak entropy solution of Problem (1)). A function \mathbf{u} of $\mathcal{N}_\omega^2(0, T; L^2(\mathbb{T}^d \times (0, 1))) \cap L^\infty(0, T; L^2(\Omega \times \mathbb{T}^d \times (0, 1)))$ is a measure-valued weak entropy solution of the stochastic scalar conservation law (1) with the initial condition $u_0 \in L^2(\mathbb{T}^d)$, if P-a.s. in Ω , for all $\eta \in \mathcal{A}$ and for all $\varphi \in \mathcal{C}$

$$\begin{aligned} & \int_{\mathbb{T}^d} \eta(u_0 - k)\varphi(x, 0)dx + \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta(\mathbf{u}(x, t, \alpha) - k)\partial_t \varphi(x, t)d\alpha dxdt \\ & + \int_0^T \int_{\mathbb{T}^d} \int_0^1 F^\eta(\mathbf{u}(x, t, \alpha), k)\mathbf{v} \cdot \nabla_x \varphi(x, t)d\alpha dxdt \\ & + \int_{\mathbb{T}^d} \int_0^T \int_0^1 \eta'(\mathbf{u}(x, t, \alpha) - k)g(\mathbf{u}(x, t, \alpha))\varphi(x, t)d\alpha dW(x, t)dx \\ & + \frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta''(\mathbf{u}(x, t, \alpha) - k)g^2(\mathbf{u}(x, t, \alpha))\varphi(x, t)Q(x, x)d\alpha dxdt \\ & \geq 0. \end{aligned} \quad (10)$$

The main result of this article is the following theorem.

Theorem 1. Suppose that the hypotheses (H) hold and that the basis functions $\{e_j\}_{j \geq 1}$ of the Q -Brownian motion satisfy the hypotheses (4) and (5), then the measure-valued weak entropy solution is unique. Moreover, it coincides with the unique weak entropy solution.

Before presenting the Kato inequality and proving the Theorem 1, we recall the Itô's formula, and show the lemma that the solution of associated parabolic equation tends to a measure-valued solution of Definition 2 in the sense of Young measure, which will be used in sequel.

Lemma 1 (Itô's formula in the case of a Q -Brownian motion, cf.⁴). We define X by:

$$X(t) = X(0) + \int_0^t \psi(s) ds + \int_0^t \theta(s) dW(s), \quad 0 \leq t \leq T,$$

where W is a Q -Brownian motion. Suppose that ψ is an $L^2(\mathbb{T}^d)$ -valued predictable Bochner integrable process,

$$\int_0^T \|\psi(s)\|_{L^2(\mathbb{T}^d)} ds < \infty \quad \text{P-a.s.},$$

where θ is an (\mathcal{F}_t) -adapted and measurable process such that

$$\text{P}\left(\int_0^T \text{Tr}[\theta^2(s)Q] ds < \infty\right) = 1.$$

Suppose that the function $\mathcal{G}(X, t) : L^2(\mathbb{T}^d) \times [0, T] \rightarrow \mathbb{R}$ is such that its partial derivative \mathcal{G}_t and Fréchet derivatives \mathcal{G}_X and \mathcal{G}_{XX} are continuous on $L^2(\mathbb{T}^d) \times [0, T]$. Then P-a.s. for all $t \in [0, T]$,

$$\begin{aligned} \mathcal{G}(X(t), t) &= \mathcal{G}(X(0), 0) + \int_0^t \mathcal{G}_t(X(s), s) + \langle \mathcal{G}_X(X(s), s), \psi(s) \rangle_{L^2(\mathbb{T}^d)} ds \\ &\quad + \int_0^t \langle \mathcal{G}_X(X(s), s), \theta(s) dW(s) \rangle_{L^2(\mathbb{T}^d)} \\ &\quad + \frac{1}{2} \int_0^t \text{Tr} [\mathcal{G}_{XX}(X(s), s) \theta^2 Q] ds. \end{aligned} \tag{11}$$

We assume that for $\epsilon > 0$, u_ϵ is the solution of the stochastic nonlinear parabolic problem

$$du - \epsilon \Delta u dt + \text{div}(\mathbf{v} f(u)) dt = g(u) dW \quad \text{in } \Omega \times \mathbb{T}^d \times [0, T] \tag{12}$$

with an initial condition $u_0^\epsilon \in H^1(\mathbb{T}^d)$. In section 4, we will give the definition of the strong solution u_ϵ of the initial value problem (12) and prove that the strong solution exists and is unique.

Lemma 2. Let u_ϵ be the solution of the associated parabolic problem in the sense of Definition 4 in section 4. Then u_ϵ tends to a measure-valued entropy solution of Problem (1) as ϵ tends to 0 in the sense of Young measures.

Proof. We define the functions

$$\begin{aligned} X(t) &= u_\epsilon(x, t), \quad \psi = \epsilon \Delta u_\epsilon(x, t) - \text{div}(\mathbf{v} f(u_\epsilon(x, t))), \quad \theta = g(u_\epsilon(x, t)) \\ \mathcal{G}(X(t), t) &= \int_{\mathbb{T}^d} \eta(X(t) - k) \varphi(x, t) dx, \end{aligned}$$

so that

$$\begin{aligned} \mathcal{G}_X(X(t), t) &= \eta'(X(t) - k) \varphi(x, t) \\ \mathcal{G}_{XX}(X(t), t) &= \eta''(X(t) - k) \varphi(x, t) \end{aligned}$$

$$\mathcal{G}_t(X(t), t) = \int_{\mathbb{T}^d} \eta(X(t) - k) \partial_t \varphi(x, t) dx$$

and apply Itô's formula, Lemma 1 to obtain

$$\begin{aligned} & \int_{\mathbb{T}^d} \eta(u_\epsilon(T) - k) \varphi(T) dx \\ &= \int_{\mathbb{T}^d} \eta(u_\epsilon(0) - k) \varphi(0) dx + \int_0^T \int_{\mathbb{T}^d} \eta(u_\epsilon(x, t) - k) \partial_t \varphi(x, t) dx dt + \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \varphi(x, t) [\epsilon \Delta u_\epsilon(x, t) - \operatorname{div}(\mathbf{v} f(u_\epsilon(x, t)))] dx dt \\ &+ \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \varphi(x, t) g(u_\epsilon(x, t)) dW(x, t) dx + \frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) \varphi(x, t) g^2(u_\epsilon(x, t)) Q(x, x) dx dt. \end{aligned}$$

We first consider the term involving Δu_ϵ . Note that for all $\varphi \in C$,

$$\begin{aligned} & \epsilon \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t)) \varphi(x, t) \Delta u_\epsilon(x, t) dx dt \\ &= -\epsilon \int_0^T \int_{\mathbb{T}^d} \nabla_x (\eta'(u_\epsilon(x, t) - k) \varphi(x, t)) \nabla_x u_\epsilon(x, t) dx dt \\ &= -\epsilon \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \nabla_x u_\epsilon(x, t) \cdot \nabla_x \varphi(x, t) dx dt - \epsilon \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) |\nabla_x u_\epsilon(x, t)|^2 \varphi(x, t) dx dt. \end{aligned}$$

As for the term involving $\operatorname{div}(\mathbf{v} f(u_\epsilon(x, t)))$, we use integral by parts, the chain rule and $\operatorname{div} v = 0$ to deduce the equalities:

$$\begin{aligned} & - \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \operatorname{div}(\mathbf{v} f(u_\epsilon(x, t))) \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} f(u_\epsilon(x, t)) \mathbf{v} \cdot \nabla (\eta'(u_\epsilon(x, t) - k) \varphi(x, t)) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt + \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) \mathbf{v} \cdot \nabla_x u_\epsilon(x, t) \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt + \int_0^T \int_{\mathbb{T}^d} \operatorname{div} \left[\int_k^{u_\epsilon(x, t)} \eta''(\sigma - k) f(\sigma) \mathbf{v} d\sigma \right] \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \left[\eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) - \int_k^{u_\epsilon(x, t)} \eta''(\sigma - k) f(\sigma) d\sigma \right] \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \left[\eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) - [\eta'(\sigma - k) f(\sigma)]_k^{u_\epsilon(x, t)} + \int_k^{u_\epsilon(x, t)} \eta'(\sigma - k) f'(\sigma) d\sigma \right] \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \left[\eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) - \eta'(u_\epsilon(x, t) - k) f(u_\epsilon(x, t)) + \eta'(k - k) f(k) + \int_k^{u_\epsilon(x, t)} \eta'(\sigma - k) f'(\sigma) d\sigma \right] \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} \left[\int_k^{u_\epsilon(x, t)} \eta'(\sigma - k) f'(\sigma) d\sigma \right] \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt + \eta'(0) f(k) \left(\int_0^T \mathbf{v} \varphi(x, t) |_{\partial \mathbb{T}^d} dt - \int_0^T \int_{\mathbb{T}^d} \operatorname{div} \mathbf{v} \varphi(x, t) dx dt \right) \end{aligned}$$

$$= \int_0^T \int_{\mathbb{T}^d} F^\eta(u_\epsilon(x, t), k) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt.$$

We therefore deduce that

$$\begin{aligned} & \int_{\mathbb{T}^d} \eta(u_\epsilon(x, T) - k) \varphi(T) dx \\ &= \int_{\mathbb{T}^d} \eta(u_\epsilon(x, 0) - k) \varphi(0) dx + \int_0^T \int_{\mathbb{T}^d} \eta(u_\epsilon(x, t) - k) \partial_t \varphi(x, t) dx dt \\ &\quad - \epsilon \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \nabla_x u_\epsilon(x, t) \cdot \nabla_x \varphi(x, t) dx dt - \epsilon \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) |\nabla_x u_\epsilon(x, t)|^2 \varphi(x, t) dx dt \\ &\quad + \int_0^T \int_{\mathbb{T}^d} F^\eta(u_\epsilon(x, t), k) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \\ &\quad + \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \varphi(x, t) g(u_\epsilon(x, t)) dW(x, t) dx + \frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) \varphi(x, t) g^2(u_\epsilon(x, t)) Q(x, x) dx dt \end{aligned}$$

for all $\varphi \in C$. Using $\varphi(T) = 0$ and that η is a nonnegative convex function and that $|\nabla_x u_\epsilon(x, t)|^2 \geq 0$, we deduce the following inequality

$$\begin{aligned} & \int_{\mathbb{T}^d} \eta(u_\epsilon^0(x) - k) \varphi(x, 0) dx + \int_0^T \int_{\mathbb{T}^d} \eta(u_\epsilon(x, t) - k) \partial_t \varphi(x, t) dx dt \\ &\quad - \epsilon \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \nabla_x u_\epsilon(x, t) \cdot \nabla_x \varphi(x, t) dx dt + \int_0^T \int_{\mathbb{T}^d} F^\eta(u_\epsilon(x, t), k) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \\ &\quad + \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) g(u_\epsilon(x, t)) \varphi(x, t) dW(x, t) dx + \frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) g^2(u_\epsilon(x, t)) \varphi(x, t) Q(x, x) dx dt \\ &\geq 0. \end{aligned} \tag{13}$$

We remark that (44) below and the strong convergence of $\{u^{\Delta t}\}$ to u^ϵ in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ [cf. Lemma 13] below imply that u^ϵ is uniformly bounded in $L^\infty(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$. Thus there exist an entropy process $\bar{\mathbf{u}} \in L^\infty(0, T; L^2(\Omega \times \mathbb{T}^d \times (0, 1)))$ and a subsequence of u_ϵ which we denote again by u_ϵ such that u_ϵ converges to $\bar{\mathbf{u}}$ in the sense of Young measures as ϵ tends to 0 [cf.², Page 668]. We multiply (13) by $\mathbf{1}_A$ where $A \in \mathcal{F}$ is arbitrary and take the expectation to obtain

$$\begin{aligned} & \mathbb{E} \left[\mathbf{1}_A \int_{\mathbb{T}^d} \eta(u_\epsilon^0(x) - k) \varphi(x, 0) dx \right] + \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \eta(u_\epsilon(x, t) - k) \partial_t \varphi(x, t) dx dt \right] \\ &\quad - \mathbb{E} \left[\epsilon \mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) \nabla_x u_\epsilon(x, t) \cdot \nabla_x \varphi(x, t) dx dt \right] + \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} F^\eta(u_\epsilon(x, t), k) \mathbf{v} \cdot \nabla_x \varphi(x, t) dx dt \right] \\ &\quad + \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x, t) - k) g(u_\epsilon(x, t)) \varphi(x, t) dW(x, t) dx \right] \\ &\quad + \frac{1}{2} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \eta''(u_\epsilon(x, t) - k) g^2(u_\epsilon(x, t)) \varphi(x, t) Q(x, x) dx dt \right] \end{aligned} \tag{14}$$

$$\geq 0.$$

The estimate (45) together with the lower semi-continuity of the convex functional $\|\cdot\|_{L^2(0,T;L^2(\Omega;L^2(\mathbb{T}^d)))}$ implies its weak lower-semi continuity by Ekeland-Temam [cf.⁶ Corollary 2.2, p.11], so that

$$\epsilon \|\nabla u^\epsilon\|_{L^2(0,T;L^2(\Omega;L^2(\mathbb{T}^d)))}^2 \leq C_2^2. \quad (15)$$

Next we pass to the limit $\epsilon \rightarrow 0$ in (14); the estimate (15) ensures that

$$-\mathbb{E} \left[\epsilon \mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \eta'(u_\epsilon(x,t) - k) \nabla u_\epsilon(x,t) \cdot \nabla \varphi(x,t) dx dt \right] \rightarrow 0$$

as $\epsilon \rightarrow 0$. For the other terms, we use similar arguments as in⁷ to deduce that

$$\begin{aligned} & \mathbb{E} \left[\mathbf{1}_A \int_{\mathbb{T}^d} \eta(u_0 - k) \varphi(x,0) dx \right] + \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta(\bar{\mathbf{u}}(x,t,\alpha) - k) \partial_t \varphi(x,t) d\alpha dx dt \right] \\ & + \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \int_0^1 F''(\bar{\mathbf{u}}(x,t,\alpha), k) \mathbf{v} \cdot \nabla_x \varphi(x,t) d\alpha dx dt \right] \\ & + \mathbb{E} \left[\mathbf{1}_A \int_{\mathbb{T}^d} \int_0^T \int_0^1 \eta'(\bar{\mathbf{u}}(x,t,\alpha) - k) g(\bar{\mathbf{u}}(x,t,\alpha)) \varphi(x,t) d\alpha dW(x,t) dx \right] \\ & + \frac{1}{2} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta''(\bar{\mathbf{u}}(x,t,\alpha) - k) g^2(\bar{\mathbf{u}}(x,t,\alpha)) \varphi(x,t) Q(x,x) d\alpha dx dt \right] \\ & \geq 0, \end{aligned}$$

which shows that $\bar{\mathbf{u}}$ is a measure-valued entropy solution of Problem (1). \square

3 | UNIQUENESS OF THE MEASURE-VALUED WEAK ENTROPY SOLUTION

The main part of this section is to prove the Kato inequality for measure-valued weak entropy solutions of Problem (1). The Kato inequality will imply the uniqueness of the measure-valued weak entropy solution and as a consequence the uniqueness of the weak entropy solution.

Proposition 1 (Kato inequality). Let $\mathbf{u}, \hat{\mathbf{u}}$ be 2 measure-valued weak entropy solutions to (1) with initial data $u_0, \hat{u}_0 \in L^2(\mathbb{T}^d)$ respectively. Then, for any nonnegative function φ with compact support belonging to $H^1(\mathbb{T}^d \times [0,T])$, there holds

$$\begin{aligned} & \int_{\mathbb{T}^d} |u_0 - \hat{u}_0| \varphi(0) dx + \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 |\mathbf{u}(x,t,\alpha) - \hat{\mathbf{u}}(x,t,\beta)| \partial_t \varphi(x,t) d\alpha d\beta dx dt \right] \\ & + \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 F(\mathbf{u}(x,t,\alpha), \hat{\mathbf{u}}(x,t,\beta)) \mathbf{v} \cdot \nabla \varphi(x,t) d\alpha d\beta dx dt \right] \\ & \geq 0. \end{aligned} \quad (16)$$

where $Q_T = \mathbb{T}^d \times (0,T)$ and $F(a,b) = \text{sgn}(a-b)[f(a) - f(b)]$ with $\text{sgn}(a) = \frac{a}{|a|}$ for all $a \neq 0$. Note that F is a Lipschitz continuous function with constant C_f .

We first prove the main result of this article, Theorem 1.

Proof of Theorem 1. We will use below the fact that the Kato inequality holds for all nonnegative test functions $\varphi \in H^1(Q_T)$. We take K , a positive constant, and denote by ψ any smooth nonincreasing function such that $\mathbf{1}_{(-\infty, K]}(\cdot) \leq \psi(\cdot) \leq \mathbf{1}_{(-\infty, K+1]}(\cdot)$. Setting $\varphi(x, t) = \psi(|x| - VC_f t)\gamma(t)$ with $\gamma(t) = \frac{T-t}{T}$ and substituting φ in (16) yields

$$\begin{aligned} & E \left[\int_{\mathbb{T}^d} |\hat{u}_0 - u_0| \psi(|x|) \right] \gamma(0) \\ & + E \left[\int_{Q_T} \int_0^1 \int_0^1 \psi(|x| - VC_f t) |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| \gamma'(t) d\alpha d\beta dx dt \right] \\ & - E \left[\int_{Q_T} \int_0^1 \int_0^1 VC_f \psi'(|x| - VC_f t) |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| \gamma(t) d\alpha d\beta dx dt \right] \\ & + E \left[\int_{Q_T} \int_0^1 \int_0^1 \psi'(|x| - VC_f t) F(\mathbf{u}(x, t, \beta), \hat{\mathbf{u}}(x, t, \alpha)) \mathbf{v} \cdot \frac{x}{|x|} \gamma(t) d\alpha d\beta dx dt \right] \\ & \geq 0. \end{aligned} \tag{17}$$

We study the signs of the last two terms in (17). Since ψ is a nonincreasing function, we have that

$$\begin{aligned} & -E \left[\int_{Q_T} \int_0^1 \int_0^1 VC_f \psi'(|x| - VC_f t) |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| \gamma(t) d\alpha d\beta dx dt \right] \\ & + E \left[\int_{Q_T} \int_0^1 \int_0^1 \psi'(|x| - VC_f t) F(\mathbf{u}(x, t, \beta), \hat{\mathbf{u}}(x, t, \alpha)) \mathbf{v} \cdot \frac{x}{|x|} \gamma(t) d\alpha d\beta dx dt \right] \\ & \leq E \left[\int_{Q_T} \int_0^1 \int_0^1 (VC_f - VC_f) \psi'(|x| - VC_f t) |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| \gamma(t) d\alpha d\beta dx dt \right] \\ & = 0, \end{aligned}$$

which combined with (17) yields

$$\begin{aligned} & E \left[\int_{\mathbb{T}^d} |\hat{u}_0 - u_0| \psi(|x|) \right] \gamma(0) \\ & + E \left[\int_{Q_T} \int_0^1 \int_0^1 \gamma'(t) \psi(|x| - VC_f t) |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| d\alpha d\beta dx dt \right] \\ & \geq 0. \end{aligned}$$

We suppose that $\hat{u}_0 = u_0$ and choose $K = \sup_{x \in \mathbb{T}^d} |x| - VC_f T$ to deduce that $\psi(|x| - VC_f t) = 1$. Also in view of the fact that $\gamma'(t) = -\frac{1}{T}$, we deduce that

$$E \left[\int_{Q_T} \int_0^1 \int_0^1 |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| d\alpha d\beta dx dt \right] = 0 \tag{18}$$

for all $\alpha, \beta \in (0, 1)$. This implies that $\mathbf{u}(x, t, \beta) = \hat{\mathbf{u}}(x, t, \alpha)$ for almost every $x \in \mathbb{T}^d$, $t \in (0, T)$, $\omega \in \Omega$, $\alpha, \beta \in (0, 1)$. Setting $\alpha = \beta$, we deduce that $\mathbf{u}(x, t, \alpha) = \hat{\mathbf{u}}(x, t, \alpha)$ for almost every $x \in \mathbb{T}^d$, $t \in (0, T)$ so that Problem (1) has a unique measure-valued weak entropy solution, say $\mathbf{u} = \mathbf{u}(x, t, \alpha)$. Then setting $\alpha \neq \beta$, we deduce that $\mathbf{u}(x, t, \alpha) = \mathbf{u}(x, t, \beta)$ which implies that the measure-valued weak entropy solution does not depend on the third parameter. Thus it is a weak entropy solution and (18) also implies the uniqueness of the weak entropy solution of Problem (1). \square

In the sequel we prove the Kato inequality, Proposition 1.

Proof. We first recall the definition of a mollifier.

Definition 3 (Mollifier). Let ρ be a nonnegative, real-valued function belonging to $C_0^\infty(\mathbb{R}^d)$ and having the properties

$$1. \rho(x) = 0 \text{ if } |x| \geq 1;$$

$$2. \int_{\mathbb{R}^d} \rho(x) dx = 1.$$

For example,

$$\rho(x) = \begin{cases} k_\rho e^{-\frac{1}{1-|x|^2}} & \text{if } |x| < 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $k_\rho > 0$ is chosen so that the condition 2 is satisfied. Let $N \geq 1$, we define the function $\rho_N(x) = N^d \rho(Nx)$ which is nonnegative, belongs to $C_0^\infty(\mathbb{R}^d)$, and satisfies

$$1. \rho_N(x) = 0 \text{ if } |x| \geq \frac{1}{N};$$

$$2. \int_{\mathbb{R}^d} \rho_N(x) dx = 1.$$

ρ_N is called a mollifier and is such that $\text{supp } \rho_N = (-\frac{1}{N}, \frac{1}{N})^d$.

We suppose that ρ_m and ρ_n are two mollifier sequences in \mathbb{T}^d and \mathbb{R} respectively, with $\text{supp } \rho_m \subset [-\frac{2}{m}, 0]^d$ and $\text{supp } \rho_n \subset [-\frac{2}{n}, 0]$. Suppose that ρ_l is a mollifier sequence with $\text{supp } \rho_l \subset [-\frac{1}{l}, \frac{1}{l}]$. For example, one can choose $\rho_m(x) = m^d \rho(mx + 1)$, $\rho_n(x) = n\rho(nx + 1)$ and $\rho_l(x) = l\rho(lx)$. Remark that $|\rho_m| \leq Cm^d$, $|\rho_n| \leq Cn$ and $|\rho_l| \leq Cl$ for some constant C and that ρ_l is an even function for all l . Let u_ϵ be the solution of the associated parabolic problem in the sense of Definition 4. We use the notations $\hat{\mathbf{u}} = \hat{\mathbf{u}}(x, t, \alpha)$ for a measure-valued solution and \hat{u}_0 for the corresponding initial condition, and set $\mathcal{A}_k^l := \rho_l(\hat{\mathbf{u}}(x, t, \alpha) - k)$ and $\mathcal{B}_k^l := \rho_l(u_\epsilon(y, s) - k)$. Next, we deduce two inequalities based on (10) and (13) respectively. We choose $\eta = \eta_\delta \in \mathcal{A}$ such that

$$\eta'_\delta(r) = \begin{cases} 1 & \text{if } r > \delta \\ \sin(\frac{\pi r}{2\delta}) & \text{if } |r| \leq \delta \\ -1 & \text{if } r < -\delta \end{cases}$$

in the inequality (10) as well as in the inequality (13). Note that η''_δ is an even Lipschitz continuous function. In the inequality (10), we replace the test function $\varphi(x, t)$ by $\varphi(y, s)\rho_m(x - y)\rho_n(t - s)$ and then multiply it by \mathcal{B}_k^l and integrate k on \mathbb{R} , y on \mathbb{T}^d and s on $(0, T)$ and write last the term coming from the stochastic integral to obtain

$$\begin{aligned} & \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \eta_\delta(\hat{u}_0(x) - k) \varphi(y, s) \rho_n(-s) \rho_m(x - y) dx \mathcal{B}_k^l dk dy ds \right] \\ & + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) \rho_n(t - s) \partial_t \varphi(y, s) \rho_m(x - y) d\alpha dx dt \mathcal{B}_k^l dk dy ds \right] \\ & + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) \varphi(y, s) \partial_t \rho_n(t - s) \rho_m(x - y) d\alpha dx dt \mathcal{B}_k^l dk dy ds \right] \\ & + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 F^{\eta_\delta}(\hat{\mathbf{u}}(x, t, \alpha), k) \mathbf{v} \cdot \rho_m(x - y) \nabla_x \varphi(y, s) \rho_n(t - s) d\alpha dx dt \mathcal{B}_k^l dk dy ds \right] \\ & + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 F^{\eta_\delta}(\hat{\mathbf{u}}(x, t, \alpha), k) \mathbf{v} \cdot \nabla_x \rho_m(x - y) \rho_n(t - s) \varphi(y - s) d\alpha dx dt \mathcal{B}_k^l dk dy ds \right] \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x-y) \rho_n(t-s) \varphi(y, s) Q(x, x) d\alpha dx dt \mathcal{B}_k^l dk dy ds \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta'_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \varphi(y, s) \rho_m(x-y) \rho_n(t-s) dx dW(x, t) \mathcal{B}_k^l dk dy ds \right] \\
:= & I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 \\
\geq & 0.
\end{aligned}$$

Next in (13) we change the variables (x, t) to (y, s) , replace $\varphi(y, s)$ by $\varphi(y, s) \rho_m(x-y) \rho_n(t-s)$ and write last the term coming from $\epsilon \Delta u_\epsilon$, then multiply by \mathcal{A}_k^l and integrate on α, k, x and t , to obtain

$$\begin{aligned}
& \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \eta_\delta(u_\epsilon^0(y) - k) \varphi(y, 0) \rho_n(t) \rho_m(x-y) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta_\delta(u_\epsilon(y, s) - k) \rho_n(t-s) \partial_s \varphi(y, s) \rho_m(x-y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta_\delta(u_\epsilon(y, s) - k) \partial_s \rho_n(t-s) \varphi(y, s) \rho_m(x-y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} F^{\eta_\delta}(u_\epsilon(y, s), k) \mathbf{v} \cdot \rho_m(x-y) \nabla_y \varphi(y, s) \rho_n(t-s) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} F^{\eta_\delta}(u_\epsilon(y, s), k) \mathbf{v} \cdot \nabla_y \rho_m(x-y) \varphi(y, s) \rho_n(t-s) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta''_\delta(u_\epsilon(y, s) - k) g^2(u_\epsilon(y, s)) \rho_m(x-y) \rho_n(t-s) \varphi(y, s) Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_0^T \int_{\mathbb{T}^d} \int_0^1 \eta'_\delta(u_\epsilon(y, s) - k) g(u_\epsilon(y, s)) \varphi(y, s) \rho_m(x-y) \rho_n(t-s) dW(y, s) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& - \epsilon \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta'_\delta(u_\epsilon(y, s) - k) \nabla_y u_\epsilon(y, s) \nabla_y [\varphi(y, s) \rho_m(x-y)] \rho_n(t-s) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
:= & J_1 + J_2 + J_3 + J_4 + J_5 + J_6 + J_7 + J_8 \\
\geq & 0.
\end{aligned}$$

We will show that the sum of $I_i + J_i$ for $i = 1, 2, \dots, 7$ and of the term J_8 converges to the left-hand-side of Kato inequality (16). More precisely, we should prove that

$$\lim_{m \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_1 + J_1] = \mathbb{E} \left[\int_{\mathbb{T}^d} |u_0 - \hat{u}_0| \varphi(0) dx \right], \quad (19)$$

$$\lim_{m \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [J_2 + J_3] = \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 |\mathbf{u}(x, t, \alpha) - \hat{\mathbf{u}}(x, t, \beta)| \partial_t \varphi(x, t) d\alpha d\beta dx dt \right], \quad (20)$$

$$I_3 + J_3 = 0, \quad (21)$$

$$\lim_{m \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_4 + J_4] = \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 F(\mathbf{u}(x, t, \beta), \hat{\mathbf{u}}(x, t, \alpha)) \mathbf{v} \cdot \nabla \varphi(x, t) d\alpha d\beta dx dt \right], \quad (22)$$

$$\lim_{m \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_5 + J_5] = 0, \quad (23)$$

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{m \rightarrow \infty} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7 + J_7] = 0, \quad (24)$$

and

$$\lim_{m \rightarrow \infty} \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} J_8 = 0. \quad (25)$$

We refer to² for the detailed proofs of (19) - (23) and of (25). Here we will only prove (24). Since we are dealing with a Q -Brownian motion instead of a one-dimensional Brownian motion, the proof of (24) is new with respect to the proofs given in². However, let us remark some simple facts. The partial derivatives $\partial_t \varphi(y, s)$ and $\nabla_x \varphi(y, s)$ in the terms I_2 and I_4 respectively vanish. Therefore $I_2 = I_4 = 0$. Moreover, since $t \in (0, T)$ and since the support of ρ_n is included in \mathbb{R}^- , the term J_1 vanishes. Let us now prove (24). We start by proving the following result.

Lemma 3. J_7 vanishes.

Proof. We set $\mathcal{G}_n^m = \rho_m(x - y)\rho_n(t - s)$, and use that $\text{supp } \rho_n \subset \mathbb{R}^-$. We have

$$\begin{aligned} J_7 &= \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_0^T \int_{\mathbb{T}^d} \eta'_\delta(u_\epsilon - k) g(u_\epsilon) \varphi(y, s) \mathcal{G}_n^m dW(y, s) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\ &= \int_{Q_T} \int_{\mathbb{R}} \mathbb{E} \left[\int_t^T \int_{\mathbb{T}^d} \eta'_\delta g(u_\epsilon)(u_\epsilon - k) \varphi(y, s) \mathcal{G}_n^m dW(y, s) dy \int_0^1 \mathcal{A}_k^l d\alpha \right] dk dx dt. \end{aligned}$$

We set

$$\bar{\alpha}(t) = \int_0^1 \mathcal{A}_k^l d\alpha$$

and

$$\bar{\gamma}(s) = g(u_\epsilon) \eta'_\delta(u_\epsilon - k) \varphi(y, s) \rho_m(x - y) \rho_n(t - s),$$

to obtain

$$\mathbb{E} \left[\int_{\mathbb{T}^d} \bar{\alpha}(t) \int_t^T \bar{\gamma}(s) dW(\cdot, s) \right] dy = \int_{\mathbb{T}^d} \sum_{j=1}^{\infty} \sqrt{\lambda_j} e_j(\cdot) \mathbb{E} \left[\bar{\alpha}(t) \int_t^T \bar{\gamma}(s) d\beta_j(s) \right] dy.$$

Then using that

$$\mathbb{E} \left[\bar{\alpha}(t) \int_t^T \bar{\gamma}(s) d\beta_j(s) \right] = \mathbb{E}[\bar{\alpha}(t)] \mathbb{E} \left[\int_t^T \bar{\gamma}(s) d\beta_j(s) \right] = 0$$

for all j , we deduce that $J_7 = 0$. \square

Next we study the limits of the sum of the Itô correction terms $I_6 + J_6$.

Lemma 4. We have the following limit properties:

$$\begin{aligned} I_6 + J_6 &\xrightarrow{n \rightarrow \infty} \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x - y) \varphi(y, t) Q(x, x) \rho_l(u_\epsilon(y, t) - k) d\alpha dk dy dx dt \right] \\ &\quad + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta(u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \rho_m(x - y) \varphi(y, t) Q(y, y) \int_0^1 \rho_l(\hat{\mathbf{u}}(x, t, \alpha) - k) d\alpha dx dt dk dy \right] \end{aligned}$$

$$\begin{aligned} & \overline{l \rightarrow \infty} \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x-y) \varphi(y, t) Q(x, x) d\alpha dx dt dy \right] \\ & + \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g^2(u_\epsilon(y, t)) \rho_m(x-y) \varphi(y, t) Q(y, y) d\alpha dx dt dy \right]. \end{aligned}$$

Proof. We justify the limit as $n \rightarrow \infty$. We set $\mathcal{G}_n^m = \rho_n(t-s) \rho_m(x-y)$ and $\mathcal{A}_k^l = \rho_l(\hat{\mathbf{u}}(x, t, \alpha) - k)$. We define A_1 as the difference of $I_6 + J_6$ and the limit term and perform a change of variables from k to $u_\epsilon(y, s) - k$ when y and s are fixed to obtain

$$\begin{aligned} A_1 &= \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x-y) \rho_n(t-s) \varphi(y, s) Q(x, x) d\alpha \rho_l(u_\epsilon(y, s) - k) dk dy ds dx dt \right] \\ &- \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x-y) \varphi(y, t) Q(x, x) \rho_l(u_\epsilon(y, t) - k) d\alpha dk dy dx dt \right] \\ &+ \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (u_\epsilon(y, s) - k) g^2(u_\epsilon(y, s)) \rho_m(x-y) \rho_n(t-s) \varphi(y, s) Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\ &- \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \rho_m(x-y) \varphi(y, t) Q(y, y) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\ &= \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, s) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \mathcal{G}_n^m \varphi(y, s) Q(x, x) dx dt d\alpha \rho_l(k) dk dy ds \right] \\ &- \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x-y) \varphi(y, t) Q(x, x) \rho_l(k) dx dt d\alpha dk dy \right] \\ &+ \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 [\eta''_\delta (u_\epsilon(y, s) - k) g^2(u_\epsilon(y, s)) - \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t))] \mathcal{G}_n^m \varphi(y, s) Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\ &+ \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \mathcal{G}_n^m [\varphi(y, s) - \varphi(y, t)] Q(y, y) \int_0^1 \mathcal{A}_k^l d\alpha dy ds dk dx dt \right] \\ &- \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \rho_m(x-y) \left[1 - \int_0^T \rho_n(t-s) ds \right] \varphi(y, t) Q(y, y) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\ &= \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 [\eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, s) + k) - \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha)) - u_\epsilon(y, t) + k] g^2(\hat{\mathbf{u}}(x, t, \alpha)) \right. \\ &\quad \times \mathcal{G}_n^m \varphi(y, s) Q(x, x) \rho_l(k) dx dt d\alpha dk dy ds \left. \right] \\ &+ \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \mathcal{G}_n^m [\varphi(y, s) - \varphi(y, t)] Q(x, x) \rho_l(k) dx dt d\alpha dk dy ds \right] \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x - y) \right. \\
& \quad \times \left. \left[1 - \int_0^T \rho_n(t - s) ds \right] \varphi(y, t) Q(x, x) \rho_l(k) dx dt d\alpha dk dy \right] \\
& + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta''_\delta (u_\epsilon(y, s) - k) [g^2(u_\epsilon(y, s)) - g^2(u_\epsilon(y, t))] G_n^m \varphi(y, s) Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} [\eta''_\delta (u_\epsilon(y, s) - k) - \eta''_\delta (u_\epsilon(y, t) - k)] g^2(u_\epsilon(y, t)) G_n^m \varphi(y, s) Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) G_n^m [\varphi(y, s) - \varphi(y, t)] Q(y, y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& - \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \varphi(y, t) Q(y, y) \left[1 - \int_0^T \rho_n(t - s) ds \right] \rho_m(x - y) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right].
\end{aligned}$$

Therefore, as η''_δ is bounded and Lipschitz-continuous, $\varphi \in C$ and $|Q(x, x)| \leq \Lambda_1$ for all $x \in \mathbb{T}^d$, g is a bounded function, we have

$$\begin{aligned}
|A_1| & \leq \frac{\Lambda_1}{2} \mathbb{E} \left[\int_{Q_T} \int_{Q_T} \int_0^1 \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) g^2(\hat{\mathbf{u}}(x, t, \alpha)) G_n^m \varphi(y, s) d\alpha dx dt dy ds \right] \\
& + \frac{c(\varphi)\Lambda_1}{2n} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x - y) d\alpha dx dt \rho_l(k) dk dy \right] \\
& + \frac{\Lambda_1}{2} \mathbb{E} \left[\int_{T-\frac{2}{n}}^T \int_{\mathbb{T}^d} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g^2(\hat{\mathbf{u}}(x, t, \alpha)) \rho_m(x - y) \varphi(y, t) \rho_l(k) d\alpha dx dt dk dy \right] \\
& + \frac{\|\eta''_\delta\|_\infty \Lambda_1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{Q_T} [g^2(u_\epsilon(y, s)) - g^2(u_\epsilon(y, t))] G_n^m \varphi(y, s) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{\Lambda_1}{2} \mathbb{E} \left[\int_{Q_T} \int_{Q_T} \int_{\mathbb{R}} \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) g^2(u_\epsilon(y, t)) G_n^m \varphi(y, s) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{c(\varphi)\Lambda_1}{2n} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \rho_m(x - y) dy ds \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& + \frac{\Lambda_1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{T-\frac{2}{n}}^T \int_{\mathbb{R}} \int_{\mathbb{T}^d} \eta''_\delta (u_\epsilon(y, t) - k) g^2(u_\epsilon(y, t)) \varphi(y, t) \rho_m(x - y) dy \int_0^1 \mathcal{A}_k^l d\alpha dk dx dt \right] \\
& \leq \frac{\Lambda_1}{2} \mathbb{E} \left[\int_{Q_T} \int_{Q_T} \int_0^1 \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) g^2(\hat{\mathbf{u}}(x, t, \alpha)) G_n^m \varphi(y, s) d\alpha dx dt dy ds \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{c(\varphi, \eta''_\delta) \Lambda_1}{2n} \mathbb{E} \left[\int_{Q_T} \int_0^1 g^2(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha dx dt \right] + \frac{c(\varphi, \eta''_\delta) \Lambda_1}{2} \mathbb{E} \left[\int_{T-2/n}^T \int_{\mathbb{T}^d} \int_0^1 g^2(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha dx dt \right] \\
& + \frac{c(\varphi) \Lambda_1 \|\eta''_\delta\|_\infty}{2} \mathbb{E} \left[\int_0^T \int_{Q_T} [g^2(u_\epsilon(y, s)) - g^2(u_\epsilon(y, t))] \rho_n(t-s) dy ds dt \right] \\
& + \frac{c(\varphi) \Lambda_1}{2} \mathbb{E} \left[\int_0^T \int_{Q_T} \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) g^2(u_\epsilon(y, t)) \rho_n(t-s) dy ds dt \right] \\
& + \frac{c(\varphi) \Lambda_1 \|\eta''_\delta\|_\infty}{2n} \mathbb{E} \left[\int_{Q_T} g^2(u_\epsilon(y, t)) dy dt \right] + \frac{\Lambda_1 \|\eta''_\delta\|_\infty}{2} \mathbb{E} \left[\int_{T-2/n}^T \int_{\mathbb{T}^d} g^2(u_\epsilon(y, t)) \varphi(y, t) dy dt \right] \\
& \leq \frac{c(m, \varphi, \Lambda_1, M_g^2)}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^T \int_0^1 \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) \rho_n(t-s) d\alpha dx dt dy ds \right] \\
& + \frac{c(\varphi, \Lambda_1, M_g^2)}{2} \mathbb{E} \left[\int_0^T \int_{Q_T} \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) \rho_n(t-s) dy ds dt \right] \\
& + \frac{c(\varphi, \eta''_\delta, \Lambda_1)}{2} \mathbb{E} \left[\int_0^T \int_{Q_T} [g^2(u_\epsilon(y, s)) - g^2(u_\epsilon(y, t))] \rho_n(t-s) dy ds dt \right] \\
& + \frac{c(\varphi, \eta''_\delta, \Lambda_1)}{2n} M_g^2 |\mathbb{T}^d| \\
& := B_1 + B_2 + B_3 + B_4.
\end{aligned}$$

We will prove the five terms tend to 0 respectively. For the term B_1 ,

$$\begin{aligned}
& \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^T \int_0^1 \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) \rho_n(t-s) dx dt d\alpha dy ds \right] \\
& = \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \int_0^1 \int_0^T \left[\int_0^T \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) \rho_n(t-s) ds \right] dx dt d\alpha dy \right].
\end{aligned}$$

We denoted by $A_n(t) = \int_0^T \min(2\|\eta''_\delta\|_\infty, |u_\epsilon(y, s) - u_\epsilon(y, t)|) \rho_n(t-s) ds$. Therefore

$$\begin{aligned}
|A_n(t)| & \leq \int_0^T |u_\epsilon(y, s) - u_\epsilon(y, t)| \rho_n(t-s) ds \\
& \leq Cn \int_{t-\frac{2}{n}}^t |u_\epsilon(y, s) - u_\epsilon(y, t)| ds
\end{aligned}$$

Since for almost all (ω, y, α) fixed, $s \mapsto u_\epsilon(\cdot, s) \in L^2(0, T) \subset L^1(0, T)$, by Lebesgue differentiation theorem, almost every t is a Lebesgue point such that

$$\frac{n}{2} \int_{t-\frac{2}{n}}^t |u_\epsilon(y, s) - u_\epsilon(y, t)| ds \rightarrow 0$$

as $n \rightarrow \infty$ ⁸. We thus deduce that $A_n(t)$ tends to 0 for almost every $t \in [0, T]$. And since $|A_n(t)| \leq 2M_g^2\|\eta''_\delta\|_\infty$, by Lebesgue's dominated convergent theorem

$$\begin{aligned} \lim_{n \rightarrow \infty} B_1 &= \frac{c(m, \varphi, \Lambda_1, M_g^2)}{2} \lim_{n \rightarrow \infty} \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \int_0^1 \int_0^T A_n(t) dx dt d\alpha dy \right] \\ &= \frac{c(m, \varphi, \Lambda_1, M_g^2)}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \int_0^T \int_0^1 \lim_{n \rightarrow \infty} A_n(t) dx dt d\alpha dy \right] \\ &= 0. \end{aligned}$$

We use the same argument to proof that $\lim_{n \rightarrow \infty} B_2 = 0$. As for the integral B_3 ,

$$|B_3| \leq \frac{c(\varphi, \eta''_\delta, \Lambda_1, M_g, C_g)}{2} \int_{Q_T}^T \int_0^1 |u_\epsilon(y, s) - u_\epsilon(y, t)| \rho_n(t-s) ds dy dt$$

which tends to 0 as n tends to ∞ . The last term B_4 tends to 0 as n tends to ∞ . We prove thus the limit as n tends to ∞ . One can also justify the limit as $l \rightarrow \infty$ in a similar way as in². \square

Next we study the stochastic integral term, namely I_7 .

Lemma 5.

$$\lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} I_7 = - \int_{Q_T} \int_{\mathbb{T}^d} \mathbb{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g(\hat{\mathbf{u}}(x, t, \alpha)) g(u_\epsilon(y, t)) d\alpha \right] \rho_m(x-y) \varphi(y, t) Q(x, y) dy dx dt.$$

Proof. Using the same type of arguments in the proof of Lemma 3, for a given s , setting $n \geq \frac{2}{s}$ and

$$\bar{\alpha}(s) = \rho_l(u_\epsilon(s - \frac{2}{n}, y) - k)$$

and

$$\bar{\gamma}(t) = \int_{\mathbb{T}^d} \int_0^1 \eta'_\delta(\hat{\mathbf{u}} - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dx,$$

we deduce that

$$\begin{aligned} &\mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dx dW(x, t) \varphi(y, s) \right. \\ &\quad \times \rho_m(x-y) \rho_l(u_\epsilon(s - \frac{2}{n}, y) - k) dk dy ds \Big] \\ &= \int_{Q_T} \int_{\mathbb{R}} \mathbb{E} \left[\bar{\alpha}(s) \int_{s-\frac{2}{n}}^s \bar{\gamma}(t) dW(x, t) \right] dk dy ds \\ &= 0. \end{aligned} \tag{26}$$

Therefore subtracting equality (26) from the expression of I_7 , we deduce that

$$\begin{aligned} &I_7 - 0 \\ &= \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \varphi(y, s) \rho_m(x-y) dx \rho_l(u_\epsilon(s - \frac{2}{n}, y) - k) dk dy ds \right] \end{aligned}$$

$$\begin{aligned}
& -\mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dx dW(x, t) \varphi(y, s) \rho_m(x-y) \rho_l(u_\epsilon(y, s - \frac{2}{n}) - k) dk dy ds \right] \\
& = \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \rho_m(x-y) dx \right. \\
& \quad \times \left. \varphi(y, s) \left(\rho_l(u_\epsilon(y, s) - k) - \rho_l(u_\epsilon(y, s - \frac{2}{n}) - k) \right) dk dy ds \right].
\end{aligned}$$

We define $A_\epsilon = \epsilon \Delta u_\epsilon - \operatorname{div}(\mathbf{v} f(u_\epsilon))$ and apply Itô's formula, Lemma 1 to the function $\mathcal{G}(X, t) = \rho_l(X)$. Setting

$$X(t) = u_\epsilon(y, t), \quad \psi = A_\epsilon, \quad \theta = g$$

then

$$\begin{aligned}
\mathcal{G}(X(t), t) &= \rho_l(X(t)) \\
\mathcal{G}_X(X(t), t) &= \rho'_l(X(t)) \\
\mathcal{G}_{XX}(X(t), t) &= \rho''_l(X(t)) \\
\mathcal{G}_t(X(t), t) &= 0
\end{aligned}$$

we deduce that for almost all $y \in \mathbb{T}^d$ and almost all $s \in (0, T)$, with $n \geq \frac{2}{s}$

$$\begin{aligned}
& \rho_l(u_\epsilon(y, s) - k) - \rho_l(u_\epsilon(y, s - \frac{2}{n}) - k) \\
& = \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma + \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \\
& \quad + \frac{1}{2} \int_{s-\frac{2}{n}}^s \rho''_l(u_\epsilon(y, \sigma) - k) g^2(u_\epsilon(y, \sigma)) Q(y, y) d\sigma \\
& = -\frac{\partial}{\partial k} \left[\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma + \int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right. \\
& \quad \left. + \frac{1}{2} \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) g^2(u_\epsilon(y, \sigma)) Q(y, y) d\sigma \right].
\end{aligned}$$

Thus

$$\begin{aligned}
I_7 & = -\mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \rho_m(x-y) dx \varphi(y, s) \right. \\
& \quad \times \left. \frac{\partial}{\partial k} \left[\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma + \int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right. \right. \\
& \quad \left. \left. + \frac{1}{2} \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) g^2(u_\epsilon(y, \sigma)) Q(y, y) d\sigma \right] dk dy ds \right]
\end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \frac{\partial}{\partial k} \left[\int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta'_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \rho_m(x-y) dx \varphi(y, s) \right] \right. \\
&\quad \times \left[\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma + \int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right. \\
&\quad \left. \left. + \frac{1}{2} \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) g^2(u_\epsilon(y, \sigma)) Q(y, y) d\sigma \right] dk dy ds \right] \\
&= - \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \rho_m(x-y) \varphi(y, s) dx \right. \\
&\quad \times \left[\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma + \int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right. \\
&\quad \left. \left. + \frac{1}{2} \int_{s-\frac{2}{n}}^s \rho'_l(u_\epsilon(y, \sigma) - k) g^2(u_\epsilon(y, \sigma)) Q(y, y) d\sigma \right] dk dy ds \right] \\
&=: \mathbb{I}_1 + \mathbb{I}_2 + \mathbb{I}_3
\end{aligned}$$

In the following, we prove the limits

$$\lim_{n \rightarrow \infty} \mathbb{I}_1 = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathbb{I}_3 = 0;$$

and

$$\begin{aligned}
&\lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{I}_2 \\
&= - \int_{Q_T} \int_{\mathbb{T}^d} \mathbb{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g(\hat{\mathbf{u}}(x, t, \alpha)) g(u_\epsilon(y, t)) d\alpha \right] \rho_m(x-y) Q(x, y) \varphi(y, t) dx dy dt.
\end{aligned}$$

We start by the estimate of \mathbb{I}_1 . Setting

$$G_{\mathbb{I}_1} := \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) \rho_m(x-y) \varphi(y, t),$$

and recalling the assumption that $\sum_{j=1}^{\infty} \lambda_j \|e_j\|_{L^\infty(\mathbb{T}^d)}^2 \leq \Lambda_1$, and using the properties that $|\rho_m| \leq Cm^d$, $|\rho_n| \leq Cn$ and $|\rho_l| \leq Cl$ for some constant C , Cauchy-Schwarz inequality and the Itô isometry, we obtain

$$\begin{aligned}
|\mathbb{I}_1| &= \left| \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbb{E} \left[\left(\int_{s-\frac{2}{n}}^s G_{\mathbb{I}_1} dW(x, t) \right) \left(\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma \right) \right] dx dk dy ds \right| \\
&\leq \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \left(\mathbb{E} \left[\left(\int_{s-\frac{2}{n}}^s G_{\mathbb{I}_1} dW(x, t) \right)^2 \right] \right)^{\frac{1}{2}} \left(\mathbb{E} \left[\left(\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma \right)^2 \right] \right)^{\frac{1}{2}} dx dk dy ds
\end{aligned}$$

$$\begin{aligned}
&\leq \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \left(\mathbb{E} \left[\left(\sum_{j=1}^{\infty} \sqrt{\lambda_j} e_j(x) \int_{s-\frac{2}{n}}^s G_{\mathbb{I}_1} d\beta_j(t) \right)^2 \right] \right)^{\frac{1}{2}} \left(\mathbb{E} \left[\left(\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) A_\epsilon(y, \sigma) d\sigma \right)^2 \right] \right)^{\frac{1}{2}} dx dk dy ds \\
&\leq \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \left(\sum_{j=1}^{\infty} \lambda_j e_j^2(x) \mathbb{E} \left[\int_{s-\frac{2}{n}}^s \int_0^1 (\eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k))^2 g^2(\hat{\mathbf{u}}) d\alpha \rho_n^2(t-s) \varphi^2 \rho_m^2(x-y) dt \right] \right)^{\frac{1}{2}} \\
&\quad \times \left(\int_{s-\frac{2}{n}}^s 1 d\sigma \mathbb{E} \left[\int_{s-\frac{2}{n}}^s \rho_l^2(u_\epsilon(y, \sigma) - k) A_\epsilon^2(y, \sigma) d\sigma \right] \right)^{\frac{1}{2}} dx dk dy ds \\
&\leq C(\varphi) \sqrt{\Lambda_1} \frac{m^d n l}{\sqrt{n}} \int_0^T \int_{\mathbb{T}^d} \int_{\|x-y\| < \frac{1}{m}} \int_{\mathbb{R}} \left(\mathbb{E} \left[\int_{s-\frac{2}{n}}^s \int_0^1 (\eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k))^2 g^2(\hat{\mathbf{u}}) d\alpha dt \right] \right)^{\frac{1}{2}} \\
&\quad \times \left(\mathbb{E} \left[\int_{s-\frac{2}{n}}^s \mathbf{1}_{\{|u_\epsilon(y, \sigma) - k| < \frac{1}{l}\}} A_\epsilon^2(y, \sigma) d\sigma \right] \right)^{\frac{1}{2}} dk dx dy \\
&\leq C(\varphi) \sqrt{\Lambda_1} m^d l \sqrt{n} \int_0^T \int_{\mathbb{T}^d} \int_{\|x-y\| < \frac{1}{m}} \int_{\mathbb{R}} \mathbb{E} \left[\int_{s-\frac{2}{n}}^s \int_0^1 (\eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k))^2 g^2(\hat{\mathbf{u}}(x, t, \alpha)) dx dt d\alpha dk dy ds \right] \\
&\quad + C(\varphi) \sqrt{\Lambda_1} l \sqrt{n} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{R}} \mathbb{E} \left[\int_{s-\frac{2}{n}}^s \mathbf{1}_{\{|u_\epsilon(y, \sigma) - k| < \frac{1}{l}\}} A_\epsilon^2(y, \sigma) d\sigma dk dy ds \right] \\
&\leq C(\varphi, \eta_\delta) \sqrt{\Lambda_1} m^d l \sqrt{n} \int_0^T \int_{\mathbb{T}^d} \int_{\|x-y\| < \frac{1}{m}} \mathbb{E} \left[\int_{s-\frac{2}{n}}^s \int_0^1 g^2(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha dt dx dy ds \right] \\
&\quad + C(\varphi) \sqrt{\Lambda_1} \sqrt{n} \int_0^T \int_{\mathbb{T}^d} \mathbb{E} \left[\int_{s-\frac{2}{n}}^s A_\epsilon^2(y, \sigma) d\sigma dy ds \right] \\
&\leq \frac{C(\varphi, \eta_\delta) l \sqrt{\Lambda_1}}{\sqrt{n}} \left[M_g^2 + \|A_\epsilon\|_{L^2((0,T) \times \Omega \times \mathbb{T}^d)}^2 \right],
\end{aligned}$$

which tends to 0 as n tends to ∞ . The Lemma 16 in section 4 ensures that $\|A_\epsilon\|_{L^2((0,T) \times \Omega \times \mathbb{T}^d)}^2$ is bounded. Using a similar idea and the fact that $|Q(x, y)| \leq \Lambda_1$ for all $x, y \in \mathbb{T}^d$ we prove that $\lim_{n \rightarrow \infty} \mathbb{I}_3 = 0$.

We now consider the term \mathbb{I}_2 . Using the Itô isometry, we obtain:

$$\begin{aligned}
\mathbb{I}_2 &= -\mathbb{E} \left[\int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_{s-\frac{2}{n}}^s \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \varphi(y, s) \rho_m(x-y) dx \right. \\
&\quad \times \left. \left(\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right) dk dy ds \right]
\end{aligned}$$

$$\begin{aligned}
&= - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\left(\int_{s-\frac{2}{n}}^s \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) dW(x, t) \right) \right. \\
&\quad \times \left. \left(\int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) dW(y, \sigma) \right) \right] \varphi(y, s) \rho_m(x-y) dx dk dy ds \\
&= - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\left(\sum_{j_1=0}^{\infty} \sqrt{\lambda_{j_1}} e_{j_1}(x) \int_{s-\frac{2}{n}}^s \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) d\beta_j(t)_B i g \right) \right. \\
&\quad \times \left. \left(\sum_{j=0}^{\infty} \sqrt{\lambda_j} e_j(y) \int_{s-\frac{2}{n}}^s \rho_l(u_\epsilon(y, \sigma) - k) g(u_\epsilon(y, \sigma)) d\beta_j(\sigma) \right) \right] \varphi(y, s) \rho_m(x-y) dx dk dy ds \\
&= - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_{s-\frac{2}{n}}^s \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) dt \right] \\
&\quad \times \varphi(y, s) \rho_m(x-y) Q(x, y) dx dk dy ds \\
&\xrightarrow{n \rightarrow \infty} - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \rho_m(x-y) Q(x, y) dx dk dy dt \\
&\xrightarrow{l \rightarrow \infty} - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g(\hat{\mathbf{u}}(x, t, \alpha)) g(u_\epsilon(y, t)) d\alpha \right] \varphi(y, t) \rho_m(x-y) Q(x, y) dx dy dt.
\end{aligned}$$

We check the limits in details. We define respectively A_1 the difference between \mathbb{I}_2 and the limit value on n ; and A_2 the difference between the limit value on n and the limit value on l . Therefore using that $\text{supp } \rho_n \subset [-\frac{2}{n}, 0]$

$$\begin{aligned}
A_1 &= - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^T \int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_n(t-s) \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) dt \right] \\
&\quad \times \varphi(y, s) \rho_m(x-y) Q(x, y) dx dk dy ds \\
&\quad + \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \rho_m(x-y) Q(x, y) dx dk dy dt \\
&= - \int_{\mathbb{T}^d} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \int_0^T \mathbf{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \left[\int_0^T \varphi(y, s) \rho_n(t-s) ds \right] \rho_m(x-y) Q(x, y) dx dk dy dt \\
&\quad + \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta(\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \rho_m(x-y) Q(x, y) dx dk dy dt
\end{aligned}$$

$$\begin{aligned}
&= \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \left[\varphi(y, t) - \int_0^T \varphi(y, s) \rho_n(t-s) ds \right] \rho_m(x-y) Q(x, y) dx dk dy dt \\
&= \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \int_0^T [\varphi(y, t) - \varphi(y, s)] \rho_n(t-s) ds \rho_m(x-y) Q(x, y) dx dk dy dt \\
&+ \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \left[1 - \int_0^T \rho_n(t-s) ds \right] \rho_m(x-y) Q(x, y) dx dk dy dt.
\end{aligned}$$

So that

$$\begin{aligned}
|A_1| &\leq \frac{C(\varphi)\Lambda_1}{n} \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \rho_m(x-y) dt dx dk dy \\
&\quad + \Lambda_1 \int_{T-\frac{2}{n}}^T \int_{\mathbb{T}^d} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \rho_m(x-y) dt dx dk dy \\
&\leq \frac{C(\varphi)\Lambda_1}{n} \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \rho_m(x-y) dt dx dk dy \\
&\quad + \Lambda_1 \int_{T-\frac{2}{n}}^T \int_{\mathbb{T}^d} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(k) g(u_\epsilon(y, t)) \right] \\
&\quad \times \varphi(y, t) \rho_m(x-y) dt dx dk dy \\
&\leq \frac{C(\varphi, \eta''_\delta)\Lambda_1}{n} \int_{Q_T} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha g(u_\epsilon(y, t)) \right] \rho_m(x-y) dt dx dy \\
&\quad + C(\eta''_\delta)\Lambda_1 \int_{T-\frac{2}{n}}^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha g(u_\epsilon(y, t)) \right] \varphi(y, t) \rho_m(x-y) dx dy dt \\
&\leq \frac{C(\varphi, \eta''_\delta)\Lambda_1}{n} \int_{Q_T} \int_{\mathbb{T}^d} \int_0^1 \mathbf{E} [g^2(\hat{\mathbf{u}}(x, t, \alpha)) + g^2(u_\epsilon(y, t))] d\alpha \rho_m(x-y) dt dx dy
\end{aligned}$$

$$\begin{aligned}
& + C(\eta''_\delta) \Lambda_1 \int_{T-\frac{2}{n}}^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \int_0^1 \mathbf{E} [g^2(\hat{\mathbf{u}}(x, t, \alpha)) + g^2(u_\epsilon(y, t))] \varphi(y, t) \rho_m(x - y) d\alpha dx dy dt \\
& \leq \frac{C(\varphi, \eta''_\delta, T)}{n} \Lambda_1 M_g^2 |\mathbb{T}^d|
\end{aligned}$$

which tends to 0 as $n \rightarrow \infty$. This completes the proof of the limit on n , we then proof the limit on l .

$$\begin{aligned}
A_2 = & - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - k) g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(u_\epsilon(y, t) - k) g(u_\epsilon(y, t)) \right] \\
& \times \varphi(y, s) \rho_m(x - y) Q(x, y) dx dk dy dt \\
& + \int_{Q_T} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g(\hat{\mathbf{u}}(x, t, \alpha)) g(u_\epsilon(y, t)) d\alpha \right] \\
& \times \varphi(y, t) \rho_m(x - y) Q(x, y) dx dy dt \\
= & - \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \left\{ \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t) + k) - \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) \right\} \right. \\
& \left. \times g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha g(u_\epsilon(y, t)) \right] \rho_l(k) \varphi(y, t) \rho_m(x - y) Q(x, y) dx dk dy dt.
\end{aligned}$$

Using the Lipschitz-continuity of η''_δ and the fact that $\text{supp } \rho_l \subset [-\frac{1}{l}, \frac{1}{l}]$,

$$\begin{aligned}
|A_2| & \leq c(\eta''_\delta) \Lambda_1 \int_{Q_T} \int_{\mathbb{R}} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 |k| g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha \rho_l(k) g(u_\epsilon(y, t)) \varphi(y, t) \rho_m(x - y) \right] dx dk dy dt \\
& \leq \frac{c(\eta''_\delta) \Lambda_1}{l} \int_{Q_T} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 g(\hat{\mathbf{u}}(x, t, \alpha)) d\alpha g(u_\epsilon(y, t)) \varphi(y, t) \rho_m(x - y) \right] dx dy dt \\
& \leq \frac{c(\eta''_\delta, \varphi)}{l} \Lambda_1 M_g^2 |\mathbb{T}^d|
\end{aligned}$$

which tends to zero as $l \rightarrow \infty$. \square

Combining Lemma 3, 4 and 5 yields

$$\begin{aligned}
& \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7 + J_7] = \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7] \\
& = \frac{1}{2} \mathbf{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^1 g^2(\hat{\mathbf{u}}(x, t, \alpha)) \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) \rho_m(x - y) \varphi(y, t) Q(x, x) d\alpha dy dx dt \right] \\
& + \frac{1}{2} \mathbf{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^1 g^2(u_\epsilon(y, t)) \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) \rho_m(x - y) \varphi(y, t) Q(y, y) d\alpha dy dx dt \right] \\
& - \int_{Q_T} \int_{\mathbb{T}^d} \mathbf{E} \left[\int_0^1 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) g(\hat{\mathbf{u}}(x, t, \alpha)) g(u_\epsilon(y, t)) d\alpha \right] \varphi(y, t) \rho_m(x - y) Q(x, y) dy dx dt \\
& = \frac{1}{2} \mathbf{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^1 \sum_{j=1}^{\infty} \lambda_j [g(\hat{\mathbf{u}}(x, t, \alpha)) e_j(x) - g(u_\epsilon(y, t)) e_j(y)]^2 \eta''_\delta (\hat{\mathbf{u}}(x, t, \alpha) - u_\epsilon(y, t)) \rho_m(x - y) \varphi(y, t) d\alpha dy dx dt \right].
\end{aligned}$$

In the following, we prove the lemma

Lemma 6. There holds

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{m \rightarrow \infty} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7] = 0.$$

Proof. We remark that

$$\begin{aligned} & \sum_{j=1}^{\infty} \lambda_j [g(\hat{\mathbf{u}}(x, t, \alpha))e_j(x) - g(u_{\epsilon}(y, t))e_j(y)]^2 \\ &= \sum_{j=1}^{\infty} \lambda_j [(g(\hat{\mathbf{u}}(x, t, \alpha)) - g(u_{\epsilon}(y, t)))e_j(x) + g(u_{\epsilon}(y, t))(e_j(x) - e_j(y))]^2 \\ &\leq 2Q(x, x) [g(\hat{\mathbf{u}}(x, t, \alpha)) - g(u_{\epsilon}(y, t))]^2 + 2M_g^2 \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2, \end{aligned}$$

so that

$$\begin{aligned} & \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7] \\ &= \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \sum_{j=1}^{\infty} \lambda_j [g(\hat{\mathbf{u}}(x, t, \alpha))e_j(x) - g(u_{\epsilon}(y, t))e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\leq \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} 2Q(x, x) [g(\hat{\mathbf{u}}(x, t, \alpha)) - g(u_{\epsilon}(y, t))]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\quad + \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} 2M_g^2 \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right]. \end{aligned} \tag{27}$$

First we estimate the first term on the right hand-side of (27), since $\text{supp } \eta''_{\delta} \subset [-\delta, \delta]$ and $0 \leq \eta''_{\delta} \leq 2\pi/\delta$,

$$\begin{aligned} & \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} 2Q(x, x) [g(\hat{\mathbf{u}}(x, t, \alpha)) - g(u_{\epsilon}(y, t))]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\leq \Lambda_1 C_g^2 \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} (\hat{\mathbf{u}} - u_{\epsilon})^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\leq \Lambda_1 C_g^2 \delta^2 \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\leq 2\pi \Lambda_1 C_g^2 \delta \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \rho_m(x - y) \varphi(y, t) dy dx dt \right] \\ &\leq 2\pi \Lambda_1 C_g^2 \delta \int_{Q_T} \varphi(y, t) dy dt, \end{aligned}$$

which tends to 0 as δ tends to 0. In what follows, we prove that the second term on the right-hand-side of (27) tends to 0 as $m \rightarrow \infty$, namely

$$\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x - y) \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt = 0. \tag{28}$$

We recall that

$$\sum_{j=1}^{\infty} \lambda_j \|e_j\|_{\infty}^2 \leq \Lambda_1.$$

Then for each $\epsilon_1 > 0$, there exists K_0 , such that for all $K > K_0$,

$$\begin{aligned} \sum_{j=K+1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 &\leq 2 \sum_{j=K+1}^{\infty} \lambda_j e_j^2(x) + 2 \sum_{j=K+1}^{\infty} \lambda_j e_j^2(y) \\ &\leq \epsilon_1, \end{aligned} \quad (29)$$

we choose $K > K_0$ to divide the infinite sum into 2 parts

$$\begin{aligned} &\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &= \lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=1}^K \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &\quad + \lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=K+1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt. \end{aligned} \quad (30)$$

We study the first term on the right-hand side of (30). Using the fact that $\text{supp } \rho_m \subset [-\frac{1}{2m}, \frac{1}{2m}]^d$, as well as the fact that $e_j \in C(\mathbb{T}^d)$, setting $x - y := z$ and $(e_j(x) - e_j(y))^2 := f_j(z)$, we deduce that

$$\begin{aligned} &\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=1}^K \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &= \sum_{j=1}^K M_g^2 \lim_{m \rightarrow \infty} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \lambda_j \rho_m(z) f_j(z) \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dz dy dt \\ &= \sum_{j=1}^K M_g^2 \lim_{m \rightarrow \infty} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \lambda_j m^d k_{\rho} e^{-\frac{1}{1-|mz+1|}} \chi_{\{z \in \mathbb{T}^d, |mz+1| < 1\}} f_j(z) \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dz dy dt \\ &= \sum_{j=1}^K M_g^2 \lim_{m \rightarrow \infty} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \lambda_j m^d k_{\rho} e^{-\frac{1}{1-|\tau|}} \chi_{\{\tau \in \mathbb{T}^d, |\tau| < 1\}} f_j(\frac{\tau-1}{m}) \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) \frac{d\tau}{m^d} dy dt \\ &= \sum_{j=1}^K M_g^2 \lim_{m \rightarrow \infty} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \lambda_j k_{\rho} e^{-\frac{1}{1-|\tau|}} \chi_{\{\tau \in \mathbb{T}^d, |\tau| < 1\}} f_j(\frac{\tau-1}{m}) \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) d\tau dy dt \\ &= 0. \end{aligned}$$

As for the second term, we use (29) to obtain,

$$\begin{aligned} &\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=K+1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &\leq \epsilon_1 M_g^2 \lim_{m \rightarrow \infty} \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &\leq \epsilon_1 M_g^2 T |\mathbb{T}^d| \|\eta''_{\delta}\|_{L^{\infty}((0,T) \times \mathbb{T}^d)} \|\varphi\|_{L^{\infty}((0,T) \times \mathbb{T}^d)}. \end{aligned}$$

In view of (30),

$$\begin{aligned} &\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt \\ &\leq \epsilon_1 M_g^2 T |\mathbb{T}^d| \|\eta''_{\delta}\|_{L^{\infty}(\mathbb{T}^d)} \|\varphi\|_{L^{\infty}(\mathbb{T}^d)} \end{aligned}$$

for all $\epsilon_1 > 0$, i.e.

$$\lim_{m \rightarrow \infty} M_g^2 \int_0^T \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \rho_m(x-y) \sum_{j=1}^{\infty} \lambda_j [e_j(x) - e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \varphi(y, t) dx dy dt = 0.$$

Recall that

$$\begin{aligned} & \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7] \\ &= \frac{1}{2} \mathbb{E} \left[\int_{Q_T} \int_{\mathbb{T}^d} \int_0^1 \sum_{j=1}^{\infty} \lambda_j [g(\hat{\mathbf{u}}(x, t, \alpha)) e_j(x) - g(u_{\epsilon}(y, t)) e_j(y)]^2 \eta''_{\delta}(\hat{\mathbf{u}}(x, t, \alpha) - u_{\epsilon}(y, t)) \rho_m(x-y) \varphi(y, t) d\alpha dy dx dt \right], \end{aligned}$$

and in view of (27), we deduce that

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \lim_{m \rightarrow \infty} \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [I_6 + J_6 + I_7] = 0.$$

which complete the proof of Lemma 6. \square

Passing to the limits in $\sum_{i=1}^7 \{I_i + J_i\} + J_8$ by n, l, δ, ϵ and m , we obtain

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\int_{\mathbb{T}^d} |u_0 - \hat{u}_0| \varphi(0) dx \right] \\ &+ \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 |\mathbf{u}(x, t, \beta) - \hat{\mathbf{u}}(x, t, \alpha)| \partial_t \varphi(x, t) d\alpha d\beta dx dt \right] \\ &+ \mathbb{E} \left[\int_{Q_T} \int_0^1 \int_0^1 F(\mathbf{u}(x, t, \beta), \hat{\mathbf{u}}(x, t, \alpha)) \mathbf{v} \cdot \nabla \varphi(x, t) d\alpha d\beta dx dt \right], \end{aligned}$$

which completes the proof of the Kato inequality. \square

4 | EXISTENCE AND UNIQUENESS OF THE STRONG SOLUTION OF THE ASSOCIATED PARABOLIC PROBLEM

In this section, we prove the existence and uniqueness of the strong solution of the associated stochastic nonlinear parabolic problem; to that purpose we apply a time discretization method.

Definition 4. We say that u_{ϵ} is a strong solution of the problem

$$(P^{\epsilon}) \quad \begin{cases} du_{\epsilon} - \epsilon \Delta u_{\epsilon} dt + \operatorname{div}(\mathbf{v} f(u_{\epsilon})) dt = g(u_{\epsilon}) dW & \text{in } \Omega \times \mathbb{T}^d \times (0, T), \\ u_{\epsilon}(x, 0) = u_0^{\epsilon}(x) & \text{for all } x \in \mathbb{T}^d \end{cases} \quad (31)$$

with initial condition $u_0^{\epsilon}(x) \in H^1(\mathbb{T}^d)$, if u_{ϵ} satisfies

1. $u_{\epsilon} \in L^{\infty}(0, T; L^2(\Omega; L^2(\mathbb{T}^d))) \cap L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$ is an $L^2(\mathbb{T}^d)$ -adapted process to the filtration (\mathcal{F}_t) ;

2. for almost all $t \in (0, T)$, P-a.s. there holds,

$$u_{\epsilon}(\cdot, t) = u_0^{\epsilon} + \epsilon \int_0^t \Delta u_{\epsilon} ds - \int_0^t \operatorname{div}(\mathbf{v} f(u_{\epsilon})) ds + \int_0^t g(u_{\epsilon}) dW(s), \quad (32)$$

in $L^2(\mathbb{T}^d)$.

We suppose that $u_0^{\epsilon} \rightarrow u_0$ strongly in $L^2(\mathbb{T}^d)$ as $\epsilon \rightarrow 0$, where u_0 is the initial condition of Problem (1). The following result is the analog of Proposition A.2 in².

Theorem 2. Let $\epsilon > 0$ be arbitrary. Problem (\mathcal{P}^ϵ) admits a unique strong solution.

We first perform an implicit time discretization with $\Delta t = T/N$. The scheme is as follows: for a given small positive constant Δt and u_n^ϵ in $L^2(\Omega; H^1(\mathbb{T}^d))$, $\mathcal{F}_{n\Delta t}$ -measurable, find u_{n+1}^ϵ in $L^2(\Omega; H^1(\mathbb{T}^d))$, $\mathcal{F}_{(n+1)\Delta t}$ -measurable, such that P-a.s. and for all v in $H^1(\mathbb{T}^d)$

$$\int_{\mathbb{T}^d} [(u_{n+1}^\epsilon - u_n^\epsilon)v + \Delta t(\epsilon \nabla u_{n+1}^\epsilon \cdot \nabla v - f(u_{n+1}^\epsilon)v \cdot \nabla v)] dx = \int_{\mathbb{T}^d} g(u_n^\epsilon)v(W_{n+1} - W_n) dx \quad (33)$$

where $W_n = W(t_n) = \sum_{j=1}^{\infty} \sqrt{\lambda_j} \beta_j(t_n) e_j(x)$; we remark that $u_0^\epsilon = u_0^\epsilon(x)$. Before pursuing the proof of the Theorem, we first show the existence of the sequence $\{u_n^\epsilon\}_{n \geq 0}$.

Lemma 7. If $\Delta t < \frac{2\epsilon}{(VC_f)^2}$, Problem (33) possesses a unique solution $u_{n+1}^\epsilon \in L^2(\Omega; H^1(\mathbb{T}^d))$.

Proof. We set $\mathbb{V} = L^2(\Omega; H^1(\mathbb{T}^d), \mathcal{F}_{(n+1)\Delta t}, dP)$, $\mathbb{H} = L^2(\Omega; L^2(\mathbb{T}^d), \mathcal{F}_{(n+1)\Delta t}, dP)$; suppose that $u_n \in \mathbb{H}$ and define the application T_a by $\bar{u} = T_a(S)$ for all $S \in \mathbb{H}$ where $u \in \mathbb{V}$ satisfies the variational problem,

$$\mathbb{E} \left[\int_{\mathbb{T}^d} (\bar{u} - u_n^\epsilon)v + \Delta t(\epsilon \nabla \bar{u} \cdot \nabla v - f(S)v \cdot \nabla v) dx \right] = \mathbb{E} \left[\int_{\mathbb{T}^d} g(u_n^\epsilon)v(W_{n+1} - W_n) dx \right] \quad (34)$$

for all $v \in \mathbb{V}$. Since $g(u_n)$ and $W_{n+1} - W_n$ are independent variables a.e. in \mathbb{T}^d and $\mathbb{E}[W_{n+1} - W_n] = 0$, so the right hand side of (34) equals to 0. We define the bilinear form

$$a(u, v) = \mathbb{E} \left[\int_{\mathbb{T}^d} uv + \Delta t \epsilon \nabla u \cdot \nabla v dx \right]$$

and the linear form

$$l(v) = \mathbb{E} \left[\int_{\mathbb{T}^d} u_n^\epsilon v + \Delta t f(S)v \cdot \nabla v dx \right].$$

We apply the Lax-Milgram theorem for $a(\cdot, \cdot)$ and $l(\cdot)$ in the space \mathbb{V} to deduce that T_a is a well-defined map. Next we write the difference of the equality (34) for a function S_1 and the corresponding solution \bar{u}_1^ϵ and for a function S_2 and the corresponding solution \bar{u}_2^ϵ , and set $v = \bar{u}_1^\epsilon - \bar{u}_2^\epsilon$ to deduce that

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} |\bar{u}_1^\epsilon - \bar{u}_2^\epsilon|^2 + \Delta t \epsilon |\nabla(\bar{u}_1^\epsilon - \bar{u}_2^\epsilon)|^2 dx \right] \\ &= \Delta t \mathbb{E} \left[\int_{\mathbb{T}^d} (\mathbf{v}f(S_1) - \mathbf{v}f(S_2)) \cdot \nabla(\bar{u}_1^\epsilon - \bar{u}_2^\epsilon) dx \right] \end{aligned}$$

which implies that

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} |T(S_1) - T(S_2)|^2 + \frac{\Delta t \epsilon}{2} |\nabla(T(S_1) - T(S_2))|^2 dx \right] \\ &\leq \frac{\Delta t}{2\epsilon} \mathbb{E} \left[\int_{\mathbb{T}^d} |\mathbf{v}f(S_1) - \mathbf{v}f(S_2)|^2 dx \right], \end{aligned}$$

where we have used Young's inequality $ab \leq \frac{1}{2\epsilon}a^2 + \frac{\epsilon}{2}b^2$ for $a, b \in \mathbb{R}$. Thus if $\Delta t < \frac{2\epsilon}{(VC_f)^2}$, T_a is a strict contraction from \mathbb{H} to \mathbb{H} and the result of the Lemma 7 follows from the Banach fixed point theorem. \square

Next, we prove some a priori estimates.

Lemma 8. We define

$$u^{\Delta t}(t) = \sum_{n=0}^{N-1} u_{n+1}^\epsilon \cdot \mathbf{1}_{[n\Delta t, (n+1)\Delta t)}(t)$$

where $N = T/\Delta t$, then we have the following estimates

$$\begin{aligned}\|u^{\Delta t}\|_{L^\infty(0,T;L^2(\Omega;L^2(\mathbb{T}^d)))} &\leq C_1, \\ \|\nabla u^{\Delta t}\|_{L^2(0,T;(L^2(\Omega;L^2(\mathbb{T}^d)))^d)} &\leq C_2,\end{aligned}$$

where the constant C_1 depends on u_0, Λ_1, M_g and T , and the constant C_2 depends on $\epsilon, u_0, \Lambda_1, M_g$ and T . Thus,

$$\|u^{\Delta t}\|_{L^2(0,T;L^2(\Omega;H^1(\mathbb{T}^d)))} \leq C$$

for a constant C which depends on $\epsilon, u_0, \Lambda_1, M_g$ and T . We define the linear interpolation function

$$\tilde{u}^{\Delta t} = \sum_{n=1}^N \left[\frac{u_n^\epsilon - u_{n-1}^\epsilon}{\Delta t} [t - (n-1)\Delta t] + u_{n-1}^\epsilon \right] \cdot \mathbf{1}_{[(n-1)\Delta t, n\Delta t]}, \quad (35)$$

so that $\tilde{u}^{\Delta t}$ is also bounded in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d))) \cap L^\infty(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$, and $\tilde{u}^{\Delta t} - u^{\Delta t}$ converges to 0 in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$.

Thus there exist a subsequence of $\{\Delta t\}$ which we denote by $\{\Delta t_j\}$ and functions $u \in L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d))), f_u \in L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ and $g_u \in L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ such that

$$u^{\Delta t_j} \rightharpoonup u$$

and

$$\tilde{u}^{\Delta t_j} \rightharpoonup u$$

weakly in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$ as $\Delta t_j \rightarrow 0$, and

$$f(u^{\Delta t_j}) \rightharpoonup f_u \quad \text{and} \quad g(u^{\Delta t_j}) \rightharpoonup g_u$$

weakly in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ as $\Delta t_j \rightarrow 0$.

Proof. Setting the test function $v = u_{n+1}^\epsilon$ in (33) and taking the expectation yields

$$\begin{aligned}&\frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (|u_{n+1}^\epsilon|^2 - |u_n^\epsilon|^2 + |u_{n+1}^\epsilon - u_n^\epsilon|^2) dx \right] + \Delta t \epsilon \mathbb{E} \left[\int_{\mathbb{T}^d} |\nabla u_{n+1}^\epsilon|^2 dx \right] - \Delta t \mathbb{E} \left[\int_{\mathbb{T}^d} f(u_{n+1}^\epsilon) \mathbf{v} \cdot \nabla u_{n+1}^\epsilon dx \right] \\&= \mathbb{E} \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) g(u_n^\epsilon) (u_{n+1}^\epsilon - u_n^\epsilon) dx \right] + \mathbb{E} \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) g(u_n^\epsilon) u_n^\epsilon dx \right].\end{aligned} \quad (36)$$

Set $F(s) = \int_0^s f(\tau) d\tau$ and use the fact that $\operatorname{div} \mathbf{v} = 0$; then for all functions $v \in H^1(\mathbb{T}^d)$

$$\begin{aligned}\int_{\mathbb{T}^d} f(v) \mathbf{v} \cdot \nabla v dx &= \int_{\mathbb{T}^d} \mathbf{v} \cdot \nabla F(v) dx \\&= - \int_{\mathbb{T}^d} F(v) \operatorname{div} \mathbf{v} dx \\&= 0.\end{aligned}$$

Next we use (8) and (9) and that $\mathbb{E} \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) g(u_n^\epsilon) u_n^\epsilon dx \right] = 0$, and the monotone convergence theorem to deduce that

$$\begin{aligned}\mathbb{E} [(W_{n+1} - W_n)^2] &= \mathbb{E} \left[\left(\sum_{j=0}^{\infty} \sqrt{\lambda_j} (\beta_j(t_{n+1}) - \beta_j(t_n)) e_j(x) \right)^2 \right] \\&= \mathbb{E} \left[\sum_{j=0}^{\infty} \lambda_j (\beta_j(t_{n+1}) - \beta_j(t_n))^2 e_j^2(x) \right]\end{aligned} \quad (37)$$

$$\begin{aligned}
&= \sum_{j=0}^{\infty} \mathbb{E} \left[\lambda_j (\beta_j(t_{n+1}) - \beta_j(t_n))^2 e_j^2(x) \right] \\
&= \Delta t Q(x, x) \leq \Lambda_1 \Delta t.
\end{aligned}$$

Applying Young's inequality and (37) yields

$$\begin{aligned}
&\mathbb{E} \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) g(u_n^\epsilon) (u_{n+1}^\epsilon - u_n^\epsilon) dx \right] \\
&\leq \frac{1}{2\bar{\epsilon}} \mathbb{E} \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n)^2 g^2(u_n^\epsilon) dx \right] + \frac{\bar{\epsilon}}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (u_{n+1}^\epsilon - u_n^\epsilon)^2 dx \right] \\
&\leq \frac{\Delta t}{2\bar{\epsilon}} \int_{\mathbb{T}^d} \mathbb{E} [g^2(u_n^\epsilon)] Q(x, x) dx + \frac{\bar{\epsilon}}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (u_{n+1}^\epsilon - u_n^\epsilon)^2 dx \right],
\end{aligned} \tag{38}$$

for all $\bar{\epsilon} > 0$. We choose $\bar{\epsilon} = \frac{1}{2}$ and deduce from (36) that

$$\begin{aligned}
&\frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (|u_{n+1}^\epsilon|^2 - |u_n^\epsilon|^2 + |u_{n+1}^\epsilon - u_n^\epsilon|^2) dx \right] + \Delta t \epsilon \mathbb{E} \left[\int_{\mathbb{T}^d} |\nabla u_{n+1}^\epsilon|^2 dx \right] \\
&\leq \Delta t \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_n^\epsilon) Q(x, x) dx \right] + \frac{1}{4} \mathbb{E} \left[\int_{\mathbb{T}^d} (u_{n+1}^\epsilon - u_n^\epsilon)^2 dx \right] \\
&\leq \Lambda_1 \Delta t \int_{\mathbb{T}^d} \mathbb{E} [g^2(u_n^\epsilon)] dx + \frac{1}{4} \mathbb{E} \left[\int_{\mathbb{T}^d} (u_{n+1}^\epsilon - u_n^\epsilon)^2 dx \right].
\end{aligned} \tag{39}$$

We denote by $\|\cdot\|$ the norm in $L^2(\mathbb{T}^d)$, change the index in (39) to k and sum from 0 to $n-1$, $n \geq 1$, to obtain,

$$\begin{aligned}
&\frac{1}{2} \mathbb{E} [\|u_n^\epsilon\|^2] + \frac{1}{4} \sum_{k=0}^{n-1} \mathbb{E} [\|u_{k+1}^\epsilon - u_k^\epsilon\|^2] + \Delta t \epsilon \sum_{k=0}^{n-1} \mathbb{E} [\|\nabla u_{k+1}^\epsilon\|^2] \\
&\leq \frac{1}{2} \|u_0^\epsilon\|^2 + \Lambda_1 \Delta t \sum_{k=0}^{n-1} \mathbb{E} [\|g(u_k^\epsilon)\|^2] \\
&\leq \frac{1}{2} \|u_0^\epsilon\|^2 + \Lambda_1 T M_g^2.
\end{aligned} \tag{40}$$

so that

$$\mathbb{E} [\|u_n^\epsilon\|^2] \leq C_1 \tag{41}$$

where the constant C_1 depends on u_0 , Λ_1 , M_g and T , and does not depend on n . Thus

$$\begin{aligned}
\mathbb{E} [\|f(u_n^\epsilon)\|^2] &= \mathbb{E} [\|f(u_n^\epsilon) - f(0)\|^2] \\
&\leq C_f^2 \mathbb{E} [\|u_n^\epsilon\|^2] \\
&\leq C_f^2 C_1,
\end{aligned} \tag{42}$$

for all n . We recall the definition

$$u^{\Delta t} = \sum_{n=0}^{N-1} u_{n+1}^\epsilon \cdot \mathbf{1}_{[n\Delta t, (n+1)\Delta t)} \tag{43}$$

where $N = T/\Delta t$. In view of (41), we obtain

$$\|u^{\Delta t}\|_{L^\infty(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))} \leq C_1 \tag{44}$$

and by (40) we deduce that

$$\sqrt{\epsilon} \|\nabla u^{\Delta t}\|_{L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))} \leq C_2 \tag{45}$$

where the constant C_2 depends on C_1 . In turn (44) and (45) imply that

$$\|\tilde{u}^{\Delta t}\|_{L^2(0,T;L^2(\Omega;H^1(\mathbb{T}^d)))} \leq C \quad (46)$$

for a constant C which depends on ϵ and T . In view of (40) and (41)

$$\begin{aligned} & \|\tilde{u}^{\Delta t} - u^{\Delta t}\|_{L^2(0,T;L^2(\Omega \times \mathbb{T}^d))}^2 \\ &= E \left[\int_{\mathbb{T}^d} \sum_{n=1}^N \int_{(n-1)\Delta t}^{n\Delta t} (u_n^\epsilon - u_{n-1}^\epsilon)^2 \left(\frac{t - n\Delta t}{\Delta t} \right)^2 \mathbf{1}_{[(n-1)\Delta t, n\Delta t]} dt dx \right] \\ &= \frac{1}{3} \Delta t \sum_{n=1}^N E \left[\|u_n^\epsilon - u_{n-1}^\epsilon\|_{L^2(\mathbb{T}^d)}^2 \right] \\ &\leq \frac{1}{3} \Delta t \left(\frac{1}{2} \|u_0^\epsilon\|^2 + \Lambda_1 T M_g^2 \right) \end{aligned} \quad (47)$$

which converges to 0 as Δt tends to 0.

Then we prove that

$$\nabla(u^{\Delta t} - \tilde{u}^{\Delta t}) \rightharpoonup 0,$$

weakly in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)^d))$.

Considering a test function $\phi(x, t) = \sum_{n=1}^N \varphi(x) \frac{n\Delta t - t}{\Delta t} \mathbf{1}_{[(n-1)\Delta t, n\Delta t]}$ with $\varphi(x) \in H^1(\mathbb{T}^d)^d$, so that $\phi \in L^2(0, T; H^1(\mathbb{T}^d)^d)$. Using the definitions of $u^{\Delta t}$ and of $\tilde{u}^{\Delta t}$,

$$\begin{aligned} & E \left[\int_{\mathbb{T}^d} \int_0^T \nabla(u^{\Delta t} - \tilde{u}^{\Delta t}) \phi(x, t) dt dx \right] \\ &= E \left[\int_{\mathbb{T}^d} \int_0^T \nabla \left(u_n^\epsilon - \sum_{n=1}^N \frac{u_n^\epsilon - u_{n-1}^\epsilon}{\Delta t} [t - (n-1)\Delta t] - u_{n-1}^\epsilon \right) \varphi(x) \mathbf{1}_{[(n-1)\Delta t, n\Delta t]} dt dx \right] \\ &= E \left[\int_{\mathbb{T}^d} \sum_{n=1}^N \int_{(n-1)\Delta t}^{n\Delta t} \left(1 - \frac{t - (n-1)\Delta t}{\Delta t} \right) (\nabla u_n^\epsilon - \nabla u_{n-1}^\epsilon) \varphi(x) dt dx \right] \\ &= E \left[\frac{\Delta t}{2} \int_{\mathbb{T}^d} \sum_{n=1}^N (\nabla u_n^\epsilon - \nabla u_{n-1}^\epsilon) \varphi(x) dx \right] \\ &= -E \left[\frac{\Delta t}{2} \sum_{n=1}^N \int_{\mathbb{T}^d} (u_n^\epsilon - u_{n-1}^\epsilon) \operatorname{div} \varphi(x) dx \right] \end{aligned}$$

since by Cauchy-Schwarz inequality

$$\left| E \left[\int_{\mathbb{T}^d} (u_n^\epsilon - u_{n-1}^\epsilon) \operatorname{div} \varphi(x) dx \right] \right| \leq E \left[\int_{\mathbb{T}^d} (u_n^\epsilon - u_{n-1}^\epsilon)^2 dx \right] E \left[\int_{\mathbb{T}^d} |\operatorname{div} \varphi(x)|^2 dx \right],$$

so that

$$\begin{aligned} \left| E \left[\int_{\mathbb{T}^d} \int_0^T \nabla(u^{\Delta t} - \tilde{u}^{\Delta t}) \phi(x, t) dt dx \right] \right| &= \left| E \left[\frac{\Delta t}{2} \sum_{n=1}^N \int_{\mathbb{T}^d} (u_n^\epsilon - u_{n-1}^\epsilon) \operatorname{div} \varphi(x) dx \right] \right| \\ &\leq \frac{\Delta t}{2} \sum_{n=1}^N E \left[\int_{\mathbb{T}^d} (u_n^\epsilon - u_{n-1}^\epsilon)^2 dx \right] E \left[\int_{\mathbb{T}^d} |\operatorname{div} \varphi(x)|^2 dx \right] \end{aligned}$$

which converges to 0 in view of (40) and this implies $u^{\Delta t} - \tilde{u}^{\Delta t}$ converges weakly to 0 in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$.

As g is a bounded function, it implies that $g(u^{\Delta t})$ is bounded in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ uniformly in Δt . A similar computation,

$$\begin{aligned} \mathbb{E} [\|f(u_n)\|^2] &= \mathbb{E} [\|f(u_n) - f(0)\|^2] \\ &\leq C_f^2 \mathbb{E} [\|u_n\|^2] \\ &\leq C_f^2 C_1, \end{aligned} \quad (48)$$

yields that $f(u^{\Delta t})$ is bounded in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ uniformly in Δt . In view of (46), there exist a function $u \in L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$ and a subsequence $\{u^{\Delta t_j}\}$ of $\{u^{\Delta t}\}$ which converges weakly to u in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$ as Δt_j tends to zero. Moreover (42) and the assumption on g imply that there exist functions f_u and g_u and a subsequence of $\{u^{\Delta t_j}\}$, which we denote again by $\{u^{\Delta t_j}\}$, such that $f(u^{\Delta t_j})$ converges weakly to f_u and $g(u^{\Delta t_j})$ converges weakly to g_u in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ as Δt_j tends to zero. This completes the proof of Lemma 8. \square

For later use, we define

$$\tilde{B}^{\Delta t_j} = \sum_{n=1}^N \left[\frac{B_n - B_{n-1}}{\Delta t_j} [t - (n-1)\Delta t_j] + B_{n-1} \right] \cdot \mathbf{1}_{[(n-1)\Delta t_j, n\Delta t_j]} \quad (49)$$

where

$$B_n = \sum_{k=0}^{n-1} (W_{k+1} - W_k) g(u_k) = \int_0^{n\Delta t_j} g(u^{\Delta t_j}(\cdot - \Delta t_j)) dW, \quad (50)$$

in which we set $u^{\Delta t_j}(s) = u^{\Delta t_j}(0) = u_0$, for all $s < 0$. One can check that $\tilde{B}^{\Delta t_j} \in L^\infty(0, T; L^2(\Omega \times \mathbb{T}^d))$. Next, we identify the weak limit of $\int_0^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s)$ when $\Delta t_j \rightarrow 0$ in $L^2(\Omega; L^2(\mathbb{T}^d))$ for all $t \in [0, T]$.

Lemma 9. We have the following limit property

$$\int_0^t g(u^{\Delta t_j}(\cdot - \Delta t_j)) dW(x, t) \rightarrow \int_0^t g_u dW(x, t) \quad (51)$$

as Δt_j , weakly in $L^2(\Omega; L^2(\mathbb{T}^d))$ for all $t \in [0, T]$.

Proof. We set $u^{\Delta t_j}(s) = u_0$, for all $s < 0$.

We define the operator $\tilde{\mathcal{T}}$ as

$$\tilde{\mathcal{T}}(\mathcal{U}) = \int_0^t \mathcal{U} dW(x, s) \quad (52)$$

where $W(x, t)$ is the Q -Brownian motion defined by (3).

In view of the Itô isometry, and the hypothesis (4), we have that

$$\begin{aligned} \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\tilde{\mathcal{T}}(\mathcal{U}) \right)^2 dx \right] &= \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_0^t \mathcal{U} dW(x, s) \right)^2 dx \right] \\ &= \mathbb{E} \left[\int_{\mathbb{T}^d} \sum_{j=1}^{\infty} \lambda_j \left(\int_0^t \mathcal{U}(x, s) d\beta_j(s) \right)^2 e_j^2(x) dx \right. \\ &\quad \left. + \int_{\mathbb{T}^d} 2dx \sum_{j_1 \neq j_2}^{\infty} \sqrt{\lambda_{j_1} \lambda_{j_2}} e_{j_1}(x) e_{j_2}(x) \int_0^t \mathcal{U}(x, s) d\beta_{j_1}(s) \int_0^t \mathcal{U}(x, s) d\beta_{j_2}(s) \right] \\ &= \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^t \left(\sum_{j=1}^{\infty} \lambda_j e_j^2(x) \right) \mathcal{U}^2(x, s) ds dx \right] + 0 \\ &\leq \Lambda_1 \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^t \mathcal{U}^2(x, s) ds dx \right] \end{aligned}$$

$$\leq \Lambda_1 \|\mathcal{U}\|_{L^2(\Omega \times Q_T)}^2.$$

This proves that $\tilde{\mathcal{T}}$ is a bounded operator from $L^2(\Omega \times Q_T)$ to $L^2(\Omega \times \mathbb{T}^d)$. Let $\tilde{\mathcal{T}}^* : L^2(\Omega \times \mathbb{T}^d) \rightarrow L^2(\Omega \times Q_T)$ be the adjoint operator of $\tilde{\mathcal{T}}$; then

$$(\psi, \tilde{\mathcal{T}}\mathcal{U})_{L^2(\Omega \times \mathbb{T}^d)} = (\tilde{\mathcal{T}}^*\psi, \mathcal{U})_{L^2(\Omega \times Q_T)}$$

for all $\psi \in L^2(\Omega \times \mathbb{T}^d)$. Next we set $\mathcal{U} = g(u^{\Delta t_j}(\cdot - \Delta t_j))$. We recall that,

$$g(u^{\Delta t_j}(\cdot - \Delta t_j)) \rightharpoonup g_u$$

weakly in $L^2(\Omega \times Q_T)$ along a subsequence as Δt_j tend to zero. Thus,

$$\begin{aligned} (\psi, \tilde{\mathcal{T}}(g(u^{\Delta t_j}(\cdot - \Delta t_j))))_{L^2(\Omega \times \mathbb{T}^d)} &= (\tilde{\mathcal{T}}^*\psi, g(u^{\Delta t_j}(\cdot - \Delta t_j)))_{L^2(\Omega \times Q_T)} \\ &\rightarrow (\tilde{\mathcal{T}}^*\psi, g_u)_{L^2(\Omega \times Q_T)} \\ &= (\psi, \tilde{\mathcal{T}}(g_u))_{L^2(\Omega \times \mathbb{T}^d)}. \end{aligned}$$

We conclude that along a subsequence

$$\tilde{\mathcal{T}}(g(u^{\Delta t_j}(\cdot - \Delta t_j))) \rightharpoonup \tilde{\mathcal{T}}(g_u) \quad \text{weakly in } L^2(\Omega \times \mathbb{T}^d),$$

as $\Delta t_j \rightarrow 0$, or in other words, in view of the definition (52) of $\tilde{\mathcal{T}}$,

$$\int_0^t g(u^{\Delta t_j}(\cdot - \Delta t_j)) dW(x, t) \rightarrow \int_0^t g_u dW(x, t)$$

weakly in $L^2(\Omega; L^2(\mathbb{T}^d))$ for all $t \in [0, T]$. \square

Lemma 10. For almost all $t \in (0, T)$, P-a.s., there holds,

$$u(t) = u_0^\epsilon + \epsilon \int_0^t \Delta u ds - \int_0^t \operatorname{div}(\mathbf{v} f_u) ds + \int_0^t g_u dW(s),$$

in $(H^1(\mathbb{T}^d))'$, where u is the weak limit of $u^{\Delta t_j}$ in $L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))$ and its weak-* limit in $L^\infty(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$.

Proof. For all $t \in [0, T]$, there exists n such that $t \in [n\Delta t_j, (n+1)\Delta t_j]$. We return to (33) and suppose that the test function $v \in L^2(\Omega; H^1(\mathbb{T}^d))$ and is \mathcal{F}_t -adapted, and sum k from 0 to n to deduce that P-a.s.

$$\begin{aligned} &\sum_{k=0}^n \int_{\mathbb{T}^d} (u_{k+1}^\epsilon - u_k^\epsilon) v dx + \Delta t_j \sum_{k=0}^n \int_{\mathbb{T}^d} (\epsilon \nabla u_{k+1}^\epsilon \cdot \nabla v - f(u_{k+1}^\epsilon) \mathbf{v} \cdot \nabla v) dx \\ &= \sum_{k=0}^n \int_{\mathbb{T}^d} [vg(u_k^\epsilon)(W_{k+1} - W_k)] dx, \end{aligned}$$

which yields

$$\begin{aligned} &\int_{\mathbb{T}^d} (u_{n+1}^\epsilon v - u_0^\epsilon v) dx + \Delta t_j \sum_{k=0}^n \int_{\mathbb{T}^d} (\epsilon \nabla u_{k+1}^\epsilon \cdot \nabla v - f(u_{k+1}^\epsilon) \mathbf{v} \cdot \nabla v) dx \\ &= \Delta t_j \sum_{k=0}^n \int_{\mathbb{T}^d} \left[vg(u_k^\epsilon) \frac{W_{k+1} - W_k}{\Delta t_j} \right] dx \end{aligned}$$

and by definition (43) it is equivalent to

$$\begin{aligned}
& \mathbb{E} \left[\int_{\mathbb{T}^d} (u^{\Delta t_j}(t)v - u_0^\epsilon v) dx \right] + \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_0^t \epsilon \nabla u^{\Delta t_j}(s) - f(u^{\Delta t_j}(s)) \mathbf{v} ds \right) \cdot \nabla v dx \right] \\
& - \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_{n\Delta t_j}^t \epsilon \nabla u^{\Delta t_j}(s) - f(u^{\Delta t_j}(s)) \mathbf{v} ds \right) \cdot \nabla v dx \right] \\
& = \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_0^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s) \right) v dx \right] - \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_{n\Delta t_j}^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s) \right) v dx \right].
\end{aligned} \tag{53}$$

Next we estimate the terms involving the time integral on $[n\Delta t_j, t]$.

$$\begin{aligned}
\left| \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{n\Delta t_j}^t \epsilon \nabla u^{\Delta t_j}(s) \cdot \nabla v ds dx \right] \right| & \leq \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{n\Delta t_j}^t |\epsilon \nabla u^{\Delta t_j}(s) \cdot \nabla v| ds dx \right] \\
& \leq \left(\mathbb{E} \left[\int_{\mathbb{T}^d} \int_{n\Delta t_j}^t |\nabla v|^2 ds dx \right] \right)^{1/2} \left(\mathbb{E} \left[\int_{\mathbb{T}^d} \int_{n\Delta t_j}^t |\epsilon \nabla u^{\Delta t_j}(s)|^2 ds dx \right] \right)^{1/2} \\
& \leq \epsilon \sqrt{\Delta t_j} \|v\|_{L^2(\Omega; H^1(\mathbb{T}^d))} \|u^{\Delta t_j}(s)\|_{L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d)))}.
\end{aligned}$$

We deduce from a similar computation that

$$\left| \mathbb{E} \left[\int_{\mathbb{T}^d} \int_{n\Delta t_j}^t f(u^{\Delta t_j}(s)) \mathbf{v} ds \cdot \nabla v dx \right] \right| \leq \epsilon V \sqrt{\Delta t_j} \|v\|_{L^2(\Omega; H^1(\mathbb{T}^d))} \|f(u^{\Delta t_j}(s))\|_{L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))}.$$

Also in view of (37),

$$\begin{aligned}
& \left| \mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_{n\Delta t_j}^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s) \right) v dx \right] \right| \\
& \leq \left(\mathbb{E} \left[\int_{\mathbb{T}^d} v^2 dx \right] \right)^{1/2} \left(\mathbb{E} \left[\int_{\mathbb{T}^d} \left(\int_{n\Delta t_j}^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s) \right)^2 dx \right] \right)^{1/2} \\
& = \|v\|_{L^2(\Omega; L^2(\mathbb{T}^d))} \left(\mathbb{E} \left[\int_{\mathbb{T}^d} \left(g(u_n^\epsilon) \int_{n\Delta t_j}^t 1 dW(s) \right)^2 dx \right] \right)^{1/2} \\
& = \|v\|_{L^2(\Omega; L^2(\mathbb{T}^d))} \left(\mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_n^\epsilon) (W(t) - W_n)^2 dx \right] \right)^{1/2} \\
& \leq \Delta t_j \Lambda_1 M_g \|v\|_{L^2(\Omega; L^2(\mathbb{T}^d))}
\end{aligned}$$

We recall that as $\Delta t_j \rightarrow 0$,

$$\begin{aligned}
u^{\Delta t_j} & \rightharpoonup u \quad \text{in } L^2(0, T; L^2(\Omega; H^1(\mathbb{T}^d))); \\
f(u^{\Delta t_j}) & \rightharpoonup f_u \quad \text{in } L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)));
\end{aligned}$$

$$\int_0^t g(u^{\Delta t_j}(s - \Delta t_j)) dW(s) \rightharpoonup \int_0^t g_u dW(s) \quad \text{in } L^2(\Omega; L^2(\mathbb{T}^d)) \text{ for all } t \in [0, T],$$

and also using that $u^{\Delta t_j}$ is bounded in $L^\infty(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$, we deduce that for a.e. $t > 0$, there exists a subsequence of $u^{\Delta t_j}(t)$ which we denote again by $u^{\Delta t_j}(t)$ which converges to $u(t)$ weakly in $L^2(\Omega; L^2(\mathbb{T}^d))$. We let $\Delta t_j \rightarrow 0$ in (53) to deduce that

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} u(t) v dx \right] - \mathbb{E} \left[\int_{\mathbb{T}^d} u_0^\epsilon v dx \right] + \epsilon \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} \nabla u \cdot \nabla v dx ds \right] - \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} f_u \mathbf{v} \cdot \nabla v dx ds \right] \\ &= \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^t g_u dW(s) v dx \right] \end{aligned} \tag{54}$$

for all $v \in L^2(\Omega; H^1(\mathbb{T}^d))$ and all $t \in [0, T]$, so that

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} \left(u(t) - \int_0^t g_u dW(s) \right) v dx \right] - \mathbb{E} \left[\int_{\mathbb{T}^d} u_0^\epsilon v dx \right] \\ &+ \epsilon \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} \nabla u \cdot \nabla v dx ds \right] - \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} f_u \mathbf{v} \cdot \nabla v dx ds \right] \\ &= 0 \end{aligned}$$

which is equivalent to

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} \left(u(t) - \int_0^t g_u dW(s) \right) v dx \right] - \mathbb{E} \left[\int_{\mathbb{T}^d} u_0^\epsilon v dx \right] \\ &+ \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} \left\langle -\epsilon \Delta u + \operatorname{div}(f_u \mathbf{v}), v \right\rangle_{(H^1)'}, H^1 dx ds \right] \\ &= 0. \end{aligned}$$

Thus for almost all $t \in (0, T)$, P-a.s., there holds,

$$u(t) = u_0^\epsilon(\cdot) + \epsilon \int_0^t \Delta u ds - \int_0^t \operatorname{div}(\mathbf{v} f_u) ds + \int_0^t g_u dW(s), \tag{55}$$

in $(H^1(\mathbb{T}^d))'$. In the following we will identify f_u and g_u .

Lemma 11. Suppose that the hypothesis (5) holds for some positive constant Λ_2 . Then $\partial_t [\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j}]$ is bounded in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$.

Proof. We set $v = u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)$ in (33) to obtain

$$\begin{aligned} & \|u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 \\ &+ \Delta t_j \epsilon \int_{\mathbb{T}^d} \nabla u_{n+1}^\epsilon \cdot \nabla [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] dx \\ &= \Delta t_j \int_{\mathbb{T}^d} f(u_{n+1}^\epsilon) \mathbf{v} \cdot \nabla [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] dx \\ &= -\Delta t_j \int_{\mathbb{T}^d} [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] \mathbf{v} f'(u_{n+1}^\epsilon) \cdot \nabla u_{n+1}^\epsilon dx \\ &\leq \frac{1}{2} \|u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 + \frac{1}{2} C(\mathbf{v}, C_f) (\Delta t_j)^2 \|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)}^2, \end{aligned}$$

so that we have

$$\begin{aligned} & \frac{1}{2} \|u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 \\ & + \Delta t_j \epsilon \int_{\mathbb{T}^d} \nabla u_{n+1}^\epsilon \cdot \nabla [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] dx \\ & \leq \frac{1}{2} C(\mathbf{v}, C_f) (\Delta t_j)^2 \|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2. \end{aligned} \quad (56)$$

Next we estimate the term $E \left[\int_{\mathbb{T}^d} \nabla u_{n+1}^\epsilon \cdot \nabla [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] dx \right]$ in (56), using the fact that $E[(W(t_{n+1}) - W(t_n))g(u_n^\epsilon)] = 0$, we obtain

$$\begin{aligned} & E \left[\int_{\mathbb{T}^d} \nabla u_{n+1}^\epsilon \cdot \nabla [u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)] dx \right] \\ & = \frac{1}{2} E \left[\|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 + \|\nabla(u_{n+1}^\epsilon - u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 - \|\nabla u_n^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right] \\ & \quad - E \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) \nabla(u_{n+1}^\epsilon - u_n^\epsilon) \cdot \nabla g(u_n^\epsilon) dx \right] \\ & \quad - E \left[\int_{\mathbb{T}^d} \nabla(W_{n+1} - W_n) \cdot \nabla(u_{n+1}^\epsilon - u_n^\epsilon) g(u_n^\epsilon) dx \right] \\ & \geq \frac{1}{2} E \left[\|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 + \frac{1}{2} \|\nabla(u_{n+1}^\epsilon - u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 - \|\nabla u_n^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right. \\ & \quad \left. - \Lambda_1 \Delta t_j \|\nabla g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 - \Lambda_2 \Delta t_j \|g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 \right], \end{aligned} \quad (57)$$

where we have used that, in view of (37),

$$\begin{aligned} & E \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n) \nabla(u_{n+1}^\epsilon - u_n^\epsilon) \cdot \nabla g(u_n^\epsilon) dx \right] \\ & \leq E \left[\int_{\mathbb{T}^d} (W_{n+1} - W_n)^2 |\nabla g(u_n^\epsilon)|^2 dx + \frac{1}{4} \int_{\mathbb{T}^d} |\nabla(u_{n+1}^\epsilon - u_n^\epsilon)|^2 dx \right] \\ & \leq \int_{\mathbb{T}^d} E[(W_{n+1} - W_n)^2] E[|\nabla g(u_n^\epsilon)|^2] dx + \frac{1}{4} E \left[\int_{\mathbb{T}^d} |\nabla(u_{n+1}^\epsilon - u_n^\epsilon)|^2 dx \right] \\ & \leq \Lambda_1 \Delta t_j \int_{\mathbb{T}^d} E[|\nabla g(u_n^\epsilon)|^2] dx + \frac{1}{4} E \left[\int_{\mathbb{T}^d} |\nabla(u_{n+1}^\epsilon - u_n^\epsilon)|^2 dx \right]. \end{aligned}$$

One can obtain an estimate for the term $E \left[\int_{\mathbb{T}^d} \nabla(W_{n+1} - W_n) \cdot \nabla(u_{n+1}^\epsilon - u_n^\epsilon) g(u_n^\epsilon) dx \right]$ in a similar fashion by using (5). This yields (57). Next we substitute (57) into (56) to deduce that,

$$\begin{aligned} & E \left[\|u_{n+1}^\epsilon - u_n^\epsilon - (W_{n+1} - W_n)g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 \right] + \Delta t_j \epsilon E \left[\|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 - \|\nabla u_n^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 + \frac{1}{2} \|\nabla(u_{n+1}^\epsilon - u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 \right] \\ & \leq \Lambda_1 (\Delta t_j)^2 \epsilon E \left[\|\nabla g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 \right] + \Lambda_2 (\Delta t_j)^2 \epsilon E \left[\|g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 \right] + C(\mathbf{v}, C_f) (\Delta t_j)^2 E \left[\|\nabla u_{n+1}^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right]. \end{aligned}$$

We sum the above inequality on the variable n from 0 to $N - 1$ and use Lemma 8 to deduce that

$$\begin{aligned} & \sum_{n=0}^{N-1} \Delta t_j \mathbb{E} \left[\left\| \frac{u_n^\epsilon - u_{n-1}^\epsilon - (W_{n+1} - W_n)g(u_{n-1}^\epsilon)}{\Delta t_j} \right\|_{L^2(\mathbb{T}^d)}^2 \right] + \epsilon \mathbb{E} \left[\|\nabla u_N^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right] + \frac{\epsilon}{2} \sum_{n=0}^{N-1} \mathbb{E} \left[\|\nabla(u_{n+1}^\epsilon - u_n^\epsilon)\|_{L^2(\mathbb{T}^d)^d}^2 \right] \\ & \leq C(\mathbf{v}, C_f, C_g, \epsilon, \Lambda_1) \Delta t_j \sum_{n=1}^N \mathbb{E} \left[\|\nabla u_n^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right] + \epsilon \mathbb{E} \left[\|\nabla u_0^\epsilon\|_{L^2(\mathbb{T}^d)^d}^2 \right] + \Lambda_2 \Delta t_j \epsilon \sum_{n=0}^N \mathbb{E} \left[\|g(u_n^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 \right] \\ & \leq C, \end{aligned} \tag{58}$$

where C depends on $\mathbf{v}, C_f, C_g, \Lambda_1, \Lambda_2, u_0$ and ϵ . We take the difference of (35) and (49) to deduce that

$$\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j} = \sum_{n=1}^N \left[\frac{u_n - u_{n-1} - (B_n - B_{n-1})}{\Delta t_j} (t - (n-1)\Delta t_j) + (u_{n-1} - B_{n-1}) \right] \mathbf{1}_{[(n-1)\Delta t_j, n\Delta t_j]},$$

which we differentiate with respect to t

$$\begin{aligned} \partial_t [\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j}] &= \sum_{n=1}^N \left[\frac{u_n^\epsilon - u_{n-1}^\epsilon - (B_n - B_{n-1})}{\Delta t_j} \right] \mathbf{1}_{[(n-1)\Delta t_j, n\Delta t_j]} \\ &= \sum_{n=1}^N \left[\frac{u_n^\epsilon - u_{n-1}^\epsilon - (W_n - W_{n-1})g(u_{n-1}^\epsilon)}{\Delta t_j} \right] \mathbf{1}_{[(n-1)\Delta t_j, n\Delta t_j]} \end{aligned} \tag{59}$$

where we have also used the definition of B_n in (50). The result of Lemma 11 then follows from (58). \square

Lemma 12. $\int_0^t g_u dW(s)$ is continuous in time and u defined as (55) is in $C([0, T]; (H^1(\mathbb{T}^d))')$.

Proof. We have proved that, for almost every $\omega \in \Omega$,

$$\begin{aligned} \tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j} &\in L^\infty(0, T; L^2(\mathbb{T}^d)); \\ \partial_t [\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j}] &\in L^2(0, T; L^2(\mathbb{T}^d)) \subset L^2(0, T; (H^1(\mathbb{T}^d))'). \end{aligned}$$

Using the fact that the embedding from $L^2(\mathbb{T}^d)$ into $(H^1(\mathbb{T}^d))'$ is compact, and applying the Theorem of Aubin-Lions-Simon [cf. ¹, Theorem II.5.16], we deduce that the sequence $\{\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j}\}$ is relatively compact in $C([0, T]; (H^1(\mathbb{T}^d))')$. Therefore

$\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j} \rightarrow u - \int_0^t g_u dW(s)$ strongly in $C([0, T]; (H^1(\mathbb{T}^d))')$ as $\Delta t_j \rightarrow 0$. If $\int_0^t g_u dW(s)$ is continuous in time, then $u(t)$ is continuous in time. We define

$$\begin{aligned} w(x, t) &= \int_0^t g_u dW(x, s) = \sum_{j=1}^{\infty} \sqrt{\lambda_j} e_j(x) \int_0^t g_u d\beta_j(s), \\ w_N(x, t) &= \sum_{j=1}^N \sqrt{\lambda_j} e_j(x) \int_0^t g_u d\beta_j(s) \end{aligned}$$

for all $N \in \mathbb{N}$. Thus for all $1 \leq N < M, M, N \in \mathbb{N}$, by Doob's inequality and Itô isometry,

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq t \leq T} (w_M(x, t) - w_N(x, t))^2 \right] &\leq 4 \mathbb{E} [(w_M(x, T) - w_N(x, T))^2] \\ &= 4 \sum_{j=N+1}^M \lambda_j e_j^2(x) \int_0^T \mathbb{E} [g_u^2] ds. \end{aligned} \tag{60}$$

Recalling the assumption that $\sum_{j=1}^{\infty} \lambda_j \|e_j\|_{L^\infty(\mathbb{T}^d)}^2 \leq \Lambda_1$ and using the fact that $g_u \in L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$, we deduce

$$\begin{aligned} \sum_{j=1}^{\infty} \lambda_j \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^T e_j^2 g_u^2 ds dx \right] &= \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^T \sum_{j=1}^{\infty} \lambda_j e_j^2 g_u^2 ds dx \right] \\ &\leq \Lambda_1 \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^T g_u^2 ds dx \right] \\ &< \infty, \end{aligned}$$

so that in view of (60)

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq t \leq T} \|w_M(t) - w_N(t)\|_{L^2(\mathbb{T}^d)}^2 \right] &\leq \int_{\mathbb{T}^d} dx \mathbb{E} \left[\sup_{0 \leq t \leq T} (w_M(t) - w_N(t))^2 \right] \\ &\leq 4 \sum_{j=N+1}^M \lambda_j \mathbb{E} \left[\int_{\mathbb{T}^d} \int_0^T e_j^2 g_u^2 ds dx \right] \\ &\rightarrow 0. \end{aligned}$$

as $N < M$ and $N \rightarrow \infty$. This implies with the help of Borel-Cantelli's lemma that $\{w_N(t)\}$ has a Cauchy subsequence $\{w_{N'}(t)\}$ in $C([0, T], L^2(\mathbb{T}^d))$, P-a.s.. Therefore since $\{w_{N'}\}$ converges to w in $C([0, T], L^2(\mathbb{T}^d))$, it follows that $w \in C([0, T], L^2(\mathbb{T}^d))$.

As P-a.s., $u - \int_0^t g_u dW(s)$ is in $C([0, T], H^1(\mathbb{T}^d)')$, we deduce that P-a.s., $u \in C([0, T], H^1(\mathbb{T}^d)')$. \square

Next we identify f_u to $f(u)$, g_u to $g(u)$ and u to u_ϵ .

Lemma 13. We have that $u^{\Delta t_j} \rightarrow u$ strongly in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ as $\Delta t_j \rightarrow 0$. In particular, this implies that that $f_u = f(u)$ and $g_u = g(u)$. And u is a strong solution of Problem \mathcal{P}^ϵ , we will denote it by u_ϵ .

Before proving the Lemma 13, we cite the Itô's formula which will be used in the proof of Lemma 13.

Lemma 14 (Itô's formula cf.^{10,11}). Let $X_0 \in L^2(\Omega, \mathcal{F}, P; L^2(\mathbb{T}^d))$ and $Y \in L^2([0, T] \times \Omega, (H^1(\mathbb{T}^d))')$, $Z \in L^2([0, T] \times \Omega, L^2(\mathbb{T}^d))$, both progressively measurable. Define the continuous $(H^1(\mathbb{T}^d))'$ -valued process X by:

$$X(t) = X(0) + \int_0^t Y(s) ds + \int_0^t Z(s) dW(s), \quad 0 \leq t \leq T,$$

and the following Itô's formula holds for the square of its L^2 -norm P-a.s.

$$\begin{aligned} \|X(t)\|_{L^2(\mathbb{T}^d)}^2 &= \|X(0)\|_{L^2(\mathbb{T}^d)}^2 + 2 \int_0^t \left\langle Y(s), X(s) \right\rangle_{(H^1(\mathbb{T}^d))', H^1(\mathbb{T}^d)} ds \\ &\quad + 2 \int_0^t \left\langle X(s), Z(s) dW(s) \right\rangle_{L^2(\mathbb{T}^d)} + \int_0^t \sum_{j=0}^{\infty} \left\langle Z^2 Q e_j, e_j \right\rangle_{L^2(\mathbb{T}^d)} ds. \end{aligned} \tag{61}$$

And we show a preliminary estimate.

Lemma 15. There holds

$$\begin{aligned} &\limsup_{\Delta t_j \rightarrow 0} \int_0^T e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] dt \\ &\leq T \|u_0\|^2 + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f_u \mathbf{v} \cdot \nabla u dx \right] ds dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g_u g(u) Q(x, x) dx \right] ds dt \end{aligned}$$

$$-\int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u) Q(x, x) dx \right] ds dt - c \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u\|^2] ds dt - 2\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] ds dt.$$

Proof. We start from (38), choosing $\bar{\epsilon} = 1$ and deduce from (36) the following counterpart of (39),

$$\begin{aligned} & \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (|u_{n+1}^\epsilon|^2 - |u_n^\epsilon|^2 + |u_{n+1}^\epsilon - u_n^\epsilon|^2) dx \right] + \Delta t_j \epsilon \mathbb{E} \left[\int_{\mathbb{T}^d} |\nabla u_{n+1}^\epsilon|^2 dx \right] \\ & \leq \frac{1}{2} \Delta t_j \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_n^\epsilon) Q(x, x) dx \right] + \frac{1}{2} \mathbb{E} \left[\int_{\mathbb{T}^d} (u_{n+1}^\epsilon - u_n^\epsilon)^2 dx \right], \end{aligned}$$

which we slightly simplify and multiply by $e^{-cn\Delta t_j}$. This implies that for all nonnegative c and integer n ,

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{T}^d} e^{-cn\Delta t_j} |u_{n+1}^\epsilon|^2 - e^{-c(n-1)\Delta t_j} |u_n^\epsilon|^2 dx \right] + 2\epsilon \Delta t_j e^{-cn\Delta t_j} \mathbb{E} \left[\int_{\mathbb{T}^d} |\nabla u_{n+1}^\epsilon|^2 dx \right] \\ & \leq \Delta t_j e^{-cn\Delta t_j} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_n^\epsilon) Q(x, x) dx \right] + [e^{-cn\Delta t_j} - e^{-c(n-1)\Delta t_j}] \mathbb{E} \left[\int_{\mathbb{T}^d} |u_n^\epsilon|^2 dx \right]. \end{aligned}$$

We sum this inequality for k from 0 to n and also use the inequality

$$e^{-ck\Delta t_j} - e^{-c(k-1)\Delta t_j} \leq -c\Delta t_j e^{-ck\Delta t_j}$$

for all fixed c and Δt_j , to obtain

$$\begin{aligned} & e^{-cn\Delta t_j} \mathbb{E} [\|u_{n+1}^\epsilon\|^2] + 2\epsilon \Delta t_j \sum_{k=0}^n e^{-ck\Delta t_j} \mathbb{E} [\|\nabla u_{k+1}^\epsilon\|^2] \\ & \leq \mathbb{E} [\|u_0\|^2] + \Delta t_j \sum_{k=0}^n e^{-ck\Delta t_j} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_k^\epsilon) Q(x, x) dx \right] - c \Delta t_j \sum_{k=0}^n e^{-ck\Delta t_j} \mathbb{E} [\|u_k^\epsilon\|^2] \end{aligned}$$

Thus, we use the definition of $u^{\Delta t_j}$ in (43) which converges weakly to its limit to deduce that

$$\begin{aligned} & e^{-cn\Delta t_j} \mathbb{E} [\|u_{n+1}^\epsilon\|^2] + 2\epsilon \int_0^{(n+1)\Delta t_j} e^{-cs} \mathbb{E} [\|\nabla u^{\Delta t_j}\|^2] ds \\ & \leq \mathbb{E} [\|u_0\|^2] + \Delta t_j \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_0) Q(x, x) dx \right] \\ & \quad + \int_0^{n\Delta t_j} e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u^{\Delta t_j}) Q(x, x) dx \right] ds - c \int_0^{n\Delta t_j} e^{-cs} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds. \end{aligned} \tag{62}$$

Next we remark that for $t \in [n\Delta t_j, (n+1)\Delta t_j]$, the expression below is nonnegative

$$\begin{aligned} 0 & \leq (e^{-cn\Delta t_j} - e^{-ct}) \mathbb{E} [\|u_{k+1}^\epsilon\|^2] + 2\epsilon \int_t^{(n+1)\Delta t_j} e^{-cs} \mathbb{E} [\|\nabla u^{\Delta t_j}\|^2] ds \\ & \quad + \int_{n\Delta t_j}^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u^{\Delta t_j}) Q(x, x) dx \right] ds + c \int_{(t-\Delta t_j)^+}^{n\Delta t_j} e^{-ct} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds. \end{aligned}$$

since each term is nonnegative; combining it with (62) yields

$$\begin{aligned}
& e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] + 2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla u^{\Delta t_j}\|^2] ds \\
& \leq \|u_0\|^2 + \Delta t_j \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u_0) Q(x, x) dx \right] + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u^{\Delta t_j}) Q(x, x) dx \right] ds \\
& \quad - c \int_0^{(t-\Delta t_j)^+} e^{-ct} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds.
\end{aligned}$$

Using the fact that $u^{\Delta t_j}$ is bounded in $L^\infty(0, T, L^2(\Omega; L^2(\mathbb{T}^d)))$, we obtain

$$\begin{aligned}
& e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] + 2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla u^{\Delta t_j}\|^2] ds \\
& \leq \|u_0\|^2 + \Delta t_j \Lambda_1 \|g(u_0)\|^2 + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u^{\Delta t_j}) Q(x, x) dx \right] ds \\
& \quad + c \left(\int_{(t-\Delta t_j)^+}^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds - \int_0^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds \right) \\
& \leq \|u_0\|^2 + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u^{\Delta t_j}) Q(x, x) dx \right] ds - c \int_0^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j}\|^2] ds + C\Delta t_j.
\end{aligned}$$

We recall that for all $v \in H^1(\mathbb{T}^d)$, there holds $\int_{\mathbb{T}^d} f(v) \mathbf{v} \cdot \nabla v dx = 0$, and also use that $a^2 = (a-b)^2 - b^2 + 2ab$ to obtain

$$\begin{aligned}
& e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] + 2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds \\
& \quad + 4\epsilon \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} \nabla u^{\Delta t_j} \nabla u dx ds \right] - 2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] \\
& \quad - 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} [\mathbf{v} f(u^{\Delta t_j}) - \mathbf{v} f(u)] \nabla [u^{\Delta t_j} - u] dx ds \right] \\
& \leq \|u_0\|^2 + C\Delta t_j + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f(u^{\Delta t_j}) \mathbf{v} \cdot \nabla u dx \right] ds \\
& \quad + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f(u) \mathbf{v} \cdot \nabla u^{\Delta t_j} dx \right] ds + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g(u^{\Delta t_j}) g(u) Q(x, x) dx \right] ds \\
& \quad - \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u) Q(x, x) dx \right] ds + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} (g(u^{\Delta t_j}) - g(u))^2 Q(x, x) dx \right] ds \\
& \quad - c \int_0^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j} - u\|^2] ds - 2c \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} u^{\Delta t_j} u dx \right] ds + c \int_0^t e^{-cs} \mathbb{E} [\|u\|^2] ds. \tag{63}
\end{aligned}$$

Note that there exists $c_{\min} = C(\mathbf{v}, f, g, \epsilon, \Lambda_1) > 0$ such that, for $c \in (c_{\min}, +\infty)$ and Δt_j small, it holds that

$$\begin{aligned}
& -2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds + 2 \int_0^t e^{-cs} \mathbb{E} [(\mathbf{v}f(u^{\Delta t_j}) - \mathbf{v}f(u)) \nabla(u^{\Delta t_j} - u)] ds \\
& + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} (g(u^{\Delta t_j}) - g(u))^2 Q(x, x) dx \right] ds - c \int_0^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j} - u\|^2] ds \\
\leq & (-2\epsilon + \epsilon) \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds + \frac{1}{\epsilon} \int_0^t e^{-cs} \mathbb{E} [\|\mathbf{v}f(u^{\Delta t_j}) - \mathbf{v}f(u)\|^2] ds \\
& + \Lambda_1 \int_0^t e^{-cs} \mathbb{E} [\|g(u^{\Delta t_j}) - g(u)\|^2] ds - c \int_0^t e^{-cs} \mathbb{E} [\|u^{\Delta t_j} - u\|^2] ds \\
\leq & -\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds.
\end{aligned}$$

Next we substitute the upper inequality in (63) and integrate from 0 to T to obtain,

$$\begin{aligned}
& \int_0^T e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] dt + \epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds dt \\
\leq & T \|u_0\|^2 + CT \Delta t_j \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f(u^{\Delta t_j}) \mathbf{v} \cdot \nabla u \right] ds dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f(u) \mathbf{v} \cdot \nabla u^{\Delta t_j} dx \right] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g(u^{\Delta t_j}) g(u) Q(x, x) dx \right] ds dt - \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u) Q(x, x) dx \right] ds dt \\
& - 2c \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} u^{\Delta t_j} u dx \right] ds dt + c \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u\|^2] ds dt \\
& - 4\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} \nabla u^{\Delta t_j} \nabla u dx \right] ds dt + 2\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] ds dt.
\end{aligned}$$

We pass to the limit as $\Delta t_j \rightarrow 0$, and also use the facts that $\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla(u^{\Delta t_j} - u)\|^2] ds dt \geq 0$

and that $\int_{\mathbb{T}^d} f(v) \mathbf{v} \cdot \nabla v dx = 0$ to obtain

$$\begin{aligned}
& \limsup_{\Delta t_j \rightarrow 0} \int_0^T e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] dt \\
\leq & T \|u_0\|^2 + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f_u \mathbf{v} \cdot \nabla u dx \right] ds dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g_u g(u) Q(x, x) dx \right] ds dt \\
& - \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g^2(u) Q(x, x) dx \right] ds dt - c \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u\|^2] ds dt - 2\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] ds dt,
\end{aligned} \tag{64}$$

which completes the proof of the Lemma 15.

Next, we use this result to prove Lemma 13. We apply the Itô's formula, Lemma 14 [cf.¹¹, Theorem 4.2.5] and also using the equality (4.2.28) in¹¹, to deduce that

$$\begin{aligned} & e^{-ct} \mathbb{E} [\|u(t)\|^2] + 2\epsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] ds - 2 \int_0^t \mathbb{E} \left[\int_{\mathbb{T}^d} e^{-cs} \mathbf{v} f_u \cdot \nabla u \right] dx ds \\ &= \mathbb{E} [\|u_0\|^2] - c \int_0^t e^{-cs} \mathbb{E} [\|u(s)\|^2] ds + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g_u^2 Q(x, x) dx \right] ds. \end{aligned} \quad (65)$$

We integrate (65) on time interval $[0, T]$ which yields

$$\begin{aligned} & T \|u_0\|^2 + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} f_u \mathbf{v} \cdot \nabla u dx \right] ds dt \\ & - 2\epsilon \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|\nabla u\|^2] ds dt - c \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u\|^2] ds dt \\ &= \int_0^T e^{-ct} \mathbb{E} [\|u(t)\|^2] dt - \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} g_u^2 Q(x, x) dx \right] ds dt. \end{aligned} \quad (66)$$

We substitute (66) into (64) to obtain

$$\limsup_{\Delta t_j \rightarrow 0} \int_0^T e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] dt + \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{T}^d} (g_u - g(u))^2 Q(x, x) dx \right] ds dt \leq \int_0^T e^{-ct} \mathbb{E} [\|u(t)\|^2] dt,$$

which yields the inequality

$$\limsup_{\Delta t_j \rightarrow 0} \int_0^T e^{-ct} \mathbb{E} [\|u^{\Delta t_j}(t)\|^2] dt \leq \int_0^T e^{-ct} \mathbb{E} [\|u(t)\|^2] dt. \quad (67)$$

In what follows, we prove that $u^{\Delta t_j}$ converges to u strongly in $L^2(0, T; L^2(\Omega \times \mathbb{T}^d))$. We have that

$$\begin{aligned} & \|e^{-ct/2}(u^{\Delta t_j} - u)\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))}^2 \\ &= \mathbb{E} \left[\int_0^T e^{-ct} \int_{\mathbb{T}^d} (u^{\Delta t_j} - u)(u^{\Delta t_j} - u) dx dt \right] \\ &= \mathbb{E} \left[\int_0^T \int_{\mathbb{T}^d} (e^{-ct/2} u^{\Delta t_j})^2 dx dt \right] - 2 \mathbb{E} \left[\int_0^T e^{-ct} \int_{\mathbb{T}^d} u u^{\Delta t_j} dx dt \right] + \mathbb{E} \left[\int_0^T \int_{\mathbb{T}^d} e^{-ct} u^2 dx dt \right]. \end{aligned}$$

Passing to the limit and using (67), we deduce that

$$\lim_{\Delta t_j \rightarrow 0} \|e^{-ct/2}(u^{\Delta t_j} - u)\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))}^2 \leq 0,$$

so that also

$$\lim_{\Delta t_j \rightarrow 0} \|u^{\Delta t_j} - u\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))}^2 \leq e^{cT} \lim_{\Delta t_j \rightarrow 0} \|e^{-ct/2}(u^{\Delta t_j} - u)\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))}^2 = 0.$$

Finally we prove that $f_u = f(u)$ and $g_u = g(u)$. Since f is Lipschitz continuous we have

$$\|f(u) - f(u^{\Delta t_j})\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))} \leq C_f \|u - u^{\Delta t_j}\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))}.$$

Using the fact that $\|u - u^{\Delta t_j}\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))} \rightarrow 0$ we deduce $\|f(u) - f(u^{\Delta t_j})\|_{L^2(0, T; L^2(\Omega \times \mathbb{T}^d))} \rightarrow 0$. Hence, $f(u^{\Delta t_j}) \rightarrow f(u)$ strongly in $L^2(0, T; L^2(\Omega \times \mathbb{T}^d))$ as $\Delta t_j \rightarrow 0$, which in turn implies that $f_u = f(u)$. A similar proof implies that $g_u = g(u)$.

Thus for almost all $t \in (0, T)$, P-a.s., there holds,

$$u(\cdot, t) = u_0^\epsilon + \epsilon \int_0^t \Delta u ds - \int_0^t \operatorname{div}(\mathbf{v} f(u)) ds + \int_0^t g(u) dW(s), \quad (68)$$

in $(H^1(\mathbb{T}^d))'$. Next we prove that this equality also holds in $L^2(\mathbb{T}^d)$. Since

$$\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j} \rightharpoonup u(\cdot, t) - \int_0^t g(u) dW(s)$$

is in $L^2(0, T; L^2(\mathbb{T}^d))$, we have that $u(\cdot, t) - \int_0^t g(u) dW(s) \in L^2(0, T; L^2(\mathbb{T}^d))$ P-a.s.. We also remark that $\int_0^t \operatorname{div}(\mathbf{v} f(u)) ds \in L^2(\mathbb{T}^d)$ for all t ; therefore using that for almost all $t \in (0, T)$, P-a.s.,

$$\epsilon \int_0^t \Delta u ds = u(\cdot, t) - \int_0^t g(u) dW(s) + \int_0^t \operatorname{div}(\mathbf{v} f(u)) ds - u_0^\epsilon,$$

we deduce that $\epsilon \int_0^t \Delta u ds \in L^2(0, T; L^2(\mathbb{T}^d))$.

This completes the proof that for almost all $t \in (0, T)$, P-a.s., there holds,

$$u(\cdot, t) = u_0^\epsilon + \epsilon \int_0^t \Delta u ds - \int_0^t \operatorname{div}(\mathbf{v} f(u)) ds + \int_0^t g(u) dW(s), \quad (69)$$

in $L^2(\mathbb{T}^d)$, therefore u is a strong solution in the sense of Definition 4, which can be denoted by u_ϵ . \square

Lemma 16. $\epsilon \Delta u_\epsilon - \operatorname{div}(\mathbf{v} f(u_\epsilon))$ belongs to $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$.

Proof. According to Lemma 11 and the facts that $\tilde{u}^{\Delta t_j} \rightarrow u_\epsilon$ weakly in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$ and $\tilde{B}^{\Delta t_j} \rightarrow \int_0^t g(u_\epsilon) dW(s)$ in $L^2(\Omega; L^2(\mathbb{T}^d))$ for all t , we obtain that $\partial_t [\tilde{u}^{\Delta t_j} - \tilde{B}^{\Delta t_j}]$ converges up to a subsequence to $\partial_t [u_\epsilon - \int_0^t g(u_\epsilon) dW(s)]$ in $L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$. As

$$\partial_t \left[u_\epsilon - \int_0^t g(u_\epsilon) dW(s) \right] = \epsilon \Delta u_\epsilon - \operatorname{div}(\mathbf{v} f(u_\epsilon)),$$

we obtain that $\epsilon \Delta u_\epsilon - \operatorname{div}(\mathbf{v} f(u_\epsilon)) \in L^2(0, T; L^2(\Omega; L^2(\mathbb{T}^d)))$. This result is used in the proof of Lemma 5 \square

The uniqueness of the strong solution

Finally, we prove the uniqueness of the strong solution of Problem (\mathcal{P}^ϵ) . Let u_1^ϵ and u_2^ϵ be two strong solutions of Problem (\mathcal{P}^ϵ) , and take the difference of the equations for u_1^ϵ and u_2^ϵ . This yields for almost all $t \in (0, T)$,

$$\begin{aligned} u_1^\epsilon(t) - u_2^\epsilon(t) &= \int_0^t \epsilon \Delta (u_1^\epsilon - u_2^\epsilon) ds - \int_0^t (\operatorname{div}(\mathbf{v} f(u_1^\epsilon)) - \operatorname{div}(\mathbf{v} f(u_2^\epsilon))) ds \\ &\quad + \int_0^t (g(u_1^\epsilon) - g(u_2^\epsilon)) dW(x, s). \end{aligned}$$

We apply Itô's formula, Lemma 1 to the function $\mathcal{G}(X, t) = \|X\|_{L^2(\mathbb{T}^d)}^2$. We define

$$X(t) = u_1^\epsilon(t) - u_2^\epsilon(t), \quad \psi = \epsilon \Delta (u_1^\epsilon - u_2^\epsilon) - (\operatorname{div}(\mathbf{v} f(u_1^\epsilon)) - \operatorname{div}(\mathbf{v} f(u_2^\epsilon))), \quad \theta = g(u_1^\epsilon) - g(u_2^\epsilon)$$

to obtain

$$\begin{aligned} \mathcal{G}(X(t), t) &= \|X(t)\|_{L^2(\mathbb{T}^d)}^2 \\ \mathcal{G}_X(X(t), t) &= 2X(t) \\ \mathcal{G}_{XX}(X(t), t) &= 2I \end{aligned}$$

$$\mathcal{G}_t(X(t), t) = 0$$

and then

$$\begin{aligned} & \|u_1^\epsilon(t) - u_2^\epsilon(t)\|_{L^2(\mathbb{T}^d)}^2 \\ &= 2\epsilon \int_0^t \langle \Delta(u_1^\epsilon - u_2^\epsilon), u_1^\epsilon - u_2^\epsilon \rangle_{L^2(\mathbb{T}^d)} ds - 2 \int_0^t \langle \operatorname{div}(\mathbf{v}f(u_1^\epsilon)) - \operatorname{div}(\mathbf{v}f(u_2^\epsilon)), u_1^\epsilon - u_2^\epsilon \rangle_{L^2(\mathbb{T}^d)} ds \\ &\quad + 2 \int_0^t \langle g(u_1^\epsilon) - g(u_2^\epsilon), u_1^\epsilon - u_2^\epsilon \rangle_{L^2(\mathbb{T}^d)} dW(\cdot, s) + \int_0^t \sum_{j=0}^{\infty} \langle [g(u_1^\epsilon) - g(u_2^\epsilon)]^2 \lambda_j e_j, e_j \rangle_{L^2(\mathbb{T}^d)} ds. \end{aligned}$$

Next we take the expectation to obtain the following estimate

$$\begin{aligned} & \mathbb{E} \left[\|u_1^\epsilon(t) - u_2^\epsilon(t)\|_{L^2(\mathbb{T}^d)}^2 \right] \\ &\leq -2\epsilon \mathbb{E} \left[\int_0^t \|\nabla(u_1^\epsilon - u_2^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 ds \right] - 2 \mathbb{E} \left[\int_0^t \langle \operatorname{div}(\mathbf{v}f(u_1^\epsilon)) - \operatorname{div}(\mathbf{v}f(u_2^\epsilon)), u_1^\epsilon - u_2^\epsilon \rangle ds \right] \\ &\quad + 2 \mathbb{E} \left[\int_0^t \langle g(u_1^\epsilon) - g(u_2^\epsilon), u_1^\epsilon - u_2^\epsilon \rangle dW(x, s) \right] + \sum_{j=0}^{\infty} \lambda_j \|e_j\|_{L^\infty(\mathbb{T}^d)}^2 \mathbb{E} \left[\int_0^t \|g(u_1^\epsilon) - g(u_2^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 ds \right], \end{aligned}$$

so that

$$\begin{aligned} & \mathbb{E} \left[\|u_1^\epsilon(t) - u_2^\epsilon(t)\|_{L^2(\mathbb{T}^d)}^2 \right] + 2\epsilon \mathbb{E} \left[\int_0^t \|\nabla(u_1^\epsilon - u_2^\epsilon)\|_{L^2(\mathbb{T}^d)}^2 ds \right] \\ &\leq 2V \mathbb{E} \left[\int_0^t \int_{\mathbb{T}^d} |\nabla(u_1^\epsilon - u_2^\epsilon)| |f(u_1^\epsilon) - f(u_2^\epsilon)| dx dt \right] + \Lambda_1 C_g \mathbb{E} \left[\int_0^t \|u_1^\epsilon - u_2^\epsilon\|_{L^2(\mathbb{T}^d)}^2 dt \right], \end{aligned}$$

which combined with Gronwall's lemma yields the uniqueness result. \square

ACKNOWLEDGMENTS

The authors would like to thank Dr. Perla El Kettani for helpful discussions and the GDRI ReaDiNet as well as the research funding of MathAM-OIL, AIST c/o AIMR Tohoku University for financial support.

References

1. F. Boyer and P. Fabrie, *Mathematical Tools for the Study of the Incompressible Navier-Stokes Equations and Related Models*, Applied Mathematical Sciences 183. New York: Springer, 2013.
2. C. Bauzet, G. Vallet and P. Wittbold, The Cauchy problem for a conservation law with a multiplicative stochastic perturbation, *Journal of Hyperbolic Differential Equations* 2012; **9**(4): 661–709.
3. C. Bauzet, G. Vallet and P. Wittbold, The Dirichlet problem for a conservation law with a multiplicative stochastic perturbation, *J. Funct. Anal.* 2014; **4**(266): 2503–2545.
4. G. Da Prato and J. Zabczyk, *Stochastic equations in infinite dimensions*, Cambridge University Press, Second edition 2014.
5. A. Debussche and J. Vovelle, Scalar conservation laws with stochastic forcing, *it J. Funct. Anal.* 2014, **259**(4): 1014–1042.
6. I. Ekeland and R. Temam, *Convex Analysis and Variational Problems*, SIAM, Philadelphia, 1999.
7. T. Funaki, Y. Gao and D. Hilhorst, Convergence of a finite volume scheme for a stochastic conservation law involving a Q -Brownian motion, *DCDS-B* 2018; **23**(4): 1461–1504.

8. M. Giaquinta and G. Modica, *Mathematical Analysis: An Introduction to Functions of Several Variables*, Springer, 2010.
9. I. Karatzas and S. E. Shreve, *Brownian Motion and Stochastic Calculus*, Springer, 1991.
10. N. V. Krylov and B. L. Rozovskii, Stochastic evolution equations, *J. Soviet Math.* 1981; **16**: 1233-1277.
11. C. Prévot and M. Röckner, *A Concise Course on Stochastic Partial Differential Equations*, Springer, Verlag Berlin Heidelberg, 2007.