



**HAL**  
open science

## Radial singular solutions for the N-Laplace equation with exponential nonlinearities

Marius Ghergu, Jacques Giacomoni, S. Prashanth

► **To cite this version:**

Marius Ghergu, Jacques Giacomoni, S. Prashanth. Radial singular solutions for the N-Laplace equation with exponential nonlinearities. *Australian Journal of Mathematical Analysis and Applications*, 2019, 475 (1), pp.668-685. 10.1016/j.jmaa.2019.02.062 . hal-02160333

**HAL Id: hal-02160333**

**<https://hal.science/hal-02160333>**

Submitted on 22 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Radial singular solutions for the $N$ - Laplace equation with exponential nonlinearities

Marius GHERGU

School of Mathematics and Statistics,  
University College Dublin,  
Belfield, Dublin 4, Ireland  
[marius.ghergu@ucd.ie](mailto:marius.ghergu@ucd.ie)

and

Institute of Mathematics Simion Stoilow of the Romanian Academy,  
21 Calea Grivitei Street,  
010702 Bucharest, Romania

Jacques GIACOMONI

LMAP (UMR E2S-UPPA CNRS 5142)  
Université de Pau et des Pays de l'Adour  
Avenue de l'Université, F-64013 Pau cedex, France  
[jacques.giacomoni@univ-pau.fr](mailto:jacques.giacomoni@univ-pau.fr)

S. PRASHANTH

TIFR-Centre for Applicable Mathematics,  
Post Bag No. 6503, Sharadar Nagar,  
GKVK Post Office,  
Bangalore 560065, India  
[pras@math.tifrbng.res.in](mailto:pras@math.tifrbng.res.in)

February 19, 2019

## Abstract

In this paper, we consider radial distributional solutions of the quasilinear equation  $-\Delta_N u = f(u)$  in the punctured open ball  $B_R \setminus \{0\} \subset \mathbb{R}^N$ ,  $N \geq 2$ . We obtain sharp conditions on the nonlinearity  $f$  for extending such solutions to the whole domain  $B_R$  by preserving the regularity. For a certain class of nonlinearity  $f$  we obtain the existence of singular solutions and deduce upper and lower estimates on the growth rate near the singularity.

**running head:** Radial Singular solutions for  $N$ -Laplace equation

**Keywords:**  $N$ -Laplace equation;  
removable singularity;  
singular solutions;  
behaviour near the singularity

**2000 Mathematics Subject Classification:** Primary 35J62, 35B09;  
Secondary 35B65

## 1 Introduction

Let  $\Omega$  be a domain in  $\mathbb{R}^N$ ,  $N \geq 2$  and  $-\Delta_N$  denote the  $N$ -Laplace operator.

**Definition 1.1** Given  $f \in L^1_{loc}(\Omega)$ , we say that a function  $u \in L^1_{loc}(\Omega)$  is a *distributional solution of the equation  $-\Delta_N u = f$  in  $\Omega$*  if

$$|\nabla u| \in L^{N-1}_{loc}(\Omega) \text{ and } \int_{\Omega} |\nabla u|^{N-2} \nabla u \cdot \nabla \phi = \int_{\Omega} f \phi, \quad \forall \phi \in C_0^\infty(\Omega).$$

We will write “ $-\Delta_N u = f$  in  $\mathcal{D}'(\Omega)$ ” as a short-hand way of saying that  $u$  is a *distributional solution in  $\Omega$* .

Let  $B_R \subset \mathbb{R}^N$  be the open ball of radius  $R$  centered at the origin in  $\mathbb{R}^N$ . Consider the problem

$$(P_R^*) \quad \begin{cases} -\Delta_N u = f(u) & \text{in } \mathcal{D}'(B_R \setminus \{0\}), \\ u \geq 0 & \text{in } B_R \setminus \{0\}, \\ u \in L^\infty_{loc}(\overline{B}_R \setminus \{0\}), & u \text{ is radial.} \end{cases}$$

Throughout the work, we assume that the function  $f : [0, \infty) \rightarrow [0, \infty)$  satisfies:

- (f<sub>1</sub>)  $f \in C^{0,\theta}([0, \infty))$  for some exponent  $\theta \in (0, 1)$ ;
- (f<sub>2</sub>)  $\liminf_{t \rightarrow \infty} f(t) t^{1-N} > 0$ ;
- (f<sub>3</sub>) the map  $t \mapsto f(t) + \kappa t^{N-1}$  is non-decreasing for some  $\kappa > 0$ .

By a result of BREZIS and LIONS [6] (in the case  $N = 2$ ) and VÉRON [23, theorem 5.10, p. 283] (in the case  $N > 2$ ), we have the following

**Theorem 1.2** *If  $u$  solves  $(P_R^*)$ , then  $u \in W^{1,p}(B_R)$  for any  $1 \leq p < N$ . Furthermore, there exists some  $\alpha \geq 0$  such that  $u$  solves the following problem :*

$$(P_{\alpha,R}) \quad \begin{cases} -\Delta_N u = f(u) + \alpha \delta_0 & \text{in } \mathcal{D}'(B_R), \\ f(u) \in L^\infty_{loc}(\overline{B}_R \setminus \{0\}) \cap L^1(B_R). \end{cases}$$

The above result leads naturally to the following questions:

(Q1) *Is there a sharp growth condition on  $f$  that determines whether a solution of  $(P_R^*)$  can or cannot be extended to be a (distributional) solution to  $(P_{0,R})$  ?*

(Q2) *If such an extension holds and the solution in the punctured domain  $B_R \setminus \{0\}$  is smooth, is the extended solution equally smooth in  $B_R$ ?*

(Q3) *If the extended solution blows up at the origin, what is its asymptotic blow-up rate?*

When  $N = 2$  and  $f$  has at most a polynomial growth, the first two questions (Q1) and (Q2) were discussed in detail in BREZIS-LIONS [6] and LIONS [18]. For a corresponding discussion involving exponential growth nonlinearities in the case  $N = 2$ , we refer to DHANYA-GIACOMONI-PRASHANTH [9].

In the quasilinear case  $N \geq 3$ , without being exhaustive, we mention the results of GUEDDA-VÉRON [13], BIDAULT-VÉRON [4] and KICHENASSAMY-VÉRON [15]. For more on the subject, we refer to the survey VÉRON [24] and the book VÉRON [23]. We cite the book GHERGUTALIAFERRO [10] for results concerning isolated singularities for partial differential inequalities.

In the present paper, we extend the results contained in [9] to the radial quasilinear case and obtain a complete answer to questions (Q1) and (Q2) above. We also provide a partial answer to (Q3).

**Definition 1.3** *We call  $f$  a sub-exponential function if there exists  $\beta > 0$  such that*

$$(1.1) \quad C := \sup_{[0, \infty)} f(t)e^{-\beta t} < \infty.$$

*We call  $f$  to be super-exponential if it is not a sub-exponential function.*

As a complete answer to question (Q1), we show that if  $f$  is sub-exponential then we can construct, by the method of monotone iterations, a solution  $U$  to  $(P_R^*)$  which solves  $(P_{\alpha,R})$  for some  $\alpha > 0$  (see Theorem 3.1). Conversely, we show in Lemma 4.1 that any solution to  $(P_R^*)$  extends to a solution to  $(P_{0,R})$  if  $f$  is super-exponential. In contrast to this result we show that the Dirac mass in  $(P_{\alpha,R})$  is not in general removable for sub-exponential functions  $f$ .

Regarding question (Q2) above, for a super-exponential  $f$ , we construct examples of radial solutions to  $(P_{0,R})$  which blows up only at the origin (see Lemma 5.2 and Theorem 5.3). That is, such solutions are smooth in  $B_R \setminus \{0\}$ , but not in  $B_R$ . We also show that if  $f$  is sub-exponential, then any solution  $u$  of  $(P_R^*)$  that extends to a solution of  $(P_{0,R})$  is regular, say in  $C_{loc}^{1,\theta}$  (see Theorem 5.1).

The question (Q3) is partially answered in Lemma 6.1. Utilising the asymptotic analysis of ATKINSON and PELETIER (see [3]), for the super-exponential nonlinearities  $f$  we derive an upper bound for singular solutions (see Lemma 6.1) and consequently obtain the following limiting behaviour for any solution  $u$  of  $(P_R^*)$  :

$$|x|^N f(u(x)) \rightarrow 0 \quad \text{as} \quad |x| \rightarrow 0.$$

We remark that this result is new even for the semilinear case. Although we do not obtain a pointwise lower bound, we derive in Lemma 7.1 an integral bound for the behavior of the solution around the isolated singularity at the origin; see also Corollaries 7.5 and 7.6. The accurate asymptotic behaviour of singular solutions is still an open question as oscillating singular solutions to  $(P_R^*)$  may exist (see Corollary 6.2).

Nevertheless, Corollaries 6.2 and 7.7 give alternatives similar to the ones available in higher dimensions for supercritical nonlinearities (see for instance Theorem 5.13 in [23]).

The semilinear case  $N = 2$  with exponential-type nonlinearities in dimensions higher than 2 is dealt with in the recent work of KIKUCHI and WEI [16]. By using the Emden-Fowler transformation and a clever analysis of the perturbation term in the asymptotic profile, the authors prove the existence and the precise asymptotic behaviour of singular solutions. In the two dimensional case, this approach as well as the one using Harnack type inequalities as in [4], fails since the coefficients of the resulting equation do not **possess** the integrability in the right spaces.

## 2 Some preliminary results

In this section we collect some useful results to our approach. The following first observation is in order:

**Proposition 2.1** *If  $u$  solves  $(P_R^*)$ , then  $u$  is radially non-increasing in  $B_R \setminus \{0\}$ .*

*Proof.* In the radial variable,  $u$  solves the ODE:

$$-r^{1-N} \left( r^{N-1} |u'|^{N-2} u' \right)' = f(u) \quad \text{in } (0, R).$$

Noting that  $f(u) \geq 0$ , it readily follows that the map  $r \mapsto ru'(r)$  is non-increasing in  $(0, R)$ . Let

$$(2.1) \quad \ell := \lim_{r \rightarrow 0^+} ru'(r).$$

If  $\ell > 0$ , by integrating (2.1) we obtain that  $u(r) \rightarrow -\infty$  as  $r \rightarrow 0^+$ , a contradiction to the nonnegativity of  $u$  near 0. Hence  $\ell \leq 0$  and the result follows.  $\blacksquare$

First, we note the following simple consequence of definition 1.3.

**Proposition 2.2** *If  $f$  is a super-exponential nonlinearity, then*

$$\int_0^\infty f(t) e^{-\epsilon t} dt = \infty \quad \text{for any } \epsilon > 0.$$

*Proof.* From (1.1) we have that for any  $\epsilon > 0$  and any positive integer  $n$

$$\sup_{[n, \infty)} f(t) e^{-\epsilon t} = \infty.$$

Therefore, we may find  $t_n \geq n$  such that

$$f(t_n) \geq n e^{\epsilon t_n}.$$

Thus,

$$\int_{t_n}^{\infty} (f(t) + \kappa t^{N-1})e^{-\epsilon t} dt \geq \int_{t_n}^{\infty} (f(t_n) + \kappa t_n^{N-1})e^{-\epsilon t} dt \geq \frac{n}{\epsilon}.$$

The assertion now follows. ■

Our next result establishes the connection between distributional and entropy solutions to  $(P_R^*)$ . We recall that for  $g \in L^1(\Omega)$ , a function  $u \in L^1_{loc}(\Omega)$  is an entropy solution of

$$(2.2) \quad -\Delta_N u = g \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

if  $T_k(u) \in W_0^{1,N}(\Omega)$  for any  $k \in \mathbb{R}^+$ ,  $|\nabla u| \in L^{N-1}_{loc}(\Omega)$  and

$$(2.3) \quad \int_{\Omega} |\nabla u|^{N-2} \nabla u \cdot \nabla T_k(u - \phi) \leq \int_{\Omega} T_k(u - \phi) g \quad \text{for all } \phi \in C_0^{\infty}(\Omega), k \in \mathbb{R}^+.$$

Here, for  $k \in \mathbb{R}^+$ ,  $T_k$  denotes the truncation map given by  $T_k(t) = t$  for  $t \in [-k, k]$  and  $T_k(t) = k \operatorname{sgn}(t)$  for  $|t| \geq k$ .

**Proposition 2.3** *Let  $u \in C^1_{loc}(\overline{B}_R \setminus \{0\}) \cap L^1_{loc}(B_R)$ ,  $f \in L^1(B_R)$  be both radial functions and  $|\nabla u|^{N-1} \in L^p(B_R)$  for some  $p > 1$ . If  $u$  is a distributional solution to  $-\Delta_N u = f$  in  $B_R$ , then  $u - u(R)$  is an entropy solution to the same problem in  $B_R$  with homogeneous Dirichlet data on  $\partial B_R$ .*

*Proof.* Since  $u$  is a distributional solution we have  $u \in L^1_{loc}(B_R)$  has "zero trace" on  $\partial\Omega$ ,  $|\nabla u| \in L^{N-1}_{loc}(B_R)$  and

$$(2.4) \quad \int_{B_R} |\nabla u|^{N-2} \nabla u \cdot \nabla \phi = \int_{B_R} \phi f \quad \text{for all } \phi \in C_0^{\infty}(B_R).$$

We may fix  $p$  close to 1 so that  $W_0^{1,p'}(B_R) \hookrightarrow C_0(\overline{B}_R)$ . We first claim that if  $|\nabla u|^{N-1} \in L^p(B_R)$  for some  $p > 1$  and  $f \in L^1(B_R)$ , we can enlarge the class of test functions  $\phi$  in (2.4) to  $W_0^{1,p'}(B_R)$ . To see this, we take a sequence  $\{\phi_n\} \subset C_0^{\infty}(B_R)$  converging to  $\phi$  in the  $W_0^{1,p'}(B_R)$  norm. Then  $\phi_n \rightarrow \phi$  in  $C_0(\overline{B}_R)$  and the claim follows by writing (2.4) for each  $\phi_n$  and passing to the limit  $n \rightarrow \infty$ .

Let  $v := u - u(R)$ . Given  $\phi \in C_0^{\infty}(B_R)$ , we have that  $T_k(v - \phi) \in W_0^{1,\infty}(B_R)$  for every  $k \in \mathbb{R}^+$ . Hence by the above observations,

$$\begin{aligned} \int_{B_R} |\nabla v|^{N-2} \nabla v \cdot \nabla T_k(v - \phi) &= \int_{B_R} |\nabla u|^{N-2} \nabla u \cdot \nabla T_k(v - \phi) \\ &= \int_{B_R} f T_k(v - \phi) \quad \forall \phi \in C_0^{\infty}(\Omega), \forall k \in \mathbb{R}^+. \end{aligned}$$

Thus,  $v$  satisfies (2.3). ■

Finally, we recall a version of BREZIS-MERLE [7] result for the  $N$ -Laplace problem.

**Proposition 2.4** (see [1, Theorem 1.6]) *Let  $u$  be the entropy solution of the problem (2.2) where  $g \in L^1(\Omega)$ . Then the following inequality holds for any  $\delta \in (0, N\omega_N^{\frac{1}{N-1}})$ :*

$$(2.5) \quad \int_{\Omega} \exp\left(\frac{(N\omega_N^{\frac{1}{N-1}} - \delta)|u|}{\|g\|_{1,\Omega}^{\frac{1}{N-1}}}\right) \leq \frac{N\omega_N^{\frac{1}{N-1}}}{\delta} |\Omega|.$$

Here,  $\|\cdot\|_{1,\Omega}$  denotes the  $L^1(\Omega)$  norm and  $|\Omega|$  the  $N$ -dimensional Lebesgue measure of  $\Omega$  and  $\omega_N$  is the volume of the unit sphere in  $\mathbb{R}^N$ .

**Corollary 2.5** (see [1, Corollary 1.7]) *Let  $u, g$  be as above. Then,  $e^{|u|} \in L^p(\Omega)$  for any  $p \geq 1$ .*

### 3 Dirac mass solution for sub-exponential $f$

In this section, we show the following:

**Theorem 3.1** *Let  $f$  be a sub-exponential function and  $\beta, C$  be given by (1.1). Then there exist  $\alpha, R_* > 0$  depending on  $\beta, C$  such that  $(P_{\alpha, R_*})$  admits a distribution solution.*

Before proving the theorem, we construct appropriate sub and super solutions in the lemmas below.

**Lemma 3.2** *Let  $f$  be a sub-exponential function with  $\beta, C$  given by (1.1). Define*

$$(3.1) \quad v_{\beta, C} = \frac{1-N}{\beta} \left[ \log r + \frac{N}{N-1} \log(1 + C\beta^{N-1}r) \right], \quad r > 0.$$

*Then, the following pointwise inequality holds :*

$$(3.2) \quad -\Delta_N v_{\beta, C} \geq f(v_{\beta, C}) \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

*Proof.* Let

$$\mu := \frac{N}{N-1}, \quad M := C\beta^{N-1}.$$

By a straightforward calculation,

$$(3.3) \quad \begin{aligned} -\Delta_N v_{\beta, C}(r) &= r^{1-N} (-r^{N-1} |v'_{\beta}(r)|^{N-2} v'_{\beta})' \\ &= \left( \frac{\mu M (N-1)^N}{\beta^{N-1}} \right) \left[ \frac{1}{r^{N-1} (1+Mr)^2} \right] \left[ \frac{1 + (1+\mu)Mr}{1+Mr} \right]^{N-2}. \end{aligned}$$

We recall the growth condition on  $f$  in definition 1.3 and use (3.1) to find that

$$(3.4) \quad f(v_{\beta, C}) \leq C e^{\beta v_{\beta, C}} = \frac{C}{r^{N-1} (1+Mr)^N}.$$

Therefore, from (3.3) and (3.4) we obtain that

$$(3.5) \quad -\Delta_N v_{\beta, C} \geq f(v_{\beta, C}) \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

■

**Remark 3.3** Note that  $v_{\beta,C}$  is a strictly decreasing function on  $(0, \infty)$  with

$$\lim_{r \rightarrow 0^+} v_{\beta,C} = \infty \quad \text{and} \quad \lim_{r \rightarrow \infty} v_{\beta,C} = -\infty.$$

**Definition 3.4** Given  $\beta, C > 0$  define  $R_* := R_*(\beta, C)$  to be the unique zero of  $v_{\beta,C}$ .

**Lemma 3.5** Let  $f, \beta, C$  be as in the previous lemma. For  $R > 0$  define

$$(3.6) \quad w_{\beta,C,R}(r) := \frac{1-N}{\beta} \left[ \log r + \frac{N}{N-1} \log(1 + C\beta^{N-1}R) \right], \quad r > 0.$$

Then

$$(3.7) \quad w_{\beta,C,R} \leq v_{\beta,C} \quad \text{in } B_R,$$

$$(3.8) \quad w_{\beta,C,R_*} \geq 0 \quad \text{in } B_{R_*},$$

$$(3.9) \quad w_{\beta,C,R}(R) = v_{\beta,C}(R),$$

$$(3.10) \quad w_{\beta,C,R}, v_{\beta,C} \in C_{loc}^\infty(\mathbb{R}^N \setminus \{0\}),$$

$$(3.11) \quad -\Delta_N w_{\beta,C,R} = \omega_N^{\frac{1}{N-1}} \left( \frac{N-1}{\beta} \right) \delta_0 \quad \text{in } \mathcal{D}'(\mathbb{R}^N).$$

The proof of Lemma 3.5 follows by straightforward calculations.

Now we are ready to proceed with the proof of Theorem 3.1. Let  $\beta, C > 0$  be given by (1.1) and let  $v_{\beta,C}, w_{\beta,C,R}$  be the corresponding functions as in (3.1) and (3.6). Choose  $R_*$  as in definition 3.4. Given  $0 < \epsilon < R_*$ , define the annular region

$$A_{\epsilon,R_*} := B_{R_*} \setminus B_\epsilon.$$

We now set up the following iteration procedure (solved in the  $W^{1,N}$ -weak sense):

$$(I_n) \quad \begin{cases} u_0 := w_{\beta,C,R_*}; \\ -\Delta_N u_n + \kappa |u_n|^{N-2} u_n = f(u_{n-1}) + \kappa |u_{n-1}|^{N-2} u_{n-1} \quad \text{in } A_{\epsilon,R_*}, \\ u_n = w_{\beta,C,R_*} \quad \text{on } \partial A_{\epsilon,R_*}; \\ u_n \in C^{1,\theta}(\overline{A_{\epsilon,R_*}}) \text{ for any } \theta \in (0, 1) \text{ and } u_n \text{ radial.} \end{cases}$$

It is standard to see that the quasilinear equation in  $(I_n)$  is solvable in the  $W^{1,N}$ -weak sense and by Hölder regularity results (see for instance TOLKSDORF [22] and LIEBERMAN [17]) the required regularity of the solution is obtained.

Using the hypothesis  $(f_3)$ , (3.2), (3.8) and an induction argument, we obtain that

$$u_0 := w_{\beta,C,R_*} \leq u_n \leq u_{n+1} \leq v_{\beta,C} \quad \text{in } A_{\epsilon,R_*} \quad \text{for all } n.$$

Since  $v_{\beta,C}$  is bounded in  $A_{\epsilon,R_*}$  for a fixed  $\epsilon > 0$ , from the above pointwise estimates we can pass to the uniform  $C^{1,\theta}(\overline{A_{\epsilon,R_*}})$  estimates for the sequence  $\{u_n\}$  by using the classical quasilinear regularity results (see for instance TOLKSDORF [22] and LIEBERMAN [17]). Therefore, we obtain a function  $u_{\epsilon,R_*}$  such that

$$u_n \rightarrow u_{\epsilon,R_*} \quad \text{in } C^1(\overline{A_{\epsilon,R_*}})$$



and  $u_{\epsilon, R_*}$  solves (in the  $W^{1, N}$ -weak sense) the following problem:

$$\begin{cases} -\Delta_N u_{\epsilon, R_*} = f(u_{\epsilon, R_*}), w_{\beta, C, R_*} \leq u_{\epsilon, R_*} \leq v_{\beta, C} \text{ in } A_{\epsilon, R_*}, \\ u_{\epsilon, R_*} = w_{\beta, C, R_*} \text{ on } \partial A_{\epsilon, R_*}, \\ u_{\epsilon, R_*} \in C^{1, \theta}(\bar{A}_{\epsilon, R_*}) \text{ and is radial.} \end{cases}$$

Choosing a positive sequence  $\{\epsilon_n\}$  tending to 0 and noting that the corresponding sequence of solutions  $\{u_{\epsilon_n, R_*}\}$  is relatively compact in  $C_{loc}^1(\bar{B}_{R_*} \setminus \{0\})$ , we obtain a distributional solution  $U$  of

$$(P_{R_*}^*) \quad \begin{cases} -\Delta_N U = f(U), \quad w_{\beta, C, R_*} \leq U \leq v_{\beta, C} \text{ in } B_{R_*} \setminus \{0\}, \\ U = w_{\beta, C, R_*} \text{ on } |x| = R_*, \\ U \in C^{1, \theta}(\bar{B}_{R_*} \setminus \{0\}) \text{ and is radial.} \end{cases}$$

Note that

$$U(r) \sim \left( \frac{1-N}{\beta} \right) \log r \text{ as } r \rightarrow 0^+, \text{ and } U, f(U) \in L_{loc}^\infty(\mathbb{R}^N \setminus \{0\}).$$

By Theorem 1.2 we obtain that

$$f(U) \in L_{loc}^1(B_{R_*}), \quad |\nabla U| \in L^p(B_{R_*}) \text{ for all } 1 \leq p < N,$$

and for some  $\alpha \geq 0$ ,

$$-\Delta_N U = f(U) + \alpha \delta_0 \text{ in } \mathcal{D}'(B_{R_*}).$$

If  $\alpha = 0$ , from Propositions 2.3-2.4, Corollary 2.5 and Corollary 2.2 in [1] we obtain that  $U$  is bounded near the origin, contradiction. Hence, necessarily  $\alpha > 0$ .  $\blacksquare$

## 4 Removable Singularity

In this section we show that if  $f$  is super-exponential then distributional solutions of  $(P_R^*)$  can be extended to (distributional) solutions of  $(P_{0, R})$ .

**Lemma 4.1** *Let  $f$  be super-exponential and let  $u$  be a solution of  $(P_R^*)$ , and hence of  $(P_{\alpha, R})$  for some  $\alpha \geq 0$  (see Theorem 1.2). Then, necessarily  $\alpha = 0$ .*

*Proof.* Let  $0 < \eta < R$  be small. Choose a nonnegative radial test function  $\phi \in C_0^\infty(B_\eta)$  with  $\phi(0) = 1$  and  $\max_{[0, \eta]} |\phi'| \leq 1/\eta$ . We then obtain (since  $u' \leq 0$  from proposition 2.1)

$$(4.1) \quad - \int_0^\eta r^{N-1} |u'|^{N-1} \phi'(r) dr = \int_0^\eta r^{N-1} f(u(r)) \phi(r) dr + \alpha \phi(0).$$

We estimate

$$(4.2) \quad \begin{aligned} \text{L.H.S. of (4.1)} &= O\left(\max_{[0, \eta]} r^{N-1} |u'|^{N-1} \int_0^\eta |\phi'| \right) \\ &= O\left(\max_{[0, \eta]} r^{N-1} |u'|^{N-1}\right). \end{aligned}$$

Since

$$\int_0^\eta r^{N-1} f(u(r)) \phi(r) dr \rightarrow 0 \text{ as } \eta \rightarrow 0^+,$$

from (4.1) and (4.2) we will obtain  $\alpha = 0$  if we can show

$$(4.3) \quad \max_{[0, \eta]} r^{N-1} |u'|^{N-1} \rightarrow 0 \text{ as } \eta \rightarrow 0^+.$$

To this aim, we utilise the Emden-Fowler transformation given by

$$(4.4) \quad t := N \log(N/r), \quad 0 < r < R; \quad y(t) := u(r), \quad N \log(N/R) < t < \infty.$$

Consequently,

$$y'(t) = -\frac{r}{N} u'(r).$$

It can then be easily checked that if  $u$  is a radial solution of  $-\Delta_N u = f(u)$  in  $B_R \setminus \{0\}$ , then  $y$  solves the following Emden-Fowler type ODE:

$$(4.5) \quad -\left(|y'|^{N-2} y'\right)' = e^{-t} f(y) \text{ in } (N \log(N/R), \infty).$$

Note that since  $u$  is non-increasing (by Proposition 2.1), one has  $y' \geq 0$ . Therefore, (4.3) holds iff  $y'(t) \rightarrow 0$  as  $t \rightarrow \infty$ . We note from (4.5) that

$$-\left(|y'|^{N-2} y'\right)' = e^{-t} f(y) \geq 0 \text{ in } (N \log(N/R), \infty).$$

This immediately implies that  $y'$  is a decreasing function on  $(N \log(N/R), \infty)$ . Let us denote

$$\ell := \lim_{t \rightarrow \infty} y'(t).$$

Noting that  $y' \geq \ell$  in  $(N \log(N/R), \infty)$ , we obtain

$$y(t) \geq y(N \log(N/R)) + \ell(t - N \log(N/R)) \quad \text{for all } t \geq N \log(N/R).$$

Hence, from the above inequality and the assumptions  $(f_2) - (f_3)$  we find

$$\begin{aligned} \infty > \int_{B_{R/2}} (f(u) + \kappa u^{N-1}) dx &= \int_{N \log(2N/R)}^\infty \left( f(y(t)) + \kappa y(t)^{N-1} \right) e^{-t} dt \\ &\geq \int_{N \log(2N/R)}^\infty f(y(N \log(N/R)) + \ell(t - N \log(N/R))) e^{-t} dt, \end{aligned}$$

which contradicts Proposition 2.2 if  $\ell > 0$ . Therefore,  $\ell = 0$  and hence  $\alpha = 0$ . ■

## 5 Existence of singular solutions

In this section we answer question (Q2). We first show that when  $f$  is sub-exponential, then any distributional solution of  $(P_{0,R})$  is regular.

**Theorem 5.1** *Let  $f$  be a sub-exponential nonlinearity. Then any solution to  $(P_{0,R})$  is regular in  $B_R$ .*

*Proof.* Let  $u$  be a solution to  $(P_{0,R})$ . Then from propositions 2.3- 2.4 and corollary 2.5, we have that  $e^u$  (and hence  $f(u)$ ) belongs to  $L^p(B_1)$  for all  $p \geq 1$ . Thus, applying corollary 2.2 in [1] (see also related results in IOKU [14] and BOCCARDO-PERAL-VAZQUEZ [5]) we obtain that  $u \in W_0^{1,N}(B_1) \cap L^\infty(\Omega)$ . Hence from Hölder regularity results in [17], we obtain that  $u \in C^{1,\theta}(\bar{\Omega}) \cap C_0(\bar{\Omega})$  for all  $0 < \theta < 1$ .  $\blacksquare$

In the following two results, we construct solutions to  $(P_{0,R})$  which blow up only at the origin for some classes of super-exponential nonlinearities.

**Lemma 5.2** *Given any  $\mu > 1$ , there exists a super-exponential nonlinearity  $f$  satisfying :*

$$\lim_{t \rightarrow \infty} f(t)e^{-t^\mu} = 0, \quad \lim_{t \rightarrow \infty} f(t)e^{-t^{\mu-\epsilon}} = \infty \quad \forall \epsilon > 0$$

*such that the corresponding problem  $(P_{0,\frac{1}{2}})$  admits a radial solution that blows up only at the origin.*

*Proof.* Given  $\mu > 1$ , define

$$f(t) := \left(\frac{N}{\mu}\right)^N (\mu-1)(N-1) \begin{cases} (N \log 2)^{\frac{(1-\mu)(N-1)-\mu}{\mu}} e^{(N \log 2)}, & 0 \leq t \leq (N \log 2)^{\frac{1}{\mu}} \\ t^{(1-\mu)(N-1)-\mu} e^{t^\mu}, & t > (N \log 2)^{\frac{1}{\mu}}. \end{cases}$$

It can be easily checked that

$$u(x) \stackrel{\text{def}}{=} \left( \log \frac{1}{|x|^N} \right)^{\frac{1}{\mu}}, \quad x \neq 0$$

solves  $(P_{\frac{1}{2}}^*)$ . Appealing to Theorem 1.2 and Lemma 4.1 we obtain that  $u$  solves  $(P_{0,\frac{1}{2}})$ . It can be also directly checked that  $f(u) \in L^1(B_{\frac{1}{2}})$ .  $\blacksquare$

In the next result, we exhibit a class of super-exponential nonlinearities whose growth rate at infinity is of critical type (in the sense of Trudinger-Moser). Although we state the result for nonlinearities in a general form, one can check that the model class of nonlinearities

$$f_0(t) := t^{-\alpha}(1+t)^m e^{t^{\frac{N}{N-1}-t^\beta}}, \quad \alpha > 0, m \geq 0, \frac{1}{N-1} < \beta < \frac{N}{N-1}$$

satisfy the required assumptions in (i).

**Theorem 5.3** (i) *Let  $f_0 : (0, \infty) \rightarrow (0, \infty)$  be a smooth (at least  $C^3$ ) nonlinearity which has the form  $f_0(t) = h(t)e^{t^{\frac{N}{N-1}}}$  where  $h(t)$  satisfies assumptions (A1)-(A5) in [11]. Then there exists  $R > 0$  such that  $-\Delta_N u = f_0(u)$  admits a radial positive distributional solution  $z^*$  in  $B_R$  blowing up only at the origin and such that  $z^* \notin W_{loc}^{1,N}(B_R)$ .*

(ii) *Let  $f(t) := f_0(\frac{R}{2})\chi_{[0,\frac{R}{2}]}(t) + f_0(t)\chi_{[\frac{R}{2},\infty)}(t)$ . Then  $(P_{0,\frac{R}{2}})$  admits a radial solution that blows up only at the origin.*

*Proof.* (i) We use the transformation (4.4) in order to cast the differential equation  $-\Delta_N u = f(u)$  in  $B_R$  (for a positive solution) into the following equivalent form :

$$(5.1) \quad \left. \begin{aligned} -(|y'|^{N-2}y')' &= e^{-t}f_0(y), \\ y \geq 0, \quad y &\in L_{loc}^\infty \end{aligned} \right\} \text{ in } (T, \infty), \quad T \stackrel{\text{def}}{=} N \log(N/R);$$

$$\int_T^\infty f_0(y(t))e^{-t}dt < \infty.$$

For our purposes, it is more convenient to consider the following “shooting from infinity” problem depending upon a parameter  $\gamma > 0$  :

$$(S_\gamma) \quad \begin{cases} -(|y'|^{N-2}y')' = e^{-t}f_0(y), \\ y(\infty) = \gamma, y'(\infty) = 0. \end{cases}$$

Let  $y(\cdot, \gamma)$  denote the unique solution of  $(S_\gamma)$ . We see that  $y(\cdot, \gamma)$  is a strictly concave function as long as it is nonnegative. Therefore, **as any concave function cannot be bounded below at  $-\infty$** , there exists a first zero of the solution  $y(\cdot, \gamma)$  denoted by  $T(\gamma)$ . In addition, the map  $\gamma \rightarrow T(\gamma)$  is continuous (see [11, Lemma 3.1]). We divide the proof into three steps:

*Step 1 : Given a sequence  $\gamma_n \rightarrow \infty$  as  $n \rightarrow \infty$ , there exist a sequence  $\{r_n\}$  of positive numbers with  $\liminf_{n \rightarrow \infty} r_n > 0$  and a sequence of nonnegative radial solutions  $\{u_n\} \subset L^\infty(B_{r_n})$  of  $-\Delta_N u_n = f_0(u_n)$  in  $\mathcal{D}'(B_{r_n})$  with  $u_n(0) = \gamma_n$ .*

Given such  $\{\gamma_n\}$ , let  $y_n = y(\cdot, \gamma_n)$  denote the solution of  $(S_{\gamma_n})$ . By [11, proposition 4.2, p. 12], we obtain that  $T^* = \limsup_{n \rightarrow \infty} T(\gamma_n) < \infty$ . Up to a subsequence of  $\{\gamma_n\}$ , we can assume that  $T^* = \lim_{n \rightarrow \infty} T(\gamma_n)$ . We fix this subsequence of  $\{y_n\}$ , which we still call  $\{y_n\}$ . Furthermore, from the asymptotic behaviour of  $f_0$  at  $\infty$  it is not difficult to show that  $T^* > -\infty$  (see also [12, lemma 4.1]).

Let us use the change of variable in (4.4) as follows:

$$r_n = Ne^{-T(\gamma_n)/N}, \quad u_n(x) = y_n(N \log(N/|x|), \quad x \in B_{r_n} \setminus \{0\}.$$

Then, we see that  $u_n$  solves the required equation in  $B_{r_n}$  with  $u_n(0) = \gamma_n$  and  $\liminf_{n \rightarrow \infty} r_n > 0$ .

*Step 2 : Let  $\{\gamma_n\}, \{y_n\}$  be as in Step 1. Extend  $y_n$  to  $[T^* - 1, T(\gamma_n))$  by 0. Then the extended sequence (still denoted as  $\{y_n\}$ ) is uniformly bounded on compact subsets of  $[T^* - 1, \infty)$ .*

Denote  $g(t) = \log(f_0(t))$ . We define the following energy functional associated to  $(S_{\gamma_n})$ :

$$E_n(t) = (y'_n)^{N-1}(t) - \frac{N-1}{N}(y'_n(t))^N g'(y_n(t)) - e^{g(y_n(t))-t}, \quad t > T(\gamma_n).$$

**An explicit calculation gives**

$$\begin{aligned} E'_n(t) &= (N-1)(y'_n)^{N-2}(t)y''_n(t) - (N-1)(y'_n(t))^{N-1}y''_n(t)g'(y_n(t)) \\ &\quad - \left(\frac{N-1}{N}\right)(y'_n(t))^{N+1}g''(y_n(t)) - e^{g(y_n(t))-t}(g'(y_n(t))y'_n(t) - 1). \end{aligned}$$

Note that the ODE satisfied by  $y_n$  can be written as  $(N-1)(y'_n)^{N-2}(t)y''_n(t) = -e^{g(y_n(t))-t}$ . Therefore, the above expression for  $E'_n$  simplifies to

$$E'_n(t) = -\left(\frac{N-1}{N}\right)(y'_n(t))^{N+1}g''(y_n(t)).$$

Given  $s_0 > 0$ , let  $t_0(\gamma_n) > T(\gamma_n)$  be defined by the relation  $y_n(t_0(\gamma_n)) = s_0 > 0$ . We may choose  $s_0$  large enough such that  $g', g'' > 0$  in  $[s_0, \infty)$ . **Indeed, by convexity of  $s \rightarrow g(s)$  for large  $s$  and the last statement of assumption (A1) in [11], there exists  $s_0 > 0$  large enough such that  $g'$  is increasing on  $(s_0, \infty)$ ,  $\lim_{t \rightarrow \infty} g'(t) = \infty$  and  $g'' > 0$  in  $(s_0, \infty)$ .** Therefore, since  $y_n$  is strictly increasing and  $g$  is convex in  $[t_0(\gamma_n), \infty)$ , we obtain  $E'_n(t) \leq 0$  for all  $t \geq t_0(\gamma_n)$ .

Since  $\lim_{t \rightarrow \infty} E_n(t) = 0$ , we obtain that  $E_n$  is a nonnegative function on  $[t_0(\gamma_n), \infty)$ . This immediately implies that

$$(5.2) \quad y'_n(t)g'(y_n(t)) \leq \frac{N}{N-1}, \quad \forall t \geq t_0(\gamma_n).$$

Now, integrating the ODE in  $(S_{\gamma_n})$ , we have

$$(y'_n(t_0))^{N-1} = \int_{t_0(\gamma_n)}^{\infty} f_0(y_n(t))e^{-t} dt.$$

Therefore, from (5.2) and recalling that  $y_n(t_0(\gamma_n)) = s_0$ , we obtain

$$(5.3) \quad \sup_n \int_{t_0(\gamma_n)}^{\infty} f_0(y_n(t))e^{-t} dt < \infty.$$

Let  $[a, b] \subset [T^* - 1, \infty)$ . Define  $A = \{n : t_0(\gamma_n) > b\}$ ,  $B = \{n : t_0(\gamma_n) \leq b\}$ . We note that  $\sup_n y_n(b) < \infty$ . Otherwise, there must be a subsequence of  $\{y_n(b)\}$  that tends to  $\infty$  and **hence** **by the nondecreasing nature of  $t \rightarrow y_n(t)$** , this subsequence converges uniformly to  $\infty$  in  $[b, b+1]$  violating (5.3).

Again by the monotonicity of  $y_n$ , we have  $\sup_{[a,b]} y_n \leq y_n(b)$ . Therefore,

$$\sup_{n \in A} \sup_{t \in [a,b]} y_n(t) \leq \sup_{n \in A} y_n(b) \leq \sup_n y_n(t_0(\gamma_n)) = s_0.$$

Similarly,

$$\sup_{n \in B} \sup_{[a,b]} y_n(t) \leq \sup_{n \in B} y_n(b) < \infty,$$

which completes the proof in Step 2.

*Step 3: Constructing the singular solution.*

From Step 2 and the fact that  $y_n$  solves the ODE in (5.1) we obtain a subsequence of  $\{y_n\}$ , which we denote again by  $\{y_n\}$ , such that  $\{y_n\}$  and  $\{y'_n\}$  are uniformly convergent in any compact sub-interval of  $(T^*, \infty)$ . By using a diagonalisation process, we can obtain a subsequence of  $\{y_n\}$ , which we will denote by  $\{y_n\}$  again, and a positive, continuous nondecreasing function  $y^*$  on  $(T^*, \infty)$  such that  $y_n \rightarrow y^*$  locally uniformly in  $(T^*, \infty)$  and  $\{y'_n\}$  also converges locally uniformly in  $(T^*, \infty)$ . Furthermore,  $y'_n \rightarrow (y^*)'$  pointwise in  $(T^*, \infty)$  (see RUDIN [20, Theorem 7.17]).

For an integer  $m \geq 0$  and any  $n$  such that  $\gamma_n > m + s_0$ , we define  $t_m(\gamma_n)$  to be the point at which  $y_n(t_m(\gamma_n)) = m + s_0$ . We claim that  $S_m \stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} t_m(\gamma_n) < \infty$ . To see this, define  $z_n(t) = y_n(t) - m - s_0$  for  $\gamma_n > m + s_0$ . Then,  $z_n$  solves the equation

$$-(|z'_n|^{N-2} z'_n)' = e^{-t} f_0(z_n + m + s_0) \stackrel{\text{def}}{=} e^{-t} \bar{f}_0(z_n).$$

Let  $T_m(\gamma_n)$  be the first zero of  $z_n(t)$  as  $t$  decreases from infinity. It can be checked that  $\bar{f}_0$  also satisfies assumptions (A1)-(A5) in [11]. Therefore, again by [11, Proposition 4.2], we get  $S_m < \infty$ .

Fix  $m$ . If necessary, by restricting to a further subsequence of  $\{y_n\}$  (depending on  $m$ ) so that  $\lim_{n \rightarrow \infty} t_m(\gamma_n) = S_m$ , we obtain

$$y^*(S_m) = \lim_{n \rightarrow \infty} y_n(S_m) = \lim_{n \rightarrow \infty} y_n(t_m(\gamma_n)) = m + s_0.$$

Since  $y^*$  is nondecreasing, we obtain that  $y^*(t) \rightarrow \infty$  as  $t \rightarrow \infty$ . Integrating the O.D.E. satisfied by  $y_n$  we find

$$(y'_n)^{N-1}(s) - (y'_n)^{N-1}(t) = \int_s^t f_0(y_n(\rho))e^{-\rho} d\rho, \quad T^* < s < t < \infty.$$

Using the convergence of  $y_n, y'_n$ , we can pass to the limit as  $n \rightarrow \infty$  on either side of the above equation to obtain that  $y^*$  also satisfies the same integral equation. That is,  $y^*$  solves the equation

$$-(((y^*)')^{N-1})' = e^{-t} f_0(y^*) \quad \text{in } (T^*, \infty).$$

Now, from (5.3),

$$(5.4) \quad \sup_n \int_{T(\gamma_n)}^{\infty} f_0(y_n) e^{-t} dt < \infty.$$

We now come to the value  $y^*$  at  $T^*$ . Since  $y^*$  is nondecreasing,  $y^*$  has a right limit at  $T^*$ . Integrating the ODE satisfied by  $y_n$  between  $t \in [T(\gamma_n), t_0(\gamma_n)]$  and  $t_0(\gamma_n)$  and using (5.2) and (5.4), we deduce that  $\{y'_n\}$  is uniformly bounded in  $[T(\gamma_n), T^* + 1]$ . Consequently, the extended sequence  $\{y_n\}$  is uniformly bounded in the Lipschitz norm on  $[T^* - 1, T^*]$ . Then, by Ascoli-Arzelà theorem, we have

$$y^*(T^*) = \lim_{n \rightarrow \infty} y_n(T^*) = 0.$$

From (5.4) and Fatou's lemma, we obtain that  $\int_{T^*}^{\infty} f_0(y^*) e^{-t} dt < \infty$ . Thus, to summarize,  $y^*$  solves the problem (5.1) with  $T = T^*$  with the additional property:  $y(T^*) = 0$ .

Let  $R = Ne^{-T^*/N}$ . We now define  $z^*(x) = y^*(N \log(N/|x|))$  for  $x \in B_R \setminus \{0\}$ . It follows that  $z^*$  solves the following problem:

$$\left\{ \begin{array}{l} -\Delta_N z^* = f_0(z^*) \\ z^* > 0 \end{array} \right\} \text{ in } B_R \setminus \{0\}, \\ z^* = 0 \text{ on } \partial B_R, \quad \lim_{|x| \rightarrow 0} z^*(x) = \infty, \quad f_0(z^*) \in L^1(B_R).$$

(ii) We note that  $f$  satisfies assumptions  $(f_1) - (f_3)$ . From Theorem 1.2,  $z^*$  satisfies the equation  $-\Delta_N z^* = f(z^*) + \alpha \delta_0$  in the sense of distributions in  $B_{\frac{R}{2}}$  for some  $\alpha \geq 0$ . Since  $f$  is super-exponential, by Lemma 4.1 we must have  $\alpha = 0$ . Thus,  $z^*$  is the required singular solution for  $(P_{0, \frac{R}{2}})$ .

If  $z^* \in W_{loc}^{1,N}(B_R)$ , by Trudinger-Moser imbedding [19], we obtain that  $f(z^*) \in L_{loc}^p(B_R)$  for all  $p \geq 1$  and hence  $z^*$  is locally bounded in  $B_R$ , a contradiction.  $\blacksquare$

## 6 Asymptotic behaviour in the super-exponential case

From the assumption  $(f_3)$ , we can fix  $\kappa \geq 0$  so that the map

$$(6.1) \quad t \mapsto F(t) := f(t) + \kappa t^{N-1} \text{ is strictly increasing for } t \geq 0.$$

For instance, if  $f(t) = e^{t^\mu}$ ,  $\mu > 0$ , then we may choose  $\kappa = 0$  and check that  $F^{-1}(t) = (\log t)^{\frac{1}{\mu}}$ . We have the following asymptotic estimate for  $u$ .

**Lemma 6.1** *Let  $f$  be a super-exponential nonlinearity and  $\kappa$  be chosen as in (6.1). Assume that*

$$(6.2) \quad (f(t))^\lambda \leq cf(\lambda t) \text{ for some } c > 0 \text{ and all } \lambda > 1, t \geq 0.$$

Then, any unbounded solution  $u$  of  $(P_R^*)$  satisfies the following properties:

(a) Given  $\epsilon > 0$ , there exists  $R_\epsilon \in (0, R)$  such that  $u(x) \leq F^{-1}(\epsilon|x|^{-N})$  for all  $x \in B_{R_\epsilon}$ .

(b) For any  $\epsilon > 0$ ,

$$\limsup_{x \rightarrow 0} \frac{u(x)}{F^{-1}(\epsilon|x|^{-N})} = 1.$$

*Proof.* We write the problem  $(P_R^*)$  in radial co-ordinates as follows (see Proposition 2.1):

$$\begin{cases} -r^{1-N} \left( \left( r^{N-1} |u'|^{N-2} u' \right)' \right) = f(u) & \text{in } \mathcal{D}'((0, R)), \\ u \text{ nonnegative} & \text{in } (0, R). \end{cases}$$

We use again the Emden-Fowler transformations as in (4.4)-(4.5) and denote by  $y$  the transformed solution corresponding to  $u$ .

Since  $f$  is super-exponential, as in the proof of Lemma 4.1, one can show that

$$(6.3) \quad \lim_{t \rightarrow \infty} y'(t) = 0.$$

Since  $y$  is a concave function, it follows that  $y'(t) > 0$  for all  $t$ . Integrating the ODE satisfied by  $y$  (see (4.5)) between the limits  $1 \ll t < t_1$ , we obtain

$$\begin{aligned} (y')^{N-1}(t) - (y')^{N-1}(t_1) &= \int_t^{t_1} (f(y(s)) + \kappa s^{N-1}) e^{-s} ds - \int_t^{t_1} \kappa s^{N-1} e^{-s} ds \\ &\geq (f(y(t)) + \kappa t^{N-1}) (e^{-t} - e^{-t_1}) - \int_t^{t_1} \kappa s^{N-1} e^{-s} ds. \end{aligned}$$

Letting  $t_1 \rightarrow \infty$  in the above inequality and using (6.3), we deduce

$$(y')^{N-1}(t) \geq (f(y(t)) + \kappa t^{N-1}) e^{-t} - \int_t^\infty \kappa s^{N-1} e^{-s} ds$$

so that

$$(f(y(t)) + \kappa t^{N-1}) e^{-t} \leq (y')^{N-1}(t) + \int_t^\infty \kappa s^{N-1} e^{-s} ds.$$

Thus, by (6.3) we find

$$\lim_{t \rightarrow \infty} F(y(t))e^{-t} = \lim_{t \rightarrow \infty} (f(y(t)) + \kappa y^{N-1}(t))e^{-t} = 0.$$

Now, (a) follows easily. Using part (a), we have that

$$(6.4) \quad \limsup_{x \rightarrow 0} \frac{u(x)}{F^{-1}(\epsilon|x|^{-N})} \leq 1.$$

Suppose for some  $\epsilon > 0$  the above inequality is strict. Then we can find  $0 < \eta < 1$  and  $R_0 > 0$  such that

$$u(x) \leq \eta F^{-1}(\epsilon|x|^{-N}) \quad \text{in } B_{R_0} \setminus \{0\}.$$

Therefore for some  $A > 0$ ,

$$\begin{aligned} f(u(x)) &\leq F(u(x)) \leq F(\eta F^{-1}(\epsilon|x|^{-N})) \quad (\text{by monotonicity of } F) \\ &= f(\eta F^{-1}(\epsilon|x|^{-N})) + \kappa (\eta F^{-1}(\epsilon|x|^{-N}))^{N-1} \\ &\leq c^\eta (f(F^{-1}(\epsilon|x|^{-N})))^\eta + \kappa (\eta F^{-1}(\epsilon|x|^{-N}))^{N-1} \quad (\text{by (6.2) with } \lambda = 1/\eta) \\ &\leq A (f(F^{-1}(\epsilon|x|^{-N})))^\eta \quad (\text{if } R_0 \text{ is small and noting } t^{N-1} \ll (f(t))^\eta \text{ for all large } t) \\ &\leq A (f(F^{-1}(\epsilon|x|^{-N})) + \kappa (F^{-1}(\epsilon|x|^{-N}))^{N-1})^\eta. \end{aligned}$$

Thus we obtain the pointwise estimate

$$f(u) \leq O(1)|x|^{-\eta N} \quad \text{in } B_{R_0} \setminus \{0\}.$$

Thus,  $f(u) \in L_{loc}^p(B_R)$  for some  $p > 1$ . From classical estimates in Serrin [21] we deduce  $u \in L_{loc}^\infty(B_R)$ , a contradiction. This shows that for any  $\epsilon > 0$  we must have equality in (6.4) which completes our proof.  $\blacksquare$

**Corollary 6.2** *Let  $f$  and  $u$  be as in Lemma 6.1. Then, one of the following alternatives holds:*

$$(i) \quad \lim_{x \rightarrow 0} \frac{u(x)}{F^{-1}(|x|^{-N})} = 1, \text{ or}$$

(ii) *for some  $0 < c_* < 1$  and any  $a \in (c_*, 1)$ , the graph of  $u$  crosses the graph of the function  $aF^{-1}(|x|^{-N})$  infinitely often in  $B_R$ .*

*Proof.* Let

$$c_* := \liminf_{x \rightarrow 0} \frac{u(x)}{F^{-1}(|x|^{-N})}.$$

If  $c_* = 1$ , the first alternative holds. Otherwise, we may take any  $a \in (c_*, 1)$  and verify that the second alternative holds for all such  $a$ .  $\blacksquare$

We point out that a similar alternative holds in the super-critical case in higher dimensions (see Theorem 5.13 in [23]). See Corollary 7.7 for a more precise version.



## 7 An integral-type lower bound for singular solutions

**Lemma 7.1** *Let  $f$  satisfy the assumptions in (6.1) and (6.2). Further assume that there exists a  $\mu > 1$  such that*

$$(7.1) \quad f(t) \geq Ce^{t^\mu} \quad \text{for all large } t > 0 \quad \text{and some } C > 0.$$

*If  $u$  is an unbounded solution of  $(P_R^*)$ , then the following holds:*

$$(7.2) \quad \int_0^R \frac{1}{r|(\log(N/r))|^{\frac{1}{\mu}}} \left( \int_{B_r(0)} f(u) dx \right)^{\frac{1}{N-1}} dr = \infty.$$

**Remark 7.2** *Note that we always have  $\int_{B_R} f(u) dx < \infty$ .*

*Proof.* Choose  $\kappa > 0$  as in (6.1) and for any  $t \geq 0$  define  $G_\kappa(t) := Ce^{t^\mu} + \kappa t$ . Since

$$F(t) \geq G_\kappa(t) \geq G_0(t), \quad \text{for all } t \geq 0,$$

we have

$$(7.3) \quad (\log t - \log C)^{\frac{1}{\mu}} = G_0^{-1}(t) \geq F^{-1}(t) \quad \text{for all large } t > 0.$$

Supposing that the conclusion of lemma does not hold and write the solution  $u$  in the form:

$$(7.4) \quad u(x) := v(x)(N \log(N/|x|))^{\frac{1}{\mu}} \quad \text{for } x \in B_R \setminus \{0\}.$$

From (7.3) and Lemma 6.1 (a), we know that

$$(7.5) \quad 0 < v(x) < 1 \quad \text{for all small } |x|.$$

Going to the radial variable  $r$  and the Emden-Fowler variable  $t$  as given in (4.4), we have that  $y(t) := u(|x|)$  and  $z(t) := v(|x|)$  satisfy:

$$(7.6) \quad y(t) := z(t)t^{\frac{1}{\mu}} \quad \text{for } t \geq T := N \log(N/R).$$

Since  $f$  is super-exponential, we have that (see the proof of Lemma 4.1)

$$\lim_{t \rightarrow \infty} y'(t) = 0.$$

Recall that  $y$  solves the differential equation in (4.5). We see then that  $y$  also solves the equation

$$(7.7) \quad y'(t) = \left( \int_t^\infty f(y(s))e^{-s} ds \right)^{\frac{1}{N-1}}, \quad \text{for } t > T.$$

Thus, we obtain the following equation for  $z$ :

$$(7.8) \quad z'(t) + \frac{1}{\mu t} z(t) = t^{-\frac{1}{\mu}} \left( \int_t^\infty f(z(s)s^{\frac{1}{\mu}})e^{-s} ds \right)^{\frac{1}{N-1}}, \quad \text{for } t > T.$$

Note that  $0 < z(t) < 1$  for all large  $t > 0$ . Making the final transformation

$$(7.9) \quad \xi := \log t, \quad \rho(\xi) = z(t), \quad \text{for } t \geq T_0 := \max\{1, T\},$$

the equation in (7.8) simplifies to:

$$(7.10) \quad \rho'(\xi) + \frac{1}{\mu}\rho(\xi) = H(\xi), \quad \xi > \log T_0$$

where

$$(7.11) \quad H(\xi) := e^{\left(\frac{\mu-1}{\mu}\right)\xi} \left( \int_{\xi}^{\infty} f(\rho(\zeta)) e^{\frac{\zeta}{\mu}} e^{-e^{\zeta}} e^{\zeta} d\zeta \right)^{\frac{1}{N-1}}.$$

Note that  $0 < \rho(\xi) < 1$  for all large  $\xi > 0$ . It can also be deduced, following the proof of Lemma 6.1(b), that

$$\limsup_{\xi \rightarrow \infty} \rho(\xi) = 1.$$

The assumption that the integral in (7.2) is finite translates to

$$(7.12) \quad H_0 := \int_{\log T_0}^{\infty} H(\xi) d\xi < \infty.$$

Integrating the equation (7.10) between the limits  $\xi > \log T_0$  and  $\xi_n > \xi$ , we obtain :

$$(7.13) \quad \rho(\xi_n) - \rho(\xi) + \frac{1}{\mu} \int_{\xi}^{\xi_n} \rho(\zeta) d\zeta = \int_{\xi}^{\xi_n} H(\zeta) d\zeta \quad \text{for all } \xi > \log T_0.$$

Letting  $n \rightarrow \infty$ , using the fact that  $0 < \rho < 1$  and (7.12),

$$(7.14) \quad \frac{1}{\mu} \int_{\xi}^{\infty} \rho(\zeta) d\zeta \leq \int_{\xi}^{\infty} H(\zeta) d\zeta + 1 \leq H_0 + 1 \quad \text{for all } \xi > \log T_0.$$

Combining with (7.10), we get that  $\rho \in W^{1,1}([\log T_0, \infty))$ . Necessarily,  $\rho(s) \rightarrow 0$  as  $s \rightarrow \infty$ , a contradiction to the fact that  $\limsup_{s \rightarrow \infty} \rho = 1$ . This proves the lemma.  $\blacksquare$

**Remark 7.3** *The example in lemma 5.2 is relevant in the context of the above result.*

**Remark 7.4** *When  $N = 2$ , by Fubini theorem, the condition (7.2) reduces to:*

$$\int_{B_R} |\log(2/|x|)|^{1-\frac{1}{\mu}} f(u) dx = \infty.$$

Define

$$w(r) := \int_r^R s^{1-N} |\log(N/s)|^{\frac{1-N}{\mu}} ds.$$

**Corollary 7.5** *Let  $f$  satisfy the assumptions in Lemma 7.1. If  $u$  is an unbounded solution of  $(P_R^*)$ , then the following holds:*

$$(7.15) \quad \int_{B_R} f(u) w(|x|) dx = \infty.$$

*Proof.* Let  $R_0 := \min\{N, R\}$ . By (7.2), Jensen's inequality and Fubini theorem,

$$\begin{aligned} \infty &= \left( \int_0^{R_0} \frac{1}{r(\log(N/r))^{\frac{1}{\mu}}} \left( \int_{B_r(0)} f(u) dx \right)^{\frac{1}{N-1}} dr \right)^{N-1} \\ &\leq R_0^{N-2} \int_0^{R_0} \frac{1}{r^{N-1}(\log(N/r))^{\frac{N-1}{\mu}}} \left( \int_{B_r(0)} f(u) dx \right) dr \\ &\leq R_0^{N-2} \int_{B_{R_0}} f(u) w(|x|) dx. \end{aligned}$$

**Corollary 7.6** *Let  $f(t) := e^{t^\mu}$ ,  $\mu > 1$ , and  $u$  be a solution of  $(P_R^*)$ . Suppose that for some  $\theta \geq (1 - \frac{1}{\mu})(N-1) + 1$ ,*

$$(7.16) \quad u(x) \leq \left( N \log(N/|x|) - \theta \log(N \log(N/|x|)) \right)^{\frac{1}{\mu}}, \text{ for all small } |x|.$$

*Then,  $u$  is bounded.*

*Proof.* It is easy to check that (7.16) implies (refer to (7.11)),

$$(7.17) \quad \int_0^\infty H(\zeta) d\zeta < \infty.$$

From proof of Lemma 7.1 we obtain that  $\lim_{t \rightarrow \infty} z(t) = 0$ . Following the arguments at the end of proof of Lemma 6.1, the conclusion follows.  $\blacksquare$

We can now refine the bound in Lemma 6.1(a) and Corollary 6.2 (ii) for the nonlinearity in the above result.

**Corollary 7.7** *Let  $f(t) := e^{t^\mu}$ ,  $\mu > 1$ , and  $\theta \geq (1 - \frac{1}{\mu})(N-1) + 1$ . Then, one of the following alternatives holds for an unbounded solution  $u$  of  $(P_R^*)$ :*

- (i)  $\left( N \log(N/|x|) - \theta \log(N \log(N/|x|)) \right)^{\frac{1}{\mu}} \leq u(x) \leq \left( N \log(N/|x|) \right)^{\frac{1}{\mu}}$ , for all small  $|x|$  or
- (ii)  $u(x) \leq \left( N \log(N/|x|) \right)^{\frac{1}{\mu}}$  for all small  $|x|$  and the graph of  $u$  crosses the graph of the function  $\left( N \log(N/|x|) - \theta \log(N \log(N/|x|)) \right)^{\frac{1}{\mu}}$  infinitely often in  $B_r$  for all small  $r > 0$ .

*Acknowledgements.* *The authors would like to thank warmly the anonymous referees for valuable comments which have improved the paper.*

## References

- [1] J. A. AGUILAR CRESPO and I. PERAL ALONSO, *Blow-up Behavior for solutions of  $-\Delta_N u = V(x)e^u$  in bounded domains in  $\mathbb{R}^N$* , *Nonlinear Anal.*, 29(4) (1997), 365–384.

- [2] M. AIZENMAN and B. SIMON, *Brownian motion and Harnack inequality for Schrodinger operators*, Comm. Pure Appl. Math. 35 (1982), 209–271.
- [3] F. V. ATKINSON and L. A. PELETIER, *Ground states and Dirichlet problems for  $-\Delta u = f(u)$  in  $\mathbb{R}^2$* , Arch. Rational Mech. Anal., 96 (1986), 147–165.
- [4] M. F. BIDAUT-VÉRON, *Local and global behavior of solutions of quasilinear equations of Emden-Fowler type*, Arch. Rational Mech. Anal., 107(4) (1989), 293–324.
- [5] L. BOCCARDO, I. PERAL ALONSO and J. L. VAZQUEZ, *The  $N$ -Laplacian elliptic equation: Variational versus entropy solutions*, J. Math. Anal. Appl., 201 (1996), 671– 688.
- [6] H. BREZIS and P.L. LIONS, *A Note on Isolated Singularities for Linear Elliptic Equations*, Mathematical Analysis and Applications, Part A Advances in Math. Suppl. Stud., 7A, 263–266.
- [7] H. BREZIS and F. MERLE, *Uniform Estimates and blow up behaviour for solutions of  $-\Delta u = V(x)e^u$  in two dimensions*, Comm. Partial Differential Equations, 16(8-9) (1991), 1223–1253.
- [8] H. BREZIS and L. VÉRON, *Removable singularities for some nonlinear elliptic equations*, Arch. Ration. Mech. Anal., 75 (1980), 1–6.
- [9] R. DHANYA, J. GIACOMONI and S. PRASHANTH, *Isolated singularities for the exponential type semilinear elliptic equation in  $\mathbb{R}^2$* , Proc. Amer. Math. Soc., 137(12) (2009), 4099–4107.
- [10] M. GHERGU and S. D. TALIAFERRO, *Isolated singularities in partial differential inequalities*, Encyclopedia of Mathematics and Its Applications, 161, Cambridge University Press, 2016.
- [11] J. GIACOMONI, S. PRASHANTH, and K. SREENADH, *Uniqueness and multiplicity results for  $N$ -Laplace equation with critical and singular nonlinearity in a ball*, Asymptotic Analysis, 65 (2009), 195–227.
- [12] J. GIACOMONI, S. PRASHANTH, and K. SREENADH, *A global multiplicity result for  $N$ -Laplacian with critical nonlinearity of concave-convex type*, J. Differential Equations, 232 (2007), 544–572.
- [13] M. GUEDDA and L. VÉRON, *Local and global properties of solutions of quasilinear elliptic equations*, J. Differential Equations, 76 (1988), 159–189.
- [14] N. IOKU, *Brezis-Merle type inequality for a weak solution to the  $N$ -Laplace equation in Lorentz-Zygmund spaces*, Differential Integral Equations, 22(5) (2009), 495–518.
- [15] S. KICHENASSAMY and L. VÉRON, *Singular solutions of the  $p$ -Laplace equation*, Math. Ann., 275 (1986), 599–615.

- [16] H. KIKUCHI and J. WEI, *A bifurcation diagram of solutions to an elliptic equation with exponential nonlinearity in higher dimensions*, Proc. Roy. Soc. Edinburgh Sect. A, 148(1) (2018), 101–122.
- [17] G. LIEBERMAN, *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal., 12(11) (1988), 1203–1219.
- [18] P. L. LIONS, *Isolated Singularities in semilinear problems*, J. Differential Equations, 38 (1980), 441–450.
- [19] J. MOSER, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. Jour., 20(11) (1971), 1077–1092.
- [20] W. RUDIN, *Principles of mathematical analysis*, 3rd Edition, McGraw Hill International Editions, 1976.
- [21] J. SERRIN, *Local behaviour of solutions of quasilinear equations*, Acta Math., 111(1964), 247–302.
- [22] P. TOLKSDORF, *On the Dirichlet problem for quasilinear equations in domains with conical boundary points*, Comm. Partial Diff. Equations, 8(7) (1983), 773–817.
- [23] L. VÉRON, *Singularities of solutions of second order quasilinear equations*, Pitman research Notes in Mathematics Series, 1996.
- [24] L. VÉRON, *Elliptic equations involving measures*, Handbook of Differential Equations, 1 (2004), 593–712.