



HAL
open science

A Global multiplicity result for a very singular critical nonlocal equation

Jacques Giacomoni, T Mukherjee, K. Sreenadh

► **To cite this version:**

Jacques Giacomoni, T Mukherjee, K. Sreenadh. A Global multiplicity result for a very singular critical nonlocal equation. *Topological Methods in Nonlinear Analysis*, In press, 10.12775/TMNA.2019.049 . hal-02161068

HAL Id: hal-02161068

<https://hal.science/hal-02161068>

Submitted on 20 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A GLOBAL MULTIPLICITY RESULT FOR A VERY SINGULAR CRITICAL NONLOCAL EQUATION

J. Giacomoni^{*}, T. Mukherjee[†] and K. Sreenadh[‡]

Abstract

In this article, we show the global multiplicity result for the following nonlocal singular problem

$$(P_\lambda) : \quad (-\Delta)^s u = u^{-q} + \lambda u^{2_s^* - 1}, \quad u > 0 \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega,$$

where Ω is a bounded domain in \mathbb{R}^n with smooth boundary $\partial\Omega$, $n > 2s$, $s \in (0, 1)$, $\lambda > 0$, $q > 0$ satisfies $q(2s - 1) < (2s + 1)$ and $2_s^* = \frac{2n}{n-2s}$. Employing the variational method, we show the existence of at least two distinct weak positive solutions for (P_λ) in X_0 when $\lambda \in (0, \Lambda)$ and no solution when $\lambda > \Lambda$, where $\Lambda > 0$ is appropriately chosen. We also prove a result of independent interest that any weak solution to (P_λ) is in $C^\alpha(\mathbb{R}^n)$ with $\alpha = \alpha(s, q) \in (0, 1)$. The asymptotic behaviour of weak solutions reveals that this result is sharp.

Key words: Fractional Laplacian, very singular nonlinearity, variational method, Hölder regularity.

2010 Mathematics Subject Classification: 35R11, 35R09, 35A15.

1 Introduction

In this article, we prove the existence, multiplicity and Hölder regularity of weak solutions to the following fractional critical and singular elliptic equation

$$(P_\lambda) : \quad (-\Delta)^s u = u^{-q} + \lambda u^{2_s^* - 1}, \quad u > 0 \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega,$$

where Ω is a bounded domain in \mathbb{R}^n with smooth boundary $\partial\Omega$, $n > 2s$, $s \in (0, 1)$, $\lambda > 0$, $q > 0$ satisfies $q(2s - 1) < (2s + 1)$ and $2_s^* = \frac{2n}{n-2s}$. The fractional Laplace operator denoted by $(-\Delta)^s$ is defined as

$$(-\Delta)^s u(x) = 2C_s^n \text{P.V.} \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy$$

^{*}Université de Pau et des Pays de l'Adour-E2S, CNRS, LMAP (UMR 5142) Bat. IPRA, Avenue de l'Université, 64013 Pau cedex, France. e-mail:jacques.giacomoni@univ-pau.fr

[†]Department of Mathematics, Indian Institute of Technology Delhi, Hauz Khaz, New Delhi-110016, India. e-mail: tulimukh@gmail.com

[‡]Department of Mathematics, Indian Institute of Technology Delhi, Hauz Khaz, New Delhi-110016, India. e-mail: sreenadh@gmail.com

where P.V. denotes the Cauchy principal value and $C_s^n = \pi^{-\frac{n}{2}} 2^{2s-1} s \frac{\Gamma(\frac{n+2s}{2})}{\Gamma(1-s)}$, Γ being the Gamma function. The fractional power of Laplacian is the infinitesimal generator of Lévy stable diffusion process and arise in anomalous diffusion in plasma, population dynamics, geophysical fluid dynamics, flames propagation, chemical reactions in liquids and American options in finance, see [3] for instance. The theory of fractional Laplacian and elliptic equations involving it as the principal part has been evolved immensely in recent years. There is a vast literature available on it, however without giving an exhaustive list we cite [7, 10, 14, 16, 19, 21, 22] for motivation to readers.

Nowadays, researchers are inspecting on various forms of singular nonlocal equations. We cite [11, 8, 9] as some contemporary works related to it. The fractional elliptic equations with singular and critical nonlinearities was first studied by Barrios et al. in [5]. The authors considered the problem

$$(-\Delta)^s u = \lambda \frac{f(x)}{u^\gamma} + Mu^p, \quad u > 0 \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega,$$

where $n > 2s$, $M \geq 0$, $0 < s < 1$, $\gamma > 0$, $\lambda > 0$, $1 < p < 2_s^* - 1$ and $f \in L^m(\Omega)$, $m \geq 1$ is a nonnegative function. Here, authors studied the existence of distributional solutions using the uniform estimates of $\{u_n\}$ which are solutions of the regularized problems with singular term $u^{-\gamma}$ replaced by $(u + \frac{1}{n})^{-\gamma}$. Motivated by their results, Sreenadh and Mukherjee in [15] studied the singular problem

$$(-\Delta)^s u = \lambda a(x)u^{-q} + u^{2_s^*-1}, \quad u > 0 \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega,$$

where $\lambda > 0$, $0 < q \leq 1$ and $\theta \leq a(x) \in L^\infty(\Omega)$, for some $\theta > 0$. They showed that although the energy functional corresponding to this problem fails to be Fréchet differentiable, making use of its Gâteaux differentiability the Nehari manifold technique can still be benefitted to obtain existence of at least two solutions over a certain range of λ . The significance of q being less than 1 is the Gâteaux differentiability of the functional corresponding to the problem. Consider now the case $q > 1$. Let

$$X \stackrel{\text{def}}{=} \left\{ u \mid u : \mathbb{R}^n \rightarrow \mathbb{R} \text{ is measurable, } u|_\Omega \in L^2(\Omega) \text{ and } \frac{(u(x) - u(y))}{|x - y|^{\frac{n}{2}+s}} \in L^2(Q) \right\},$$

where $Q \stackrel{\text{def}}{=} \mathbb{R}^{2n} \setminus (\mathcal{C}\Omega \times \mathcal{C}\Omega)$ and $\mathcal{C}\Omega := \mathbb{R}^n \setminus \Omega$ endowed with the norm

$$\|u\|_X \stackrel{\text{def}}{=} \|u\|_{L^2(\Omega)} + [u]_X,$$

where

$$[u]_X \stackrel{\text{def}}{=} \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy \right)^{\frac{1}{2}}.$$

Let $J : X_0 \rightarrow \mathbb{R}$ be the functional defined by

$$J(u) \stackrel{\text{def}}{=} \frac{C_s^n}{2} \|u\|_{X_0}^2 - \frac{1}{1-q} \int_\Omega |u|^{1-q} dx - \frac{\lambda}{2_s^*} \int_\Omega |u|^{2_s^*} dx$$

for any u in the Hilbert space $X_0 \stackrel{\text{def}}{=} \{u \in X : u = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}$ equipped with the inner product

$$\langle u, v \rangle \stackrel{\text{def}}{=} \int_Q \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+2s}} dx dy.$$

J may not be defined on the whole space nor it is even continuous on $D(I) \equiv \{u \in X_0 : I(u) < \infty\}$ and the approach used for $q < 1$ can not be extended. Besides this, one has that the interior of $D(I) = \emptyset$ because of the singular term. But we notice that if we enforce the condition $q > 1$ satisfies $q(2s - 1) < (2s + 1)$ then we can prove that $D(I)$ is non empty and Gâteaux differentiable on a suitable convex cone of X_0 .

The existence of weak solutions to (P_λ) when $\lambda \in (0, \Lambda)$ and no solution when $\lambda > \Lambda$ has been already obtained by Giacomoni et al. in [12]. But here the multiplicity of solutions has been achieved in $L^1_{loc}(\Omega)$ only, by using non smooth critical point theory, so the questions of existence of solutions in the energy space and of Hölder regularity were still pending. This article is bringing answers to these two issues. For that, we followed the approach of [13] but we notify that the adversity and novelty of this article lies in extending Haitao's technique in a nonlocal framework. The regularity of weak solution of the purely singular problem

$$(-\Delta)^s u = u^{-q}, \quad u > 0, \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega$$

plays a vital role in our study. This has been obtained by Adimurthi, Giacomoni and Santra in [1] in recent times. In the present paper we extend the Hölder regularity results proved in [1, Theorem 1.4] in our framework of weak solutions (see definition 1.1 below) rather than the more restricted classical solutions framework (defined in [1]). It requires additional L^∞ -estimates and the use of the weak comparison principle.

Our paper has been organized as follows. Section 2 contains some preliminary results used in the subsequent sections. Section 3 and 4 contain the proof of existence of first and second weak solution to (P_λ) respectively (Theorem 1.2). The proof of the Hölder regularity result (Theorem 1.3) is done in Section 4 based on a priori estimates proved in Proposition 4.1. Now we state the main results proved in the paper. First we define the notion of weak solutions.

Definition 1.1 *A function $u \in X_0$ is said to be a weak solution of (P_λ) if there exists a $m_K > 0$ such that $u > m_K$ in every compact subset K of Ω , and it satisfies*

$$C_s^n \int_Q \frac{(u(x) - u(y))(\phi(x) - \phi(y))}{|x - y|^{n+2s}} dx dy = \int_\Omega (u^{-q} + u^{2^*_s-1})\phi dx, \text{ for all } \phi \in X_0.$$

Given any $\phi \in C_0(\overline{\Omega})$ such that $\phi > 0$ in Ω we define

$$C_\phi(\Omega) \stackrel{\text{def}}{=} \{u \in C_0(\overline{\Omega}) \mid \exists c \geq 0 \text{ such that } |u(x)| \leq c\phi(x), \forall x \in \Omega\}$$

with the usual norm $\left\| \frac{u}{\phi} \right\|_{L^\infty(\Omega)}$ and the associated positive cone. We define the following open convex subset of $C_\phi(\Omega)$ as

$$C_\phi^+(\Omega) \stackrel{\text{def}}{=} \left\{ u \in C_\phi(\Omega) \mid \inf_{x \in \Omega} \frac{u(x)}{\phi(x)} > 0 \right\}.$$

In particular, C_ϕ^+ contains all those functions $u \in C_0(\Omega)$ with $k_1\phi \leq u \leq k_2\phi$ in Ω for some $k_1, k_2 > 0$. Let $\phi_{1,s}$ be the first positive normalized eigenfunction ($\|\phi_{1,s}\|_{L^\infty(\Omega)} = 1$) of $(-\Delta)^s$ in X_0 . We recall that $\phi_{1,s} \in C^s(\mathbb{R}^N)$ and $\phi_{1,s} \in C_{\delta^s}^+(\Omega)$ where $\delta(x) = \text{dist}(x, \partial\Omega)$ (see for instance Proposition 1.1 and Theorem 1.2 in [17]). We then define the barrier function ϕ_q as follows:

$$\phi_q \stackrel{\text{def}}{=} \begin{cases} \phi_{1,s} & \text{if } 0 < q < 1, \\ \phi_{1,s} \left(\ln \left(\frac{2}{\phi_{1,s}} \right) \right)^{\frac{1}{q+1}} & \text{if } q = 1, \\ \phi_{1,s}^{\frac{2}{q+1}} & \text{if } q > 1. \end{cases}$$

We prove the following as the main results:

Theorem 1.2 *There exists $\Lambda > 0$ such that*

- (i) (P_λ) admits at least two solutions in $X_0 \cap C_{\phi_q}^+(\Omega)$ for every $\lambda \in (0, \Lambda)$;
- (ii) (P_λ) admits no solution for $\lambda > \Lambda$;
- (iii) (P_Λ) admits at least one positive solution $u_\Lambda \in X_0 \cap C_{\phi_q}^+(\Omega)$.

Theorem 1.3 *Let $\lambda \in (0, \Lambda]$, $q > 0$ satisfies $q(2s - 1) < (2s + 1)$ and $u \in X_0$ is any positive weak solution of (P_λ) then*

- (i) $u \in C^s(\mathbb{R}^n)$ when $0 < q < 1$;
- (ii) $u \in C^{s-\epsilon}(\mathbb{R}^n)$ for any small enough $\epsilon > 0$ when $q = 1$;
- (iii) $u \in C^{\frac{2s}{q+1}}(\mathbb{R}^n)$ when $q > 1$.

Remark 1.4 *Here, the Hölder regularity for the weak solutions of (P_λ) obtained is optimal because of the behavior of the solution near $\partial\Omega$ since we showed that any weak solution of (P_λ) lies in $C_{\phi_q}^+(\Omega)$.*

Remark 1.5 *It follows from Theorem 1.3 that the extremal solution (when $\lambda = \Lambda$), in case of critical growth nonlinearities is a classical solution which extends the results in [1] where in this regard only subcritical nonlinearities are considered.*

2 Preliminaries

We start by some preliminary results. The energy functional corresponding to (P_λ) is given by $I_\lambda : X_0 \rightarrow \mathbb{R}$ defined as

$$I_\lambda(u) \stackrel{\text{def}}{=} \begin{cases} \frac{C_s^n \|u\|_{X_0}^2}{2} - \frac{1}{1-q} \int_\Omega |u|^{1-q} dx - \frac{1}{2_s^*} \int_\Omega |u|^{2_s^*} dx & \text{if } q \neq 1, \\ \frac{C_s^n \|u\|_{X_0}^2}{2} - \int_\Omega \ln |u| dx - \frac{1}{2_s^*} \int_\Omega |u|^{2_s^*} dx & \text{if } q = 1. \end{cases}$$

Let $q > 0$ satisfies $q(2s - 1) < (2s + 1)$. Then for any $\varphi \in X_0$ and $u \in C_{\phi_q}^+(\Omega)$, by Hardy's inequality (see [23, Lemma 3.2.6.1, p. 259]) we obtain

$$\int_{\Omega} u^{-q} \varphi \leq \left(\int_{\Omega} \frac{dx}{(\delta(x))^{\frac{2s(q-1)}{q+1}}} \right)^{\frac{1}{2}} \left(\frac{\varphi^2}{(\delta(x))^{2s}} \right)^{\frac{1}{2}} < K \|\varphi\| < +\infty \quad (2.1)$$

where $K > 0$ is a constant. If we define $D(I) = \{u \in X_0 : I_{\lambda}(u) < \infty\}$ then by virtue of (2.1) we get that $D(I) \neq \emptyset$. This gives an importance of the inequality $q(2s - 1) < (2s + 1)$. From the proof of [1, Theorem 1.2], we know that if $0 < q < 1$ and $u \in X_0$ satisfies $u \geq c\delta^s$ then I_{λ} is Gâteaux differentiable at u . In the proposition below, we show the same property of I_{λ} when $q \geq 1$ satisfies $q(2s - 1) < (2s + 1)$.

Proposition 2.1 *If $M = \{u \in X_0 : u_1 \leq u \leq u_2\}$ where $u_1 \in C_{\phi_q}^+(\Omega)$ and $u_2 \in X_0$ then I_{λ} is Gâteaux differentiable at u in the direction $(v - u)$ where $v, u \in M$.*

Proof. We need to show that

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{I_{\lambda}(u + t(v - u)) - I_{\lambda}(u)}{t} &= C_s^n \int_Q \frac{(v(x) - v(y))((v - u)(x) - (v - u)(y))}{|x - y|^{n+2s}} dx dy \\ &\quad - \int_{\Omega} u^{-q}(v - u) dx - \lambda \int_{\Omega} u^{2s^*-1}(v - u) dx. \end{aligned}$$

It is enough to show this for the singular term; for the rest two terms, the proof is standard. For any $t \in (0, 1)$, $u + t(v - u) \in M$ since M is convex. Consider $F(u) = \frac{1}{1 - q} \int_{\Omega} u^{1-q} dx$ then using Mean Value Theorem we get

$$\begin{aligned} \frac{F(u + t(v - u)) - F(u)}{t} &= \frac{1}{t(1 - q)} \int_{\Omega} ((u + t(v - u))^{1-q} - u^{1-q})(x) dx \\ &= \int_{\Omega} (u + t\theta(v - u))^{-q}(x)(v - u)(x) dx \end{aligned}$$

for some $\theta \in (0, 1)$. Since $(u + t\theta(v - u)) \in M$ and (2.1), we have

$$\int_{\Omega} (u + t\theta(v - u))^{-q}(v - u) dx \leq \int_{\Omega} u_1^{-q}(v - u) dx < +\infty.$$

So using Lebesgue Dominated convergence theorem we pass through the limit $t \rightarrow 0$ and get

$$\lim_{t \rightarrow 0} \frac{F(u + t(v - u)) - F(u)}{t} = \int_{\Omega} u^{-q}(v - u) dx.$$

This completes the proof. \square

Let $L(u) := (-\Delta)^s u - u^{-q}$. Then L forms a monotone operator. So we have the following comparison principle:

Lemma 2.2 *Let $u_1, u_2 \in X_0 \cap C_{\phi_q}^+(\Omega)$ are weak solutions to*

$$L(u_1) = g_1 \text{ in } \Omega, \quad L(u_2) = g_2 \text{ in } \Omega$$

with $g_1, g_2 \in L^2(\Omega)$ such that $g_1 \leq g_2$ a.e. in Ω . Then $u_1 \leq u_2$ a.e. in Ω . Moreover if $g \in L^\infty(\Omega)$ then the problem

$$L(u) = g \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega$$

has a unique solution in X_0 .

3 Existence result

Let us define

$$\Lambda := \sup\{\lambda > 0 : (P_\lambda) \text{ has a weak solution}\}.$$

Also let $\underline{w} \in C_0(\overline{\Omega})$ solves the purely singular problem

$$(-\Delta)^s \underline{w} = \underline{w}^{-q}, \quad \underline{w} > 0 \text{ in } \Omega, \quad \underline{w} = 0 \text{ in } \mathbb{R}^n \setminus \Omega.$$

Then [1, Theorems 1.2 and 1.4] give us that \underline{w} is unique, $\underline{w} \in X_0 \cap C_{\phi_q}^+(\Omega)$ and $\underline{w} \in C^{s_q}(\mathbb{R}^n)$

$$\text{where } s_q \stackrel{\text{def}}{=} \begin{cases} s \text{ if } q < 1, \\ s - \epsilon \text{ for any } \epsilon > 0 \text{ if } q = 1, \\ \frac{2s}{q+1} \text{ if } 1 < q \text{ and } q(2s-1) < 2s+1. \end{cases}$$

For sake of clarity we basically focus on the case $1 \leq q$ and $q(2s-1) < (2s+1)$. Indeed, when $q \in (0, 1)$, the case follows easily along the same line. In this context, the next result is an important lemma for Λ .

Lemma 3.1 *It holds $0 < \Lambda < +\infty$.*

Proof. First we prove that $\Lambda < +\infty$. Using $\phi_{1,s}$ as the test function in (P_λ) we get

$$\int_{\Omega} (u^{-q} \phi_{1,s} + \lambda u^{2s^*-1} \phi_{1,s}) \, dx = \int_{\mathbb{R}^n} \phi_{1,s} (-\Delta)^s u \, dx = \int_{\mathbb{R}^n} u (-\Delta)^s \phi_{1,s} \, dx = \lambda_{1,s} \int_{\Omega} u \phi_{1,s} \, dx. \quad (3.1)$$

If we choose a $\lambda > 0$ which satisfies $t^{-q} + \lambda t^{2s^*-1} > 2\lambda_{1,s} t$ for all $t > 0$ then we get a contradiction to (3.1). Therefore it must be $\Lambda < +\infty$. Now to prove $\Lambda > 0$ we need sub and supersolution for (P_λ) . It is easy to see that $\underline{u}_\lambda = \underline{w}$ forms a subsolution of (P_λ) and $\overline{u}_\lambda = \underline{u}_\lambda + Mz$ for $\lambda > 0$ small enough and for a $M = M(\lambda) > 0$ forms a supersolution of (P_λ) , where $0 < z \in X_0$ solves $(-\Delta)^s z = 1$ in Ω . Now we define the closed convex subset M_λ of X_0 as

$$M_\lambda := \{u \in X_0 : \underline{u}_\lambda \leq u \leq \overline{u}_\lambda\}.$$

Consider the iterative scheme ($k \geq 1$):

$$(P_{\lambda,k}) \begin{cases} (-\Delta)^s u_k - u_k^{-q} = \lambda u_{k-1}^{2s^*-1}, & u_k > 0 \text{ in } \Omega \\ u_k = 0 & \text{in } \mathbb{R}^n \setminus \Omega \end{cases}$$

with $u_0 = \underline{u}_\lambda$. The existence of $\{u_k\}$ in $X_0 \cap M_\lambda \cap C_{\phi_q}^+(\Omega)$ can be proved by considering the approximated problem corresponding to $(P_{\lambda,k})$, for instance we refer [6, Proposition 2.3]. From Lemma 2.2, it follows that $\{u_k\}$ is increasing and $u_k \in M_\lambda$ for all k . Let $\lim_{k \uparrow \infty} u_k = u_\lambda$. Then testing $(P_{\lambda,k})$ by u_k we get

$$\|u_k\|^2 \leq 2 \int_{\Omega} \overline{u_\lambda}^2 dx + \lambda \int_{\Omega} \overline{u_\lambda}^{-2_s^*} dx + \int_{\Omega} \overline{u_\lambda} \underline{u_\lambda}^{-q} \leq K_\lambda$$

where $K_\lambda > 0$ is a constant depending on λ . So, up to a subsequence, $u_k \rightharpoonup u_\lambda$ in X_0 . Finally using Lebesgue dominated convergence Theorem we pass through the limit in $(P_{\lambda,k})$ to obtain u_λ solves (P_λ) weakly and obviously, $u_\lambda \in M_\lambda$. This proves that $\Lambda > 0$. \square

In the next result, we prove the existence of a weak solution for (P_λ) whenever $\lambda \in (0, \Lambda)$. In the proof, we use a minimization on a conical shell argument similar as in [2, Lemma 4.1] and in [6, Proposition 3.5]. But here we take advantage of the existence of a strict positive subsolution to control the singular nonlinearity.

Proposition 3.2 *For each $\lambda \in (0, \Lambda)$, (P_λ) admits a weak solution $w \in C_{\phi_q}^+(\Omega)$.*

Proof. The proof goes along the line of Perron's method adapted over a nonlocal framework (see [13, Lemma 2.2]). Let $\lambda \in (0, \Lambda)$ and $\lambda' \in (\lambda, \Lambda)$ then it is easy to see that $u_{\lambda'}$, a weak solution of $(P_{\lambda'})$, forms a supersolution for (P_λ) . Such a λ' exists because of the definition of Λ and Lemma 3.1. Let \underline{u}_λ be the same function as defined in Lemma 3.1 and consider the closed convex subset W_λ of X_0 as

$$W_\lambda = \{u \in X_0 : \underline{u}_\lambda \leq u \leq u_{\lambda'}\}$$

Then for each $u \in W_\lambda$, because of fractional Sobolev embedding I_λ satisfies

$$I_\lambda(u) \geq \begin{cases} \frac{C_s^n \|u\|^2}{2} - \frac{C}{2_s^*} \|u\|^{2_s^*} & \text{if } q > 1, \\ \frac{C_s^n \|u\|^2}{2} - \frac{C}{2_s^*} \|u\|^{2_s^*} - C(\lambda') & \text{if } q \leq 1 \end{cases}$$

where $C(\lambda')$ is a positive constant depending solely on λ' . Then I_λ is bounded from below and coercive over W_λ . If $\{u_k\} \subset W_\lambda$ be such that $u_k \rightharpoonup u_0$ in X_0 as $k \rightarrow \infty$ then since for each k , $u_k \geq \underline{u}_\lambda$ for $q > 1$ and $u_k \leq u_{\lambda'}$ for $q \in (0, 1]$, $\int_{\Omega} u_k^{1-q} dx \leq \int_{\Omega} \underline{u}_\lambda^{1-q} dx$, we can use Lebesgue Dominated convergence theorem to get that

$$\int_{\Omega} u_k^{1-q} dx \rightarrow \int_{\Omega} u_0^{1-q} dx \text{ as } k \rightarrow \infty.$$

Hence from weak lower semicontinuity of norms, it follows that I_λ is weakly lower semicontinuous over W_λ . Moreover, W_λ is weakly sequentially closed subset of X_0 . Therefore there exists a $w \in W_\lambda$ such that

$$\inf_{u \in W_\lambda} I_\lambda(u) = I_\lambda(w). \tag{3.2}$$

Claim- w is a weak solution of (P_λ) .

Let $\varphi \in X_0$ and $\epsilon > 0$ then we define

$$v_\epsilon = \min\{u_{\lambda'}, \max\{\underline{u}_\lambda, w + \epsilon\varphi\}\} = w + \epsilon\varphi - \varphi^\epsilon + \varphi_\epsilon$$

where $\varphi^\epsilon = \max\{0, w + \epsilon\varphi - u_{\lambda'}\}$ and $\varphi_\epsilon = \max\{0, \underline{u}_\lambda - w - \epsilon\varphi\}$. By construction $v_\epsilon \in W_\lambda$ and $\varphi^\epsilon, \varphi_\epsilon \in X_0 \cap L^\infty(\Omega)$. Since $w + t(v_\epsilon - w) \in W_\lambda$ for each $t \in (0, 1)$, using (3.2) and Proposition 2.1 we get that

$$\begin{aligned} 0 &\leq \lim_{t \rightarrow 0^+} \frac{I_\lambda(w + t(v_\epsilon - w)) - I_\lambda(w)}{t} \\ &= \int_Q (v_\epsilon - w)(-\Delta)^s w \, dx - \int_\Omega w^{-q}(v_\epsilon - w) \, dx - \int_\Omega w^{2^*_s-1}(v_\epsilon - w) \, dx. \end{aligned}$$

This on simplification gives

$$\int_{\mathbb{R}^n} \varphi(-\Delta)^s w \, dx - \int_\Omega (w^{-q} + \lambda w^{2^*_s-1})\varphi \, dx \geq \frac{1}{\epsilon}(E^\epsilon - E_\epsilon) \quad (3.3)$$

where

$$\begin{aligned} E^\epsilon &= \int_{\mathbb{R}^n} \varphi^\epsilon(-\Delta)^s w \, dx - \int_\Omega (w^{-q} + \lambda w^{2^*_s-1})\varphi^\epsilon \, dx \\ &= \int_{\mathbb{R}^n} \varphi^\epsilon(-\Delta)^s (w - u_{\lambda'}) \, dx + \int_{\mathbb{R}^n} \varphi^\epsilon(-\Delta)^s u_{\lambda'} \, dx - \int_\Omega (w^{-q} + \lambda w^{2^*_s-1})\varphi^\epsilon \, dx \\ E_\epsilon &= \int_{\mathbb{R}^n} \varphi_\epsilon(-\Delta)^s w \, dx - \int_\Omega (w^{-q} + \lambda w^{2^*_s-1})\varphi_\epsilon \, dx \\ &= \int_{\mathbb{R}^n} \varphi_\epsilon(-\Delta)^s (w - \underline{u}_\lambda) \, dx + \int_{\mathbb{R}^n} \varphi_\epsilon(-\Delta)^s \underline{u}_\lambda \, dx - \int_\Omega (w^{-q} + \lambda w^{2^*_s-1})\varphi_\epsilon \, dx. \end{aligned}$$

We define $\Omega^\epsilon = \{x \in \Omega : (w + \epsilon\varphi)(x) \geq u_{\lambda'} > w(x)\}$ so that $\mathcal{L}(\Omega^\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0^+$ and also $\mathcal{C}\Omega^\epsilon := \Omega \setminus \Omega^\epsilon \subset \{x \in \Omega : (w + \epsilon\varphi)(x) < u_{\lambda'}(x)\}$ which implies that $\mathcal{L}(\Omega^\epsilon \times \mathcal{C}\Omega^\epsilon) \rightarrow 0$ as

$\epsilon \rightarrow 0^+$. Now we consider the term

$$\begin{aligned}
& \int_{\mathbb{R}^n} \varphi^\epsilon (-\Delta)^s (w - u_{\lambda'}) dx \\
&= \int_Q \frac{((w - u_{\lambda'})(x) - (w - u_{\lambda'})(y))(\varphi^\epsilon(x) - \varphi^\epsilon(y))}{|x - y|^{n+2s}} dx dy \\
&= \int_{\Omega^\epsilon} \int_{\Omega^\epsilon} \frac{|(w - u_{\lambda'})(x) - (w - u_{\lambda'})(y)|^2}{|x - y|^{n+2s}} dx dy \\
&\quad + \epsilon \int_{\Omega^\epsilon} \int_{\Omega^\epsilon} \frac{((w - u_{\lambda'})(x) - (w - u_{\lambda'})(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy \\
&\quad + 2 \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})^2(x)}{|x - y|^{n+2s}} dx dy + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})(x)\varphi(x)}{|x - y|^{n+2s}} dx dy \\
&\quad - 2 \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})(x)(w - u_{\lambda'})(y)}{|x - y|^{n+2s}} dx dy + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})(y)\varphi(x)}{|x - y|^{n+2s}} dx dy \\
&\quad + 2 \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega} \frac{(w - u_{\lambda'})^2(x)}{|x - y|^{n+2s}} dx dy + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega} \frac{(w - u_{\lambda'})(x)\varphi(x)}{|x - y|^{n+2s}} dx dy \\
&\geq \epsilon \int_{\Omega^\epsilon} \int_{\Omega^\epsilon} \frac{((w - u_{\lambda'})(x) - (w - u_{\lambda'})(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy \\
&\quad + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})(x)\varphi(x)}{|x - y|^{n+2s}} dx dy - 2\epsilon^2 \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{\varphi(x)\varphi(y)}{|x - y|^{n+2s}} dx dy \\
&\quad + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega^\epsilon} \frac{(w - u_{\lambda'})(y)\varphi(x)}{|x - y|^{n+2s}} dx dy + 2\epsilon \int_{\Omega^\epsilon} \int_{\mathcal{C}\Omega} \frac{(w - u_{\lambda'})(x)\varphi(x)}{|x - y|^{n+2s}} dx dy
\end{aligned}$$

where to obtain the last inequality, we use the fact that if $(x, y) \in \Omega^\epsilon \times \mathcal{C}\Omega^\epsilon$ then $(w - u_{\lambda'})(x)(w - u_{\lambda'})(y) \leq \epsilon^2 \varphi(x)\varphi(y)$. Therefore we get

$$\frac{1}{\epsilon} \int_{\mathbb{R}^n} \varphi^\epsilon (-\Delta)^s (w - u_{\lambda'}) dx \geq o(1) \text{ as } \epsilon \rightarrow 0^+.$$

Moreover using the fact that $u_{\lambda'}$ is a supersolution of (P_λ) , the other terms of $\frac{1}{\epsilon} E^\epsilon$ can be estimated as

$$\begin{aligned}
& \frac{1}{\epsilon} \int_{\mathbb{R}^n} \varphi^\epsilon (-\Delta)^s u_{\lambda'} dx - \frac{1}{\epsilon} \int_{\Omega} (w^{-q} + \lambda w^{2^*s-1}) \varphi^\epsilon dx \\
&\geq \frac{1}{\epsilon} \int_{\Omega^\epsilon} (u_{\lambda'}^{-q} - w^{-q}) \varphi^\epsilon dx + \frac{1}{\epsilon} \int_{\Omega^\epsilon} (u_{\lambda'}^{2^*s-1} - w^{2^*s-1}) \varphi^\epsilon dx \\
&\geq - \int_{\Omega^\epsilon} |u_{\lambda'}^{-q} - w^{-q}| |\varphi| dx = o(1) \text{ as } \epsilon \rightarrow 0^+.
\end{aligned}$$

Altogether we get

$$\frac{1}{\epsilon} E^\epsilon \geq o(1) \text{ as } \epsilon \rightarrow 0^+$$

and similarly we obtain

$$\frac{1}{\epsilon} E_\epsilon \leq o(1) \text{ as } \epsilon \rightarrow 0^+.$$

Hence (3.3) gives that for all $\varphi \in X_0$

$$\int_{\mathbb{R}^n} \varphi(-\Delta)^s w \, dx - \int_{\Omega} (w^{-q} + \lambda w^{2_s^*-1}) \varphi \, dx \geq o(1) \text{ as } \epsilon \rightarrow 0^+$$

but since φ was arbitrary, this implies that w is a weak solution of (P_λ) . This establishes the proof. \square

We now prove a special property of w , the weak solution of (P_λ) obtained in Proposition 3.2 following the proof in [6, Proposition 3.5]. We also refer [2, Proposition 5.2] where similar ideas were already used.

Lemma 3.3 *Let $\lambda \in (0, \Lambda)$ and w denotes the weak solution of (P_λ) obtained in Proposition 3.2. Then w forms a local minimum of the functional I_λ .*

Proof. We argue by contradiction, so suppose w is not a local minimum of I_λ . Then there exists a sequence $\{u_k\} \subset X_0$ satisfying

$$\|u_k - w\| \rightarrow 0 \text{ as } k \rightarrow \infty \text{ and } I_\lambda(u_k) < I_\lambda(w). \quad (3.4)$$

We define $\underline{u} = \underline{u}_\lambda$ and $\bar{u} = u_{\lambda'}$ as sub and supersolution of (P_λ) as defined in the proof of Proposition 3.2. Also we define

$$v_k = \max\{\underline{u}, \min\{u_k, \underline{u}\}\} = \begin{cases} \underline{u}, & \text{if } u_k < \underline{u}, \\ u_k, & \text{if } \underline{u} \leq u_k \leq \bar{u}, \\ \bar{u}, & \text{if } u_k > \bar{u}, \end{cases}$$

and $\underline{w}_k = (u_k - \underline{u})^-$, $\bar{w}_k = (u_k - \bar{u})^+$. Correspondingly, we define the sets $\underline{S}_k = \text{Supp}(\underline{w}_k)$ and $\bar{S}_k = \text{Supp}(\bar{w}_k)$. Then $u_k = v_k - \underline{w}_k + \bar{w}_k$ and $v_k \in W_\lambda$ where W_λ has been defined in Proposition 3.2. The main idea of the proof is to establish that the measures of \bar{S}_k and \underline{S}_k tend to 0 as $k \rightarrow \infty$ which together with $v_k \in W_\lambda$ force $I_\lambda(u_k)$ to be beyond $I_\lambda(w)$. First, we have that

$$\begin{aligned} \int_{\Omega} (u_k^+)^{1-q} dx &= \int_{\underline{S}_k} (u_k^+)^{1-q} dx + \int_{\bar{S}_k} (u_k^+)^{1-q} dx + \int_{\underline{u} \leq v_k \leq \bar{u}} (v_k)^{1-q} dx \\ &= \int_{\underline{S}_k} ((u_k^+)^{1-q} - \underline{u}^{1-q}) dx + \int_{\bar{S}_k} ((u_k^+)^{1-q} - \bar{u}^{1-q}) dx + \int_{\Omega} (v_k)^{1-q} dx \\ \text{and } \int_{\Omega} (u_k^+)^{2_s^*} dx &= \int_{\underline{S}_k} (u_k^+)^{2_s^*} dx + \int_{\bar{S}_k} (u_k^+)^{2_s^*} dx + \int_{\underline{u} \leq v_k \leq \bar{u}} (v_k)^{2_s^*} dx \\ &= \int_{\underline{S}_k} ((u_k^+)^{2_s^*} - \underline{u}^{2_s^*}) dx + \int_{\bar{S}_k} ((u_k^+)^{2_s^*} - \bar{u}^{2_s^*}) dx + \int_{\Omega} (v_k)^{2_s^*} dx. \end{aligned}$$

Then we can express $I_\lambda(u_k)$ as

$$\begin{aligned} I_\lambda(u_k) &= I_\lambda(v_k) + \frac{J_0}{2} - \frac{1}{1-q} \left(\int_{\underline{S}_k} ((u_k^+)^{1-q} - \underline{u}^{1-q}) dx + \int_{\bar{S}_k} ((u_k^+)^{1-q} - \bar{u}^{1-q}) dx \right) \\ &\quad - \frac{\lambda}{2_s^*} \left(\int_{\underline{S}_k} ((u_k^+)^{2_s^*} - \underline{u}^{2_s^*}) dx + \int_{\bar{S}_k} ((u_k^+)^{2_s^*} - \bar{u}^{2_s^*}) dx \right) \end{aligned} \quad (3.5)$$

where $J_0 = C_s^n (\|u_k\|^2 - \|v_k\|^2)$. While denoting $S_k = \{x \in \Omega : \underline{u} \leq v_k \leq \bar{u}\}$ and $h_k(x, y) = (u_k(x) - u_k(y))^2 - (v_k(x) - v_k(y))^2$, we get

$$\begin{aligned} J_0 &= \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy + \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy + 2 \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy \\ &+ 2 \int_{\underline{S}_k} \int_{S_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy + 2 \int_{\overline{S}_k} \int_{S_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy. \end{aligned}$$

Since $u_k = \overline{w}_k + \bar{u}$ and $v_k = \bar{u}$ in \overline{S}_k and $u_k = \underline{u} - \underline{w}_k$ and $v_k = \underline{u}$ in \underline{S}_k we get that

$$\begin{aligned} \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy &= \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{(\underline{w}_k(x) - \underline{w}_k(y))^2}{|x - y|^{n+2s}} dx dy \\ &- 2 \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{(\underline{w}_k(x) - \underline{w}_k(y))(\underline{u}(x) - \underline{u}(y))}{|x - y|^{n+2s}} dx dy \\ \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy &= \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{(\overline{w}_k(x) - \overline{w}_k(y))^2}{|x - y|^{n+2s}} dx dy \\ &+ 2 \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{(\overline{w}_k(x) - \overline{w}_k(y))(\bar{u}(x) - \bar{u}(y))}{|x - y|^{n+2s}} dx dy. \end{aligned}$$

Also similarly we obtain

$$\begin{aligned} \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy &= \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(\underline{w}_k(x) + \overline{w}_k(y))^2}{|x - y|^{n+2s}} dx dy \\ &- 2 \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(\underline{w}_k(x) + \overline{w}_k(y))(\underline{u}(x) - \bar{u}(y))}{|x - y|^{n+2s}} dx dy, \\ \int_{\underline{S}_k} \int_{S_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy &= \int_{\underline{S}_k} \int_{S_k} \frac{\underline{w}_k^2(x)}{|x - y|^{n+2s}} dx dy - 2 \int_{\underline{S}_k} \int_{S_k} \frac{\underline{w}_k(x)(\underline{u}(x) - u_k(y))}{|x - y|^{n+2s}} dx dy, \\ \int_{\overline{S}_k} \int_{S_k} \frac{h_k(x, y)}{|x - y|^{n+2s}} dx dy &= \int_{\overline{S}_k} \int_{S_k} \frac{\overline{w}_k^2(x)}{|x - y|^{n+2s}} dx dy + 2 \int_{\overline{S}_k} \int_{S_k} \frac{\overline{w}_k(x)(\bar{u}(x) - u_k(y))}{|x - y|^{n+2s}} dx dy. \end{aligned}$$

Since $\mathcal{C}\underline{S}_k = \overline{S}_k \cup S_k$, $\mathcal{C}\overline{S}_k = \underline{S}_k \cup S_k$ and

$$\begin{aligned} \|\underline{w}_k\|^2 &= \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{(\underline{w}_k(x) - \underline{w}_k(y))^2}{|x - y|^{n+2s}} dx dy + 2 \int_{\underline{S}_k} \int_{\mathcal{C}\underline{S}_k} \frac{\underline{w}_k^2(x)}{|x - y|^{n+2s}} dx dy \\ \|\overline{w}_k\|^2 &= \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{(\overline{w}_k(x) - \overline{w}_k(y))^2}{|x - y|^{n+2s}} dx dy + 2 \int_{\overline{S}_k} \int_{\mathcal{C}\overline{S}_k} \frac{\overline{w}_k^2(x)}{|x - y|^{n+2s}} dx dy, \end{aligned}$$

using all above estimates, we can express J_0 as

$$\begin{aligned}
J_0 &= C_s^n (\|\underline{w}_k\|^2 + \|\overline{w}_k\|^2) + 2 \left(\int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(w_k(x) + \overline{w}_k(y))^2}{|x-y|^{n+2s}} dx dy - \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{w_k^2(x)}{|x-y|^{n+2s}} \right. \\
&\quad \left. - \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k^2(x)}{|x-y|^{n+2s}} dx dy \right) - 2 \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{(w_k(x) - w_k(y))(u(x) - u(y))}{|x-y|^{n+2s}} dx dy \\
&\quad + 2 \int_{\overline{S}_k} \int_{\overline{S}_k} \frac{(\overline{w}_k(x) - \overline{w}_k(y))(\overline{u}(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy - 4 \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(w_k(x) + \overline{w}_k(y))(u(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy \\
&\quad - 4 \int_{\underline{S}_k} \int_{\underline{S}_k} \frac{w_k(x)(u(x) - u_k(y))}{|x-y|^{n+2s}} dx dy + 4 \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k(x)(\overline{u}(x) - u_k(y))}{|x-y|^{n+2s}} dx dy.
\end{aligned}$$

Now we notice that if $(x, y) \in \underline{S}_k \times \underline{S}_k$ then $(u(x) - u_k(y)) \leq (u(x) - u(y))$, if $(x, y) \in \overline{S}_k \times \underline{S}_k$ then $(\overline{u}(x) - u_k(y)) \geq (\overline{u}(x) - \overline{u}(y))$ and

$$\begin{aligned}
&\int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(w_k(x) + \overline{w}_k(y))^2}{|x-y|^{n+2s}} dx dy - \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{w_k^2(x)}{|x-y|^{n+2s}} dx dy - \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k^2(x)}{|x-y|^{n+2s}} dx dy \\
&= 2 \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k(x)w_k(y)}{|x-y|^{n+2s}} dx dy.
\end{aligned}$$

Also using change of variables, we have

$$\begin{aligned}
&\int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(w_k(x) + \overline{w}_k(y))(u(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy \\
&= \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{w_k(x)(u(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy - \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k(x)(\overline{u}(x) - u(y))}{|x-y|^{n+2s}} dx dy.
\end{aligned}$$

Therefore altogether we obtain

$$\begin{aligned}
J_0 &\geq C_s^n (\|\underline{w}_k\|^2 + \|\overline{w}_k\|^2) + 4 \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k(x)w_k(y)}{|x-y|^{n+2s}} dx dy + 2 \int_{\mathbb{R}^n} \overline{w}_k(-\Delta)^s \overline{u} dx - 2 \int_{\mathbb{R}^n} w_k(-\Delta)^s u dx \\
&\quad - 4 \int_{\overline{S}_k} \int_{\underline{S}_k} \frac{\overline{w}_k(x)(\overline{u}(x) - u(y))}{|x-y|^{n+2s}} dx dy + 4 \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{w_k(x)(u(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy \\
&\quad - 4 \int_{\underline{S}_k} \int_{\overline{S}_k} \frac{(w_k(x) + \overline{w}_k(y))(u(x) - \overline{u}(y))}{|x-y|^{n+2s}} dx dy \\
&\geq C_s^n (\|\underline{w}_k\|^2 + \|\overline{w}_k\|^2) + 2 \int_{\mathbb{R}^n} \overline{w}_k(-\Delta)^s \overline{u} dx - 2 \int_{\mathbb{R}^n} w_k(-\Delta)^s u dx
\end{aligned}$$

where we used the fact that if $(x, y) \in \overline{S}_k \times \underline{S}_k$ then $\overline{w}_k(x)w_k(y) \geq 0$. Now recalling that u and \overline{u} forms sub and supersolution of (P_λ) respectively, inserting the above inequality in (3.5)

we obtain

$$\begin{aligned}
I_\lambda(u_k) &\geq I_\lambda(v_k) + \frac{C_s^n \|w_k\|^2}{2} + \frac{C_s^n \|\overline{w_k}\|^2}{2} + \int_{\underline{S_k}} \left(\frac{\overline{u}^{1-q} - (\overline{u} + \overline{w_k})^{1-q}}{1-q} + \overline{u}^{-q} \overline{w_k} \right) dx \\
&\quad + \int_{\underline{S_k}} \left(\frac{\underline{u}^{1-q} - (\underline{u} - \underline{w_k})^{1-q}}{1-q} - \underline{u}^{-q} \underline{w_k} \right) dx + \lambda \int_{\underline{S_k}} \left(\frac{\overline{u}^{2_s^*} - (\overline{u} + \overline{w_k})^{2_s^*}}{2_s^*} + \overline{u}^{2_s^*-1} \overline{w_k} \right) dx \\
&\quad + \lambda \int_{\underline{S_k}} \left(\frac{\underline{u}^{2_s^*} - (\underline{u} - \underline{w_k})^{2_s^*}}{2_s^*} - \underline{u}^{2_s^*-1} \underline{w_k} \right) dx.
\end{aligned}$$

Now from Mean Value Theorem it follows that there exists $\theta \in (0, 1)$ (where θ may change its value for different function below) such that

$$\begin{aligned}
I_\lambda(u_k) &\geq I_\lambda(v_k) + \frac{C_s^n \|w_k\|^2}{2} + \frac{C_s^n \|\overline{w_k}\|^2}{2} - \int_{\underline{S_k}} ((\overline{u} + \theta \overline{w_k})^{-q} - \overline{u}^{-q}) \overline{w_k} dx \\
&\quad - \int_{\underline{S_k}} (\underline{u}^{-q} - (\underline{u} + \theta \underline{w_k})^{-q}) \underline{w_k} dx - \lambda \int_{\underline{S_k}} ((\overline{u} + \theta \overline{w_k})^{2_s^*-1} - \overline{u}^{2_s^*-1}) \overline{w_k} dx \\
&\quad - \lambda \int_{\underline{S_k}} (\underline{u}^{2_s^*-1} - (\underline{u} + \theta \underline{w_k})^{2_s^*-1}) \underline{w_k} dx \\
&\geq I_\lambda(v_k) + \frac{C_s^n \|w_k\|^2}{2} + \lambda \int_{\underline{S_k}} ((\overline{u} + \theta \overline{w_k})^{2_s^*-1} - \overline{u}^{2_s^*-1}) \overline{w_k} dx \\
&\quad - \lambda \int_{\underline{S_k}} (\underline{u}^{2_s^*-1} - (\underline{u} + \theta \underline{w_k})^{2_s^*-1}) \underline{w_k} dx.
\end{aligned} \tag{3.6}$$

Now since $2_s^* > 2$, there exists constant $C > 0$ such that (3.6) reduces to

$$\begin{aligned}
I_\lambda(u_k) &\geq I_\lambda(v_k) + \frac{C_s^n \|w_k\|^2}{2} + \frac{C_s^n \|\overline{w_k}\|^2}{2} - C \int_{\underline{S_k}} (\underline{u}^{2_s^*-2} - \underline{w_k}^{2_s^*-2}) \underline{w_k}^2 dx \\
&\quad - C \int_{\underline{S_k}} (\overline{u}^{2_s^*-2} - \overline{w_k}^{2_s^*-2}) \overline{w_k}^2 dx \\
&\geq I_\lambda(v_k) + \frac{C_s^n \|w_k\|^2}{2} + \frac{C_s^n \|\overline{w_k}\|^2}{2} - C \left(\int_{\underline{S_k}} |\underline{u}|^{2_s^*} \right)^{\frac{2_s^*-2}{2_s^*}} \|w_k\|^2 \\
&\quad - C \left(\int_{\underline{S_k}} |\overline{u}|^{2_s^*} \right)^{\frac{2_s^*-2}{2_s^*}} \|\overline{w_k}\|^2.
\end{aligned} \tag{3.7}$$

Claim- $\lim_{k \rightarrow \infty} |\overline{S_k}| = 0$ and $\lim_{k \rightarrow \infty} |\underline{S_k}| = 0$.

Let $\alpha > 0$ and define

$$\begin{aligned}
A_k &= \{x \in \Omega : u_k \geq \overline{u} \text{ and } \overline{u} > w + \alpha\}, \hat{A}_k = \{x \in \Omega : u_k \leq \underline{u} \text{ and } \underline{u} < w - \alpha\} \\
B_k &= \{x \in \Omega : u_k \geq \overline{u} \text{ and } \overline{u} \leq w + \alpha\}, \hat{B}_k = \{x \in \Omega : u_k \leq \underline{u} \text{ and } \underline{u} \geq w - \alpha\}.
\end{aligned}$$

Since

$$0 = \mathcal{L}(\{x \in \Omega : \overline{u} < w\}) = \mathcal{L}(\cap_{j=1}^{\infty} \{x \in \Omega : \overline{u} < w + \frac{1}{j}\})$$

so there exists $j_0 \geq 1$ large enough and $\alpha < 1/j_0$ such that $\mathcal{L}(\{x \in \Omega : \bar{u} < w + \alpha\}) \leq \epsilon/2$. This implies that $\mathcal{L}(B_k) \leq \epsilon/2$ and similarly, we obtain $\mathcal{L}(\hat{B}_k) \leq \epsilon/2$. From (3.4) we already have $|u_k - w|_2 \rightarrow 0$ as $k \rightarrow \infty$. So for $k \geq k_0$ large enough we get that

$$\frac{\alpha^2 \epsilon}{2} \geq \int_{\Omega} |u_k - w|^2 dx \geq \int_{A_k} |u_k - w|^2 dx \geq \alpha^2 \mathcal{L}(A_k)$$

which implies that $\mathcal{L}(A_k) \leq \frac{\epsilon}{2}$ for $k \geq k_0$. Similarly we obtain $\mathcal{L}(\hat{A}_k) \leq \frac{\epsilon}{2}$ for $k \geq k_0$. Now since $\overline{S_k} \subset A_k \cap B_k$ and $\underline{S_k} \subset \hat{A}_k \cap \hat{B}_k$ we get that $\mathcal{L}(\overline{S_k}) \leq \epsilon$ and $\mathcal{L}(\underline{S_k}) \leq \epsilon$ for $k \geq k_0$. This proves the claim. Thus

$$\left(\int_{\overline{S_k}} |\bar{u}|^{2_s^*} \right)^{\frac{2_s^* - 2}{2_s^*}} \leq o(1) \text{ and } \left(\int_{\underline{S_k}} |\underline{u}|^{2_s^*} \right)^{\frac{2_s^* - 2}{2_s^*}} \leq o(1)$$

which imposing in (3.7) gives that for large enough k

$$I_{\lambda}(u_k) \geq I_{\lambda}(v_k) \geq I_{\lambda}(w)$$

which is a contradiction to (3.4). Therefore w must be a local minimum of I_{λ} over X_0 . \square

Theorem 3.4 *There exists a positive weak solution to (P_{Λ}) .*

Proof. Let $\lambda_m \uparrow \Lambda$ as $m \rightarrow \infty$ and $\{u_{\lambda_m}\}$ be a sequence of positive weak solutions to (P_{λ_m}) , such that u_{λ_m} forms the local minimum of I_{λ_m} as seen in Lemma 3.3. Since we consider the minimal solutions, we get $u_m \leq u_{m+1}$ for each m . Then, it is easy to see that $I_{\lambda_m} < 0$ in the case $0 < q < 1$ whereas there exists a constant K independent of m such that $I_{\lambda_m} \leq K$ for all m when $q > 1$ but $q(2s - 1) < (2s + 1)$. This implies that $\{u_{\lambda_m}\}$ is uniformly bounded in X_0 . Therefore, up to a subsequence there exists $u_{\Lambda} \in X_0$ such that $u_{\lambda_m} \rightharpoonup u_{\Lambda}$ weakly and pointwise a.e. in X_0 as $m \rightarrow \infty$. Also by construction $u_{\lambda_m} \geq \underline{u}_{\lambda_1}$ as defined in Lemma 3.1. Therefore, u_{Λ} is a positive weak solution of (P_{Λ}) . \square

4 Multiplicity result

We have already obtained the first solution for (P_{λ}) in the previous section when $\lambda \in (0, \Lambda)$ in X_0 -topology. We fix $\lambda \in (0, \Lambda)$ and let w denotes the first weak solution of (P_{λ}) obtained in Proposition 3.2. In this section, we prove the existence of second solution of (P_{λ}) using the machinery of mountain pass Lemma and with the help of Ekeland variational principle. Precisely, we extend the approach used in [13] to the non local setting and for $q \geq 1$. This can be done by using the asymptotic boundary behavior of $w \in C_{\phi_q}^+(\Omega)$ and the Hardy's inequality to control the singular nonlinearity in the cone T defined below:

$$T \stackrel{\text{def}}{=} \{x \in X_0 : u \geq w \text{ a.e. in } \Omega\}$$

and since w forms a local minimizer of I_{λ} we get that $I_{\lambda}(u) \geq I_{\lambda}(w)$ whenever $\|u - w\| \leq \sigma_0$, for some constant $\sigma_0 > 0$. Then one of the following cases holds

(ZA) (Zero Altitude) $\inf\{I_\lambda(u) \mid u \in T, \|u - w\| = \sigma\} = I_\lambda(w)$ for all $\sigma \in (0, \sigma_0)$.

(MP) (Mountain Pass) There exists a $\sigma_1 \in (0, \sigma_0)$ such that $\inf\{I_\lambda(u) \mid u \in T, \|u - w\| = \sigma_1\} > I_\lambda(w)$.

Before investigating the two distinguished cases (ZA) and (MP), we prove the following regularity result for weak solutions to (P_λ) :

Proposition 4.1 *Any weak solution to (P_λ) for $\lambda \in (0, \Lambda]$ belongs to $L^\infty(\Omega) \cap C_{\phi_q}^+(\Omega)$.*

Proof. Let $u \in X_0$ denotes a weak solution to (P_λ) . We know that $\underline{u}_\lambda \in X_0 \cap C_{\phi_q}^+(\Omega)$ (defined in Lemma 3.1) forms a subsolution to (P_λ) satisfying $(-\Delta)^s \underline{u}_\lambda = \underline{u}_\lambda^{-q}$ in Ω . We first have

Claim : $\underline{u}_\lambda \leq u$ a.e. in Ω .

Let us prove the above claim. Suppose it is not true. First it is easy to see that for any $v \in X_0$ it holds

$$(v(x) - v(y))(v^+(x) - v^-(y)) \geq |v^+(x) - v^+(y)|^2, \text{ for any } x, y \in \mathbb{R}^n.$$

Therefore using $(\underline{u}_\lambda - u)^+$ as the test function in

$$(-\Delta)^s(\underline{u}_\lambda - u) \leq \underline{u}_\lambda^{-q} - u^{-q} \text{ in } \Omega$$

we get

$$\begin{aligned} 0 &\leq C_s^n \int_Q \frac{|(\underline{u}_\lambda - u)^+(x) - (\underline{u}_\lambda - u)^+(y)|^2}{|x - y|^{n+2s}} dx dy \\ &\leq C_s^n \int_Q \frac{((\underline{u}_\lambda - u)^+(x) - (\underline{u}_\lambda - u)^+(y))((\underline{u}_\lambda - u)(x) - (\underline{u}_\lambda - u)(y))}{|x - y|^{n+2s}} dx dy \\ &\leq \int_\Omega (\underline{u}_\lambda^{-q} - u^{-q})(\underline{u}_\lambda - u)^+ dx \leq 0. \end{aligned}$$

Hence it must be that $\text{meas}\{x \in \Omega : \underline{u}_\lambda(x) \geq u(x)\} = 0$ which gives a contradiction. Therefore $\underline{u}_\lambda \leq u$ a.e. in Ω . Let us now prove that $u \in L^\infty(\Omega)$. We follow the approach in [4, Proposition 2.2]. By virtue of the above claim and Hardy's inequality, we know that $\int_\Omega u^{-q} \phi dx < \infty$ for any $\phi \in X_0$. We aim to show that $(u - 1)^+$ belongs to $L^\infty(\Omega)$ which will imply that $u \in L^\infty(\Omega)$. If $f(t) = (t - 1)^+$ for $t \in \mathbb{R}$ and $\psi(t) \in C^\infty(\mathbb{R})$ be a convex and increasing function such that $\psi'(t) \leq 1$ when $t \in [0, 1]$ and $\psi'(t) = 1$ when $t \geq 1$ then we can define

$$\psi_\epsilon(t) = \epsilon \psi(t/\epsilon)$$

so that $\psi_\epsilon \rightarrow f$ uniformly as $\epsilon \rightarrow 0$. Also since ψ_ϵ is smooth, by regularity results and the uniform convergence of ψ_ϵ to f we get that

$$(-\Delta)^s \psi_\epsilon(u) \rightarrow (-\Delta)^s (u - 1)^+ \text{ as } \epsilon \rightarrow 0.$$

Moreover because ψ_ϵ is convex and differentiable, we know that

$$(-\Delta)^s \psi_\epsilon(u) \leq \psi'_\epsilon(u) (-\Delta)^s u \leq \chi_{\{u > 1\}} (-\Delta)^s u$$

where $\chi_{\{u>1\}}$ denotes the characteristic function over the set $\{x \in \Omega : u(x) > 1\}$. Then passing on the limits $\epsilon \rightarrow 0$ in above equation, we obtain

$$(-\Delta)^s(u-1)^+ \leq \chi_{\{u>1\}}(-\Delta)^s u \leq \chi_{\{u>1\}}(u^{-q} + \lambda u^{2_s^*-1}) \leq C(1 + ((u-1)^+)^{2_s^*-1})$$

for some constant $C > 0$. Therefore using [4, Proposition 2.2], we conclude that $(u-1)^+ \in L^\infty(\Omega)$. Finally we show that $u \in C_{\phi_q}^+(\Omega)$. Let z_λ be the unique solution (refer to [1, Theorem 1.1] with $\delta = q$ and $\beta = 0$) to

$$(-\Delta)^s z_\lambda = z_\lambda^{-q} + \lambda c, \quad u > 0 \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega,$$

with $c = \|u\|_\infty^{2_s^*-1}$ then similarly we can prove that $u \leq z_\lambda$. Therefore using local regularity results in [17, Propositions 2.2 and 2.3] derived from [20, Propositions 2.8 and 2.9], $u \in C_{\phi_q}^+(\Omega)$. This completes the proof. \square

Lemma 4.2 *Let (ZA) holds then for any $\sigma \in (0, \sigma_0)$ there exists a solution $v \in C_\phi^+(\Omega)$ to (P_λ) such that $0 < w < v$ in Ω and $\|v - w\| = \sigma$.*

Proof. We follow the proof of Lemma 2.6 of [13] in a nonlocal framework. We fix $\sigma \in (0, \sigma_0)$ and $r > 0$ such that $\sigma - r > 0$ and $\sigma + r < \sigma_0$. Let us define the set

$$W = \{u \in T \mid 0 < \sigma - r \leq \|u - w\| \leq \sigma + r\}$$

which is closed in X_0 and by (ZA), $\inf_{u \in W} I_\lambda(u) = I_\lambda(w)$. So using Ekeland variational principle, for any minimizing sequence $\{u_k\} \subset X_0$ satisfying $\|u_k\| = \sigma$ and $I_\lambda(u_k) \leq I_\lambda(w) + \frac{1}{k}$, we get another sequence $\{v_k\} \subset W$ such that

$$\begin{cases} I_\lambda(v_k) \leq I_\lambda(u_k) \leq I_\lambda(w) + \frac{1}{k} \\ \|u_k - v_k\| \leq \frac{1}{k} \\ I_\lambda(v_k) \leq I_\lambda(z) + \frac{1}{k}\|z - v_k\|, \text{ for all } z \in W. \end{cases} \quad (4.1)$$

We can choose $\epsilon > 0$ small enough so that $v_k + \epsilon(z - v_k) \in W$ for $z \in T$. So from (4.1) we obtain

$$\frac{I_\lambda(v_k + \epsilon(z - v_k)) - I_\lambda(v_k)}{\epsilon} \geq -\frac{1}{k}\|z - v_k\|.$$

Letting $\epsilon \rightarrow 0^+$ and using the fact that $v_k \geq w$ for each k , for $z \in T$ we get

$$\int_{\mathbb{R}^n} (-\Delta)^s v_k(z - v_k) - \int_{\Omega} v_k^{-q}(z - v_k) dx - \lambda \int_{\Omega} v_k^{2_s^*-1}(z - v_k) dx \geq -\frac{1}{k}\|z - v_k\|. \quad (4.2)$$

Now since $\{v_k\}$ forms a bounded sequence in X_0 , we get that there exists a $v \in X_0$ such that, up to a subsequence, $v_k \rightharpoonup v$ weakly in X_0 and pointwise a.e. in Ω as $k \rightarrow \infty$. Since $v_k \geq w$ for each k , we get $v \geq w$ a.e. in Ω . In what follows, we will prove that v is a weak

solution of (P_λ) . For $\phi \in X_0$ and $\epsilon > 0$, we set $\phi_{k,\epsilon} = (v_k + \epsilon\phi - w)^- \in X_0$ which implies that $(v_k + \epsilon\phi + \phi_{k,\epsilon}) \in T$. Putting $z = v_k + \epsilon\phi + \phi_{k,\epsilon}$ in (4.2) we get

$$\begin{aligned} C_s^n \int_Q \frac{(v_k(x) - v_k(y))((\epsilon\phi + \phi_{k,\epsilon})(x) - (\epsilon\phi + \phi_{k,\epsilon})(y))}{|x - y|^{n+2s}} dx dy - \int_\Omega v_k^{-q} (\epsilon\phi + \phi_{k,\epsilon}) dx \\ - \lambda \int_\Omega v_k^{2_s^*-1} (\epsilon\phi + \phi_{k,\epsilon}) dx \geq \frac{-1}{k} \|(\epsilon\phi + \phi_{k,\epsilon})\|. \end{aligned} \quad (4.3)$$

We define the sets $\Omega_{k,\epsilon} = \text{Supp } \phi_{k,\epsilon}$, $\Omega_\epsilon = \text{Supp } \phi_\epsilon$ and $\Omega_0 = \{x \in \Omega : v(x) = w(x)\}$. Then we get that $\mathcal{L}(\Omega_\epsilon \setminus \Omega_0) \rightarrow 0$ as $\epsilon \rightarrow 0$ and $\mathcal{L}(\Omega_{k,\epsilon} \setminus \Omega_\epsilon) + \mathcal{L}(\Omega_\epsilon \setminus \Omega_{k,\epsilon}) \rightarrow 0$ as $k \rightarrow \infty$. Also since $|\phi_{k,\epsilon}| \leq w + \epsilon|\phi|$, using Lebesgue Dominated convergence theorem we get $\phi_{k,\epsilon} \rightarrow \phi_\epsilon = (v + \epsilon\phi - w)^-$ in $L^m(\Omega)$ for all $m \in [1, 2_s^*]$. Moreover $\phi_{k,\epsilon} \rightharpoonup \phi_\epsilon$ weakly in X_0 and pointwise a.e. in Ω as $k \rightarrow \infty$. Now we estimate the following integral

$$\begin{aligned} \int_Q \frac{(v_k(x) - v_k(y))(\phi_{k,\epsilon}(x) - \phi_{k,\epsilon}(y))}{|x - y|^{n+2s}} dx dy \\ = \int_Q \frac{(v_k(x) - v_k(y))(\phi_\epsilon(x) - \phi_\epsilon(y))}{|x - y|^{n+2s}} dx dy \\ + \int_Q \frac{(v_k(x) - v_k(y))((\phi_{k,\epsilon} - \phi_\epsilon)(x) - (\phi_{k,\epsilon} - \phi_\epsilon)(y))}{|x - y|^{n+2s}} dx dy := I_1 + I_2. \end{aligned} \quad (4.4)$$

We show that $I_2 \leq o_k(1)$ for which we split the integrals and estimate them separately. Let $H_k = \Omega_{k,\epsilon} \cap \Omega_\epsilon$ and $G_k = \Omega_{k,\epsilon} \setminus \Omega_\epsilon \cup \Omega_\epsilon \setminus \Omega_{k,\epsilon}$. Then

$$\begin{aligned} \int_\Omega \int_{\mathcal{C}\Omega} \frac{(v_k(x) - v_k(y))((\phi_{k,\epsilon} - \phi_\epsilon)(x) - (\phi_{k,\epsilon} - \phi_\epsilon)(y))}{|x - y|^{n+2s}} \\ \leq \int_{H_k} \int_{\mathcal{C}\Omega} \frac{v(x)(v - v_k)(x)}{|x - y|^{n+2s}} + \int_{G_k} \int_{\mathcal{C}\Omega} \frac{v_k(x)(\phi_{k,\epsilon} - \phi_\epsilon)(x)}{|x - y|^{n+2s}} \\ \leq \int_{H_k} \int_{\mathcal{C}\Omega} \frac{v(x)(v - v_k)(x)}{|x - y|^{n+2s}} + \int_{G_k} \int_{\mathcal{C}\Omega} \frac{v_k(x)\phi_{k,\epsilon}(x)}{|x - y|^{n+2s}} \\ = \int_{H_k} \int_{\mathcal{C}\Omega} \frac{v(x)(v - v_k)(x)}{|x - y|^{n+2s}} + o_k(1) \end{aligned} \quad (4.5)$$

using the fact that $\mathcal{L}(\Omega_{k,\epsilon} \setminus \Omega_\epsilon) + \mathcal{L}(\Omega_\epsilon \setminus \Omega_{k,\epsilon}) \rightarrow 0$ as $k \rightarrow \infty$ and Lebesgue Dominated convergence theorem. Similarly

$$\begin{aligned} \int_\Omega \int_\Omega \frac{(v_k(x) - v_k(y))((\phi_{k,\epsilon} - \phi_\epsilon)(x) - (\phi_{k,\epsilon} - \phi_\epsilon)(y))}{|x - y|^{n+2s}} \\ \leq \int_{H_k} \int_{H_k} \frac{(v(x) - v(y))((v - v_k)(x) - (v - v_k)(y))}{|x - y|^{n+2s}} \\ + 2 \int_{H_k} \int_{G_k} \frac{(v_k(x) - v_k(y))((\phi_{k,\epsilon} - \phi_\epsilon)(x) - (\phi_{k,\epsilon} - \phi_\epsilon)(y))}{|x - y|^{n+2s}} \\ + \int_{G_k} \int_{G_k} \frac{(v_k(x) - v_k(y))((\phi_{k,\epsilon} - \phi_\epsilon)(x) - (\phi_{k,\epsilon} - \phi_\epsilon)(y))}{|x - y|^{n+2s}} \\ \leq \int_{H_k} \int_{H_k} \frac{(v(x) - v(y))((v - v_k)(x) - (v - v_k)(y))}{|x - y|^{n+2s}} + o_k(1) \end{aligned} \quad (4.6)$$

using again the Lebesgue Dominated convergence theorem with the fact that $v_k - v \rightarrow 0$ and $\phi_{k,\epsilon} - \phi_\epsilon \rightarrow 0$ pointwise as $k \rightarrow \infty$. Combining (4.5) and (4.6) we obtain that

$$I_2 \leq \int_{H_k} \int_{H_k \cup C\Omega} \frac{(v(x) - v(y))((v - v_k)(x) - (v - v_k)(y))}{|x - y|^{n+2s}} + o_k(1) = o_k(1).$$

Therefore using this in (4.4), we obtain

$$\int_Q \frac{(v_k(x) - v_k(y))(\phi_{k,\epsilon}(x) - \phi_{k,\epsilon}(y))}{|x - y|^{n+2s}} dx dy \leq \int_Q \frac{(v_k(x) - v_k(y))(\phi_\epsilon(x) - \phi_\epsilon(y))}{|x - y|^{n+2s}} dx dy + o_k(1).$$

Moreover, we have that $|v_k^{-q}(\epsilon\phi + \phi_{k,\epsilon})| \leq w^{-q}(w + 2\epsilon\phi) \in L^1(\Omega)$ using the Hardy's inequality. Thus using Lebesgue Dominated convergence theorem and passing on the limits $k \rightarrow \infty$ in (4.3) we get

$$0 \leq C_s^n \int_Q \frac{(v_k(x) - v_k(y))((\epsilon\phi + \phi_\epsilon)(x) - (\epsilon\phi + \phi_\epsilon)(y))}{|x - y|^{n+2s}} dx dy - \int_\Omega (v^{-q} + \lambda v^{2_s^*-1})(\epsilon\phi + \phi_\epsilon) dx.$$

Using the fact that w is a weak solution of (P_λ) and $v \geq w$, the above inequality implies that

$$\begin{aligned} & C_s^n \int_Q \frac{(v(x) - v(y))(\phi(x) - \phi(y))}{|x - y|^{n+2s}} dx dy - \int_\Omega v^{-q} \phi dx - \lambda \int_\Omega v^{2_s^*-1} \phi dx \\ & \geq -\frac{1}{\epsilon} \left(C_s^n \int_Q \frac{(v(x) - v(y))(\phi_\epsilon(x) - \phi_\epsilon(y))}{|x - y|^{n+2s}} dx dy - \int_\Omega v^{-q} \phi_\epsilon dx - \lambda \int_\Omega v^{2_s^*-1} \phi_\epsilon dx \right) \\ & \geq \frac{1}{\epsilon} \left(C_s^n \int_Q \frac{((w - v)(x) - (w - v)(y))(\phi_\epsilon(x) - \phi_\epsilon(y))}{|x - y|^{n+2s}} dx dy + \int_\Omega (v^{-q} - w^{-q}) \phi_\epsilon dx \right) \\ & \geq C_s^n \int_{\Omega_\epsilon} \int_{\Omega_\epsilon} \frac{((v - w)(x) - (v - w)(y))(\phi(x) - \phi(y))}{|x - y|^{n+2s}} dx dy \\ & \quad + 2C_s^n \int_{\Omega_\epsilon} \int_{\{w \leq v + \epsilon\phi\}} \frac{((v - w)(x) - (v - w)(y))\phi(x)}{|x - y|^{n+2s}} dx dy \\ & \quad + 2C_s^n \int_{\Omega_\epsilon} \int_{C\Omega} \frac{(v - w)(x)\phi(x)}{|x - y|^{n+2s}} dx dy + \int_{\Omega_\epsilon} (v^{-q} - w^{-q}) \phi dx \\ & = o(1) \text{ as } \epsilon \rightarrow 0^+ \end{aligned}$$

using the fact that $|\Omega_\epsilon \setminus \Omega_0| \rightarrow 0$ as $\epsilon \rightarrow 0^+$. From this, we get that

$$C_s^n \int_Q \frac{(v(x) - v(y))(\phi(x) - \phi(y))}{|x - y|^{n+2s}} dx dy - \int_\Omega v^{-q} \phi dx - \lambda \int_\Omega v^{2_s^*-1} \phi dx = 0 \text{ for all } \phi \in X_0.$$

Claim- The sequence $v_k \rightarrow v$ strongly in X_0 as $k \rightarrow \infty$.

From Brezis Lieb lemma we have

$$\begin{aligned} & \|v_k\|^2 - \|v_k - v\|^2 = \|v\|^2 + o(1) \\ & \int_\Omega |v_k|^{2_s^*} dx - \int_\Omega |v_k - v|^{2_s^*} dx = \int_\Omega |v|^{2_s^*} dx + o(1). \end{aligned}$$

Since $v_k, v \geq w$ a.e. in Ω , we get

$$\int_{\Omega} |v_k|^{1-q} dx - \int_{\Omega} |v|^{1-q} dx = \int_{\Omega} (v_k + \theta v)^{-q} (v_k - v) dx, \text{ for } \theta \in [0, 1].$$

We know that $(v_k + \theta v)^{-q} (v_k - v) \rightarrow 0$ pointwise a.e. in Ω and $v_k, v \geq w \in C_{\phi}^+(\Omega)$. Therefore for any $E \subset \Omega$, we have

$$\int_{\Omega} (v_k + \theta v)^{-q} (v_k - v) dx \leq C \|\delta^{\frac{(1-q)s}{1+q}}(x)\|_{L^2(E)} \|v_k - v\|, \text{ using Hardy's inequality.} \quad (4.7)$$

Since $q(2s-1) < (2s+1)$, for any $\epsilon > 0$, there exists a $\rho > 0$ such that $\|\delta^{\frac{(1-q)s}{1+q}}(x)\|_{L^2(E)} < \epsilon$ whenever $\mathcal{L}(E) < \rho$. Hence from (4.7) and Vitali's convergence theorem we obtain

$$\int_{\Omega} (v_k + \theta v)^{-q} (v_k - v) dx \rightarrow 0 \text{ as } k \rightarrow \infty$$

that is

$$\int_{\Omega} |v_k|^{1-q} dx \rightarrow \int_{\Omega} |v|^{1-q} dx \text{ as } k \rightarrow \infty.$$

Taking v as the testing function in (4.2), we deduce

$$C_s^n \|v_k - v\|^2 \leq \lambda \|v_k - v\|_{L^{2_s^*}(\Omega)}^{2_s^*} + o_k(1). \quad (4.8)$$

In the other hand, taking $z = 2v_k$ in (4.2) we infer

$$C_s^n \|v_k\|^2 - \int_{\Omega} v_k^{1-q} dx - \lambda \|v_k\|_{L^{2_s^*}(\Omega)}^{2_s^*} \geq o_k(1). \quad (4.9)$$

Since v is a weak solution, we have that

$$C_s^n \|v\|^2 - \int_{\Omega} v^{1-q} dx - \lambda \|v\|_{L^{2_s^*}(\Omega)}^{2_s^*} = 0. \quad (4.10)$$

From (4.9) and (4.10),

$$C_s^n \|v_k - v\|^2 \geq \lambda \|v_k - v\|_{L^{2_s^*}(\Omega)}^{2_s^*} + o_k(1). \quad (4.11)$$

From (4.8) and (4.11), we have that

$$C_s^n \|v_k - v\|^2 = \lambda \|v_k - v\|_{L^{2_s^*}(\Omega)}^{2_s^*} + o_k(1). \quad (4.12)$$

Without loss of generality, we can assume that $I_{\lambda}(w) \leq I_{\lambda}(v)$. Then, we easily get

$$I_{\lambda}(v_k) - I_{\lambda}(v) \leq I_{\lambda}(w) - I_{\lambda}(v) + o_k(1) \leq o_k(1)$$

from which it follows that

$$\frac{C_s^n}{2} \|v_k - v\|^2 - \frac{\lambda}{2_s^*} \|v_k - v\|_{L^{2_s^*}(\Omega)}^{2_s^*} \leq o_k(1). \quad (4.13)$$

From (4.12) and (4.13), we infer that $v_k \rightarrow v$ strongly in X_0 . This proves the claim. Since $v_k \in W$ we conclude that $v \in W$ and $v \not\equiv w$. Next we prove that $w < v$ in Ω . For that, we first observe that from Proposition 4.1 $w, v \in L^\infty(\Omega) \cap C_{\phi_q}^+(\Omega)$. Now suppose that there exists $x_0 \in \Omega$ such that $v(x_0) = w(x_0)$. Then, since $v \geq w$, $v, w \in C(\mathbb{R}^n)$ and $v \not\equiv w$, we get

$$0 > C_s^n \int_{\mathbb{R}^n} \frac{(v-w)(x_0) - (v-w)(y)}{|x_0 - y|^{n+2s}} = v^{-q}(x_0) + \lambda v^{2s^*-1}(x_0) - (w^{-q}(x_0) + \lambda w^{2s^*-1}(x_0)) = 0$$

from which we get a contradiction. Therefore $v > w$ in Ω . \square

We define

$$S_s = \inf_{u \in X_0 \setminus \{0\}} \frac{\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy}{\left(\int_\Omega |u|^{2s^*} dx\right)^{2/2s^*}}$$

as the best constant for the embedding $X_0 \hookrightarrow L^{2s^*}(\Omega)$. Consider the family of minimizers $\{U_\epsilon\}$ of S_s (refer [18]) defined as

$$U_\epsilon(x) = \epsilon^{-\frac{(n-2s)}{2}} u^* \left(\frac{x}{\epsilon} \right), \quad x \in \mathbb{R}^n$$

where $u^*(x) = \bar{u} \left(\frac{x}{S_s^{1/2s}} \right)$, $\bar{u}(x) = \frac{\tilde{u}(x)}{|u|_{2s^*}}$ and $\tilde{u}(x) = \alpha(\beta^2 + |x|^2)^{-\frac{n-2s}{2}}$ with $\alpha \in \mathbb{R} \setminus \{0\}$ and $\beta > 0$ are fixed constants. Then for each $\epsilon > 0$, U_ϵ satisfies

$$(-\Delta)^s u = |u|^{2s^*-2} u \quad \text{in } \mathbb{R}^n.$$

Let $\nu > 0$ be such that $B_{4\nu} \subset \Omega$ and let $\zeta \in C_c^\infty(\mathbb{R}^n)$ be such that $0 \leq \zeta \leq 1$ in \mathbb{R}^n , $\zeta \equiv 0$ in $\mathbb{R}^n \setminus B_{2\nu}$ and $\zeta \equiv 1$ in B_ν . For each $\epsilon > 0$ and $x \in \mathbb{R}^n$, we define $\Phi_\epsilon(x) := \zeta(x)U_\epsilon(x)$. From [12, Lemma 4.13], we have the following.

Lemma 4.3 $\sup\{I_\lambda(u + t\Phi_\epsilon) : t \geq 0\} < I_\lambda(u) + \frac{s(C_s^n S_s)^{\frac{n}{2s}}}{n\lambda^{\frac{n-2s}{2s}}}$, for any sufficiently small $\epsilon > 0$.

Now we prove the existence of second solution if (MP) holds.

Lemma 4.4 *Let (MP) holds then there exists a $v \in X_0 \cap C_\phi^+(\Omega)$, verifying $w < v$ in Ω , which solves (P_λ) weakly.*

Proof. From Lemma 4.3, it follows that there exists $\epsilon > 0$ and $R_0 \geq 1$ such that

$$(i) \quad I_\lambda(w + R U_\epsilon) < I_\lambda(w) \quad \text{for } \epsilon \in (0, \epsilon_0) \text{ and } R \geq R_0.$$

$$(ii) \quad I_\lambda(w + t R_0 U_\epsilon) < I_\lambda(w) + \frac{s(C_s^n S_s)^{\frac{n}{2s}}}{n\lambda^{\frac{n-2s}{2s}}} \quad \text{for } \epsilon \in (0, \epsilon_0) \text{ and } t \in [0, 1].$$

We define the complete metric space

$$\Gamma := \{\eta \in C([0, 1], T) : \eta(0) = w, \|\eta(1) - w\| > \sigma_1, I_\lambda(\eta(1)) < I_\lambda(w)\}$$

with metric defined as $d(\eta', \eta) = \max_{t \in [0,1]} \{\|\eta'(t) - \eta(t)\|\}$ for all $\eta, \eta' \in \Gamma$. From (i) above, we get that $\eta(t) = w + tR_0U_\epsilon \in \Gamma$ for large enough $R_0 > 0$. This gives that $\Gamma \neq \emptyset$. Let $\gamma_0 = \inf_{\eta \in \Gamma} \max_{t \in [0,1]} I_\lambda(\eta(t))$ then by virtue of (ii) above and condition (MP), we get

$$I_\lambda(w) < \gamma_0 \leq I_\lambda(w) + \frac{s(C_s^n S_s)^{\frac{n}{2s}}}{n\lambda^{\frac{n-2s}{2s}}}.$$

Now let $\Psi(\eta) = \max_{t \in [0,1]} I_\lambda(\eta(t))$ for $\eta \in \Gamma$. Then using Ekeland's variational principle, we get a sequence $\{\eta_k\} \subset \Gamma$ such that

$$\Psi(\eta_k) < \gamma_0 + \frac{1}{k} \text{ and } \Psi(\eta_k) < \Psi(\eta) + \frac{1}{k} \|\Psi(\eta) - \eta(\eta_k)\|_\Gamma, \forall \eta \in \Gamma. \quad (4.14)$$

We define

$$\Lambda_k = \{t \in [0, 1] : I_\lambda(\eta_k(t)) = \max_{x \in [0,1]} I_\lambda(\eta_k(x))\}$$

. **Claim:** There exists a $t_k \in \Lambda_k$ such that if $v_k = \eta_k(t_k)$ and $z \in T$ then

$$\int_{\mathbb{R}^n} (-\Delta)^s v_k (z - v_k) - \int_{\Omega} (v_k^{-q} + \lambda v_k^{2_s^*-1})(z - v_k) dx \geq -\frac{1}{k} \max\{1, \|z - v_k\|\}.$$

We prove it by contradiction, so assume that for every $t \in \Lambda_k$ there exists a $z_t \in T$ such that

$$\begin{aligned} \int_{\mathbb{R}^n} (-\Delta)^s \eta_k(t) \left(\frac{z_t - \eta_k(t)}{\max\{1, \|z_t - \eta_k(t)\|\}} \right) dx \\ - \int_{\Omega} ((\eta_k(t))^{-q} + \lambda(\eta_k(t))^{2_s^*-1}) \left(\frac{z_t - \eta_k(t)}{\max\{1, \|z_t - \eta_k(t)\|\}} \right) dx < -\frac{1}{k}. \end{aligned} \quad (4.15)$$

Since I_λ is locally Lipschitz in T , z_t can be chosen to be locally constant on Λ_t . Therefore for each $t \in \Lambda_k$ there exists a neighborhood N_t of t in $(0, 1)$ such that for each $r \in N_t \cap \Gamma_k$, (4.15) holds that is

$$\begin{aligned} \int_{\mathbb{R}^n} (-\Delta)^s \eta_k(r) \left(\frac{z_t - \eta_k(r)}{\max\{1, \|z_t - \eta_k(r)\|\}} \right) dx \\ - \int_{\Omega} ((\eta_k(r))^{-q} + \lambda(\eta_k(r))^{2_s^*-1}) \left(\frac{z_t - \eta_k(r)}{\max\{1, \|z_t - \eta_k(r)\|\}} \right) dx < -\frac{1}{k}. \end{aligned} \quad (4.16)$$

It is possible to choose a finite set $\{r_1, r_2, \dots, r_m\} \subset \Lambda_k$ such that $\Lambda_k \subset \cup_{i=1}^m J_{r_i}$. For notational convenience, we set $z_i = z_{r_i}$ and denote $\{\kappa_1, \kappa_2, \dots, \kappa_m\}$ as the partition of unity associated with covering $\{J_{r_1}, J_{r_2}, \dots, J_{r_m}\}$ of Λ_k . Now if we define $z(r) = \sum_{i=1}^m \kappa_i(r) z_i$ for $r \in [0, 1]$ then $z(r) \in T$ for each $r \in [0, 1]$. Therefore from (4.16) we deduce that for all $r \in [0, 1]$

$$\begin{aligned} \int_{\mathbb{R}^n} (-\Delta)^s \eta_k(r) \left(\frac{z(r) - \eta_k(r)}{\max\{1, \|z(r) - \eta_k(r)\|\}} \right) dx \\ - \int_{\Omega} ((\eta_k(r))^{-q} + \lambda(\eta_k(r))^{2_s^*-1}) \left(\frac{z(r) - \eta_k(r)}{\max\{1, \|z(r) - \eta_k(r)\|\}} \right) dx < -\frac{1}{k}. \end{aligned}$$

Let $h : [0, 1] \rightarrow [0, 1]$ be a continuous function such that $h(t) = 1$ in a neighborhood of Λ_k and $h(0) = h(1) = 0$. Also we set $\mu_k(t) = \max\{1, \|z(t) - \eta_k(t)\|\}$ and

$$\eta(t) = \eta_k(t) + \frac{h(t)\epsilon}{\mu_k(t)}(z(t) - \eta_k(t)).$$

For $\epsilon \in (0, 1)$, $\eta(t) \in T$ for all $t \in [0, 1]$. Hence (4.14) gives us that

$$\max_{t \in [0, 1]} I_\lambda(\eta_k(t)) \leq \max_{t \in [0, 1]} I_\lambda(\eta(t)) + \frac{\epsilon}{k} \max_{t \in [0, 1]} \left(h(t) \frac{\|z(t) - \eta_k(t)\|}{\mu_k(t)} \right). \quad (4.17)$$

If $t_{k,\epsilon} \in [0, 1]$ denotes the value such that $I_\lambda(\eta(t_{k,\epsilon})) = \max_{t \in [0, 1]} I_\lambda(\eta(t))$ then we can assume that $t_{k,\epsilon_j} \rightarrow t_k$ for some $t_k \in [0, 1]$, where ϵ_j is a sequence such that $\epsilon_j \rightarrow 0$. Using the continuity of η , we deduce that

$$\eta(t_{k,\epsilon_j}) \rightarrow \eta_k(t_k) \text{ as } \epsilon_j \rightarrow 0.$$

Hence from (4.17) we obtain that $\max_{t \in [0, 1]} I_\lambda(\eta_k(t)) \leq \max_{t \in [0, 1]} I_\lambda(\eta_k(t_k))$ which implies $I_\lambda(\eta_k(t_k)) = \max_{t \in [0, 1]} I_\lambda(\eta_k(t))$. So $t_k \in \Gamma_k$ and $h(t_{k,\epsilon_j}) = 1$ for $j > 0$ large enough, by definition.

If we set $v_k = \eta_k(t_k)$, $v_{k,j} = \eta_k(t_{k,\epsilon_j})$ and $\mu_{k,j} = \max\{1, \|z(t_{k,\epsilon_j}) - v_{k,j}\|\}$ then for large enough j we obtain

$$I_\lambda(v_{k,j}) \leq I_\lambda(v_k) \leq I_\lambda \left(v_{k,j} + \frac{\epsilon_j}{\mu_{k,j}}(z(t_{k,\epsilon_j}) - v_{k,j}) \right) + \frac{\epsilon_j}{k}. \quad (4.18)$$

It is easy to see that $\mu_{k,j} \rightarrow \theta_k := \max\{1, \|z(t_k) - v_k\|\}$ and $\|v_k - v_{k,j}\| \rightarrow 0$ as $j \rightarrow \infty$. Let $p_j = v_{k,j} - v_k$ and

$$k_j = p_j + \epsilon_j \left(\frac{z(t_{k,\epsilon_j}) - v_{k,j}}{\mu_{k,j}} - \frac{z(t_k) - v_k}{\theta_k} \right) = p_j + o(1).$$

Then from (4.18), we obtain

$$\frac{1}{\epsilon_j} \left(I_\lambda \left(v_k + \epsilon_j \left(\frac{z(t_k) - v_k}{\theta_k} \right) + k_j \right) + I_\lambda(v_k + p_j) \right) \geq -\frac{1}{k} \text{ as } j \rightarrow \infty.$$

But since $v_k + \epsilon_j \left(\frac{z(t_k) - v_k}{\theta_k} \right) \geq w$ using the fact that $z(t_k) \in T$, from Proposition 2.1 and the above inequality we get

$$\int_{\mathbb{R}^n} (-\Delta)^s v_k \left(\frac{z(t_k) - v_k}{\theta_k} \right) dx - \int_{\Omega} (v_k^{-q} + \lambda v_k^{2_s^* - 1}) \left(\frac{z(t_k) - v_k}{\theta_k} \right) dx \geq -\frac{1}{k}.$$

This is a contradiction to (4.15). Thus, the claim holds. So there exists a sequence $\{v_k\}$ satisfying

$$\begin{cases} \int_{\mathbb{R}^n} (-\Delta)^s v_k(z - v_k) - \int_{\Omega} (v_k^{-q} + \lambda v_k^{2_s^* - 1})(z - v_k) dx \geq -\frac{c}{k}(1 + \|z\|) \text{ for all } z \in T \\ I_\lambda(v_k) \rightarrow \gamma_0 \text{ as } k \rightarrow \infty \end{cases} \quad (4.19)$$

where $c > 0$ is some constant. Setting $z = 2v_k$ in (4.14) and using (4.19) we get

$$\gamma_0 + o(1) \geq \frac{sC_s^n}{n} \|v_k\|^2 - \frac{2_s^* - 1 + q}{2_s^*(1-q)} \int_{\Omega} |v_k|^{1-q} dx - \frac{c}{2_s^* k} (1 + 2\|v_k\|).$$

Now this implies that $\{v_k\}$ must be bounded in X_0 , thus up to a subsequence, $v_k \rightharpoonup v$ weakly in X_0 as $k \rightarrow \infty$. Using similar ideas as in (ZA) case, it can be shown that v is a weak solution of (P_λ) . Then the remaining part of the proof is similar as in [12, Proposition 4.12] (see also [13, Lemma 2.7] in the local setting) and consists of proving the strong convergence of the sequence $\{v_k\}$ to v . To this aim we use that the energy $I_\lambda(v_k)$ is strictly below the first critical level $I_\lambda(w) + \frac{s(C_s^n S_s)^{\frac{n}{2s}}}{n\lambda^{\frac{n-2s}{2s}}}$ which implies

$$\frac{C_s^n}{2} \|v_k - v\|^2 - \frac{\lambda}{2_s^*} \|v_k - v\|_{L^{2_s^*}(\Omega)}^{2_s^*} < \frac{s(C_s^n S_s)^{\frac{n}{2s}}}{n\lambda^{\frac{n-2s}{2s}}}. \quad (4.20)$$

Now (4.12) and (4.20) and the fact that $S_s \|v_k - v\|_{L^{2_s^*}(\Omega)}^2 \leq \|v_k - v\|^2$ force $\|v_k - v\| \rightarrow 0$ as $k \rightarrow \infty$. Thus, we infer that $I_\lambda(v) = \gamma_0$ and $v \neq w$ and the proof of $w < v$ in Ω can be performed as in the proof of lemma 4.2. \square

Proof of Theorem 1.2: The proof follows from Lemma 4.2, Lemma 4.4, Proposition 4.1 and Proposition 3.4 along with Proposition 3.2. \square

Proof of Theorem 1.3: The proof follows directly from Proposition 4.1 and [1, Theorem 1.2] with $\delta = q$ and $\beta = 0$. To see that the regularity result falls into the scope of [1, Theorem 1.2], note that u is a classical solution as defined in [1, Definition 1]. \square

Acknowledgements. The authors would like to thank warmly the anonymous referee for valuable comments which have improved the paper.

References

- [1] Adimurthi, J. Giacomoni and S. Santra, *Positive solutions to a fractional equation with singular nonlinearity*, J. Differential Equations, 265 (4) (2018), 1191-1226.
- [2] S. Alama, *Semilinear elliptic equations with sublinear indefinite nonlinearities*, Adv. Diff. Equations 4 (1999) 813–842.
- [3] D. Applebaum, *Lévy process-from probability to finance and quantum groups*, Notices Amer. Math. Soc., 51 (2004) 1336-1347.
- [4] B. Barrios, E. Colorado, R. Servadei and F. Soria, *A critical fractional equation with concave-convex power nonlinearities*, Ann. I. H. Poincaré, 32 (2015) 875-900.
- [5] B. Barrios, I. De Bonis, M. Medina and I. Peral, *Semilinear problems for the fractional laplacian with a singular nonlinearity*, Open Math., 13 (2015) 390–407.

- [6] B. Barrios, M. Medina and I. Peral, *Some remarks on the solvability of non-local elliptic problems with the Hardy potential*, Communications in Contemporary Mathematics, 16(4) (2014), 1350046 (29 pages).
- [7] C. Bucur and E. Valdinoci, *An Introduction to the Fractional Laplacian*, Nonlocal Diffusion and Applications, Lecture Notes of the Unione Matematica Italiana, 20 (2016), Springer, Cham.
- [8] Z. Cai, C. Chu and C. Lei, *Existence of positive solutions for a fractional elliptic problems with the Hardy-Sobolev-Maz'ya potential and critical nonlinearities*, Boundary Value Problems, (2017), DOI 10.1186/s13661-017-0912-8.
- [9] W. Chen, *Fractional elliptic problems with two critical sobolev-hardy exponents*, Electronic Journal of Differential Equations, 22 (2018), 1–12.
- [10] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math., 136(5) (2012), 521–573.
- [11] A. Fiscella and P. Pucci, *On certain nonlocal Hardy-Sobolev critical elliptic Dirichlet problems*, Adv. Differential Equations, 21(5/6) (2016), 571–599.
- [12] J. Giacomoni, T. Mukherjee and K. Sreenadh, *Positive solutions of fractional elliptic equation with critical and singular nonlinearity*, Advances in Nonlinear Analysis, 6(3) (2017), 327–354.
- [13] Y. Haitao, *Multiplicity and asymptotic behavior of positive solutions for a singular semi-linear elliptic problem*, J. Differential Equations, 189 (2003) 487–512.
- [14] T. Kato, *Fractional powers of dissipative operators*, J. Math. Soc. Japan, 13 (1961), 246–274.
- [15] T. Mukherjee and K. Sreenadh, *Critical growth fractional elliptic problem with singular nonlinearities*, Electronic Journal of differential equations, 54 (2016) 1–23.
- [16] N. Landkof, *Foundations of modern potential theory*, Die Grundlehren der mathematischen Wissenschaften, Band 180. Springer-Verlag, New York-Heidelberg, 1972.
- [17] X. Ros-Oton and J. Serra, *The Dirichlet problem for the fractional Laplacian: Regularity up to the boundary*, J. Math. Pures Appl., 101 (2014), 275–302.
- [18] R. Servadei and E. Valdinoci, *The Brezis-Nirenberg result for the fractional laplacian*, Trans. Amer. Math. Soc., 367 (2015) 67–102.
- [19] L. Silvestre, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Thesis (Ph.D.)-The University of Texas at Austin, 2005.

- [20] L. Silvestre, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Comm. Pure Appl. Math., 60 (1) (2007), 67–112.
- [21] E.M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [22] E.M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30 Princeton University Press, Princeton, N.J., 1970.
- [23] H. Triebel, *Interpolation theory, function spaces, differential operators*, John Ambrosius Barth, Heidelberg, second edition, 1995.