



HAL
open science

The colloidal stabilization of young red wine by Acacia senegal gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins

Michael Nigen, Rafael Apolinar Valiente, Nerea Iturmendi, Pascale Williams, Thierry Doco, Virginie Moine, Arnaud Massot, Isabelle Jaouen, Christian Sanchez

► To cite this version:

Michael Nigen, Rafael Apolinar Valiente, Nerea Iturmendi, Pascale Williams, Thierry Doco, et al.. The colloidal stabilization of young red wine by Acacia senegal gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins. *Food Hydrocolloids*, 2019, 97, 10.1016/j.foodhyd.2019.105176 . hal-02164154

HAL Id: hal-02164154

<https://hal.science/hal-02164154>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Distributed under a Creative Commons Attribution 4.0 International License

Accepted Manuscript

The colloidal stabilization of young red wine by *Acacia senegal* gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins

Michaël Nigen, Rafael Apolinar Valiente, Nerea Iturmendi, Pascale Williams, Thierry Doco, Virginie Moine, Arnaud Massot, Isabelle Jaouen, Christian Sanchez



PII: S0268-005X(19)30393-5

DOI: <https://doi.org/10.1016/j.foodhyd.2019.105176>

Article Number: 105176

Reference: FOOHYD 105176

To appear in: *Food Hydrocolloids*

Received Date: 18 February 2019

Revised Date: 21 May 2019



Accepted Date: 18 June 2019

Please cite this article as: Nigen, Michaël., Valiente, R.A., Iturmendi, N., Williams, P., Doco, T., Moine, V., Massot, A., Jaouen, I., Sanchez, C., The colloidal stabilization of young red wine by *Acacia senegal* gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins, *Food Hydrocolloids* (2019), doi: <https://doi.org/10.1016/j.foodhyd.2019.105176>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comment citer ce document :

Nigen, M., Apolinar Valiente, R., Iturmendi, N., Williams, P., Doco, T., Moine, Massot, Jaouen, I., Sanchez, C. (2019). The colloidal stabilization of young red wine by *Acacia senegal* gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins. *Food Hydrocolloids*. 97. . DOI : 10.1016/i.foodhyd.2019.105176



Graphical Abstract.

Comment citer ce document :

Nigen, M., Apolinar Valiente, R., Iturmendi, N., Williams, P., Doco, T., Moine, Massot, Jaouen, I., Sanchez, C. (2019). The colloidal stabilization of young red wine by Acacia senegal gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins. Food Hydrocolloids. 97. . DOI : 10.1016/j.foodhyd.2019.105176

1 The colloidal stabilization of young red wine by *Acacia*
2 *senegal* gum: The involvement of the protein backbone from
3 the protein-rich arabinogalactan-proteins

4
5
6 Michaël Nigen^{a*}, Rafael Apolinar Valiente^a, Nerea Iturmendi^c, Pascale Williams^b, Thierry
7 Doco^b, Virginie Moine^c, Arnaud Massot^c, Isabelle Jaouen^d and Christian Sanchez^a

8
9 ^a UMR1208 Ingénierie des Agropolymères et Technologies Emergentes, INRA-Montpellier
10 SupAgro-CIRAD-Université Montpellier, 2 Place Pierre Viala F-34060 Montpellier, France

11
12 ^b UMR 1083 Science Pour l'Œnologie, INRA-Montpellier SupAgro-Université Montpellier, 2
13 Place Pierre Viala F-34060 Montpellier, France

14
15 ^c BIOLAFFORT, 126 quai de la Souys, 33100 Bordeaux, France

16
17 ^d ALLAND & ROBERT, 9 rue Saintonge, 75003 Paris, France

18
19
20
21
22 * corresponding author. Tel.: 33 4 99 61 26 15. Fax: 33 4 99 61 30 76. E-mail:
23 michael.nigen@umontpellier.fr

24

25 **Abstract**

26 The arabinogalactan-proteins (AGPs) ~~type biopolymers~~ from *Acacia senegal* gum are used as
27 protective colloid in young red wines to prevent the precipitation of the coloring matter.
28 Usually, *Acacia senegal* gum is chosen based on its efficiency to stabilize iron
29 hexacyanoferrate salts in hydro-alcoholic solution. In this study, the protective activity of
30 *Acacia senegal* gum and its ~~three~~ macromolecular fractions (HIC-F1, HIC-F2 and HIC-F3),
31 separated according to their hydrophobicity, towards the precipitation of iron
32 hexacyanoferrate salts and polyphenols in hydro-alcoholic solutions are reported. ~~The~~ AGPs
33 prevent the colloidal instability of both iron hexacyanoferrate salts and polyphenols in
34 “model” hydro-alcoholic solutions and young red wine with a good correlation between
35 results obtained on both systems. This result ~~then~~ strengthens the use of iron hexacyanoferrate
36 salts (hydro-alcoholic – mineral test) for the evaluation of the protective activity of *A. senegal*
37 gum in young red wine. The precipitation of iron hexacyanoferrate salts is avoided by the
38 electrostatic binding of Ca^{2+} , the driver of the instability, with ~~the~~ negatively charged carried
39 by amino-acids from the protein backbone or both by amino acids and uronic acid
40 monosaccharides localized close to the protein backbone of AGPs. The protective activity
41 closely depends on the protein content of AGPs in both iron hexacyanoferrate salts and
42 polyphenols hydro-alcoholic solutions: the more they are rich in proteins, the more their
43 colloidal stabilizing efficiency is (HIC-F3>HIC-F2>HIC-F1). The differences observed in the
44 protective activity between AGPs from the three HIC fractions are relied not only to their
45 protein content but also to their related rate of glycosylation that modulates the protein
46 accessibility to its environment, then their physicochemical properties as their hydrophobic
47 behavior.

48

49 **Keywords:** Acacia gum, Arabinogalactan proteins, colloidal stability, young red wine,
50 coloring matter

51 1. Introduction

52 Red wine is a mixture of water, alcohols, organic acid and of complex molecules at the origin
53 of its organoleptic properties and in constant evolution during ageing. The color as well as the
54 clarity of red wines are ones of the qualities required by the consumers. Red wines must
55 present a colloidal stability not only at the time of bottling but also during ageing and storage
56 until its consumption (Ribéreau-Gayon, Glories, Maujean & Dubourbieu, 2006). The phenolic
57 compounds, responsible of the organoleptic perceptions, are also unstable molecules able to
58 aggregate and precipitate both in young and aged red wines (Ribéreau-Gayon et al., 2006;
59 Alcade-Eon, Garcia-Estévez, Puente, Rivas-Gonzalo & Escribano-Bailon, 2014). Among
60 others, the phenolic compounds can react with ferric iron to form soluble complexes that may
61 then flocculate and precipitate during ageing (Ribéreau-Gayon et al., 2006). The colloidal
62 stability in young red wines may be ensured by several techniques as a cold treatment, fining
63 or addition of protective colloids such as carboxymethyl cellulose, metatartaric acid,
64 mannoproteins and Acacia gum (Waters, Pellerin & Brillouet, 1994; Waters, Pellerin &
65 Brillouet, 1994a; Dupin, Stockdale, Williams, Jones, Markides & Waters, 2000; Escot,
66 Feuillat, Dulau & Charpentier, 2001; Riou, Vernhet, Doco & Moutounet, 2002; Riberau-
67 Gayon et al., 2006; Poncet-Legrand, Doco, Williams & Vernhet, 2007; Pérez
68 Lamela, García-Falcón, Simal-Gándara & Orriols-Fernández, 2007; Gerbaud, Gabas,
69 Blouin & Crachereau, 2010; Teissedre, 2012; Gordillo, Cejudo-Bastante, Rodríguez-Pulido,
70 Jara-Palacios, Ramírez-Pérez, González-Miret & Heredia, 2014). The cold and fining
71 treatments cause the precipitation of existing particles responsible of the turbidity as well as
72 those potentially formed during ageing. Therefore, both treatments have a clarifying and a
73 stabilizing action (Riberau-Gayon et al., 2006). In young red wine, Acacia gum is used as a

74 protective colloid in order to prevent or limit the aggregation and precipitation of unstable
75 colloids as the coloring matter (Ribereau-Gayon et al., 2006; Teissedre, 2012). In particular,
76 Acacia gum macromolecules bind to polyphenols by the involvement of both hydrogen
77 bonding and hydrophobic interactions that prevents their interaction and aggregation with
78 proteins (de Freitas, Carvalho & Mateus, 2003; Mateus, Carvalho, Luis & de Freitas, 2004;
79 Soares, Gonçalves, Fernandes, Mateus & de Freitas, 2009; Soares, Mateus & de Freitas, 2012;
80 Chung, Rojanasasithara, Mutilangi & McClements, 2016). Hence, Acacia gum is mainly used
81 in young red wine for its colloidal stabilizing properties. It stabilizes young red wine but does
82 not clarify it.

83 Acacia gum is a natural exudate obtained from the trunk and branches of *Acacia senegal* and
84 *Acacia seyal* trees (Williams & Phillips, 2000). Acacia gum macromolecules are highly
85 glycosylated hydroxyproline-rich arabinogalactan-proteins (AGPs) belonging to the
86 glycoprotein superfamily (Akiyama, Eda & Kato, 1984). AGPs from Acacia gum are mainly
87 composed by sugars (D-galactose, L-arabinose, L-rhamnose, D-glucuronic acid, and 4-O-
88 methyl-D-glucuronic acid) with a small fraction of proteins and minerals (Idris, Williams &
89 Phillips, 1998; Lopez-Torrez, Nigen, Williams, Doco & Sanchez, 2015). The sugars are
90 organized into hyperbranched polysaccharide blocks covalently linked to the polypeptide
91 backbone in serine- and hydroxyproline-rich domains (Lopez-Torrez et al., 2015). The highly
92 branched polysaccharide structure is formed by main chains of 1,3-linked β -D-
93 galactopyranose substituted by side chains in O-6 position. Units of α -L-arabinofuranosyl and
94 α -L-rhamnopyranosyl are distributed in the main and side chains while β -D-
95 glucuronopyranosyl and 4-O-methyl- β -D-glucuronopyranosyl are mostly end-units
96 (Anderson, Hirst & Stoddart, 1966; Anderson & Stobbart, 1966; Lopez-Torrez et al., 2015).

97 *A. senegal* gum can be defined as a continuum of AGPs differing by their sugar, amino acid
98 and mineral content and composition, sugar to amino acid ratio, polarity, number of charges,

99 molar mass, size and shape (Randall, Phillips & Williams, 1989; Islam, Phillips, Sljivo,
100 Snowden & Williams, 1997; Renard, Lavenant-Gourgeon, Ralet & Sanchez, 2006; Mejia
101 Tamayo, Nigen, Apolinar-Valiente, Doco, Williams, Renard & Sanchez, 2018). These AGPs
102 can be separated according to their polarity using hydrophobic interaction chromatography
103 (HIC) with the recovery of three fractions (Randall et al., 1989; Renard et al., 2006). These
104 three fractions were historically named arabinogalactan (AG), arabinogalactan-protein (AGP)
105 and glycoprotein (GP) according to their protein content and elution order. ~~However, all the~~
106 ~~ability of these three fractions to bind the β -glucosyl Yariv's reagent suggests that they all~~
107 ~~belong to the AGP family because of their reactivity to the Yariv's reactant~~ (Osman et al.
108 1993). Hence, to avoid any confusion, we named them HIC-F1 (arabinogalactan, AG), HIC-
109 F2 (arabinogalactan-protein, AGP) and HIC-F3 (glycoprotein, GP) in their order of elution,
110 reflecting also an increasing hydrophobic index. HIC-F1 is the most abundant fraction (85-
111 92% of the whole gum) as compared to HIC-F2 (6-16% of the whole gum) and HIC-F3 (1-3%
112 of the whole gum). The sugar composition was similar between the three fractions, with
113 however a larger content of arabinose in HIC-F2 and HIC-F3 and a larger content of charged
114 sugars in HIC-F1. HIC-F3 was the richest fraction in proteins with values around 25-40%,
115 while the amount of proteins was around 8-10% and 0.5-1% for HIC-F2 and HIC-F1,
116 respectively. These three HIC fractions differed also by their mean molar mass (M_w) and high
117 M_w AGPs content (AGPs with M_w upper than 10^6 g.mol⁻¹). HIC-F1 was mainly composed of
118 low M_w AGPs, while HIC-F2 and HIC-F3 were richest in high M_w AGPs.

119 In young red wines, it is advised to use *Acacia senegal* gum at a maximal concentration of
120 300 mg/L (Code International des Pratiques Œnologiques, Fiche Code OIV (12/72) – Edition
121 01/2015 II.3.3-7) even if no legal limit exists in the EC 606-2009. Before its addition in young
122 red wines, the efficiency of *A. senegal* gum towards wine colloidal instability is evaluated
123 according to an “efficacy test” of the International Oenological Codex (International

124 Oenological Codex: COEI-1-GOMARA, 2000). This “efficacy test” consists in determining
125 the quantity of *A. senegal* gum required to prevent the flocculation of a colloidal iron
126 hexacyanoferrate solution in hydro-alcoholic medium by calcium salt. Hence, the *A. senegal*
127 gum efficiency towards wine colloidal instability is usually evaluated using a hydro-alcoholic
128 – mineral solution which seems in appearance far enough from wine matrices on their
129 biochemical and physicochemical properties. The colloidal stabilization of two so different
130 matrices by *A. senegal* gum raises some questions on the colloidal stability mechanism and
131 especially on the AGPs at the origin of this colloidal stabilization.

132 The aim of this study was to investigate the colloidal stabilizing properties of AGPs from *A.*
133 *senegal* gum in three hydro-alcoholic solutions: a hydro-alcoholic – mineral solution, “wine-
134 like” solution and young red wine. The AGPs responsible of the colloidal stability in the three
135 hydro-alcoholic solutions were identified. Furthermore, the colloidal stabilization mechanism
136 involving AGPs from *A. senegal* gum in hydro-alcoholic – mineral solution was
137 characterized.

138

139 **2. Materials and Methods**

140 *2.1. Materials*

141 Instant spray dried Acacia *senegal* gum was provided by ALLAND & ROBERT Company -
142 Natural and organic gums (Port Mort, France). *A. senegal* gum was fractionated by
143 hydrophobic interaction chromatography (HIC) according to Renard et al. (Renard et al.,
144 2006). Three HIC fractions were isolated ~~that were~~ and named HIC-F1, HIC-F2 and HIC-F3
145 according to their elution order, and then their hydrophobic behaviour. The biochemical
146 composition and structural properties (mean molar mass) of *A. senegal* gum and HIC
147 fractions were described by Mejia-Tamayo (Mejia-Tamayo et al., 2018) and presented in
148 supplementary data (Table S1).

149 The pronase was from *streptomyces griseus* (lot 10165921001, Roche). The grape marc
150 powder rich in polyphenols (64%) was provided by Grap'Sud Company (Cruviers-Lascours,
151 France). The unstable young red wine (Cabernet sauvignon/Merlot, 2015) was provided by
152 BioLaffort Company (Bordeaux, France). All other reagents were of analytical grade from
153 Sigma-Aldrich (USA).

154

155 2.2. Hydrolysis of *Acacia senegal* gum and HIC-F1 fraction by pronase

156 *A. senegal* gum and HIC-F1 fraction were dissolved at a concentration of 10 mg.ml⁻¹ in ultra
157 pure deionised water (18.2 mΩ resistivity) containing 0.02% NaN₃ and stirred overnight at
158 room temperature. The pH was adjusted at 6.5 using a small aliquot of NaOH solution before
159 to add pronase at a final concentration of 0.14 mg.ml⁻¹. The enzymatic hydrolysis occurred
160 during 48h at 35°C. The enzymatic reaction was stopped by removing the pronase from the
161 samples by centrifugation at 3000 rpm using a centricon Vivaspin 20 (cut-off of 50 000 Da).
162 The hydrolysis products were also removed by the centrifugation step. The samples were then
163 washed four times with ultra pure deionised water (18.2 mΩ resistivity) by centrifugation at
164 3000 rpm using a centricon Vivaspin 20 (cut-off of 50 000 Da), before their freeze-drying.

165

166 2.3. Preparation of *Acacia senegal* gum and its HIC fractions

167 Stock solution of *A. senegal* gum and HIC fractions were dissolved in ~~ultra-pure~~ultra-pure
168 deionised water (18.2 mΩ resistivity) and stirred overnight at room temperature. The
169 solutions were centrifuged at 12 500 rpm for 30 min at 20°C to remove insoluble materials
170 and air bubbles.

171

172 2.4. Electrophoretic mobility measurements

173 The electrophoretic mobility of *A. senegal* gum prepared in hydro-alcoholic solutions (tartaric
174 acid: 3 g.L⁻¹; potassium sulfate: 1 g.L⁻¹; ethanol: 12% v/v) at pH 3.1, 3.5 and 4 was
175 determined using a Zetasizer 1000 (Malvern, United Kingdom). The measurements were
176 performed at 25°C. The data are the average of three measurements.

177

178 2.5. Multi-detector high performance size exclusion chromatography

179 HPSEC experiments were performed using a Shimadzu HPLC system (Shimadzu, Kyoto,
180 Japan) coupled to one Shodex OHpak SB-G pre-column followed by four Shodex OHpak SB
181 columns in series (SB 806 HQ, SB 805 HQ, SB 804 HQ and SB 803 HQ). 50 µl of *A. senegal*
182 gum and HIC fractions solutions (1 mg.ml⁻¹) were injected. The elution was performed with
183 0.1 mol.L⁻¹ LiNO₃ solution containing 0.02% NaN₃ at a flow rate of 1 ml.min⁻¹ and 30°C.

184 The HPSEC system was fitted in series to a Dawn Heleos II multi-angle laser light scattering
185 (Wyatt Technology Corp., Santa Barbara, Ca, USA) and an Optilab T-rEX refractometer
186 (Wyatt Technology Corp., Santa Barbara, Ca, USA). The molar mass distribution and the
187 weight-average molar mass (M_w) were calculated using ASTRA software 6.1.1.17 (Wyatt
188 Technologies, Santa Barbara, CA). The data were analyzed using the Zimm's model (1st
189 order) and a refractive index increment (dn/dc) of 0.155, 0.162, 0.160 and 0.145 ml.g⁻¹ for *A.*
190 *senegal* gum, HIC-F1, HIC-F2 and HIC-F3, respectively, as determined experimentally.

191

192 2.6. Colloidal stability of hydro-alcoholic - mineral solution

193 The hydro-alcoholic – mineral solution was prepared with ~~ultra-pure~~ultra-pure deionised
194 water (18.2 mΩ resistivity) according to the ~~International~~ Oenological International Codex
195 (O.I.V.). This solution ~~It~~was composed of tartaric acid (3 g.L⁻¹), potassium sulfate (1 g.L⁻¹),
196 ethanol (12% v/v), potassium ferrocyanide (60 mg.L⁻¹) and iron ~~solution~~(5 mg.L⁻¹~~‰~~).
197 ~~When~~The addition of iron (Fe³⁺) ~~was added~~ to the hydro-alcoholic – mineral solution

198 ~~containing potassium ferrocyanide ($K_4[Fe(CN)_6]$)~~, induced the formation of a soluble
199 “Prussian blue” colloidal solution of $K[Fe^{III}Fe^{II}(CN)_6]$ ~~is obtained~~. The hydro-alcoholic –
200 mineral solution was stirred during 2 h at room temperature before to adjust ~~the~~ its pH at 3.1
201 using NaOH solution (1 mol.L^{-1}). The hydro-alcoholic – mineral solution was then filtered
202 through a $0.45 \mu\text{m}$ regenerated cellulose membrane (GE Healthcare Life Sciences, Germany).
203 For the studies according to pH, the hydro-alcoholic – mineral solution was prepared at pH
204 3.1, 3.5 and 4.0.

205 The hydro-alcoholic – mineral solution was destabilized by the addition of CaCl_2 (22 g.L^{-1})
206 prepared in ultra-pure deionised water ($18.2 \text{ m}\Omega$ resistivity). The added Ca^{2+} bound to
207 $K[Fe^{III}Fe^{II}(CN)_6]$ inducing the formation of “Prussian blue” precipitates ($\text{Ca}[Fe^{III}Fe^{II}(CN)_6]_2$).
208 Most of experiments were performed at a final CaCl_2 concentration of 0.67 g.L^{-1} . Some
209 experiments were performed at a final concentration of 0.33, 0.5, 1 and 1.33 g.L^{-1} to
210 investigate the influence of calcium. Each measurement was triplicated.

211 The efficiency of *A. senegal* gum and its HIC fraction to prevent the colloidal mineral
212 destabilization was studied by varying their concentration in the hydro-alcoholic – mineral
213 solution. The final concentration of *A. senegal* gum or its HIC fractions in the hydro-alcoholic
214 – mineral solution ranged between 0 and 1 g.L^{-1} . *A. senegal* gum or HIC fraction solutions
215 were added to the hydro-alcoholic – mineral solution before the addition of CaCl_2 solution.
216 Each measurement was triplicated.

217

218 2.7. Colloidal stability of hydro-alcoholic – grape marc solution and young red wine

219 The hydro-alcoholic – grape marc solution (wine-like medium) was prepared with ultra-pure
220 deionised water ($18.2 \text{ m}\Omega$ resistivity). It contained tartaric acid (2.7 g.L^{-1}), potassium sulfate
221 (0.9 g.L^{-1}), grape marc powder (3 g.L^{-1}) and ethanol (12% v/v). The solution was gently
222 stirred during 2 h at room temperature before to adjust the pH at 3.5 using NaOH solution (1

223 mol.L⁻¹). The solution was then filtered through a 0.45 µm regenerated cellulose membrane
 224 (GE Healthcare Life Sciences, Germany).

225 The colloidal destabilization of the hydro-alcoholic – grape marc solution and the young red
 226 wine was induced by cooling the solutions to 10°C. The efficiency of *A. senegal* gum and its
 227 HIC fraction to prevent the destabilization of the hydro-alcoholic – grape marc solution and
 228 the young red wine was studied at 10°C by varying their concentration between 0 and 2 g.L⁻¹.
 229 Each measurement was triplicated.

230

231 2.8. Characterization of the colloidal stability of hydro-alcoholic model solutions and young 232 red wine

233 The colloidal stability of the hydro-alcoholic – mineral and hydro-alcoholic – grape marc
 234 solutions, and the young red wine was investigated using a Turbiscan Tower equipped with a
 235 pulsed near infrared light source ($\lambda = 880$ nm) and two synchronous detectors, a transmission
 236 (T) and a backscattering (BS) detectors (Formulacion, France). The samples were loaded into
 237 cylindrical glass tubes and scanned throughout its entire height. The transmittance (T, in %)
 238 was measured every 1 min and 50 sec during 24 h at 25°C for the hydro-alcoholic – mineral
 239 solutions and during 48 h at 10°C for the hydro-alcoholic – grape marc solutions and young
 240 red wine.

241 The transmittance (T) profiles were analyzed using the TowerSoft software, version 1.1.0.36
 242 (Formulacion, France). ΔT (%) corresponded to the difference in transmittance between the
 243 scan_i and the first scan. The Turbiscan Stability Index (TSI) was also determined as the
 244 following:

$$245 \quad TSI = \sum_i \frac{\sum h |scan_i(h) - scan_{i-1}(h)|}{H}$$

246 The TSI corresponded to the transmittance variation at all measured position (h) throughout
247 the entire height (H) of the sample between the scan_i and the scan_{i-1}. The plot of TSI
248 according to time corresponds to the kinetic behavior of the samples that takes into account all
249 the physical phenomena (especially destabilization mechanism) occurring during the
250 measurement.

251 The TSI kinetics were used to plot the TSI values at 24h in hydro alcoholic – mineral solution
252 and 48h in hydro alcoholic – grape marc solution and young red wine according to AGPs
253 concentration. This representation assimilated to a “stability curve” was used to
254 experimentally determine the AGPs concentration, named critical concentration, necessary to
255 obtain the colloidal stability of the solutions.

256

257 3. Results

258 3.1. Colloidal stabilization mechanism of the hydro-alcoholic – mineral solution by *A.* 259 *senegal* gum

260 The colloidal stability of the hydro-alcoholic – mineral solution (pH 3.1) containing *A.*
261 *senegal* gum was followed by measuring at 25°C the transmittance (T) of the solutions during
262 24h using a light scattering equipment. The figure 1 showed the temporal change in
263 transmittance (ΔT) of three hydro-alcoholic – mineral solutions without and with 0.09 and
264 0.14 g.L⁻¹ of *A. senegal* gum (Figure 1A, 1B and 1C). After 24h, the transmittance (T) of the
265 hydro-alcoholic – mineral solutions without and with 0.09 g.L⁻¹ of *A. senegal* gum increased
266 respectively by 60% and 35% in the middle of the sample (cell height: 2 – 36 mm) reflecting
267 a clarification of the solution. This phenomenon was confirmed by the loss of the blue color
268 of the solutions after 24h (Figure 1A and 1B). The clarification resulted from the formation of
269 a salt precipitate ($\text{Ca}[\text{Fe}^{\text{III}}\text{Fe}^{\text{II}}(\text{CN})_6]_2$) between $\text{K}[\text{Fe}^{\text{III}}\text{Fe}^{\text{II}}(\text{CN})_6]$ and Ca^{2+} that pelleted, as
270 evidenced by the decrease of T by 5% and 2% after 24h in the bottom of the cell (cell height:

271 1 – 1.5 mm) for the solutions without and with 0.09 g.L⁻¹ of *A. senegal* gum, respectively
272 (Figure 1A and 1B). On the other hand, the transmittance of the hydro-alcoholic – mineral
273 solution containing 0.14 g.L⁻¹ of *A. senegal* gum was constant during the 24h of
274 measurement, reflecting the colloidal stability of the solution. The influence of *A. senegal*
275 gum concentration (0 – 0.14 g.L⁻¹) on the colloidal stability kinetics of the hydro-alcoholic –
276 mineral solution was analyzed by plotting the TSI (Turbiscan Stability Index) according to
277 time (Figure 2A). The increase of *A. senegal* gum concentration reduced and delayed the
278 instability mechanism occurring in the hydro-alcoholic – mineral solution. The colloidal
279 instability was totally prevented for an *A. senegal* gum concentration above around 0.11 g.L⁻¹.
280 The minimal *A. senegal* gum concentration ([AG]_{critical}) necessary to obtain the hydro-
281 alcoholic – mineral colloidal stability was determined by plotting the TSI values after 24h
282 according to *A. senegal* gum concentration (stability curve) (Figure 2B). Below an *A. senegal*
283 gum concentration of 0.05 g.L⁻¹, the TSI values at 24h was constant and similar to the control
284 sample without *A. senegal* gum. In these experimental conditions, the concentration of *A.*
285 *senegal* gum was too low. Above 0.05 g.L⁻¹ of *A. senegal* gum, the TSI values decreased
286 progressively before to reach a constant TSI value (≤ 2) similar to the control without Ca²⁺ for
287 *A. senegal* gum concentration upper than 0.11 g.L⁻¹. Based on the stability curve, the
288 [AG]_{critical} was 0.114 ± 0.002 g.L⁻¹ (Figure 2B).
289 The colloidal stabilization mechanism was further investigated by modifying the
290 physicochemical condition as the pH of the solutions and the amount of Ca²⁺ added to induce
291 the colloidal instability mechanism. The stability curves of hydro-alcoholic – mineral
292 solutions prepared at pH 3.1, 3.5 and 4.0 are shown in Figure 3A. As the pH increased from
293 3.1 to 4.0, the stability curves were shifted towards lower *A. senegal* gum concentrations
294 necessary to prevent the colloidal instability. The [AG]_{critical} were 0.114 ± 0.002 , $0.094 \pm$
295 0.003 and 0.064 ± 0.002 g.L⁻¹ at pH 3.1, 3.5 and 4.0, respectively. Hence, the colloidal

296 stabilizing properties of *A. senegal* gum were greatly influenced by pH and enhanced with its
297 increase in this pH range. AGPs from *A. senegal* gum are weakly charged glycoproteins
298 characterized by a global negative charge in aqueous solution for pH ranging from 2.0 to 10.0
299 (Burgess & Carless, 1984). Moreover, the negative AGPs μ_E increased from pH 2.0 to ~ 5.0
300 in aqueous solutions reflecting the increase of the global negative charge (Burgess & Carless,
301 1984; Schmitt, Sanchez, Thomas & Hardy, 1999). Thus, the electrophoretic mobility (μ_E) of
302 *A. senegal* gum in hydro-alcoholic solutions prepared at pH 3.1, 3.5 and 4.0 was measured
303 and plotted according to $[AG]_{critical}$ (Figure 3B). A good correlation was evidenced between
304 *A. senegal* gum μ_E and its concentration necessary to stabilize the hydro-alcoholic – mineral
305 solutions. The more the negative *A. senegal* gum μ_E was, the less the *A. senegal* gum quantity
306 was required to prevent the colloidal instability. Hence, negative charges carried by AGPs
307 were certainly involved in this colloidal stabilization mechanism.

308 Since the colloidal instability of the hydro-alcoholic – mineral solution is induced by the
309 addition of Ca^{2+} , it was hypothesized that electrostatic interactions between Ca^{2+} and negative
310 charges of AGPs could play an important role in the stabilization mechanism. Therefore, we
311 determined the $[AG]_{critical}$ in hydro-alcoholic – mineral solutions (pH 3.1) containing different
312 final Ca^{2+} concentrations (0.33 to 1.33 g.L⁻¹). A close linear relationship between the
313 $[AG]_{critical}$ and the amount of Ca^{2+} added in the hydro-alcoholic – mineral solutions was
314 obtained (Figure 4). The increase of Ca^{2+} concentration induced also the increase of
315 $[AG]_{critical}$. Based on its structural and physicochemical properties, hyperbranched and
316 negatively charged, AGPs from *A. senegal* gum appear as a natural mineral carrier that
317 contains ~2–3.4 wt% of ash for this Acacia gum sample, and especially Ca^{2+} (Anderson,
318 Douglas, Morrison & Weiping, 1990; Debon & Tester, 2001; Mhinzi, 2003). Lamport and
319 Varnai evidenced that Ca^{2+} bound to AGPs with a fairly strong binding constant of 6.5×10^{-6}
320 mol.L⁻¹ (Lamport & Varnai, 2013). They also showed that the Ca^{2+} binding sites of AGPs

321 were not naturally fully saturated by Ca^{2+} . Hence, as expected, we can easily suggest that
322 AGPs stabilized the hydro-alcoholic – mineral solutions by its electrostatic binding with Ca^{2+} ,
323 the driver of the instability mechanism.

324

325 3.2. Identification AGPs from *A. senegal* gum involved in the colloidal stability mechanism of 326 the hydro-alcoholic – mineral solution

327 As previously mentioned, *A. senegal* gum is a continuum of AGPs differing by their
328 biochemical, structural and physicochemical properties (Randall et al., 1989; Renard et al.,
329 2006). To better identify the macromolecules involved in this colloidal stabilization
330 mechanism, *A. senegal* gum was fractionated using Hydrophobic Interaction Chromatography
331 (HIC). The three fractions obtained were named HIC-F1, HIC-F2 and HIC-F3 according to
332 their elution order and consequently to their increasing hydrophobic index. The biochemical
333 composition and some structural properties of *A. senegal* gum and its HIC fractions were
334 previously described (Mejia-Tamayo et al., 2018) and presented in supplementary data (table
335 1). The three HIC fractions are composed of the same sugars (D-galactose, L-arabinose, D-
336 glucuronic acid and 4-O-methyl-D-glucuronic acid) with some differences in their
337 proportions. The arabinose to galactose (Ara/Gal) molar ratio was 0.69, 1.03 and 1.29 for
338 HIC-F1, HIC-F2 and HIC-F3, respectively. HIC-F1 was also richer in uronic acid (glucuronic
339 and 4-O-methyl-glucuronic acids) than HIC-F2 and HIC-F3 fractions (21.7 wt% vs. 16.2 wt%
340 and 14.4 wt%, respectively). Hence, the carbohydrate blocks of HIC-F1 were supposed to
341 carry more negative charges than HIC-F2 and HIC-F3 ones. Biochemically, the main
342 difference between HIC fractions was the protein content and consequently the protein/sugar
343 ratio. HIC-F3 showed a higher amount of proteins (13.8 wt%) and then protein/sugar ratio
344 (0.169) as compared to HIC-F2 (6.3 wt% and 0.069) and HIC-F1 (0.5 wt% and 0.005).
345 Structurally, the three HIC fractions differed mainly by their mean molar mass (M_w) and high

346 molar mass (HM_w) macromolecules content ($M_w > 10^6 \text{ g}\cdot\text{mol}^{-1}$). HIC-F2 and HIC-F3 were
347 richer in protein-rich high molar mass (HM_w) AGPs than HIC-F1 (73, 56 and 3%,
348 respectively) (supplementary table 1).

349 The stability curves of the hydro-alcoholic – mineral solution in the presence of the three HIC
350 fractions are shown in the Figure 5. They can be classified from the most effective to the least
351 in terms of stabilization as: HIC-F3 > HIC-F2 > HIC-F1. Thus, the critical concentrations of
352 HIC-F1, HIC-F2 and HIC-F3 fractions were 0.570, 0.090 and 0.012 $\text{g}\cdot\text{L}^{-1}$, respectively.
353 Though HIC-F3 was the minor fraction in weight (around 1 wt% of whole *A. senegal* gum),
354 its efficiency for the colloidal stabilization of the hydro-alcoholic – mineral solution was 48,
355 10 and 7.5 times larger than those of HIC-F1, *A. senegal* gum and HIC-F2, respectively.
356 Hence, the colloidal stability properties of *A. senegal* gum towards the hydro-alcoholic –
357 mineral solution were mainly due to its two minor fractions in weight, HIC-F2 and HIC-F3
358 (around 11 wt% of whole *A. senegal* gum), that were especially richer in protein-rich HM_w
359 AGPs than HIC-F1 (Table 1).

360 HIC-F1 (89 wt% of whole *A. senegal* gum) that was characterized by poor colloidal
361 stabilizing property contained also a low amount (3%) of protein-rich HM_w AGPs. To precise
362 whether the macromolecular origin of the colloidal stabilizing properties is due to this
363 fraction, its HM_w AGPs were removed by an enzymatic hydrolysis treatment using the
364 pronase, a protease mixture. Several studies showed the specific hydrolysis activity of pronase
365 for protein-rich HM_w AGPs from *A. senegal* gum without degrading and affecting the
366 structural properties of low M_w (LM_w) AGPs (Connolly, Fenyo & Vandeveld, 1988; Randall
367 et al., 1989; Flindt, Al-Assaf, Phillips & Williams, 2005; Mahendran, Williams, Phillips, Al-
368 Assaf & Baldwin, 2008; Renard, Lavenant-Gourgeon, Lapp, Nigen & Sanchez, 2014). The
369 refractive index (RI) signal and the M_w distribution of HIC-F1 before and after the enzymatic
370 treatment are shown in Figure 6A. The pronase hydrolyzed the protein-rich AGPs decreasing

371 the HM_w AGPs content from 3 to 1.7%, without affecting the quantity and the structural
372 properties of LM_w AGPs (Figure 6A). The colloidal stabilizing properties of the hydrolyzed
373 HIC-F1 towards the hydro-alcoholic – mineral solution were characterized and compared to
374 HIC-F1 (Figure 6C). The hydro-alcoholic – mineral solutions was totally unstable until 1.2
375 $g.L^{-1}$ of hydrolyzed HIC-F1, a concentration twice more than the minimal HIC-F1
376 concentration necessary to totally prevent the colloidal instability ($[HIC-F1]_{critical} = 0.570 g.L^{-1}$).
377 Hence, the removal of less than 1.5% of AGPs from HIC-F1, corresponding to HM_w
378 AGPs, induced the loss of its colloidal stabilizing properties. The main contribution of
379 protein-rich HM_w AGPs was also confirmed by studying the colloidal stabilizing properties of
380 hydrolyzed *A. senegal* gum by pronase. After the enzymatic treatment, the content of protein-
381 rich HM_w ($M_w > 10^6 g.mol^{-1}$) AGPs decreased from 14 to 6% (Figure 6B). As observed for
382 HIC-F1, *A. senegal* gum also lost its stabilizing properties after the pronase treatment: the
383 hydro-alcoholic – mineral solutions containing $0.12 g.L^{-1}$ of hydrolyzed *A. senegal* gum was
384 totally unstable while $[AG]_{critical}$ was found to be $0.114 g.L^{-1}$. These results highlighted the
385 major role of the protein-rich AGPs in the colloidal stabilizing mechanism of the hydro-
386 alcoholic – mineral solution. Hence, HIC-F2 and HIC-F3 fractions prevented the colloidal
387 instability of the hydro-alcoholic – mineral solution (i.e. formation of salt precipitates) by
388 their electrostatic binding with Ca^{2+} . The key role of the protein-rich AGPs on the functional
389 properties of *A. senegal* gum was also evidenced by studying its interfacial properties
390 (Elmaman, Al-Assaf, Phillips & Williams, 2008). In this study, the removal of the protein-
391 rich AGPs by a pronase treatment resulted in the loss of the interfacial properties of *A.*
392 *senegal* gum.

393

394 *3.3. Colloidal stabilization of a hydro-alcoholic – grape marc solution and an unstable young*

395 *red wine by AGPs from A. senegal gum*

396 In order to correlate the results obtained on the hydro-alcoholic – mineral solution to the wine
397 matrix, the stabilizing properties of *A. senegal* gum and its HIC fractions were characterized
398 on an unstable synthetic hydro-alcoholic – grape marc solution (wine-like medium) and an
399 unstable young red wine.

400 The hydro-alcoholic – grape marc solution and the young red wine that were stable at 25°C
401 (control sample) were destabilized upon cooling to 10°C that induced the precipitation of the
402 unstable colloidal substances as the coloring matter (Ribereau-Gayon et al., 2006; Alcade-Eon
403 et al., 2014). The colloidal stability kinetics of the hydro-alcoholic – grape marc solution and
404 the young red wine supplemented with *A. senegal* gum and HIC fractions were followed by
405 measuring the transmittance of the solution during 48 h at 10°C. The increase of *A. senegal*
406 gum concentration from 0 to 0.5 g.L⁻¹ led to the colloidal stabilization of the hydro-alcoholic
407 – grape marc solution (Figure 7A). The solution supplemented with 0.5 g.L⁻¹ of *A. senegal*
408 gum presented a similar kinetic behavior as the control sample at 25°C suggesting that AGPs
409 from *A. senegal* gum totally prevented the colloidal instability. This result was in agreement
410 with the use of *A. senegal* gum, and more globally the AGPs, as protective colloids in wines
411 and model beverages to prevent the aggregation and precipitation of polyphenols and proteins
412 (Ribereau-Gayon et al., 2006; Waters et al., 1994; Riou et al., 2002; Mateus et al., 2004;
413 Luck, Liao, Murray, Grimmer, Warminski, Williamson, Lilley & Haslam, 1994; Liang, Liu,
414 Qi, Su, Yu, Wang & He, 2013; Soares et al., 2009; de Freitas et al., 2003), and the chemical
415 modifications of anthocyanins (Chung et al., 2016). The efficiency of *A. senegal* gum and its
416 HIC fractions towards the colloidal instability of this hydro-alcoholic – grape marc solution
417 were characterized by plotting the stability curves after 48h of kinetics (Figure 7B). The
418 critical concentrations of *A. senegal* gum, HIC-F1, HIC-F2 and HIC-F3 fractions were found
419 to be 0.245 ± 0.010 , 0.996 ± 0.090 , 0.118 ± 0.014 and 0.027 ± 0.003 g.L⁻¹, respectively. The
420 colloidal stabilizing properties of *A. senegal* gum, HIC-F2 and HIC-F3 fractions were also

421 confirmed on an unstable young red wine (Figure 7C). They totally prevented the young red
422 wine colloidal instability at 10°C for concentrations of 0.086 ± 0.007 , 0.045 ± 0.006 and
423 0.015 ± 0.002 g.L⁻¹, respectively. Unlike the other two HIC fractions, HIC-F1 was not
424 efficient for the colloidal stabilization of the young red wine even at a concentration of 2 g.L⁻¹
425 (data not shown).

426

427 4. Discussion

428 The colloidal stabilizing properties of *A. senegal* gum and its HIC fractions isolated from
429 Hydrophobic Interaction Chromatography were investigated on three colloidal hydro-
430 alcoholic matrices: a hydro-alcoholic – mineral matrix, a hydro-alcoholic – grape marc
431 solution (“wine-like” matrix) and an unstable young red wine. Whatever the hydro-alcoholic
432 matrix tested, the HIC fractions can be classified in the same following order according to
433 their colloidal stabilizing efficiency: HIC-F1 << HIC-F2 < HIC-F3. *A. senegal* gum that is a
434 mixture of the HIC fractions presented some intermediate properties between those of HIC-F1
435 and HIC-F2. Before its addition in young red wines, ~~Usually, *A. senegal* gum added in young~~
436 ~~red wines is usually chosen based on its ability to stabilize a hydro-alcoholic – mineral matrix.~~
437 When we plot the critical concentrations of *A. senegal* gum and its HIC fractions determined
438 in hydro-alcoholic – mineral solution as a function of those determined in hydro-alcoholic –
439 grape marc solution or young red wine (Figure 8), a good correlation between both systems is
440 obtained. The more the AGPs from *A. senegal* gum are efficient to stabilize the hydro-
441 alcoholic – mineral solution, the better they are to stabilize the polyphenols in young red
442 wines. Hence, these results confirmed and strengthened the relevance and the use of the
443 hydro-alcoholic – mineral test for the evaluation of *A. senegal* gum colloidal stabilizing
444 properties before its addition in young red wine.

445 In the hydro-alcoholic – mineral solution, the colloidal stability was assessed by the
446 electrostatic binding of Ca^{2+} , the “positively charged” driver of the colloidal instability, with
447 the negatively charged AGPs. According to the monosaccharide and amino acid compositions
448 of *Acacia senegal* gum (Mejia-Tamayo et al., 2018), the negative charges of AGPs are carried
449 out both by the carbohydrate moiety through the carboxylic groups of glucuronic and 4-O-
450 methyl glucuronic acid monosaccharides, and the polypeptidic backbone by the carboxylic
451 groups of aspartic and glutamic acid amino acid side chains. The content and proportion of
452 these negatively charged uronic acids and amino acids are fraction dependent with a large
453 excess of uronic acids in the three fractions. HIC-F1 has a higher uronic acid content (20.9
454 wt% of whole fraction) as compared to HIC-F2 and HIC-F3 (14.9 wt% and 11.7 wt%,
455 respectively) (supplementary table 1). Recently, Grein-Iankovski et al. (2018) characterized
456 an acid dissociation constant (pK_a) between 3 and 4 for the AGPs from *A. senegal* gum that
457 was attributed to the carboxylic group of glucuronic acids. This chemical property suggests a
458 partial ionization of the carboxylic groups in the range of the pH studied in the hydro-
459 alcoholic solutions, and moreover the increase of the negative charge of AGPs with the
460 increase of pH from 3.1 to 4.0. This result is in well accordance with the decrease of the
461 critical concentration of *A. senegal* gum when the pH is increased from 3.1 to 4.0. Recently,
462 †The binding of Ca^{2+} with the carboxyl group of two uronic acids of a conserved O-Hyp-
463 linked arabinogalactan polysaccharide was demonstrated using molecular dynamic
464 simulations (Lamport & Varnai, 2013). Hence, based on the excess of uronic acid in the three
465 HIC fractions and the molecular dynamic simulations it could be expected as a first
466 approximation that the mineral colloidal stability occurred mainly by the binding of Ca^{2+} to
467 the uronic acids of carbohydrate blocks. However, our results showed that HIC-F1, the richest
468 fraction in uronic acids, had the weakest colloidal stabilizing properties, while HIC-F3, the
469 one with the least uronic acids content, was the most effective fraction for the colloidal

470 stabilization of the hydro-alcoholic – mineral solution (Figure 5). Hence, the stabilizing
471 properties of AGPs could not be only explained by their uronic acids content. We also showed
472 a significant decrease of the stabilizing properties of HIC-F1 and *A. senegal* gum after the
473 hydrolysis of their protein-rich AGPs by a pronase treatment. The pronase is a cocktail of
474 proteases hydrolyzing the accessible polypeptidic backbone of protein-rich AGPs without
475 degrading the low molar mass (LM_w) AGPs poor in proteins (Renard et al. 2014). This
476 enzymatic treatment gave rise to macromolecules presenting similar molar masses (M_w from
477 1.8 to 6.0×10^5 $g \cdot mol^{-1}$) and conformational properties as low molar mass (LM_w) AGPs
478 naturally found in *A. senegal* gum (Connolly et al., 1988, Randall et al., 1989, Aoki, Al-
479 Assaf, Katayama & Phillips, 2007, Renard et al., 2014). Since the conformational properties,
480 and probably the sugar organization, of the released carbohydrate blocks of AGPs was
481 seemingly not disturbed by the pronase treatment, this suggested that the uronic acids of the
482 carbohydrate blocks were certainly not directly involved in the electrostatic Ca^{2+} binding in
483 this colloidal stability test. Similar results were obtained on AGPs purified from a red wine by
484 Waters et al. These authors showed that the greatest protein haze protective activity in
485 Chardonnay wine was reached with AGPs presenting the highest protein content. Moreover,
486 this protective activity was not affected by the sugar hydrolysis of AGPs using α -
487 arabinofuranosidase and endo-(1 \rightarrow 6)- β -D-galactanase, while it was totally lost when the
488 protein moiety was altered with the loss of 75% of initial protein after smith degradation
489 treatment (Waters et al., 1994). These results revealed the key role of the accessible
490 polypeptidic backbone of AGPs in their colloidal stabilizing properties. In the hydro-alcoholic
491 – mineral solution, it could be hypothesized that the electrostatic binding of Ca^{2+} occurred
492 with only the negatively charged amino acids (aspartic and glutamic acids) or with the
493 involvement of both negatively charged amino acids and uronic acid monosaccharides
494 localized close to the protein backbone.

495 As observed in the hydro-alcoholic – mineral solution, the protein-rich AGPs from *A. senegal*
496 gum, HIC-F2 and HIC-F3 fractions, were also found to be the most effective for the colloidal
497 stabilization of the hydro-alcoholic – grape marc solution and the unstable young red wine.
498 The surface and colloidal stabilizing properties of the protein-rich AGPs were previously
499 demonstrated in other type of matrices. After *A. senegal* gum fractionation by size exclusion
500 chromatography (SEC) and hydrophobic interaction chromatography (HIC), Ray et al.
501 showed that the protein-rich AGPs made the best emulsions in model beverage (Ray, Bird,
502 Iacobucci & Clark, 1995). Protein-rich AGPs were also found to be the most effective in
503 decreasing the interfacial tension of n-hexadecane-water interface (Castellani, Al-Assaf,
504 Axelos, Phillips & Anton, 2010), to preferentially adsorb at the interface of latex dispersions
505 (Snowden, Phillips & Williams, 1987) and oil droplets (Randall, Phillips & Williams, 1988),
506 and to stabilize carbon nanotube dispersions (Li, Zhang, Jin & Cai, 2018).

507 The stabilizing properties of the protein-rich AGPs towards such a chemical diversity of
508 molecules (minerals, polyphenols, proteins, oils, latex, etc...) could be relied to their intrinsic
509 properties. Structurally, the three HIC fractions seems not so different. AGPs from these
510 fractions are hyperbranched glycoproteins adopting more or less extended ellipsoidal
511 conformations (Sanchez, Schmitt, Kolodziejczyk, Lapp, Gaillard & Renard, 2008; Renard,
512 Garnier, Lapp, Schmitt & Sanchez, 2012; Renard, Garnier, Lapp, Schmitt & Sanchez, 2013;
513 Renard et al., 2014; Lopez-Torrez et al., 2015). AGPs from HIC-F2 and HIC-F3 fractions also
514 appeared as more “flexible” than those of HIC-F1 (Mejia-Tamayo et al., 2018).
515 Biochemically, HIC fractions presented some more marked differences, especially in their
516 protein content and consequently in their protein to sugar ratio (supplementary table 1) that
517 influenced their hydrophobic and hydration properties (Mejia-Tamayo et al., 2018). HIC-F1,
518 HIC-F2 and HIC-F3 were eluted in that order according to their protein content and
519 increasing hydrophobic behavior. From our results, the higher the HIC fraction is rich in

520 protein, the more effective it is for the colloidal stabilization of the three hydro-alcoholic
521 solutions. In the hydro-alcoholic – grape marc solution and young red wine, these results were
522 in accordance with the putative stabilization of polyphenols by their binding with AGPs via
523 the establishment of hydrophobic interactions and hydrogen bonding (de Freitas et al., 2003;
524 Mateus et al., 2004; Soares et al., 2009; Soares et al., 2012; Chung et al., 2016). Hence, the
525 protein content and certainly the amino acids composition and distribution in the protein
526 backbone of AGPs from *A. senegal* gum appeared as the main factor for the expression of
527 their colloidal stabilizing properties in hydro-alcoholic matrices. This behavior agrees with
528 the work performed by Dickinson et al. dealing with the surface properties of Acacia gums.
529 Based on the surface activity of various Acacia gums with different protein content, these
530 authors previously evidenced a good correlation between the Acacia gum protein content and
531 its interfacial properties (Dickinson, Murray, Stainsby & Anderson, 1988; Dickinson, 2003).
532 The protein content appeared also as essential for the colloidal stabilizing properties of
533 mannoproteins, other glycoproteins considered as protective colloids in wine. The addition of
534 mannoproteins containing a high proportion of proteins (10 to 30%) to red wine prevented the
535 coloring matter precipitation and protein haze in white wine (Waters et al., 1994a; Escot et al.,
536 2001; Charpentier, Escot, Gonzalez, Dulau & Feuillat, 2004; Poncet et al., 2007), while the
537 mannoproteins with a low protein content were devoid of protective activity (Charpentier et
538 al., 2004; Guadalupe, Palacios & Ayestaran, 2007; Poncet et al., 2007; Guadalupe &
539 Ayestaran, 2008).

540 It is useful to remember that AGPs are constituted by a protein backbone covalently linked to
541 hyperbranched carbohydrate blocks that partially hindered it from its environment. Therefore,
542 if we consider the protein as the major component for the colloidal stabilizing properties of
543 AGPs, it would seem appropriate to consider not only the protein content but also its
544 accessibility to its environment that is closely link to the rate of glycosylation. HIC-F1,

545 presenting the weakest colloidal stabilizing properties, is highly glycosylated as compared to
546 HIC-F2 and HIC-F3: their protein to sugar ratios (w/w) were 0.005, 0.069 and 0.169,
547 respectively. The high glycosylation rate of HIC-F1 cause the protein less accessible to its
548 environment as demonstrated by its resistance to the protein hydrolysis assay (Flindt et al.,
549 2005; Renard et al., 2014). Moreover, the rate of glycosylation also influenced the structural
550 and physicochemical properties of HIC fractions. The more the AGPs were glycosylated, then
551 hydrated, the less their flexibility and hydrophobic behavior were marked (Mejia Tamayo et
552 al., 2018). Therefore, we can advance that the surface properties of AGPs are mainly due to
553 the physicochemical properties of their protein backbone which are subtly modulated and
554 controlled by their rate of glycosylation.

555

556 **5. Conclusion**

557 In this research, the colloidal stabilizing properties of arabinogalactan-proteins from *A.*
558 *senegal* gum in hydro-alcoholic – mineral and hydro-alcoholic – polyphenols solutions were
559 investigated. The AGPs prevented the colloidal instability of both calcium iron
560 hexacyanoferrate salts in “model” hydro-alcoholic solution and polyphenols in young red
561 wine. A good relationship was evidenced between the stabilizing properties of AGPs
562 determined in these two hydro-alcoholic solutions. The protein moiety of the AGPs appeared
563 to be essential for these functional properties whatever the hydro-alcoholic solutions. The
564 more the AGPs were rich in proteins, the more their colloidal stabilizing efficiency were. In
565 the hydro-alcoholic – mineral solution, the AGPs avoided the precipitation of potassium
566 ferrocyanide salts by their electrostatic binding with Ca^{2+} , the driver of the instability.

567

568 **Acknowledgements**

569 The authors would like to thank BioLaffort Company (Bordeaux, France) for the financial
570 support and Alland&Robert Company (Port-Mort, France) for the gift of *Acacia senegal* gum
571 sample.

572

573 **References**

574 Akiyama, Y., Eda, S., & Kato K. (1984). Gum Arabic is a kind of arabinogalactan protein.
575 *Agricultural and Biological Chemistry*, 48, 235-237.

576 Alcade-Eon, C., Garcia-Estévez, I., Puente, V., Rivas-Gonzalo, J.C., & Escribano-Bailon,
577 M.T. (2014). Color stabilization of red wines. A chemical and colloidal approach. *Journal of*
578 *Agricultural and Food Chemistry*, 62, 6984-6994.

579 Anderson, D.M.W., Hirst, E., & Stoddart, J.F. (1966). Studies on uronic acid materials Part
580 XVII some structural features of *Acacia senegal* gum (gum Arabic). *Journal of the Chemical*
581 *Society C: Organic*, 1959-1966.

582 Anderson, D.M.W., & Stoddart, J.F. (1966). Studies on uronic acid materials Part XV. The
583 use of molecular-sieve chromatography in studies on *Acacia senegal* gum (Gum Arabic).
584 *Carbohydrate Research*, 2, 104-114.

585 Anderson, D.M.W., Douglas, D.M.B., Morrison, N.A., & Weiping, W. (1990). Specifications
586 for gum Arabic (*Acacia senegal*); analytical data for samples collected between 1904 and
587 1989. *Food Additives and Contaminants*, 7, 303-321.

588 Aoki, H., Al-Assaf, S., Katayama, T., & Phillips, G.O. (2007). Characterization and properties
589 of *Acacia senegal* (L.) Willd. Var. *senegal* with enhanced properties (*Acacia* (sen) SUPER
590 GUM (TM): Part 2- Mechanism of the maturation process. *Food Hydrocolloids*, 21, 329-337.

591 Burgess, D.J., & Carless, J.E. (1984). Microelectrophoretic studies of gelatin and *Acacia* for
592 the prediction of complex coacervation. *Journal of Colloid and Interface Science*, 98, 1-8.

- 593 Castellani, O., Al-Assaf, S., Axelos, M., Phillips, G.O., & Anton, M. (2010). Hydrocolloids
594 with emulsifying capacity. Part-2 Adsorption properties at the n-hexadecane-water interface.
595 *Food Hydrocolloids*, 24, 121-130.
- 596 Charpentier, C., Escot, S., Gonzalez, E., Dulau, L., & Feuillat, M. (2004). Influence of yeast
597 glycosylated proteins on tannins aggregation in model solution. *Journal International des*
598 *Sciences de la Vigne et du Vin*, 38, 209-218.
- 599 Chung, C., Rojanasasithara, T., Mutilangi, W., & McClements, D.J. (2016). Enhancement of
600 colour stability of anthocyanins in model beverages by gum Arabic addition. *Food Chemistry*,
601 201, 14-22.
- 602 Connolly, S., Fenyo, J.C., & Vandeveld, M.C. (1988). Effect of a proteinase on the
603 macromolecular distribution of *Acacia senegal* gum. *Carbohydrate Polymers*, 8, 23-32.
- 604 Debon, S.J.J., & Tester, R.F. (2001). In vitro binding of calcium, iron and zinc by non-starch
605 polysaccharides. *Food Chemistry*, 73, 401-410.
- 606 Dickinson, E., Murray, B.S., Stainsby, G., & Anderson, D.M.W. (1988). Surface activity and
607 emulsifying behavior of some *Acacia* gums. *Food Hydrocolloids*, 2, 477-490.
- 608 Dickinson, E. (2003). Hydrocolloids at interfaces and the influence on the properties of
609 dispersed systems. *Food Hydrocolloids*, 17, 25-39.
- 610 de Freitas, V., Carvalho, E., & Mateus, N. (2003). Study of carbohydrate influence on protein-
611 tannin aggregation by nephelometry. *Food Chemistry*, 81, 503-509.
- 612 Dupin, I.V.S., Stockdale, V.J., Williams, P.J., Jones, G.P., Markides, A.J., & Waters, E.J.
613 (2000). *Saccharomyces cerevisiae* mannoproteins that protect wine from protein haze:
614 evaluation of extraction methods and immunolocalization. *Journal of Agricultural and Food*
615 *Chemistry*, 48, 1086-1095.

- 616 Elmaman, M., Al-Assaf, S., Phillips, G.O., & Williams, P.A. (2008). Studies on Acacia
617 exudate gums: Part VI. Interfacial rheology of *Acacia senegal* and *Acacia seyal*. *Food*
618 *Hydrocolloids*, 22, 682-689.
- 619 Escot, S., Feuillat, M., Dulau, L., & Charpentier, C. (2001). Release of polysaccharides by
620 yeasts and the influence of released polysaccharides on colour stability and wine astringency.
621 *Australian Journal of Grape and Wine research*, 7, 153-159.
- 622 Flindt, C., Al-Assaf, S., Phillips, G.O., & Williams, P.A. (2005). Studies on Acacia exudate
623 gums: Part V. Structural features of *Acacia seyal*. *Food Hydrocolloids*, 19, 687-701.
- 624 Gerbaud, V., Gabas, N., Blouin, J., & Crachereau, J.C. (2010). Study of wine acid salt
625 stabilization by addition of carboxymethylcellulose (CMC) : comparison with the « protective
626 colloids » effect. *Journal International des Sciences de la Vigne et du Vin*, 44, 231-242.
- 627 Gordillo, B., Cejudo-Bastante, M.J., Rodríguez-Pulido, F.J., Jara-Palacios, M.J., Ramírez-
628 Pérez, P., González-Miret, M.L., & Heredia, F.J. (2014) Impact of adding white pomace to
629 red grapes on the phenolic composition and color stability of syrah wines from a warm
630 climate. *Journal of Agricultural and Food Chemistry*, 62, 2663-2671.
- 631 Grein-Iankovski, A., Ferreira, J.G.L., Orth, E.S., Sierakowski, M.R., Cardoso, M.B., Simas,
632 F.F., & Riegel-Vidotti, I.C. (2018). A comprehensive study of the relation between structural
633 and physical chemical properties of acacia gums. *Food Hydrocolloids*, 85, 167-175.
- 634 Guadalupe, Z., Palacios, A., & Ayestaran, B. (2007). Maceration enzymes and
635 mannoproteins: A possible strategy to increase colloidal stability and color extraction in red
636 wines. *The Journal of Agricultural and Food Chemistry*, 55, 4854-4862.
- 637 Guadalupe, Z., & Ayestaran, B. (2008). Effect of commercial mannoprotein addition on
638 polysaccharide, polyphenolic, and color composition in red wines. *The Journal of*
639 *Agricultural and Food Chemistry*, 56, 9022-9029.


- 640 Idris, O.H.M., Williams, P.A., & Phillips, G.O. (1998). Characterization of gum from Acacia
641 senegal trees of different age and location using multidetection gel permeation
642 chromatography. *Food Hydrocolloids*, 12, 379-388.
- 643 International Oenological Codex: COEI-1-GOMARA, 2000.
- 644 Islam, A., Phillips, G., Sljivo, A., Snowden, M., & Williams, P. (1997). A review of recent
645 developments on the regulatory, structural and functional aspects of gum Arabic. *Food*
646 *Hydrocolloids*, 11, 493-505.
- 647 Lamport, D.T.A., & Varnai, P. (2013). Periplasmic arabinogalactan glycoproteins act as a
648 calcium capacitor that regulates plant growth and development. *New Phytologist*, 197, 58-64.
- 649 Li, X., Zhang, H., Jin, Q., & Cai, Z. (2018). Contribution of arabinogalactan protein to the
650 stabilization of single-walled carbon nanotubes in aqueous solution of gum Arabic. *Food*
651 *Hydrocolloids*, 78, 55-61.
- 652 Liang, M., Liu, R., Qi, W., Su, R., Yu, Y., Wang, L., & He, Z. (2013). Interaction between
653 lysozyme and procyanidin: Multilevel structural nature and effect of carbohydrates. *Food*
654 *Chemistry*, 138, 1596-1603.
- 655 Lopez-Torrez, L., Nigen, M., Williams, P., Doco, T., & Sanchez, C. (2015). Acacia *senegal*
656 vs. Acacia *seyal* gums – Part 1: Composition and structure of hyperbranched plant exudates.
657 *Food Hydrocolloids*, 51, 41-53.
- 658 Luck, G., Liao, H., Murray, N.J., Grimmer, H.R., Warminski, E.E., Williamson, M.P., Lilley,
659 T.H., & Haslam, E. (1994). Polyphenols, astringency and proline-rich proteins.
660 *Phytochemistry*, 37, 357-371.
- 661 Mahendran, T., Williams, P.A., Phillips, G.O., Al-Assaf, S., & Baldwin, T.C. (2008). New
662 insights into the structural characteristics of the arabinogalactan-protein (AGP) fraction of
663 gum Arabic. *Journal of Agricultural and Food Chemistry*, 56, 9269-9276.

- 664 Mateus, N., Carvalho, E., Luis, C., & de Freitas, V. (2004). Influence of the tannin structure
665 on the disruption effect of carbohydrates on protein-tannin aggregates. *Analytica Chimica*
666 *Acta*, 513, 135-140.
- 667 Mejia-Tamayo, V., Nigen, M., Apolinar-Valiente, R., Doco, T., Williams, P., Renard, D., &
668 Sanchez, C. (2018). Flexibility and hydration of amphiphilic hyperbranched arabinogalactan-
669 protein from plant exudate: a volumetric perspective. *Colloids and Interface*, 2, 1-24.
- 670 Mhinzi, G.S. (2003). Intra-species variation of the properties of gum exudates from *Acacia*
671 *senegal var. senegal* and *Acacia seyal var. fistula* from Tanzania. *Bulletin of the Chemical*
672 *Society of Ethiopia*, 17, 67-74.
- 673 Osman, M.E., Menzies, A.R., Williams, P.A., Phillips, G.O., & Baldwin, T.C. (1993). The
674 molecular characterization of the polysaccharide gum from *Acacia senegal*. *Carbohydrate*
675 *Research*, 246, 303-318.
- 676 Pérez-Lamela, C., García-Falcón, M.S., Simal-Gándara, J., & Orriols-Fernández,
677 I. (2007). Influence of grape variety, wine system and enological treatments on the colour
678 stability of young red wines. *Food Chemistry*, 101, 601-606.
- 679 Poncet-Legrand, C., Doco, T., Williams, P., & Vernhet, A. (2007). Inhibition of grape seed
680 tannin aggregation by wine mannoproteins: effect of polysaccharide molecular weight.
681 *American Journal of Enology and Viticulture*, 58, 87-91.
- 682 Randall, R.C., Phillips, G.O., & Williams, P.A. (1988). The role of the proteinaceous
683 component on the emulsifying properties of gum arabic. *Food Hydrocolloids*, 2, 131-140.
- 684 Randall, R.C., Phillips, G.O., & Williams, P.A. (1989). Fractionation and characterization of
685 gum from *Acacia senegal*. *Food Hydrocolloids*, 3, 65-75.
- 686 Ray, A.K., Bird, P.B., Iacobucci, A.I., & Clark, B.C. (1995). Functionality of gum Arabic.
687 Fractionation, characterization and evaluation of gum fractions in citrus oil emulsions and
688 model beverages. *Food Hydrocolloids*, 9, 123-131.

- 689 Renard, D., Lavenant-Gourgeon, L., Ralet, M.C., & Sanchez C. (2006). Acacia *senegal* gum:
690 continuum of molecular species differing by their protein to sugar ratio, molecular weight,
691 and charges. *Biomacromolecules*, 7, 2637-2649.
- 692 Renard, D., Garnier, C., Lapp, A., Schmitt, C., & Sanchez, C. (2012). Structure of
693 arabinogalactan-protein from Acacia gum: From porous ellipsoids to supramolecular
694 architectures. *Carbohydrate Polymers*, 90, 322-332.
- 695 Renard, D., Garnier, C., Lapp, A., Schmitt, C., & Sanchez C. (2013). Corrigendum to
696 “Structure of arabinogalactan-protein from Acacia gum: From porous ellipsoids to
697 supramolecular architectures” [*Carbohydrate Polymers* 90 (2012) 322-332]. *Carbohydrate*
698 *Polymers*, 97, 864-867.
- 699 Renard, D., Lavenant-Gourgeon, L., Lapp, A., Nigen, M., & Sanchez, C. (2014). Enzymatic
700 hydrolysis studies of arabinogalactan-protein structure from Acacia gum: The self-similarity
701 hypothesis of assembly from a common building block. *Carbohydrate Polymers*, 112, 648-
702 661.
- 703 Ribéreau-Gayon, P., Glories, Y., Maujean, A., & Dubourbieu, D. (2006). *Handbook of*
704 *Enology. Vol. 2. The Chemistry of Wine Stabilization and Treatments*, 2nd ed.; Wiley:
705 Chichester, UK.
- 706 Riou, V., Vernhet, A., Doco, T., & Moutounet, M. (2002). Aggregation of grape seed tannins
707 in model wine – effect of wine polysaccharides. *Food Hydrocolloids*, 16, 17-23.
- 708 Sanchez, C., Schmitt, C., Kolodziejczyk, E., Lapp, A., Gaillard, C., & Renard, D. (2008). The
709 Acacia gum arabinogalactan fraction is a thin oblate ellipsoid: A new model based on small-
710 angle neutron scattering and ab initio calculation. *Biophysical Journal*, 94, 629-639.
- 711 Schmitt, C., Sanchez, C., Thomas, F., & Hardy, J. (1999). Complex coacervation between β -
712 lactoglobulin and acacia gum in aqueous medium. *Food Hydrocolloids*, 13, 483-496.

- 713 Snowden, M.L., Phillips, G.O., & Williams, P.A. (1987). Functional characteristics of gum
714 arabic. *Food Hydrocolloids*, 1, 291-300.
- 715 Soares, S.I., Gonçalves, R.M., Fernandes, I., Mateus, N., & de Freitas, V. (2009). Mechanistic
716 approach by which polysaccharides inhibit α -amylase/procyanidin aggregation. *Journal of*
717 *Agricultural and Food Chemistry*, 57, 4352-4358.
- 718 Soares, S., Mateus, N., & de Freitas, V. (2012). Carbohydrates inhibit salivary proteins
719 precipitation by condensed tannins. *Journal of Agricultural and Food Chemistry*, 60, 3966-
720 3972.
- 721 Teissedre, P.L. (2012). Arabic Gum in Oenology. In: Kennedy J.F, Phillips G.O., Williams
722 P.A. (eds) Arabic gum. RSC Publishing, Cambridge.
- 723 Waters, E.J., Pellerin, P., & Brillouet, J-M. (1994). A wine arabinogalactan-protein that
724 reduces heat-induced wine protein haze. *Bioscience Biotechnology and Biochemistry*, 58, 43-
725 48.
- 726 Waters, E.J., Pellerin, P., & Brillouet, J-M. (1994a). A *Saccharomyces* mannoprotein that
727 protects wine from protein haze. *Carbohydrate polymers*, 23, 185-191.
- 728 Williams, P.A., & Phillips, G.O. (2000). Gum Arabic. In G.O. Phillips, & P.A. Williams
729 (Eds.), *Handbook of hydrocolloids*, Boca Raton: CRC Press.
- 730

731 Figure 1. Transmittance changes of hydro-alcoholic – mineral solutions without (A) and with
 732 0.09 (B) and 0.14 g.L⁻¹ (C) of *A. senegal* gum. The transmittance was registered at 25°C
 733 during 24h. Pictures of the solutions before and after 24h of kinetic are presented on the right.




734

735

736

737 Figure 2. (A) Colloid stability kinetics of hydro-alcoholic – mineral solutions containing
 738 various *A. senegal* gum concentrations: 0 (●), 0.05 (●), 0.08 (●), 0.09 (●), 0.10 (●), 0.11 (●)
 739 and 0.14 (●) g.L⁻¹. (B) Colloidal stability curve of hydro-alcoholic – mineral solution in
 740 presence of *A. senegal* gum after 24h of kinetic. The experiments were performed at 25°C.
 741 The line (figure 2B) is shown to guide the eyes. The experiments were triplicated.
 742




743

744

745

746

747 Figure 3. (A) Colloidal stability curves of hydro-alcoholic – mineral solutions in presence of
 748 *A. senegal* gum (AG) at pH 3.1 (○), 3.5 (○) and 4.0 (○) after 24h of kinetic. (B) Relationship
 749 between the electrophoretic mobility of *A. senegal* gum (AG) and its critical concentration in
 750 hydro-alcoholic – mineral solutions. The lines (figure 3A) are shown to guide the eyes. The
 751 experiments were triplicated.



752

753


754

755

756 Figure 4. Relationship between the concentration of Ca^{2+} in the hydro-alcoholic – mineral
757 solution and the critical stabilizing concentration in *A. senegal* gum ($[\text{AG}]_{\text{crit}}$). The
758 experiments were triplicated.

759

760



761

762

763

764

765

766

767

768

769


770

771

772

773

774 Figure 5. Colloidal stability curves of hydro-alcoholic – mineral solution in presence of A.
 775 *senegal* gum (○), HIC-F1 (○), HIC-F2 (○) and HIC-F3 (○) fractions after 24h of kinetic. The
 776 experiments were performed at 25°C. The lines are shown to guide the eyes. The inset figure
 777 is a zoom of the left part. The experiments were triplicated.
 778




779

780

781

782 Figure 6. Molar mass distribution (thick line) and normalized refractive index signal (thin
 783 line) of HIC-F1 (A) and *A. senegal* gum (B) before (black) and after (red) pronase treatment.
 784 Colloidal stability curves of hydro-alcoholic – mineral solution in presence of HIC-F1 (C) and
 785 *A. senegal* gum (D) before (○) and after (●) pronase treatment after 24h of kinetic. The lines
 786 (figures 6C and 6D) are shown to guide the eyes. The colloidal stability experiments were
 787 triplicated.
 788
 789




790

791

792 Figure 7. (A) Colloid stability kinetic of hydro-alcoholic – grape marc solutions according to
793 *A. senegal* gum (AG) concentration at 10°C. The concentrations of *A. senegal* gum were 0
794 (●), 0.01 (●), 0.025 (●), 0.05 (●), 0.1 (●) and 0.5 (●) g.L⁻¹. The colloidal stability kinetic of
795 the hydro-alcoholic – grape marc solution at 25°C without *A. senegal* gum is shown as a
796 control (●) (B) Colloidal stability curves of hydro-alcoholic – grape marc solution in presence
797 of *A. senegal* gum (○), HIC-F1 (○), HIC-F2 (○) and HIC-F3 (○) fractions after 48h of kinetic
798 at 10°C. (C) Colloidal stability curves of a young red wine in presence of *A. senegal* gum (○),
799 HIC-F2 (○) and HIC-F3 (○) fractions after 48h of kinetic at 10°C. The lines (figures 7B and
800 7C) are shown to guide the eyes. The inset figure is a zoom of the left part. The experiments
801 were triplicated.

802

803




804

805

806 Figure 8. Relationship between the critical concentrations of *A. senegal* gum (AG) and its
 807 HIC-fractions determined in hydro-alcoholic – mineral solution and those determined in
 808 hydro-alcoholic grape marc solution (A) and young red wine (B). The experiments were
 809 triplicated.

810



811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826 Supplementary table 1. Biochemical and structural properties of *Acacia senegal* gum (AG),

827 and its HIC fractions.

828

	AG	HIC-F1	HIC-F2	HIC-F3
Total dry matter (mg.g ⁻¹)	893.4	921.6	926.2	921.9
Sugars (mg.g ⁻¹)	944.4	961.3	918.3	813.0
Galactose	363.6	374.9	315.9	270.7
Arabinose	281.4	257.6	326.9	311.4
Rhamnose	120.9	120.2	125.8	113.0
Glucuronic acid	168.1	195.1	143.3	111.4
4-O-Me-Glucuronic acid	9.4	13.5	5.5	5.7
Proteins (mg.g ⁻¹)	21.5	4.9	63.1	137.7
Ash (mg.g ⁻¹)	34.1	30.5	19.3	49.3
M _w (g.mol ⁻¹)	6.8 × 10 ⁵	3.5 × 10 ⁵	1.5 × 10 ⁶	1.6 × 10 ⁶
High M _w content (%) ^a	14	3	73	56

829

830 ^a The high M_w corresponded to macromolecules with a M_w upper than 10⁶ g.mol⁻¹.

831

832

Highlights.

- AGPs from *Acacia senegal* gum prevent the coloring matter precipitation
- The more AGPs are rich in proteins, the more their stabilizing efficiency are
- Stabilizing properties of AGPs are correlated in “synthetic” solutions and red wine
- In “synthetic” mineral solution, AGPs avoid the precipitation by Ca²⁺ binding

ACCEPTED MANUSCRIPT

Comment citer ce document :

Nigen, M., Apolinar Valiente, R., Iturmendi, N., Williams, P., Doco, T., Moine, Massot, Jaouen, I., Sanchez, C. (2019). The colloidal stabilization of young red wine by *Acacia senegal* gum: The involvement of the protein backbone from the protein-rich arabinogalactan-proteins. *Food Hydrocolloids*. 97. . DOI : 10.1016/i.foodhvd.2019.105176