



MAMP and PRR Interaction at the Centre of Probiotic/Host Interaction

Gwénaël Jan

► To cite this version:

Gwénaël Jan. MAMP and PRR Interaction at the Centre of Probiotic/Host Interaction. 13. International Scientific Conference on Probiotics, Prebiotics, Gut Microbiota and Health – IPC2019, Jun 2019, Prague, Czech Republic. , 2019. hal-02164170

HAL Id: hal-02164170

<https://hal.science/hal-02164170>


Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MAMPs and PRRs interaction at the center of probiotic/host interaction



Gwénaël JAN
UMR 1253 STLO
INRA Agrocampus Ouest, Rennes, France






Microbiota / host interaction, an old story




Coevolution and dependence



Davenport et al. BMC Biology (2017)

→ Our microbiota evolved with us




→ And we adapted to its products




A key role of the first colonizers



Good bugs



Main inhabitants and main actors



Milani et al. Microbiol.Mol.Biol.Rev. (2017)

Turroni et al. Cell.Mol.Life Sci. (2014)

Dramatic effects of the absence

Bifidobacterium longum subspecies *infantis*: champion colonizer of the infant gut

Mark A. Underwood^{1,2}, J. Bruce German^{3,4}, Carlito B. Lebrilla^{3,4} and David A. Mills^{2,3,5}

Oligosaccharides are abundant in human milk. Production of these highly diverse structures requires significant energy expenditure by the mother and yet these human milk oligosaccharides offer no direct nutritive value to her infant. A primary function of human milk oligosaccharides is to shape the intestinal microbiota with the *Bifidobacterium* spp.

B. infantis displays anti-inflammatory activity

result of a large repertoire of bacterial genes encoding an array of glycosidases and oligosaccharide transporters not found in other bacterial species. *In vitro*, *B. infantis* grows better than other bacterial strains in the presence of human milk oligosaccharides, displays anti-inflammatory activity in premature

hypothesis that human milk contained "bifidogenic factors" that stimulated the growth of these bifidobacteria (1).

Prebiotics are dietary supplements that promote health benefits by stimulating the growth and/or activity in the gut lumen of commensal microbes (ideally without stimulating potential pathogens). Probiotics are live microorganisms and a variety of mechanisms and a variety of volition of

a collection of complex prebiotic oligosaccharides found in abundance in human milk and a single bacterial subspecies, *Bifidobacterium longum* subsp. *infantis* (*B. infantis*) unique in its capacity to consume these oligosaccharides. Note that the species *B. longum* has two subspecies: *B. longum* subsp. *infantis*

B. infantis decreases the risk of necrotizing enterocolitis

rotating enterocolitis in premature infants. Colonization with *B. infantis* is also associated with increased vaccine responses. Probiotic organisms have historically been selected based on ease of production and stability. The advantages of *B. infantis*, selected through coevolution with human milk glycans, present an opportunity for focused manipulation of the infant intestinal microbiota.

The colonization of the fetal gut begins *in utero* with swallowing of amniotic fluid. At that point, infants begin a lifelong relationship!

community of microflora. *B. infantis* increases vaccine response

gut microbiome) and presumably use negative consequences of the phenotype of the gut microbiota occur with rupture of the fetal membranes, birth, initiation of feeding, addition of solid foods, weaning, and interventions such as antibiotics, acid-suppression, and prebiotic or probiotic dietary supplements. The predominance of "bifid" microbes in the stools of healthy infants was described more than 100 years ago, prompting the

INTESTINAL MICROBIOTA

Humans stand at the end of the long evolution of mammals, with differences in milk conspicuous for the volume, number of structures, and complexity of milk oligosaccharides (Figure 1) (2–4). Human milk oligosaccharides (HMOs) are the third largest solid component of human milk (after lactose and fat) even in times of famine (5), and yet these free glycans are not digestible by the infant as the human gut does not produce the glycosidases necessary to cleave the HMO linkages. The obvious evolutionary question is: What benefit is provided to the infant that justifies the mother's tremendous expenditure of energy?

complex molecules to answer this question. The capacity of select gut microbes to deconstruct and consume HMOs (6,7). Among multiple microbial species studied, only two genera, *Bifidobacterium* and *Bacteroides*, are able to comprehensively utilize HMOs as a primary food source (Table 1) (8,9). This relative resistance to microbial consumption allows the HMOs to arrive intact in the distal small bowel and the colon where the largest numbers of commensal bacteria thrive.

¹Department of Pediatrics, University of California, Davis, Sacramento, California; ²Foods for Health Institute, University of California, Davis, Davis, California; ³Department of Food Science and Technology, University of California, Davis, Davis, California; ⁴Department of Chemistry, University of California, Davis, Davis, California; ⁵Department of Viticulture and Enology, University of California, Davis, Davis, California. Correspondence: Mark A. Underwood (mark.underwood@ucdmc.ucdavis.edu)

Received 14 April 2014; accepted 4 August 2014; advance online publication 5 November 2014; doi:10.1038/px.2014.156

Copyright © 2015 International Pediatric Research Foundation, Inc.

Volume 77 | Number 1 | January 2015

Pediatric RESEARCH 229



The Journal of Clinical Investigation

RESEARCH ARTICLE

Commensal *Propionibacterium* strain UF1 mitigates intestinal inflammation via Th17 cell regulation

Natalia C. Gómez¹, Daniel J. Sartor¹, Michael J. Owen³, Josef Neu⁴, and Sourour Mohamadzadeh^{1,2}

¹Department of Infectious Diseases and Immunology, ²Division of Gastroenterology, Hepatology & Nutrition, Department of Medicine, ³Department of Physiological Sciences, ⁴Division of Neonatology, Department of Pediatrics, Loyola University Chicago, Maywood, Illinois. *Propionibacterium* mitigate induced necrotizing enterocolitis

Consul Propionibacterium regulate Th17 cells leading to NEC, which remains a mystery. It is known that this diminution correlates with alterations in the gut microbiota, particularly enrichment of *Propionibacterium* species. Transfaunation of microbiota from HBM-fed preterm infants or a newly identified and cultured *Propionibacterium* strain, P. UF1, to germfree mice conferred protection against pathogen infection and correlated with profound increases in intestinal Th17 cells. The induction of Th17 cells was dependent on bacterial dihydroxyacetone acetyltransferase (DlaT), a major protein expressed on the P. UF1 surface layer (S-layer). Binding of P. UF1 to its cognate receptor, SIGNR1, on dendritic cells resulted in the regulation of intestinal phagocytes. Importantly, transfer of P. UF1 profoundly mitigated induced NEC-like injury in neonatal mice. Together, these results mechanistically elucidate the protective effects of HBM and P. UF1-induced immunoregulation, which safeguard against proinflammatory diseases, including NEC.



Healing effect of probiotics



nature publishing group

Clinical Investigation

Article



Microbiology and Molecular
Biology Reviews®

Probiotics (*Lactobacillus acidophilus* and *Bifidobacterium bifidum*) prevent NEC in VLBW infants fed breast milk but not formula


Andreas Repa¹, Margarita Thanhaeuser¹, David Endress², Michael Weber³, Alexandra Kreissl¹, Christoph Binder¹, Angelika Berger¹ and Nadja Haiden¹

The First Microbial Colonizers of the Human Gut: Composition, Activities, and Health Implications of the Infant Gut Microbiota



Christian Milani,^a Sabrina Duranti,^a Francesca Bottacini,^b Eoghan Casey,^b Francesca Turroni,^{a,c} Jennifer Mahony,^b Clara Belzer,^d Susana Delgado Palacio,^e Silvia Arboleya Montes,^e Leonardo Mancabelli,^a Gabriele Andrea Lugli,^a Juan Miguel Rodriguez,^f Lars Bode,^g Willem de Vos,^{d,h} Miguel Gueimonde,^e Abelardo Margolles,^e Douwe van Sinderen,^b Marco Ventura^{a,c}



The gut microbiota and inflammation



Pro- and anti-inflammatory bugs








Microbial Associated Molecular Patterns / Pattern Recognition Receptors






MAMPs : Gram-positive surface



→ Decorated with many motives

→ Macromolecules and glycobiome

PRRs: host receptors




Lebeau et al. Nature Reviews (2010)


→ Immune or epithelial cells

Nature Reviews | Microbiology

→ Toll-like receptors (TLRs), NOD-like receptors (NLRs), C-type lectin receptors (CLRs)



Nature Reviews | Microbiology



Bailey et al. Front. Immunol. (2014)

→ Different immunomodulatory effects

Example of *Lactobacillus salivarius* PG

Inflammatory bowel disease

Anti-inflammatory capacity of selected lactobacilli in experimental colitis is driven by NOD2-mediated recognition of a specific peptidoglycan-derived muropeptide

Elise Macho Fernandez,¹ Véronique Valenti,¹ Christoph Rockel,² Corinna Hermann,² Bruno Pot,¹ Ivo Gomperts Boneca,^{3,4} Corinne Granette¹

Results The protective capacity of strain *Lactobacillus salivarius* Ls33 was correlated with a local IL-10 production and was abolished in *Nod2*-deficient mice. PGN purified from Ls33 rescued mice from colitis in an IL-10-dependent manner and favoured the development of CD103⁺ DCs and CD4⁺Foxp3⁺ regulatory T cells. In







Figure 4 Peptidoglycan (PGN) from selected strains rescues mice from trinitrobenzene sulfonic acid (TNBS)-induced colitis. BALB/c mice were injected intraperitoneally with phosphate-buffered saline (PBS; n=10), 50 µg (n=5) or 200 µg (n=10) of PGN from Ls33 (Ls33 PGN) or NCFM (NCFM PGN) before induction of colitis. (A) Wallace score evaluated 2 days after induction of colitis. (B) Ameho scores illustrated with representative sections of the distal colon of healthy mice or mice with TNBS-induced colitis, injected intraperitoneally with PBS or 200 µg of PGNs (original magnification, ×20). Data represent mean values with the SEM (*p<0.05; **p<0.01; ***p<0.001 vs PBS-treated mice).

Example of *Lactobacillus plantarum* LTA



Mutation of Dlt reduces D-Ala in TA and makes *L. plantarum* induce IL-10 in PBMCs




TA from WT induce pro-inflammatory cytokines, while TA from Dlt mutant does not

Enhanced antiinflammatory capacity of a *Lactobacillus plantarum* mutant synthesizing modified teichoic acids

Corinne Grangette*,†, Sophie Nutten*,‡, Emmanuelle Palumbo§, Siegfried Morath¶, Corinna Hermann¶, Joelle Dewulf*, Bruno Pot*, Thomas Hartung*, Pascal Hols§, and Annick Mercenier*,‡



*Laboratoire de Bactériologie des Écosystèmes, Institut Pasteur de Lille-Institut de Biologie de Lille, 59019 Lille Cedex, France; †Unité de Génétique, Institut des Sciences de la Vie, Université Catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium; and ¶Department of Biochemical Pharmacology, Konstanz University, 78457 Konstanz, Germany

Communicated by Todd R. Klaenhammer, North Carolina State University, Raleigh, NC, May 18, 2005 (received for review November 19, 2004)



Mutation of Dlt mutant protects more than WT *plantarum* against colitis

Example of *Lactobacillus rhamnosus* pili




The pili- mutant induces more inflammatory IL-8



Functional Analysis of *Lactobacillus rhamnosus* GG Pili in Relation to Adhesion and Immunomodulatory Interactions with Intestinal Epithelial Cells

Sarah Lebeer,^{a,b} Ingmar Claes,^a Hanne L. P. Tytgat,^a Tine L. A. Verhoeven,^a Erya Marien,^a Ingemar von Ossowski,^c Justus Reunanan,^c Alri Palva,^c Willem M. de Vos,^{c,d} Stigrid C. J. De Keersmaecker,^a and Jos Vanderleyden^a


^aCentre of Microbial and Plant Genetics, K.U. Leuven, Leuven, Belgium; ^bDepartment of Bioscience Engineering, University of Antwerp, Antwerp, Belgium; ^cDepartment of Veterinary Biosciences, University of Helsinki, Helsinki, Finland; ^dLaboratory of Microbiology, Wageningen University, Wageningen, The Netherlands^d



Probiotic Gut Microbiota Isolate Interacts with Dendritic Cells via Glycosylated Heterotrimeric Pili

Hanne L. P. Tytgat^{1,2,3}, Nienke H. van Teijlingen⁴, Ruby May A. Sullan^{5a}, François P. Douillard⁶, Pia Rasinkangas⁶, Marcel Messing⁷, Justus Reunanan⁶, Reetta Satokari⁶, Jos Vanderleyden¹, Yves F. Dufrêne³, Teunis B. H. Geijtenbeek⁴, Willem M. de Vos^{3,6,7}, Sarah Lebeer^{1,2,*}

Example of *Bifidobacterium bifidum* CSPS




SCIENCE IMMUNOLOGY | RESEARCH ARTICLE




MICROBIOME

Cell surface polysaccharides of *Bifidobacterium bifidum* induce the generation of Foxp3⁺ regulatory T cells


Ravi Verma¹, Changhon Lee², Eun-Ji Jeun², Jaeu Yi², Kwang Soon Kim¹, Ambarnil Ghosh¹, Seohyun Byun², Choong-Gu Lee¹, Hye-Ji Kang¹, Gi-Cheon Kim¹, Chang-Duk Jun³, Gwenaël Jan⁴, Chang-Hee Suh⁵, Ju-Yang Jung⁵, Jonathan Sprent^{2,6}, Dipayan Rudra^{1,2}, Cristina De Castro^{7,8}, Antonio Molinaro^{8,9}, Charles D. Surh^{1,2}, Sin-Hyeog Im^{1,2,*}




B. bifidum monocolonization enhances Treg population in the colon Lamina Propria




B. bifidum surface PS mediates Tcell induction




Example of *Lactobacillus acidophilus* Slp




L. acidophilus induces IL-10 in immature dendritic cells





L. acidophilus binds to DC-SIGN receptors, not the slpA- mutant.



SlpA mediates *L. acidophilus* binding to DC-SIGN receptors



Purified SlpA binds to DC-SIGN receptors



S layer protein A of *Lactobacillus acidophilus* NCFM regulates immature dendritic cell and T cell functions


Sergey R. Konstantinova^{a,b,1}, Hauke Smidt^a, Willem M. de Vos^{a,c}, Sven C. M. Bruijns^b, Satwinder Kaur Singh^b, Florence Valence^d, Daniel Molle^d, Sylvie Lortal^d, Eric Altermann^e, Todd R. Klaenhammer^{f,2}, and Yvette van Kooyk^{b,2}

^aLaboratory of Microbiology, Wageningen University, Dreijenplein 10, 6703 HB, Wageningen, The Netherlands; ^bVU University Medical Center Amsterdam, v.d. Boechorstraat 7, 1081 BT Amsterdam, The Netherlands; ^cDepartment of Basic Veterinary Sciences, Helsinki University, FIN-00014, Helsinki, Finland;


^dInstitut National de la Recherche Agronomique, 65 rue de Saint Brieuc, 35000 Rennes, France; and ^eDepartment of Microbiology and ^fSoutheast Dairy Foods Research Center, North Carolina State University, Raleigh, NC 27695

Contributed by Todd R. Klaenhammer, October 15, 2008 (sent for review July 21, 2008)




Propionibacterium freudenreichii SlpB



Untreated Guanidine treated








Surface protein SlpB induces production of IL-10 in PBMC cells.



Inactivation of SlpB suppresses IL-10 induction

P. freudenreichii cheese in a murine model of colitis



5



Food/commensal/probiotic bacteria have various MAMPs



Various immune responses, may be strain-dependent



We can make fermented foods with these strains



Houem Rabah

Fillipe Luis Rosa do Carmo

Song Huang

Floriane Gaucher
20th June, 11H55,
Probiotics for health

