



HAL
open science

Combinatorial descriptions of the crystal structure on certain PBW bases

Ben Salisbury, Adam Schultze, Peter Tingley

► **To cite this version:**

Ben Salisbury, Adam Schultze, Peter Tingley. Combinatorial descriptions of the crystal structure on certain PBW bases. 28-th International Conference on Formal Power Series and Algebraic Combinatorics, Simon Fraser University, Jul 2016, Vancouver, Canada. 10.46298/dmtcs.6377 . hal-02166334

HAL Id: hal-02166334

<https://hal.science/hal-02166334>

Submitted on 26 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Combinatorial descriptions of the crystal structure on certain PBW bases

Ben Salisbury^{1†}, Adam Schultze^{2‡}, and Peter Tingley^{3§}

¹*Department of Mathematics, Central Michigan University, Mount Pleasant, MI 48859, USA*

²*Department of Mathematics and Statistics, State University of New York at Albany, Albany, NY 12222, USA*

³*Department of Mathematics and Statistics, Loyola University of Chicago, Chicago, IL 60660, USA*

Abstract. Lusztig’s theory of PBW bases gives a way to realize the crystal $B(\infty)$ for any simple complex Lie algebra where the underlying set consists of Kostant partitions. In fact, there are many different such realizations, one for each reduced expression for the longest element of the Weyl group. There is an algorithm to calculate the actions of the crystal operators, but it can be quite complicated. For ADE types, we give conditions on the reduced expression which ensure that the corresponding crystal operators are given by simple combinatorial bracketing rules. We then give at least one reduced expression satisfying our conditions in every type except E_8 , and discuss the resulting combinatorics. Finally, we describe the relationship with more standard tableaux combinatorics in types A and D .

Résumé. La théorie de Lusztig de bases PBW donne un moyen de réaliser le cristal $B(\infty)$ pour toute algèbre de Lie complexe-simple où l’ensemble sous-jacent est constitué de partitions Kostant. En fait, il existe de nombreuses réalisations différentes, une pour chaque expression réduite de l’élément le plus long du groupe de Weyl. Il existe un algorithme pour calculer les actions des opérateurs de cristal, mais elle peut être assez compliqué. Pour les types ADE, nous donnons des conditions sur l’expression réduite qui assurent que les opérateurs de cristal correspondants sont donnés par une simple règle de d’appariement combinatoire. Nous donnons ensuite au moins une expression réduite satisfaisant nos conditions pour chaque type, sauf E_8 , et discutons la combinatoire résultant. Enfin, nous décrivons les relations avec la combinatoire de tableaux standards dans les types A et D .

Keywords. canonical basis, crystal basis, PBW basis, Kostant partition, multisegment, tableaux

1 Introduction

We consider the crystal $B(\infty)$ for a simple Lie algebra over \mathbf{C} of simply-laced type. This is a combinatorial object that contains a great deal of information about the algebra and its finite-dimensional representations. It is usually defined by a complicated algebraic construction, but it can often be realized in quite simple ways.

[†]Email: ben.salisbury@cmich.edu. Partially supported by CMU Early Career grant #C62847.

[‡]Email: aschultze@albany.edu. Partially supported by NSF grant DMS-1265555.

[§]Email: ptingley@luc.edu. Partially supported by NSF grant DMS-1265555.

Lusztig's early algebraic construction of the canonical basis of the quantum group $U_q(\mathfrak{g})$ (see [13, Chapters 41 and 42] and references therein) can be interpreted as giving a number of parameterizations of $B(\infty)$, one for each reduced expression for the longest word w_0 in the Weyl group. For each of these realizations, at least one of the crystal operators is very simple, but others may be complicated. However, Lusztig explicitly describes how the realizations are related for reduced expressions that differ by a braid move. This gives a way to realize the whole crystal: Fix a reduced expression for w_0 . Then an element of the crystal is simply a PBW monomial. To apply a crystal operator, modify the reduced expression via a sequence of braid moves until that operator is simple, then apply the operator, then modify it back. For more on this point of view, see [19].

This procedure is algorithmic, but can be complicated. In type A_n , there is a simpler realization using multisegments. As discussed in [2], this is precisely Lusztig's crystal structure for the reduced expression $w_0 = (s_1 s_2 \cdots s_n)(s_1 s_2 \cdots s_{n-1}) \cdots (s_1 s_2) s_1$. This is not true for other reduced expressions (unless they are related to this one by trivial braid moves).

In the current work, we give a set of conditions on a reduced expression that ensure Lusztig's crystal structure on Kostant partitions is given by a simple bracketing procedure, similar to the type A_n structure on multisegments. There is at least one such reduced expression in every simply-laced type except E_8 .

We then discuss the type A_n and D_n situation in some detail. There is already a nice combinatorial realization of $B(\infty)$ in these types, due to J. Hong and H. Lee [4], where the underlying set consists of marginally large tableaux. We explicitly describe the unique crystal isomorphism between marginally large tableaux and Kostant partitions in these types. The isomorphism in type A_n is essentially the same as the one given in [2], and naturally factors through the realization of $B(\infty)$ in terms of multisegments. The type D_n isomorphism is new; in particular, it does not agree with the bijection used in [10].

The type D_n situation has one interesting new feature: it is not possible to find a reduced expression that is adapted both for calculating both the ordinary crystal operators and the $*$ -operators (i.e., the crystal operators twisted by Kashiwara's involution). Thus, while the ordinary crystal operators can be described with a bracketing procedure, the $*$ -operators are more complicated. This may explain why the embeddings $B(\lambda) \hookrightarrow B(\infty)$ of the finite crystals into the infinity crystal seem more difficult to understand in type D_n , since the conditions describing which elements of $B(\infty)$ are present in a fixed $B(\lambda)$ reference the $*$ -crystal structure. See [5] for some recent work discussing these embeddings.

Reineke [15] has also given an explicit description of the crystal operators on Lusztig's PBW basis for certain reduced expressions of w_0 , using quiver representation theory. There the reduced expression must be adapted to an orientation of the Dynkin diagram. Our construction works nicely when the reduced expression is " i -semi-adapted for all i " (see §3). These conditions do not coincide. For instance, the reduced expression we use in §4.2 for type D_n is not adapted to any orientation of the Dynkin diagram.

The results here will appear with full proofs in two upcoming papers. See [17] for the general results and [18] for the connection to marginally large tableaux in type D_n .

2 Background

2.1 General

Let \mathfrak{g} be a complex-simple Lie algebra of type ADE. Let I be the index set of \mathfrak{g} , $A = (a_{ij})$ the Cartan matrix, $\{\alpha_i\}_{i \in I}$ the positive simple roots, $\{\alpha_i^\vee\}_{i \in I}$ the simple coroots, Φ^+ the set of positive roots, P the weight lattice, P^\vee the dual weight lattice, W the Weyl group with longest element w_0 , and $\{s_i\}_{i \in I}$ the

generating reflections. Let $(-|-)$ denote a symmetric bilinear form on P satisfying $(\alpha_i|\alpha_j) = a_{ij}$ and let $\langle -, - \rangle: P^\vee \times P \rightarrow \mathbf{Z}$ be the canonical pairing. Let $R(w_0)$ denote the set of reduced expressions for the longest element w_0 of the Weyl group.

Let $U_q(\mathfrak{g})$ be the quantized universal enveloping algebra of \mathfrak{g} , which is a $\mathbf{Q}(q)$ -algebra generated by E_i, F_i , and q^h , for $i \in I$ and $h \in P^\vee$, subject to certain relations (see, for example, [3]). Let $U_q^-(\mathfrak{g})$ be the subalgebra generated by the F_i 's.

2.2 Crystals

Let e_i, f_i be the Kashiwara operators on $U_q^-(\mathfrak{g})$ defined in [7], $\mathcal{A} \subset \mathbf{Q}(q)$ be the subring of functions regular at $q = 0$ and define $L(\infty)$ to be the \mathcal{A} -lattice spanned by

$$S = \{f_{i_1} f_{i_2} \cdots f_{i_t} \cdot 1 \in U_q^-(\mathfrak{g}) : t \geq 0, i_k \in I\}.$$

Theorem/Definition 2.1 ([7])

1. Let $\pi: L(\infty) \rightarrow L(\infty)/qL(\infty)$ be the natural projection and set $B(\infty) = \pi(S)$. Then $B(\infty)$ is a \mathbf{Q} -basis of $L(\infty)/qL(\infty)$.
2. The operators e_i and f_i act on $L(\infty)/qL(\infty)$ for each $i \in I$. Moreover, $e_i: B(\infty) \rightarrow B(\infty) \sqcup \{0\}$ and $f_i: B(\infty) \rightarrow B(\infty)$ for each $i \in I$. For $b, b' \in B(\infty)$, we have $f_i b = b'$ if and only if $e_i b' = b$.

2.3 Reduced expressions, convex orders, and PBW bases

Definition 2.2 A total order \prec on Φ^+ is called convex if, for all triples of roots β, β', β'' with $\beta' = \beta + \beta''$, we have either $\beta \prec \beta' \prec \beta''$ or $\beta'' \prec \beta' \prec \beta$.

Theorem 2.3 ([14]) There is a one-to-one correspondence between $R(w_0)$ and convex orders on Φ^+ . Explicitly, if $w_0 = s_{i_1} s_{i_2} \cdots s_{i_N}$, then the corresponding convex order \prec is defined by

$$\beta_1 = \alpha_{i_1} \prec \beta_2 = s_{i_1}(\alpha_{i_2}) \prec \cdots \prec \beta_N = s_{i_1} s_{i_2} \cdots s_{i_{N-1}}(\alpha_{i_N}).$$

For $c \in \mathbf{Z}_{>0}$ define

$$F_i^{(c)} := \frac{F_i^c}{[c]!}, \text{ where } [c]! := \prod_{j=1}^c \frac{q^j - q^{-j}}{q - q^{-1}}.$$

Given $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$ and $\mathbf{c} = (c_\beta^{\mathbf{i}} \in \mathbf{Z}_{\geq 0}^N : \beta \in \Phi^+)$, define

$$F_{\mathbf{i}}^{\mathbf{c}} = F_{\beta_1}^{(c_{\beta_1}^{\mathbf{i}})} F_{\beta_2}^{(c_{\beta_2}^{\mathbf{i}})} \cdots F_{\beta_N}^{(c_{\beta_N}^{\mathbf{i}})} \text{ where } F_{\beta_k}^{(c_{\beta_k}^{\mathbf{i}})} = T_{i_1} T_{i_2} \cdots T_{i_{k-1}}(F_{i_k}^{(c_{\beta_k}^{\mathbf{i}})}), \tag{2.4}$$

and T_i is the Lusztig automorphism of $U_q(\mathfrak{g})$ defined in [13, §37.1.3] (where it is denoted $T_{i,-1}''$). Then the set

$$\mathcal{B}_{\mathbf{i}} = \{F_{\mathbf{i}}^{\mathbf{c}} : \mathbf{c} \in \mathbf{Z}_{\geq 0}^N\} \tag{2.5}$$

forms a $\mathbf{Q}(q)$ -basis of $U_q^-(\mathfrak{g})$ called the PBW basis.

Theorem 2.6 ([16]) For $\mathbf{i} \in R(w_0)$, $\text{Span}_{\mathcal{A}}(\mathcal{B}_{\mathbf{i}}) = L(\infty)$ and $\mathcal{B}_{\mathbf{i}} + qL(\infty) = B(\infty)$.

Since $\mathcal{B}_i + qL(\infty) = B(\infty)$ for every $\mathbf{i} \in R(w_0)$, there is a parametrization of the elements of $B(\infty)$ dependent on a chosen \mathbf{i} . Given $b \in B(\infty)$, denote the bijection which takes $b \mapsto F_i^c \text{ mod } qL(\infty) \mapsto \mathbf{c}$ by \mathbf{c}^i , and call $\mathbf{c} = \mathbf{c}^i(b)$ the *i-Lusztig data* of b . The crystal structure on $B(\infty)$ can be interpreted using these parameterizations.

Proposition 2.7 ([1, 13]) Fix some $i \in I$ and let $\mathbf{i} \in R(w_0)$ be such that $i_1 = i$. Suppose $b \in B(\infty)$ and $\mathbf{c}^i(b) = (c_1, c_2, \dots, c_N)$. Then $\mathbf{c}^i(f_i b) = (c_1 + 1, c_2, \dots, c_N)$. Moreover, if $c_1 = 0$, then $e_i b = 0$. Otherwise, $\mathbf{c}^i(e_i b) = (c_1 - 1, c_2, \dots, c_N)$.

If \mathbf{i} is such that $i_1 = i$, then the operator f_i is easily understood on $B(\infty)$, as indexed by \mathcal{B}_i . To understand the whole crystal structure, it is necessary to understand how the Lusztig data $\mathbf{c}^i(b)$ are related for different \mathbf{i} . All reduced expressions for w_0 are related by a sequence of braid moves, so it is enough to understand what happens to $\mathbf{c}^i(b)$ when \mathbf{i} changes by a single braid move. The following is due to Lusztig.

Lemma 2.8 ([12, §2.1]) Fix $\mathbf{i} \in R(w_0)$ and $b \in \mathcal{B}_i$. Let $\{\beta_1 \prec \dots \prec \beta_N\}$ be the order on Φ^+ corresponding to \mathbf{i} .

1. If $\mathbf{i}' \in R(w_0)$ is such that \mathbf{i} and \mathbf{i}' differ by replacing two consecutive entries (i_k, i_{k+1}) , such that $a_{i_k, i_{k+1}} = 0$ by (i_{k+1}, i_k) , then $\mathbf{c}_{\beta}^{\mathbf{i}'}(b) = \mathbf{c}_{\beta}^{\mathbf{i}}(b)$ for all β (although the order has changed).
2. If $\mathbf{i}' \in R(w_0)$ is such that \mathbf{i} and \mathbf{i}' differ by replacing three consecutive (i_k, i_{k+1}, i_{k+2}) such that $a_{i_k, i_{k+1}} = -1$ and $i_{k+2} = i_k$ by (i_{k+1}, i_k, i_{k+1}) , then
 - $\mathbf{c}_{\beta_k}^{\mathbf{i}'}(b) = \max\{c_{\beta_{k+1}}^{\mathbf{i}}(b), c_{\beta_k}^{\mathbf{i}}(b) + c_{\beta_{k+1}}^{\mathbf{i}}(b) - c_{\beta_{k+2}}^{\mathbf{i}}(b)\}$,
 - $\mathbf{c}_{\beta_{k+1}}^{\mathbf{i}'}(b) = \min\{c_{\beta_k}^{\mathbf{i}}(b), c_{\beta_{k+2}}^{\mathbf{i}}(b)\}$,
 - $\mathbf{c}_{\beta_{k+2}}^{\mathbf{i}'}(b) = \max\{c_{\beta_{k+1}}^{\mathbf{i}}(b), c_{\beta_{k+1}}^{\mathbf{i}}(b) + c_{\beta_{k+2}}^{\mathbf{i}}(b) - c_{\beta_k}^{\mathbf{i}}(b)\}$,

and for all other β , $\mathbf{c}_{\beta}^{\mathbf{i}'}(b) = \mathbf{c}_{\beta}^{\mathbf{i}}(b)$.

Caution: β_k and β_{k+2} are actually the $(k + 2)^{nd}$ and k^{th} roots respectively in the order for \mathbf{i}' . See Example 2.10.

Lemma 2.9 Let $\mathbf{i}, \mathbf{i}' \in R(w_0)$ and suppose $\{\beta_1 \prec \dots \prec \beta_N\}$ and $\{\beta_1 \prec' \dots \prec' \beta_N\}$ are the convex orderings on Φ^+ determined by \mathbf{i} and \mathbf{i}' , respectively.

1. The reduced expressions \mathbf{i}, \mathbf{i}' are related by a 2-term braid move $(i_k, i_{k+1}) \rightarrow (i_{k+1}, i_k)$ if and only if $(\beta_k | \beta_{k+1}) = 0$. In this case, $\beta_k \prec \beta_{k+1}$ is replaced by $\beta_{k+1} \prec' \beta_k$ after the braid move.
2. The reduced expressions \mathbf{i}, \mathbf{i}' are related by a 3-term braid move $(i_k, i_{k+1}, i_{k+2}) \rightarrow (i_{k+1}, i_k, i_{k+1})$, with $i_k = i_{k+2}$, if and only if $\{\beta_k, \beta_{k+1}, \beta_{k+2}\}$ form a root system of type \mathfrak{sl}_3 . In this case, $\beta_k \prec \beta_{k+1} \prec \beta_{k+2}$ is replaced by $\beta_{k+2} \prec' \beta_{k+1} \prec' \beta_k$ after the braid move.

For any two reduced expressions, we can understand the map $R_{\mathbf{i}'}^{\mathbf{i}} : \mathbf{Z}_{\geq 0}^N \rightarrow \mathbf{Z}_{\geq 0}^N$ sending $\mathbf{c}^i(b)$ to $\mathbf{c}^{i'}(b)$ by finding a way to move from \mathbf{i} to \mathbf{i}' by a sequence of braid moves, and composing the maps above. Putting this together gives a realization of $B(\infty)$ where the underlying set is \mathcal{B}_i for some fixed \mathbf{i} , and the f_i are calculated as in the following example.

Example 2.10 Let \mathfrak{g} be of type D_4 , $\mathbf{i} = 123421234234$, where α_2 is the simple root at the trivalent node of the Dynkin diagram. The corresponding order on positive roots is

$$1 \prec 12 \prec 123 \prec 124 \prec 1234 \prec 12234 \prec 2 \prec 24 \prec 23 \prec 234 \prec 3 \prec 4,$$

where 1 is identified with α_1 , 12 with $\alpha_1 + \alpha_2$, and so on. Consider

$$b = F_1^{(2)} F_{12}^{(1)} F_{123}^{(4)} F_{124}^{(2)} F_{1234}^{(1)} F_{12234}^{(3)} F_2^{(3)} F_{24}^{(1)} F_{23}^{(2)} F_{234}^{(1)} F_3^{(2)} F_4^{(0)} \in \mathcal{B}_{\mathbf{i}}, \quad (2.11)$$

Calculating $f_1 b$ is easy: the exponent of F_1 simply increases by 1. The calculation of $f_4 b$, goes as follows:

$$\begin{aligned} b &= F_1^{(2)} F_{12}^{(1)} F_{123}^{(4)} F_{124}^{(2)} F_{1234}^{(1)} F_{12234}^{(3)} F_2^{(3)} F_{24}^{(1)} F_{23}^{(2)} F_{234}^{(1)} F_3^{(2)} F_4^{(0)} \\ &\simeq F_1^{(2)} F_{12}^{(1)} F_{124}^{(2)} F_{123}^{(4)} F_{1234}^{(1)} F_{12234}^{(3)} F_2^{(3)} F_{24}^{(1)} F_{23}^{(2)} F_{234}^{(1)} F_4^{(0)} F_3^{(2)} \\ &\simeq F_1^{(2)} F_{12}^{(1)} F_{124}^{(2)} F_{123}^{(4)} F_{1234}^{(1)} F_{12234}^{(3)} F_2^{(3)} F_{24}^{(1)} F_4^{(1)} F_{234}^{(0)} F_{23}^{(3)} F_3^{(2)} \\ &\quad \vdots \\ &\simeq F_4^{(2)} F_1^{(2)} F_{124}^{(1)} F_{12}^{(2)} F_{1234}^{(1)} F_{123}^{(4)} F_{12234}^{(3)} F_{24}^{(1)} F_2^{(3)} F_{234}^{(0)} F_{23}^{(3)} F_3^{(2)} \\ f_4 b &\simeq F_4^{(3)} F_1^{(2)} F_{124}^{(1)} F_{12}^{(2)} F_{1234}^{(1)} F_{123}^{(4)} F_{12234}^{(3)} F_{24}^{(1)} F_2^{(3)} F_{234}^{(0)} F_{23}^{(3)} F_3^{(2)} \\ &\quad \vdots \\ &\simeq F_1^{(2)} F_{12}^{(1)} F_{123}^{(3)} F_{124}^{(2)} F_{1234}^{(2)} F_{12234}^{(3)} F_2^{(3)} F_{24}^{(1)} F_{23}^{(2)} F_{234}^{(1)} F_3^{(2)} F_4^{(0)}. \end{aligned}$$

The first step performs two 2-term braid moves and the corresponding (trivial) piecewise linear bijections. The second step is the piecewise-linear bijection for a 3-term braid move. Specifically, by Lemma 2.8,

$$\begin{aligned} c_{23}^{\mathbf{i}'}(b) &= \max\{c_{234}^{\mathbf{i}}(b), c_{23}^{\mathbf{i}}(b) + c_{234}^{\mathbf{i}}(b) - c_4^{\mathbf{i}}(b)\} = \max\{1, 2 + 1 - 0\} = 3, \\ c_{234}^{\mathbf{i}'}(b) &= \min\{c_{23}^{\mathbf{i}}(b), c_4^{\mathbf{i}}(b)\} = \min\{2, 0\} = 0, \\ c_4^{\mathbf{i}'}(b) &= \max\{c_{234}^{\mathbf{i}}(b), c_{234}^{\mathbf{i}}(b) + c_4^{\mathbf{i}}(b) - c_{23}^{\mathbf{i}}(b)\} = \max\{1, 1 + 0 - 2\} = 1. \end{aligned}$$

This, along with the reordering of the roots by Lemma 2.9, gives the third line above. Continue making braid moves and applying the corresponding piecewise-linear bijections to get a PBW monomial with F_4 as the leftmost factor. (To do this, recall that, by Lemma 2.9, two roots can be interchanged with a 2-term braid move exactly if they are perpendicular, and that a three-term braid move applies to three consecutive roots β, β', β'' if and only if $\beta' = \beta + \beta''$.) Then increase that exponent by 1. Then do braid moves and the corresponding piecewise-linear bijections to get back to the original order (not shown). The result is $f_4 b$, expressed as an element of $\mathcal{B}_{\mathbf{i}}$.

3 Semi-adapted words and bracketing

In general, calculating f_i as in Example 2.10 can be computationally involved. However, for some words, the application of f_i can be calculated by a simple bracketing procedure. We now discuss those words.

3.1 Semi-adapted words

Definition 3.1 Fix a reduced expression \mathbf{i} for w_0 , and $i \in I$. We say that \mathbf{i} is adapted for i if $i_1 = i$. We say that \mathbf{i} is semi-adapted for i if one can perform a sequence of braid moves to \mathbf{i} to get to a word \mathbf{i}' with $i'_1 = i$, and each of these is either

- a 2-term braid move, or
- a 3-term braid move such that the corresponding roots before the move are $(\beta, \beta + \alpha_i, \alpha_i)$, in that order, for some β .

Definition 3.2 Fix an i -semi-adapted reduced expression \mathbf{i} . Let η_1, \dots, η_k be the roots prior to α_i in the corresponding order such that $(\alpha_i | \eta_j) < 0$, in the order they appear left to right. Let $\nu_j = \eta_j + \alpha_i$, which is a root.

3.2 Bracketing rules

Recall that a Kostant partition is a formal $\mathbf{Z}_{\geq 0}$ -linear combination of positive roots, which we denote by $(c_\beta)_{\beta \in \Phi^+}$. Therefore, an \mathbf{i} -Lusztig datum $\mathbf{c}^{\mathbf{i}}(b)$ may be identified with a Kostant partition.

Definition 3.3 For any Kostant partition $\mathbf{c} = (c_\beta)_{\beta \in \Phi^+}$, define $S_i^{\mathbf{i}}(\mathbf{c})$ to be the string of brackets

$$c_{\nu_1} \quad c_{\eta_1} \quad \quad \quad c_{\nu_k} \quad c_{\eta_k} \quad c_{\alpha_i} \\) \cdots) (\cdots (\cdots) \cdots) (\cdots (\quad) \cdots) .$$

We sequentially pair adjacent brackets $()$ until the remaining brackets are a subsequence of the form $) \cdots) (\cdots ($. A bracket in $S_i^{\mathbf{i}}(\mathbf{c})$ is called uncanceled if it is not paired in this procedure.

Theorem 3.4 Fix $i \in I$ and an i -semi-adapted word \mathbf{i} . Let $b \in \mathcal{B}_{\mathbf{i}}$, and let $\mathbf{c} = (\mathbf{c}^{\mathbf{i}}_\beta(b))$ be the corresponding Kostant partition.

- If the leftmost uncanceled $'($ in $S_i^{\mathbf{i}}(\mathbf{c})$ corresponds to η_j then

$$c_{\nu_j}^{\mathbf{i}}(f_i b) = c_{\nu_j}^{\mathbf{i}}(b) + 1, \quad c_{\eta_j}^{\mathbf{i}}(f_i b) = c_{\eta_j}^{\mathbf{i}}(b) - 1, \quad \text{and} \quad c_{\beta}^{\mathbf{i}}(f_i b) = c_{\beta}^{\mathbf{i}}(b) \text{ for all other } \beta.$$

If there is no uncanceled $'($, then $c_{\alpha_i}^{\mathbf{i}}(b)$ increases by 1 and all other $c_{\beta}^{\mathbf{i}}(b)$ are unchanged.

- If the rightmost uncanceled $')$ in $S_i^{\mathbf{i}}(\mathbf{c})$ corresponds to ν_j then

$$c_{\nu_j}^{\mathbf{i}}(e_i b) = c_{\nu_j}^{\mathbf{i}}(b) - 1, \quad c_{\eta_j}^{\mathbf{i}}(e_i b) = c_{\eta_j}^{\mathbf{i}}(b) + 1, \quad \text{and} \quad c_{\beta}^{\mathbf{i}}(e_i b) = c_{\beta}^{\mathbf{i}}(b) \text{ for all other } \beta.$$

If there is no uncanceled $')$ in $S_i^{\mathbf{i}}(\mathbf{c})$, then $e_i b = 0$.

Thus, if $\mathbf{i} \in R(w_0)$ is i -semi-adapted for all i , then the crystal structure on $\mathcal{B}_{\mathbf{i}}$ can be described completely using bracketing rules. The proof of Theorem 3.4 will appear in [17]. It relies on the following lemma.

Lemma 3.5 In any sequence of braid moves allowed by Definition 3.1, the 3-term braid moves happen in the order $(\eta_k, \nu_k, \alpha_i)$, then $(\eta_{k-1}, \nu_{k-1}, \alpha_i)$, and so on.


3.3 Existence of semi-adapted words

In [11], Littelmann details specific reduced expressions for the longest element of the Weyl group for which the conditions describing the associated string cone can be expressed. These reduced expressions are called “nice decompositions” and correspond to “good enumerations” of the underlying Dynkin diagram. At least one such nice decomposition is given for each finite type except E_8 and F_4 . As will be shown in [17], the reverse of these nice decompositions are i -semi-adapted for all i . In particular, the words i^A and i^D described in the subsequent sections are precisely the reverse of Littelmann’s nice decompositions in types A and D , respectively.

4 Explicit descriptions

4.1 Type A

The Lie algebra of type A_n has Dynkin diagram



Write the positive roots as $\{\alpha_{i,j} : 1 \leq i \leq j \leq n\}$, where $\alpha_{i,j} = \alpha_i + \alpha_{i+1} + \cdots + \alpha_j$. The word i^A corresponding to the reduced expression $w_0 = (s_1 s_2 \cdots s_n)(s_1 s_2 \cdots s_{n-1}) \cdots (s_1 s_2) s_1$ is i -semi-adapted for all i . Therefore, the crystal structure on \mathcal{B}_{i^A} is given by bracketing rules.

In [2], they show that the crystal structure on \mathcal{B}_{i^A} essentially gives the realization of $B(\infty)$ using the multisegments from [6, 9]. That is, they show that the map which takes each root $\alpha_{i,j}$ to the segment $[i, j]$ is a crystal isomorphism. This isomorphism can also be understood using tableaux, which we discuss in more detail in the next section.

To understand why this map is a crystal isomorphism, consider the corresponding order on positive roots. In this case, $\alpha_{i,j} \prec \alpha_{i',j'}$ if and only if $i < i'$ or $i = i'$ and $j < j'$. Then, given $i \in I$ and a Kostant partition $c = (c_\beta)_{\beta \in \Phi^+}$, the string of brackets $S_i^A(c)$ is

$$\underbrace{\cdots)}_{c_{\alpha_{1,i}}} \underbrace{\cdots(}_{c_{\alpha_{1,i-1}}} \underbrace{\cdots)}_{c_{\alpha_{2,i}}} \underbrace{\cdots(}_{c_{\alpha_{2,i-1}}} \cdots \underbrace{\cdots)}_{c_{\alpha_{i-i,i}}} \underbrace{\cdots(}_{c_{\alpha_{i-i,i-1}}} \underbrace{\cdots)}_{c_{\alpha_{i,i}}}$$

The order on roots agrees with the order on segments used to define the crystal operators in [6].

Example 4.1 Consider type A_3 with $i = 123121$ and $i = 2$. The corresponding order on positive roots is $1 \prec \frac{2}{1} \prec \frac{3}{1} \prec 2 \prec \frac{3}{2} \prec 3$, where we identify 1 with α_1 , $\frac{2}{1}$ with $\alpha_1 + \alpha_2$, and so on. If $b \in B(\infty)$ is such that $c^i(b) = (2, 3, 1, 3, 3, 2)$, then the corresponding Kostant partition is

$$1 \quad 1 \quad \frac{2}{1} \quad \frac{2}{1} \quad \frac{2}{1} \quad \frac{3}{1} \quad 2 \quad 2 \quad 2 \quad \frac{3}{2} \quad \frac{3}{2} \quad \frac{3}{2} \quad 3 \quad 3.$$

Placing the parts/roots in the order prescribed by Definition 3.2, we get

$$\frac{2}{1} \frac{2}{1} \frac{2}{1} 1 1 2 2 2 \xrightarrow{f_2} \frac{2}{1} \frac{2}{1} \frac{2}{1} 1 1 2 2 2 2.$$

$$))) \text{ (((())) }$$

Hence, applying f_2 to the given partition above yields $1 1 \frac{2}{1} \frac{2}{1} \frac{2}{1} \frac{3}{1} 2 2 2 2 \frac{3}{2} \frac{3}{2} \frac{3}{2} 3 3$.

$\beta_{i,k} = \alpha_i + \cdots + \alpha_k, \quad 1 \leq i \leq k \leq n-1$
$\gamma_{i,k} = \alpha_i + \cdots + \alpha_{n-2} + \alpha_n + \alpha_{n-1} + \cdots + \alpha_k, \quad 1 \leq i < k \leq n$

Tab. 4.1: Positive roots of type D_n .


However, in type A there are many other good enumerations, and hence many other reduced expressions where the crystal structure on the corresponding PBW monomials is given by a bracketing rule. For example, in type A_4 ,



is a good enumeration. It could be very interesting to understand the corresponding combinatorics.

4.2 Type D

The Lie algebra of type D_n has Dynkin diagram



The list of positive roots in type D_n is given in Table 4.1.

Lemma 4.2 Define \mathbf{i}^D to be the word associated to the reduced expression

$$w_0 = (s_1 s_2 \cdots s_{n-1} s_n s_{n-2} \cdots s_2 s_1)(s_2 \cdots s_{n-1} s_n s_{n-2} \cdots s_2) \cdots (s_{n-2} s_{n-1} s_n s_{n-2}) s_{n-1} s_n.$$

Then \mathbf{i}^D is semi-adapted for all i .

The order of the positive roots corresponding to the subword $(i, i + 1, \dots, n, n - 2, \dots, i)$ of \mathbf{i}^D , for $1 \leq i \leq n - 2$, as

$$\begin{cases} \beta_{i,i} \prec \beta_{i,i+1} \prec \cdots \prec \beta_{i,n-2} \prec \beta_{i,n-1} \prec \gamma_{i,n} \prec \gamma_{i,n-1} \prec \cdots \prec \gamma_{i,i+1} & \text{if } i \equiv 1 \pmod 2, \\ \beta_{i,i} \prec \beta_{i,i+1} \prec \cdots \prec \beta_{i,n-2} \prec \gamma_{i,n} \prec \beta_{i,n-1} \prec \gamma_{i,n-1} \prec \cdots \prec \gamma_{i,i+1} & \text{if } i \equiv 0 \pmod 2. \end{cases}$$

The ordering on the roots corresponding to the suffix $(n - 1, n)$ of \mathbf{i}^D is

$$\begin{cases} \beta_{n-1,n-1} \prec \gamma_{n-1,n} & \text{if } n \equiv 0 \pmod 2, \\ \gamma_{n-1,n} \prec \beta_{n-1,n-1} & \text{if } n \equiv 1 \pmod 2. \end{cases}$$

It follows that, for a Kostant partition $\mathbf{c} = (c_\beta)_{\beta \in \Phi^+}$, the string of brackets $S_i^D(\mathbf{c})$ needed to compute f_i is obtained by canceling brackets in

$$\begin{cases} \underbrace{\left(\cdots \right)}_{c_{\beta_{1,i}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{1,i-1}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{1,i}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{1,i+1}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{i-1,i}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{i-1,i-1}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{i-1,i}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{i-1,i+1}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{i,i}}}, & \text{if } i \neq n, \\ \underbrace{\left(\cdots \right)}_{c_{\gamma_{1,n}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{1,n-2}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{1,n-1}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{1,n-1}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{n-2,n}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{n-2,n-2}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{n-2,n-1}}} \underbrace{\left(\cdots \right)}_{c_{\beta_{n-2,n-1}}} \underbrace{\left(\cdots \right)}_{c_{\gamma_{n-1,n}}}, & \text{if } i = n. \end{cases}$$

Computing f_3 on this tableaux gives

$$f_2\Theta^{-1}(b) = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & 3 & 3 & 3 & 4 & & & \\ \hline 2 & 2 & 2 & 2 & 3 & 3 & 3 & 3 & 4 & 4 & 4 & & & & & & & & & & \\ \hline 3 & 4 & 4 & & & & & & & & & & & & & & & & & & \\ \hline \end{array} .$$

One can verify that this agrees with the calculation in Example 4.1.

5.2 Type D

As in [4], let $\mathcal{T}(\infty)$ be the set of marginally large semistandard tableaux on the alphabet

$$J(D_n) := \left\{ 1 \prec \dots \prec n-1 \prec \frac{n}{\bar{n}} \prec \overline{n-1} \prec \dots \prec \bar{1} \right\},$$

with $n-1$ rows such that the contents of the i th row are less than or equal to \bar{i} (for each $i = 1, \dots, n-1$) and entries n and \bar{n} do not appear in the same row.

As in type A , $\mathcal{T}(\infty)$ has a crystal structure. To compute the action of the Kashiwara operators e_i and f_i on $T \in \mathcal{T}(\infty)$, form a sequence of brackets by reading boxes in the tableau in rows from right to left, starting with the top row. Add ‘)’ under any letter for which there is an i -colored arrow entering the corresponding box in Figure 5.1, and add ‘(’ under any letter for which there is an i -colored arrow leaving the corresponding box in Figure 5.1. Sequentially cancel all $()$ -pairs to obtain a sequence of the form $(\dots)(\dots)(\dots)$. The brackets that remain are called *uncanceled*.




Fig. 5.1: The fundamental crystal of type D_n .

Definition 5.2 Let $T \in \mathcal{T}(\infty)$ and $i \in I$.

1. Let x be the letter in T corresponding to the right-most uncanceled ‘)’ . Then $e_i T$ is the tableau obtained from T by replacing the box containing x by the box containing the letter at the other end of the i -arrow from x in Figure 5.1. If the result is not marginally large, then delete exactly one column containing the elements $1, \dots, i$ so that the result is marginally large. If there is no uncanceled ‘)’ , then $e_i T = 0$.
2. Let y be the letter in T corresponding to the left-most uncanceled ‘(’ . Then $f_i T$ is the tableau obtained from T by replacing the box containing y by the box containing the letter at the other end of the i -arrow from y in Figure 5.1. If the result is not marginally large, then insert exactly one column containing the elements $1, \dots, i$ so that the result is marginally large.

Definition 5.2 uses the middle-Eastern reading, as defined in [3], whereas in [4] they use the far-Eastern reading. However, the resulting crystal graphs are identical.

Proposition 5.3 *The crystal graphs obtained from $\mathcal{T}(\infty)$ using the far-Eastern reading and the middle-Eastern reading, respectively, are identical.*

Remark 5.4 *In contrast to type A_n , the crystal structure on irreducible highest weight crystals of type D_n modeled by Kashiwara-Nakashima tableaux [8] using these two readings are not in fact the same. This is only a property of marginally large tableaux.*

Definition 5.5 *Define a map Ψ from $\mathcal{T}(\infty)$ to Kostant partitions as follows. For a tableaux $T \in \mathcal{T}(\infty)$, let R_1, \dots, R_{n-1} denote the rows of T starting at the top. Set $\Psi(T) = (c_\beta)_{\beta \in \Phi^+}$, where (c_β) is obtained from the trivial data $(0, 0, \dots, 0)$ in the following way:*

1. if $j \neq n - 1$, each \bar{j} in R_j increases both $c_{\beta_{j,j}}$ and $c_{\gamma_{j,j+1}}$ by 1;
2. if $j = n - 1$, each \bar{j} in R_j increases both $c_{\beta_{n-1,n-1}}$ and $c_{\gamma_{n-1,n}}$ by 1;
3. for each pair k, \bar{k} in R_j , where $k \neq n - 1$, increase both $c_{\beta_{j,k}}$ and $c_{\gamma_{j,k+1}}$ by 1;
4. for each pair $n - 1, \overline{n - 1}$ in R_j , increase both $c_{\beta_{j,n-1}}$ and $c_{\gamma_{j,n}}$ by 1;
5. each remaining $k \in \{j, j + 1, \dots, n\}$ in R_j increases $c_{\beta_{j,k-1}}$ by 1;
6. each remaining $\bar{k} \in \{\bar{n}, \overline{n - 1}, \dots, \overline{j + 1}\}$ in R_j increases $c_{\gamma_{j,k}}$ by 1.

Example 5.6 *Let $n = 4$ and*

$$T = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & \bar{3} & \bar{1} & \bar{1} & \bar{1} & \\ \hline 2 & 2 & 2 & 2 & 3 & \bar{4} & \bar{3} & \bar{3} & & & & & & & & \\ \hline 3 & \bar{4} & \bar{3} & & & & & & & & & & & & & \\ \hline \end{array} .$$

The sequence of $\bar{1}$'s in the first row increases both $c_{\beta_{1,1}}$ and $c_{\gamma_{1,2}}$ by 3, the $\bar{3}$ in the first row increases $c_{\gamma_{1,3}}$ by 1 and the pair of 2's in the first row increases $c_{\beta_{1,1}}$ by 2. In the second row, the pair $(3, \bar{3})$ increases both $c_{\beta_{2,3}}$ and $c_{\gamma_{2,4}}$ by 1, the $\bar{4}$ increases $c_{\gamma_{2,3}}$ by 1, and the 3 increases $c_{\gamma_{2,4}}$ by 1. Finally, the $\bar{3}$ in the third row increases both $c_{\beta_{3,3}}$ and $c_{\gamma_{3,4}}$ by 1 and the $\bar{4}$ increases $c_{\gamma_{3,4}}$ by 1. In summary,

$$\Psi(T) = 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ \begin{array}{c} 3\bar{4} \\ 2 \\ 1 \end{array} \ \begin{array}{c} 2 \\ 3\bar{4} \\ 2 \\ 1 \end{array} \ \begin{array}{c} 2 \\ 3\bar{4} \\ 2 \\ 1 \end{array} \ \begin{array}{c} 2 \\ 3\bar{4} \\ 2 \\ 1 \end{array} \ 4 \ 4 \ 3 \ 3\bar{4} \ 3 \ 4 \ 4.$$

Recall that Kostant partitions correspond to elements of \mathcal{B}_{iD} . The following is the main result of [18]:

Theorem 5.7 *The map $\Psi: \mathcal{T}(\infty) \rightarrow \mathcal{B}_{iD}$ defined above is a crystal isomorphism.*

References

[1] Arkady Berenstein and Andrei Zelevinsky, *Tensor product multiplicities, canonical bases and totally positive varieties*, Invent. Math. **143** (2001), no. 1, 77–128. MR 1802793 (2002c:17005)

[2] John Claxton and Peter Tingley, *Young tableaux, multisegments, and PBW bases*, Sém. Lothar. Combin. **73** (2014/2015), Art. B73c, 21 pp.

- [3] Jin Hong and Seok-Jin Kang, *Introduction to quantum groups and crystal bases*, Graduate Studies in Mathematics, vol. 42, American Mathematical Society, Providence, RI, 2002. MR 1881971 (2002m:17012)
- [4] Jin Hong and Hyeonmi Lee, *Young tableaux and crystal $\mathcal{B}(\infty)$ for finite simple Lie algebras*, J. Algebra **320** (2008), no. 10, 3680–3693. MR 2457716 (2009j:17008)
- [5] ———, *Crystal $\mathcal{B}(\lambda)$ as a subset of the tableau description of $\mathcal{B}(\infty)$ for the classical Lie algebra types*, Algebr. Represent. Theory **18** (2015), no. 1, 137–160. MR 3317847
- [6] Nicolas Jacon and Cédric Lecouvey, *Kashiwara and Zelevinsky involutions in affine type A*, Pacific J. Math. **243** (2009), no. 2, 287–311. MR 2552260 (2010k:20012)
- [7] Masaki Kashiwara, *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516. MR 1115118 (93b:17045)
- [8] Masaki Kashiwara and Toshiki Nakashima, *Crystal graphs for representations of the q -analogue of classical Lie algebras*, J. Algebra **165** (1994), no. 2, 295–345. MR 1273277 (95c:17025)
- [9] Bernard Leclerc, Jean-Yves Thibon, and Eric Vasserot, *Zelevinsky’s involution at roots of unity*, J. Reine Angew. Math. **513** (1999), 33–51. MR 1713318 (2001f:20011)
- [10] Kyu-Hwan Lee and Ben Salisbury, *Young tableaux, canonical bases, and the Gindikin-Karpelevich formula*, J. Korean Math. Soc. **51** (2014), no. 2, 289–309.
- [11] P. Littelmann, *Cones, crystals, and patterns*, Transform. Groups **3** (1998), no. 2, 145–179. MR 1628449 (99e:17009)
- [12] George Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498. MR 1035415 (90m:17023)
- [13] ———, *Introduction to quantum groups*, Modern Birkhäuser Classics, Birkhäuser/Springer, New York, 2010, Reprint of the 1994 edition. MR 2759715 (2011j:17028)
- [14] Paolo Papi, *A characterization of a special ordering in a root system*, Proc. Amer. Math. Soc. **120** (1994), no. 3, 661–665. MR 1169886 (94e:20056)
- [15] Markus Reineke, *On the coloured graph structure of Lusztig’s canonical basis*, Math. Ann. **307** (1997), no. 4, 705–723. MR 1464139
- [16] Yoshihisa Saito, *PBW basis of quantized universal enveloping algebras*, Publ. Res. Inst. Math. Sci. **30** (1994), no. 2, 209–232. MR 1265471 (95e:17021)
- [17] Ben Salisbury, Adam Schultze, and Peter Tingley, *Combinatorial descriptions of the crystal structure on certain PBW bases*, 2016, in preparation.
- [18] ———, *PBW bases and marginally large tableaux in type D*, 2016, in preparation.
- [19] Peter Tingley, *Elementary construction of Lusztig’s canonical basis*, AMS Contemporary Math volume on Groups, Rings, Group Rings and Hopf Algebras, dedicated to Don Passman, To appear, arXiv:1602.04895.