



**HAL**  
open science

## Links in the complex of weakly separated collections

Suho Oh, David Speyer

► **To cite this version:**

Suho Oh, David Speyer. Links in the complex of weakly separated collections. 28-th International Conference on Formal Power Series and Algebraic Combinatorics, Simon Fraser University, Jul 2016, Vancouver, Canada. 10.46298/dmtcs.6389 . hal-02166342

**HAL Id: hal-02166342**

**<https://hal.science/hal-02166342>**

Submitted on 26 Jun 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Links in the complex of weakly separated collections (Extended Abstract)

Suho Oh<sup>1†</sup> and David Speyer<sup>2‡</sup>

<sup>1</sup>San Marcos, TX

<sup>2</sup>Ann Arbor, MI

**Abstract.** Plabic graphs are combinatorial objects used to study the totally nonnegative Grassmannian. Faces of plabic graphs are labeled by  $k$ -element sets of positive integers, and a collection of such  $k$ -element sets are the face labels of a plabic graph if that collection forms a maximal weakly separated collection. There are moves that one can apply to plabic graphs, and thus to maximal weakly separated collections, analogous to mutations of seeds in cluster algebras. In this short note, we show if two maximal weakly separated collections can be mutated from one to another, then one can do so while freezing the face labels they have in common. In particular, this provides a new, and we think simpler, proof of Postnikov’s result that any two reduced plabic graphs with the same decorated permutations can be mutated to each other.

**Résumé.** Les graphes “plabic” sont des objets combinatoires utilisés pour l’étude de la Grassmannienne totalement positive. Les faces des graphes “plabic” sont étiquetées par des ensembles de  $k$  entiers positifs et une collection de tels ensembles correspond à un graphe “plabic” si cette collection est une collection maximale faiblement séparée. Certaines opérations peuvent être effectuées sur les graphes “plabic”, et donc sur les collections faiblement séparées, analogues aux mutations de graines dans les algèbres amassées. Dans cet article, nous montrons que si deux collections maximales faiblement séparées peuvent être transformées l’une en l’autre par ces opérations, alors il est possible de le faire en gelant les étiquettes faciales qui sont communes aux deux collections. En particulier, ceci fournit une nouvelle (et de notre point de vue plus simple) preuve du résultat de Postnikov qui affirme que deux graphes “plabic” réduits avec les mêmes permutations faciales peuvent être transformés l’un en l’autre par ces opérations.

**Keywords.** weak separation, maximal weakly separated collection, plabic graphs, plabic tiling, total positivity, Grassmannian

## 1 Introduction

Fix two positive integers  $k \leq n$ . Let  $[n] := \{1, \dots, n\}$ . We will generally consider  $[n]$  as cyclically ordered. We will say that  $i_1, i_2, \dots, i_r$  in  $[n]$  are *cyclically ordered* if  $i_s < i_{s+1} < \dots < i_r < i_1 < i_2 < \dots < i_{s-1}$  for some  $s \in [r]$ .

Fix positive integers  $k < n$ . Let  $I$  and  $J$  be two  $k$ -element subsets of  $\{1, 2, \dots, n\}$ . The following definition is due to Leclerc and Zelevinsky [2], see also [9] and [6]: The sets  $I$  and  $J$  are called *weakly*

<sup>†</sup>Email: suhooh@txstate.edu

<sup>‡</sup>Email: speyer@umich.edu

*separated* if there do **not** exist  $a, b, c$  and  $d$  cyclically ordered with  $a, c \in I \setminus J$  and  $b, d \in J \setminus I$ . Graphically,  $I$  and  $J$  are weakly separated if we can draw a chord across the circle separating  $I \setminus J$  from  $J \setminus I$ . We write  $I \parallel J$  to indicate that  $I$  and  $J$  are weakly separated.

Write  $\binom{[n]}{k}$  for the set of  $k$  element subsets of  $[n]$ . We will use the term *collection* to refer to a subset of  $\binom{[n]}{k}$ . We define a *weakly separated collection* to be a collection  $\mathcal{C} \subset \binom{[n]}{k}$  such that, for any  $I$  and  $J$  in  $\mathcal{C}$ , the sets  $I$  and  $J$  are weakly separated. We define a *maximal weakly separated collection* to be a weakly separated collection which is not contained in any other weakly separated collection.

Following Leclerc and Zelevinsky [2], Scott [9] observed the following:

**Proposition 1.1.** [9], cf. [2] *Let  $S \in \binom{[n]}{k-2}$  and let  $a, b, c, d$  be cyclically ordered elements of  $[n] \setminus S$ . Suppose that a maximal weakly separated collection  $\mathcal{C}_1$  contains  $S \cup \{a, b\}$ ,  $S \cup \{b, c\}$ ,  $S \cup \{c, d\}$ ,  $S \cup \{d, a\}$  and  $S \cup \{a, c\}$ . Then  $\mathcal{C}_2 := (\mathcal{C}_1 \setminus \{S \cup \{a, c\}\}) \cup \{S \cup \{b, d\}\}$  is also a maximal weakly separated collection.*

When  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are related as in this proposition, we will say that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are *mutations* of each other. Relying on results of [7], in [6] the authors proved that any two maximal weakly separated collections are linked by a sequence of mutations. As a corollary, any two maximal weakly separated collections have the same cardinality – namely  $k(n - k) + 1$ .

In other words, if we form a simplicial complex whose vertices are indexed by  $\binom{[n]}{k}$ , and whose faces are the maximal weakly separated sets, then this complex is pure of dimension  $k(n - k)$  and is connected in codimension 1. This complex was further studied in [3].

In this paper, we will study the links of faces in this complex. Namely, our main result is:

**Theorem 1.2.** *Let  $\mathcal{B} \subset \binom{[n]}{k}$  be a weakly separated collection. Let  $\mathcal{C}$  and  $\mathcal{C}'$  be two maximal weakly separated collections containing  $\mathcal{B}$ . Then  $\mathcal{C}$  and  $\mathcal{C}'$  are linked by a chain of mutations  $\mathcal{C} = \mathcal{C}_1 \rightarrow \mathcal{C}_2 \rightarrow \dots \rightarrow \mathcal{C}_r = \mathcal{C}'$  where all the  $\mathcal{C}_i$  contain  $\mathcal{B}$ .*

In other words, if  $\sigma$  is a face of the simplicial complex described above, with codimension greater than 1, then the link of  $\sigma$  is connected in codimension 1.

Even the case  $\mathcal{B} = \emptyset$ , where this result is due to Postnikov [7], our proof is new and independent of Postnikov’s.

## 2 Notations

We will use the following notations through out the paper: We write  $(a, b)$  for the open cyclic interval from  $a$  to  $b$ . In other words, the set of  $i$  such that  $a, i, b$  is cyclically ordered. We write  $[a, b]$  for the closed cyclic interval,  $[a, b] = (a, b) \cup \{a, b\}$ , and use similar notations for half open intervals.

If  $S$  is a subset of  $[n]$  and  $a$  an element of  $[n]$ , we may abbreviate  $S \cup \{a\}$  and  $S \setminus \{a\}$  by  $Sa$  and  $S \setminus a$ .

In this paper, we need to deal with three levels of objects: elements of  $[n]$ , subsets of  $[n]$ , and collections of subsets of  $[n]$ . For clarity, we will denote these by lower case letters, capital letters, and calligraphic letters, respectively.

The use of the notation  $I \setminus J$  does not imply  $J \subseteq I$ .

## 3 Positroids

Postnikov and the authors more generally studied weakly separated collections within combinatorial objects known as positroids. We review this material briefly here; see [7] and [6] for more. A *Grassmann*

**necklace** is a sequence  $\mathcal{I} = (I_1, \dots, I_n)$  of  $k$ -element subsets of  $[n]$  such that, for  $i \in [n]$ , the set  $I_{i+1}$  contains  $I_i \setminus \{i\}$ . (Here the indices are taken modulo  $n$ .) If  $i \notin I_i$ , then we should have  $I_{i+1} = I_i$ .

Define a linear order  $<_i$  on  $[n]$  by

$$i <_i i + 1 <_i i + 2 <_i \dots <_i i - 1.$$

We extend  $<_i$  to  $k$  element sets, as follows. For  $I = \{i_1, \dots, i_k\}$  and  $J = \{j_1, \dots, j_k\}$  with  $i_1 <_i i_2 <_i \dots <_i i_k$  and  $j_1 <_i j_2 <_i \dots <_i j_k$ , define the partial order

$$I \leq_i J \text{ if and only if } i_1 \leq_i j_1, \dots, i_k \leq_i j_k.$$

Given a Grassmann necklace  $\mathcal{I} = (I_1, \dots, I_n)$ , define the **positroid**  $\mathcal{M}_{\mathcal{I}}$  to be

$$\mathcal{M}_{\mathcal{I}} := \left\{ J \in \binom{[n]}{k} \mid I_i \leq_i J \text{ for all } i \in [n] \right\}.$$

Fix a Grassmann necklace  $\mathcal{I} = (I_1, \dots, I_n)$ , with corresponding positroid  $\mathcal{M}_{\mathcal{I}}$ . Then  $\mathcal{C}$  is called a *weakly separated collection inside  $\mathcal{M}_{\mathcal{I}}$*  if  $\mathcal{C}$  is a weakly separated collection and  $\mathcal{I} \subseteq \mathcal{C} \subseteq \mathcal{M}_{\mathcal{I}}$ . We call  $\mathcal{C}$  a *maximal weakly separated collection inside  $\mathcal{M}_{\mathcal{I}}$*  if it is maximal among weakly separated collections inside  $\mathcal{M}_{\mathcal{I}}$ .

Our actual main result is

**Theorem 3.1.** *Let  $\mathcal{I}$  be a Grassmann necklace and let  $\mathcal{B}$  be a weakly separated collection in  $\mathcal{M}_{\mathcal{I}}$ . Let  $\mathcal{C}$  and  $\mathcal{C}'$  be two maximal weakly separated collections in  $\mathcal{M}_{\mathcal{I}}$  containing  $\mathcal{B}$ . Then  $\mathcal{C}$  and  $\mathcal{C}'$  are linked by a chain of mutations  $\mathcal{C} = \mathcal{C}_1 \rightarrow \mathcal{C}_2 \rightarrow \dots \rightarrow \mathcal{C}_r = \mathcal{C}'$  where all the  $\mathcal{C}_i$  contain  $\mathcal{B}$  and are in  $\mathcal{M}_{\mathcal{I}}$ .*

The case  $I_i = \{i, i + 1, \dots, i + k - 1\}$  corresponds to taking  $\mathcal{M}_{\mathcal{I}}$  to be all of  $\binom{[n]}{k}$ . This result also implies the main result of [1]. (Just take  $\mathcal{B}$  to be any collection of boundary labels.)

## 4 Plabic tilings

In this section, we review the plabic tiling construction from [6]. The motivation for this construction is as follows: The main result of [6] is that maximal weakly separated collections are in bijection with certain planar bipartite graphs called “reduced plabic graphs”. The planar dual of a reduced plabic graph is thus a bi-colored CW complex, homeomorphic to a two-dimensional disc. The plabic tiling construction assigns a bi-colored two-dimensional CW complex to any weakly separated collection, maximal or not. For the purposes of this paper, we only need plabic tilings, not plabic graphs.


Let us fix  $\mathcal{C}$ , a weakly separated collection in  $\mathcal{M}_{\mathcal{I}}$ . For  $I$  and  $J \in \mathcal{M}_{\mathcal{I}}$ , say that  $I$  **neighbors**  $J$  if

$$|I \setminus J| = |J \setminus I| = 1.$$

Let  $K$  be any  $(k - 1)$  element subset of  $[n]$ . We define the **white clique**  $\mathcal{W}(K)$  to be the set of  $I \in \mathcal{C}$  such that  $K \subset I$ . Similarly, for  $L$  a  $(k + 1)$  element subset of  $[n]$ , we define the **black clique**  $\mathcal{B}(L)$  for the set of  $I \in \mathcal{C}$  which are contained in  $L$ . We call a clique **nontrivial** if it has at least three elements. Observe that, if  $\mathcal{X}$  is a nontrivial clique, then it cannot be both black and white.

Observe that a white clique  $\mathcal{W}(K)$  is of the form  $\{Ka_1, Ka_2, \dots, Ka_r\}$  for some  $a_1, a_2, \dots, a_r$ , which we take to be cyclically ordered. Similarly,  $\mathcal{B}(L)$  is of the form  $\{L \setminus b_1, L \setminus b_2, \dots, L \setminus b_s\}$ , with the  $b_i$ ’s cyclically ordered. If  $\mathcal{W}(K)$  is nontrivial, we define the boundary of  $\mathcal{W}(K)$  to be the cyclic graph

$$(Ka_1) \rightarrow (Ka_2) \rightarrow \dots \rightarrow (Ka_r) \rightarrow (Ka_1).$$



**Fig. 1:** Example of a plabic tiling

Similarly, the boundary of a nontrivial  $\mathcal{B}(L)$  is

$$(L \setminus b_1) \rightarrow (L \setminus b_2) \rightarrow \dots \rightarrow (L \setminus b_s) \rightarrow (L \setminus b_1).$$

If  $(J, J')$  is a two element clique, then we define its boundary to be the graph with a single edge  $(J, J')$ ; we define an one element clique to have empty boundary.

We now define a two dimensional CW-complex  $\Sigma(\mathcal{C})$ . The vertices of  $\Sigma(\mathcal{C})$  will be the elements of  $\mathcal{C}$ . There will be an edge  $(I, J)$  if

1.  $\mathcal{W}(I \cap J)$  is nontrivial and  $(I, J)$  appears in the boundary of  $\mathcal{W}(I \cap J)$  or
2.  $\mathcal{B}(I \cup J)$  is nontrivial and  $(I, J)$  appears in the boundary of  $\mathcal{B}(I \cup J)$  or
3.  $\mathcal{W}(I \cap J) = \mathcal{B}(I \cup J) = \{I, J\}$ .

There will be a two-dimensional face of  $\Sigma(\mathcal{C})$  for each nontrivial clique  $\mathcal{X}$  of  $\mathcal{C}$ . The boundary of this face will be the boundary of  $\mathcal{X}$ . We will refer to each face of  $\Sigma(\mathcal{C})$  as **black** or **white**, according to the color of the corresponding clique. We call a CW-complex of the form  $\Sigma(\mathcal{C})$  a **plabic tiling**. An implicit claim here is that, if  $\mathcal{W}(I \cap J)$  and  $\mathcal{B}(I \cup J)$  are both nontrivial, then  $(I, J)$  is a boundary edge of both, so that 2-dimensional faces of  $\Sigma(\mathcal{C})$  are glued along boundary edges. This is not obvious, but it is true; see [6, Lemma 9.2].

So far,  $\Sigma(\mathcal{C})$  is an abstract CW-complex. Our next goal is to embed it in a plane.

Fix  $n$  points  $v_1, v_2, \dots, v_n$  in  $\mathbb{R}^2$ , at the vertices of a convex  $n$ -gon in clockwise order. We write  $e_1, e_2, \dots, e_n$  for the standard basis of  $\mathbb{R}^n$ . We define a linear map  $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^2$  by  $e_a \mapsto v_a$ . For  $I \in \binom{[n]}{t}$ , set  $e_I = \sum_{a \in I} e_a$ . We abbreviate  $\pi(e_I)$  by  $\pi(I)$ .

We extend the map  $\pi$  to a map from  $\Sigma(\mathcal{C})$  to  $\mathbb{R}^2$  as follows: Each vertex  $I$  of  $\Sigma(\mathcal{C})$  is sent to  $\pi(I)$  and each face of  $\Sigma(\mathcal{C})$  is sent to the convex hull of the images of its vertices. We encourage the reader to consult Figure 1 and see that the vector  $\pi(Si) - \pi(Sj)$  is a translation of  $v_i - v_j$ .<sup>(i)</sup>

We define  $\pi(\mathcal{I})$  to be the closed polygonal curve whose vertices are, in order,  $\pi(I_1), \pi(I_2), \dots, \pi(I_n), \pi(I_1)$ . For example in Figure 1, we consider the closed polygonal curve given by

$$\pi(\{1, 2, 3\}), \pi(\{2, 3, 4\}), \dots, \pi(\{7, 2, 1\}), \pi(\{1, 2, 3\}),$$

<sup>(i)</sup> This figure is extremely similar to [6, Figure 9], we have redrawn it to avoid issues of figure reuse.

coming from the Grassmann necklace

$$(\{1, 2, 3\}, \{2, 3, 4\}, \{3, 4, 5\}, \{4, 5, 6\}, \{5, 6, 7\}, \{6, 7, 1\}, \{7, 2, 1\}).$$

We now summarize the main results of [6] concerning plabic tilings:

**Proposition 4.1** ([6, Prop. 9.4, Prop. 9.8, Prop 9.10, Theorem 11.1]). *With the above notation,  $\pi(\mathcal{I})$  is a simple closed curve, except that if  $\mathcal{I}$  has repeated elements then  $\pi(\mathcal{I})$  may touch itself at those vertices, in a manner which can be perturbed to a simple closed curve. If  $\mathcal{C}$  is a weakly separated collection in  $\mathcal{M}_{\mathcal{I}}$ , then the map  $\pi : \Sigma(\mathcal{C}) \rightarrow \mathbb{R}^2$  is injective, and its image lands inside the curve  $\pi(\mathcal{I})$*

*The collection  $\mathcal{C}$  is maximal among weakly separated collections in  $\mathcal{M}_{\mathcal{I}}$  if and only if  $\Sigma(\mathcal{C})$  fills the entire interior of the curve  $\pi(\mathcal{I})$ .*

*If  $J$  is weakly separated from all elements of  $\mathcal{I}$ , then  $J \in \mathcal{M}_{\mathcal{I}}$  if and only if  $\pi(J)$  is inside the curve  $\pi(\mathcal{I})$ .*

We will sometimes speak of **triangulating**  $\Sigma(\mathcal{C})$ , meaning to take each 2-cell of  $\Sigma(\mathcal{C})$  and divide it into triangles. Coloring these triangles with the color of the corresponding 2-cells, the vertices of a white triangle are of the form  $(Sa, Sb, Sc)$  for some  $k - 1$  element set  $S$  and some  $a, b, c \in [n] \setminus S$ . The vertices of a black triangle are of the form  $(S \setminus a, S \setminus b, S \setminus c)$  for some  $k + 1$  element set  $S$  and some  $a, b, c \in S$ . Note that the image of a triangle under  $\pi$  is a translate of  $\text{Hull}(v_a, v_b, v_c)$  or  $\text{Hull}(-v_a, -v_b, -v_c)$  respectively. The triangle is oriented clockwise if  $(a, b, c)$  are cyclically ordered.

## 5 A lemma regarding mutations

We will need the following lemma.

**Lemma 5.1.** *Let  $H$  be a subset of  $[n]$  of cardinality  $k - 2$ ; let  $a, b, c, d$  be circularly ordered elements in  $[n] \setminus H$ . Let  $J$  be another  $k$  element subset of  $[n]$ . Suppose that  $Hac$  and  $Hbd$  are weakly separated with  $J$ . Then  $Hab, Hbc, Hcd$  and  $Hda$  are weakly separated with  $J$ .*

The relevance of this lemma is as follows: Suppose that  $\mathcal{C}$  is a weakly separated collection which contains  $Hac$  and all of whose elements other than  $Hac$  are weakly separated from  $Hbd$ . Then the lemma shows that  $\mathcal{C}' := \mathcal{C} \cup \{Hab, Hbc, Hcd, Had\}$  is weakly separated. Extending  $\mathcal{C}'$  to some maximal weakly separated collection  $\mathcal{C}_{\max}$ , we can mutate  $\mathcal{C}_{\max}$  to change  $Hac$  to  $Hbd$ . So the lemma shows that, if a weakly separated collections looks like it should be mutable in a certain manner, then it can be extended to a maximal weakly separated collection which is mutable in that manner.

*Proof.* We will show  $Hab \parallel J$ , the cases of  $Hbc, Hcd$  and  $Had$  are similar due to cyclic symmetry. Assume for the sake of contradiction that  $Hab$  and  $J$  are not weakly separated. Set  $J' = J \setminus \{a, b, c, d\}$ .

**Case 1:**  $J' = H$ . In this case,  $J$  is one of  $Hab, Hac, Had, Hbc, Hbd$  and  $Hcd$ . In each case,  $Hab \parallel J$ .

**Case 2:**  $J' \subsetneq H$ . Then  $|J'| - |H| \leq |J| - |H| = 2$ , so  $J' \setminus H$  has either one or two elements.

**Case 2a:**  $J' = H \cup \{p\}$  for some  $p \notin H$ . The equation  $|Hab| = |J|$  shows that  $J \setminus J'$  has three elements, which are among  $\{a, b, c, d\}$ . Checking all four possibilities for  $J \setminus J'$  and all four possible positions for  $p$  relative to the circularly ordered set  $\{a, b, c, d\}$  checks the claim.

**Case 2b:**  $J' = H \cup \{p, q\}$  for some  $p$  and  $q \notin H$ . Then  $|J \setminus J'| = 4$ , so  $a, b, c$  and  $d$  are in  $J$ . Checking all possible positions for  $\{p, q\}$  among the circularly oriented set  $\{a, b, c, d\}$  proves the claim.

**Case 3:**  $H \subsetneq J'$ . This is extremely similar to Case 2, we omit the details.

**Case 4:**  $H \not\subseteq J'$  and  $J' \not\subseteq H$ . We must have  $H \parallel J'$  as subsets of the circularly ordered set  $[n] \setminus \{a, b, c, d\}$ . Let  $p, q, r$  and  $s$  be the circularly ordered elements of  $[n] \setminus \{a, b, c, d\}$  so that  $\{p, q\} \subseteq H \setminus J' \subseteq [p, q]$  and  $\{r, s\} \subseteq J' \setminus H \subseteq [r, s]$ . Using that  $Hac \parallel J$  and  $Hbd \parallel J$ , we see that it is impossible for any of  $\{a, b, c, d\}$  to lie in  $(p, q) \cap J$ , or in  $(r, s) \setminus J$ . Thus there are no elements of  $J \setminus H$  in  $[p, q]$  and no elements of  $Habcd \setminus J$  in  $[r, s]$ .

Suppose for the sake of contradiction that  $Hab \not\parallel J$ . Thinking about how this can be compatible with the above restrictions on  $J \setminus H$  and  $Habcd \setminus J$ , we see that we are in one of the following two cases:

**Case 4a:** There are  $u$  and  $v$ , with  $(q, u, v, r)$  circularly ordered, with  $u \in J \setminus Hab$  and  $v \in Hab \setminus J$ . We must have  $u \in \{c, d\}$  and  $v \in \{a, b\}$ . If  $(u, v) = (d, a)$ , then  $Hac \not\parallel J$ . If  $(u, v) = (c, a)$ , then  $Hac \not\parallel J$  implies  $d \in H$  and  $Hbd \parallel J$  implies  $d \notin H$ . If  $(u, v) = (d, b)$ , then  $Hac \parallel J$  implies  $a \notin H$  and  $Hbd \parallel J$  implies  $a \in H$ . Finally, if  $(u, v) = (c, b)$ , then  $Hbd \not\parallel J$ .

**Case 4b:** There are  $u$  and  $v$ , with  $(s, u, v, p)$  circularly ordered, with  $u \in Hab \setminus J$  and  $v \in J \setminus Hab$ . This case is very similar to case 4a, and we omit the details. □

## 6 Main result

In this section, we will prove our main result: if  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are maximal weakly separated collections of some positroid  $\mathcal{M}_{\mathcal{I}}$ , then  $\mathcal{C}_1$  can be mutated to  $\mathcal{C}_2$  while preserving the sets they have in common. Throughout this section, we will fix a positroid  $\mathcal{M}$  and its Grassmann necklace  $\mathcal{I}$ .

Let  $\mathcal{B}$  be a weakly separated collection contained in  $\mathcal{W}$ . We will say that two maximal weakly separated collections  $\mathcal{C}$  and  $\mathcal{C}'$  are  **$\mathcal{B}$ -equivalent within  $\mathcal{M}_{\mathcal{I}}$**  if there is a chain of maximal weakly separated collections  $\mathcal{C} = \mathcal{C}^1 \rightarrow \mathcal{C}^2 \rightarrow \dots \rightarrow \mathcal{C}^{q-1} \rightarrow \mathcal{C}^q = \mathcal{C}'$ , such that:

- $\mathcal{B} \subseteq \mathcal{C}^1, \dots, \mathcal{C}^q$ ,
- $\mathcal{C}^{i+1}$  is obtained from  $\mathcal{C}^i$  by one mutation move.
- All the  $\mathcal{C}^i$  obey  $\mathcal{I} \subseteq \mathcal{C}^i \subseteq \mathcal{M}_{\mathcal{I}}$ .

**Theorem 6.1.** *If  $\mathcal{C}$  and  $\mathcal{C}'$  are maximal weakly separated collections within  $\mathcal{M}_{\mathcal{I}}$  containing  $\mathcal{B}$ , then  $\mathcal{C}$  and  $\mathcal{C}'$  are  $\mathcal{B}$ -equivalent within  $\mathcal{M}_{\mathcal{I}}$ .*

*Proof.* Since weakly separated collections in  $\mathcal{M}_{\mathcal{I}}$  contain  $\mathcal{I}$  by definition, we may assume that  $\mathcal{I} \subset \mathcal{B}$ . So the condition  $\mathcal{I} \subseteq \mathcal{C}^i$  will follow from  $\mathcal{B} \subseteq \mathcal{C}^i$ , and we will only need to check the other conditions.

Let  $\Sigma(\mathcal{B})$  be the 2-dimensional CW-complex defined in the previous section associated to  $\mathcal{B}$ . We fix a map  $\pi$  as in the previous section. So  $\pi(\Sigma(\mathcal{B}))$  is a closed region of  $\mathbb{R}^2$ , whose exterior boundary is  $\pi(\mathcal{I})$ . Let  $A(\mathcal{B})$  be the area of the bounded regions of  $\mathbb{R}^2 \setminus \pi(\Sigma(\mathcal{B}))$ . Let  $\delta = \min \text{Area}(\text{Hull}(v_a, v_b, v_c))$  where the minimum is over  $1 \leq a < b < c \leq n$ . So  $\delta$  is the smallest possible area of triangle appearing in a triangulation of some  $\Sigma(\mathcal{C})$ . Our proof is by induction on  $\lceil A(\mathcal{B})/\delta \rceil$ . If  $t := \lceil A(\mathcal{B})/\delta \rceil = 0$  then  $A(\mathcal{B}) = 0$  and  $\Sigma(\mathcal{B})$  fills the entire interior of  $\pi(\mathcal{I})$ , so  $\mathcal{B}$  is maximal in  $\mathcal{M}_{\mathcal{I}}$  and  $\mathcal{B}$  is the only maximal weakly separated collection in  $\mathcal{M}_{\mathcal{I}}$  containing  $\mathcal{B}$ , so the Theorem is vacuously true.

Now, suppose that  $A(\mathcal{B}) > 0$ . So there is some hole (connected region inside  $\pi(\Sigma(\mathcal{B}))$  that is not covered by the tiles) within  $\pi(\Sigma(\mathcal{B}))$ . Let  $K$  and  $L$  be the  $k$ -element sets labeling two consecutive elements on the boundary of the hole. Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be two maximal weakly separated collections in  $\mathcal{M}_{\mathcal{I}}$  containing  $\mathcal{B}$ . Then  $K$  and  $L$  lie in a common face of  $\Sigma(\mathcal{C}_r)$  (for  $r = 1, 2$ .) Triangulate  $\Sigma(\mathcal{C}_r)$  using the edge

$(K, L)$ . Let  $J_r$  be the third vertex of the triangle of  $\Sigma(\mathcal{C}_r)$  containing  $(K, L)$  and lying on the hole side. Let  $T_r$  be the triangle  $(J_r, K, L)$ . We now divide into 2 cases depending on the colors of the triangles  $T_r$ .

**Case 1:**  $T_1$  and  $T_2$  are both white or both black. We present the case that the triangles are white; the other case is very similar. Set  $H = K \cap L$ . Then the  $J_r$  are of the forms  $He_r$  for some  $e_1$  and  $e_2$ . From this we can compute that  $J_1$  and  $J_2$  are weakly separated from each other. Also, by hypothesis,  $\mathcal{B} \cup \{J_1\}$  and  $\mathcal{B} \cup \{J_2\}$  are weakly separated. So  $\mathcal{B} \cup \{J_1, J_2\}$  is weakly separated; complete  $\mathcal{B} \cup \{J_1, J_2\}$  to a maximal weakly separated collection  $\mathcal{C}'$  in  $\mathcal{M}_{\mathcal{I}}$ .

Set  $\mathcal{B}_r = \mathcal{B} \cup \{J_r\}$ . Then  $\Sigma(\mathcal{B}_r)$  is  $\Sigma(\mathcal{B})$  with an extra triangle added on, so  $A(\mathcal{B}_r) \leq A(\mathcal{B}) - \delta$ . Now,  $\mathcal{C}'$  and  $\mathcal{C}_r$  contain  $\mathcal{B}_r$ . So, by induction,  $\mathcal{C}_r$  is  $\mathcal{B}_r$ -equivalent to  $\mathcal{C}'$  within  $\mathcal{M}_{\mathcal{I}}$ . Connecting the chains  $\mathcal{C}_1 \rightarrow \dots \rightarrow \mathcal{C}' \rightarrow \dots \rightarrow \mathcal{C}_2$ , we see that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are  $\mathcal{B}$ -equivalent within  $\mathcal{M}_{\mathcal{I}}$ .

**Case 2:**  $T_1$  is white and  $T_2$  is black: Then we can write  $(J_1, J_2, K, L)$  as  $(Hac, Hbd, Hab, Had)$ . Since  $(J_1, K, L)$  and  $(J_2, K, L)$  are oriented the same way, the triples  $(c, b, d)$  and  $(a, d, b)$  are cyclicly oriented the same way, which shows that  $(a, b, c, d)$  are cyclicly oriented. By Lemma 5.1,  $Hab, Hac, Had, Hbc, Hbd$  and  $Hcd$  are weakly separated from  $\mathcal{B}$ . Set  $\mathcal{B}_1 = \mathcal{B} \cup \{Hac, Hab, Had, Hbc, Hcd\}$  and  $\mathcal{B}_2 = \mathcal{B} \cup \{Hbd, Hab, Had, Hbc, Hcd\}$ , so the  $\mathcal{B}_r$  are weakly separated. Moreover, in  $\Sigma(\mathcal{B}_r)$ , the new vertices that we have added lie immediately adjacent to the edge  $(Hab, Had)$  of  $\Sigma(\mathcal{B})$ , inside the hole of  $\Sigma(\mathcal{B})$ , and hence lie inside  $\pi(\mathcal{I})$ . So, by Lemma 4.1, these new vertices lie in  $\mathcal{M}_{\mathcal{I}}$ , so  $\mathcal{B}_1$  and  $\mathcal{B}_2$  are weakly separated collections in  $\mathcal{M}_{\mathcal{I}}$ .

Complete  $\mathcal{B}_1$  to a maximal weakly separated collection  $\mathcal{C}'_1$  within  $\mathcal{M}_{\mathcal{I}}$ ; define  $\mathcal{C}'_2$  to be the mutation of  $\mathcal{C}'_1$  where we replace  $Hac$  by  $Hbd$ . Then  $\mathcal{C}_1 \cap \mathcal{C}'_1 \supseteq \mathcal{B} \cup \{Hac\}$  and  $\mathcal{C}_2 \cap \mathcal{C}'_2 \supseteq \mathcal{B} \cup \{Hbd\}$ . The complexes  $\Sigma(\mathcal{B} \cup \{Hac\})$  and  $\Sigma(\mathcal{B} \cup \{Hbd\})$  are  $\Sigma(\mathcal{B})$  with one added triangle. So, by induction,  $\mathcal{C}_1$  and  $\mathcal{C}'_1$  are  $(\mathcal{B} \cup \{Hac\})$ -equivalent within  $\mathcal{M}_{\mathcal{I}}$ , and  $\mathcal{C}_2$  and  $\mathcal{C}'_2$  are  $(\mathcal{B} \cup \{Hbd\})$ -equivalent within  $\mathcal{M}_{\mathcal{I}}$ . Chaining together the mutations  $\mathcal{C}_1 \rightarrow \dots \rightarrow \mathcal{C}'_1 \rightarrow \mathcal{C}'_2 \rightarrow \dots \rightarrow \mathcal{C}_2$ , we see that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are  $\mathcal{B}$ -equivalent. □

## 7 Implications of the main result

In this section, we go over the direct implications of Theorem 6.1. Each maximal weakly separated collection corresponds to a reduced plabic graph and the mutation of maximal weakly separated collections corresponds to *square moves* of plabic graphs [6].

We can define a **plabic complex** of a positroid  $\mathcal{M}$ . Consider a simplicial complex where the vertices are labeled with Plücker coordinates and the facets are given by plabic graphs (maximal weakly separated collections) of  $\mathcal{M}$ . A special case of this complex, when  $\mathcal{M}$  is the uniform matroid  $\binom{[n]}{k}$ , was studied in [3]. Hess and Hirsch also conjectured that the complex is a pseudomanifold with boundary.

A simplicial complex is a **pseudomanifold with boundary** if it satisfies the following properties [8]:

- (pure) The facets have the same dimension.
- (non-branching) Each codimension 1 face is a face of one or two facets.
- (strongly connected) Any two facets can be joined by a chain of facets in which each pair of neighboring facets have a common codimension 1 face.

Therefore, another way to interpret Theorem 6.1 is:

**Corollary 7.1.** *Let  $\mathcal{M}$  be a positroid. The plabic complex of  $\mathcal{M}$  is a pseudomanifold with a boundary.*



We end with some questions.

**Question 7.2.** *The simplicial complex coming from both weakly separated collections and strongly separated collections happen to share this property. Is there some way to come up with a uniform theory on separation of sets that exhibit this behavior?*

We have shown that one can mutate a (reduced) plabic graph into another plabic graph while preserving the set of facet labels they have in common. Is this the optimal (in terms of number of mutations needed) way to transform a plabic graph into another?

**Question 7.3.** *Let  $\mathcal{C}$  and  $\mathcal{C}'$  be two different maximal weakly separated collections of a same positroid  $\mathcal{M}_{\mathcal{I}}$ . Consider all possible chains of mutation from  $\mathcal{C}$  to  $\mathcal{C}'$ . Among all the chains that have shortest length (least number of mutations used), is there one that preserves  $\mathcal{C} \cap \mathcal{C}'$ ?*

## Acknowledgements

The first author would like to thank Sergey Fomin for useful discussions.

## References

- [1] Vladimir Danilov, Alexander Karzanov and Gleb Koshevoy. Combined tilings and the purity phenomenon on separated set-systems. arXiv:1401.6418.
- [2] Bernard Leclerc and Andrei Zelevinsky. Quasicommuting families of quantum Plücker coordinates. In *Advances in Math. Sciences (Kirillov's seminar)*, *AMS Translations*, **181** 85–108, 1998.
- [3] Daniel Hess and Benjamin Hirsch. On the topology of weakly and strongly separated set complexes. *Topology and its Applications*, **160(2)**, 328 – 336, 2013.
- [4] Suho Oh. Positroids and Schubert matroids. *Journal of Combinatorial Theory A*, **118(8)**, 2426 – 2435, 2011.
- [5] Suho Oh. Chambers of wiring diagrams. Submitted to *Discrete Mathematics*.
- [6] Suho Oh, Alexander Postnikov and David Speyer. Weak Separation and Plabic Graphs. To appear in *Proceedings of London Mathematical Society*.
- [7] Alexander Postnikov. Total positivity, grassmannians, and networks, 2006. <http://math.mit.edu/~apost/papers/tpgrass.pdf>
- [8] Edwin Spanier *Algebraic Topology*, McGraw-Hill series in higher mathematics, **55(1)**, Springer, 1994.
- [9] Joshua Scott. Quasi-commuting families of quantum minors. *Journal of Algebra*, **290(1)**, 204 – 220, 2005.