



HAL
open science

Yang-Baxter basis of Hecke algebra and Casselman's problem (extended abstract)

Maki Nakasuji, Hiroshi Naruse

► **To cite this version:**

Maki Nakasuji, Hiroshi Naruse. Yang-Baxter basis of Hecke algebra and Casselman's problem (extended abstract). 28-th International Conference on Formal Power Series and Algebraic Combinatorics, Simon Fraser University, Jul 2016, Vancouver, Canada. 10.46298/dmtcs.6370 . hal-02166345

HAL Id: hal-02166345

<https://hal.science/hal-02166345>

Submitted on 26 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Yang-Baxter basis of Hecke algebra and Casselman's problem (extended abstract)

Maki Nakasuji^{1†} and Hiroshi Naruse^{2‡}

¹*Sophia University, Japan*

²*University of Yamanashi, Japan*

Abstract. We generalize the definition of Yang-Baxter basis of type A Hecke algebra introduced by A.Lascoux, B.Leclerc and J.Y.Thibon (Letters in Math. Phys., 40 (1997), 75–90) to all the Lie types and prove their duality. As an application we give a solution to Casselman's problem on Iwahori fixed vectors of principal series representation of p -adic groups.

Résumé. Nous généralisons la définition de la base de l'algèbre de Hecke de Yang-Baxter de type A introduit par A.Lascoux, B.Leclerc and J.Y.Thibon (Letters in Math. Phys., 40 (1997), 75–90) à tous les types de Lie et prouvons la dualité. Comme application nous donnons une solution du problème de Casselman sur les vecteurs fixés par le sous-groupe d'Iwahori de la série principale des groupes p -adiques.

Keywords. Yang-Baxter basis, Iwahori fixed vector, Casselman's problem, Demazure-Lusztig operator

1 Introduction

Yang-Baxter basis of Hecke algebra of type A was defined in the paper of Lascoux-Leclerc-Thibon [LLT]. There is also a modified version in [Las]. First we generalize the latter version to all the Lie types. Then we will solve the Casselman's problem on the basis of Iwahori fixed vectors using Yang-Baxter basis and Demazure-Lusztig type operator. This paper is an extended abstract and the detailed proofs will appear in [NN].

2 Generic Hecke algebra

2.1 Root system, Weyl group and generic Hecke algebra

Let $\mathcal{R} = (\Lambda, \Lambda^*, R, R^*)$ be a (reduced) semisimple root data cf. [Dem]. More precisely $\Lambda \simeq \mathbb{Z}^r$ is a weight lattice with $\text{rank } \Lambda = r$. There is a pairing $\langle \cdot, \cdot \rangle: \Lambda^* \times \Lambda \rightarrow \mathbb{Z}$. $R \subset \Lambda$ is a root system with simple roots $\{\alpha_i\}_{1 \leq i \leq r}$ and positive roots R^+ . $R^* \subset \Lambda^*$ is the set of coroots, and there is a bijection $R \rightarrow R^*$, $\alpha \mapsto \alpha^*$. We also denote the coroot $\alpha^* = h_\alpha$. The Weyl group W of \mathcal{R} is generated by

[†]Email: nakasuji@sophia.ac.jp. KAKENHI 15K17519.

[‡]Email: hnaruse@yamanashi.ac.jp. KAKENHI 25400041.

simple reflections $S = \{s_i\}_{1 \leq i \leq r}$. The action of W on Λ is given by $s_i(\lambda) = \lambda - \langle \alpha_i^*, \lambda \rangle \alpha_i$ for $\lambda \in \Lambda$. We define generic Hecke algebra $H_{t_1, t_2}(W)$ over $\mathbb{Z}[t_1, t_2]$ with two parameters t_1, t_2 as follows. Generators are $h_i = h_{s_i}$, with relations $(h_i - t_1)(h_i - t_2) = 0$ for $1 \leq i \leq r$ and the braid relations $\underbrace{h_i h_j \cdots}_{m_{i,j}} = \underbrace{h_j h_i \cdots}_{m_{i,j}}$, where $m_{i,j}$ is the order of $s_i s_j$ for $1 \leq i < j \leq r$. We need to extend the coefficients

to the quotient field of the group algebra $\mathbb{Z}[\Lambda]$. An element of $\mathbb{Z}[\Lambda]$ is denoted as $\sum_{\lambda \in \Lambda} c_\lambda e^\lambda$. The Weyl group acts on $\mathbb{Z}[\Lambda]$ by $w(e^\lambda) = e^{w\lambda}$. We extend the coefficient ring $\mathbb{Z}[t_1, t_2]$ of $H_{t_1, t_2}(W)$ to

$$Q_{t_1, t_2}(\Lambda) := \mathbb{Z}[t_1, t_2] \otimes Q(\mathbb{Z}[\Lambda])$$

where $Q(\mathbb{Z}[\Lambda])$ is the quotient field of $\mathbb{Z}[\Lambda]$.

$$H_{t_1, t_2}^{Q(\Lambda)}(W) := Q_{t_1, t_2}(\Lambda) \otimes_{\mathbb{Z}[t_1, t_2]} H_{t_1, t_2}(W).$$

For $w \in W$, an expression of $w = s_{i_1} s_{i_2} \cdots s_{i_\ell}$ with minimal number of generators $s_{i_k} \in S$ is called a reduced expression in which case we write $\ell(w) = \ell$ and call it the length of w . Then $h_w = h_{i_1} h_{i_2} \cdots h_{i_\ell}$ is well defined and $\{h_w\}_{w \in W}$ forms a $Q_{t_1, t_2}^{Q(\Lambda)}$ -basis of $H_{t_1, t_2}^{Q(\Lambda)}(W)$.

2.2 Yang-Baxter basis and its properties

Yang-Baxter basis was introduced in the paper [LLT] to investigate the relation with Schubert calculus. There is also a variant in [Las] for type A case. We generalize that results to all Lie types.

For $\lambda \in \Lambda$, we define $E(\lambda) = e^{-\lambda} - 1$. Then $E(\lambda + \nu) = E(\lambda) + E(\nu) + E(\lambda)E(\nu)$. In particular, if $\lambda \neq 0$, $\frac{1}{E(\lambda)} + \frac{1}{E(-\lambda)} = -1$.

Proposition 1 For $\lambda \in \Lambda$, if $\lambda \neq 0$, let $h_i(\lambda) := h_i + \frac{t_1 + t_2}{E(\lambda)}$. Then these satisfy the **Yang-Baxter relations**, i.e. if we write $[p, q] := p\lambda + q\nu$ for fixed $\lambda, \nu \in \Lambda$, the following equations hold. We assume all appearance of $[p, q]$ is nonzero.

$$\begin{aligned} h_i([1, 0])h_j([0, 1]) &= h_j([0, 1])h_i([1, 0]) && \text{if } m_{i,j} = 2 \\ h_i([1, 0])h_j([1, 1])h_i([0, 1]) &= h_j([0, 1])h_i([1, 1])h_j([1, 0]) && \text{if } m_{i,j} = 3 \\ h_i([1, 0])h_j([1, 1])h_i([1, 2])h_j([0, 1]) &= h_j([0, 1])h_i([1, 2])h_j([1, 1])h_i([1, 0]) && \text{if } m_{i,j} = 4 \\ h_i([1, 0])h_j([1, 1])h_i([2, 3]) &= h_j([0, 1])h_i([1, 3])h_j([1, 2]) \\ \times h_j([1, 2])h_i([1, 3])h_j([0, 1]) &= \times h_i([2, 3])h_j([1, 1])h_i([1, 0]) && \text{if } m_{i,j} = 6 \end{aligned}$$

Proof: We can prove these equations by direct calculations. □

Remark 1 In [Che] I. Cherednik treated Yang-Baxter relation in more general setting. There is also a related work [Kat] by S. Kato and the proof of Theorem 2.4 in [Kat] suggests a uniform way to prove Yang-Baxter relations without direct calculations.

We use the Bruhat order $x \leq y$ on elements $x, y \in W$ (cf.[Hum]). Following [Las] we define the Yang-Baxter basis Y_w for $w \in W$ recursively as follows.

$$Y_e := 1, Y_w := Y_{w'}(h_i + \frac{t_1+t_2}{w'E(\alpha_i)}) \text{ if } w = w's_i > w'.$$

Using the Yang-Baxter relation above it is easy to see that Y_w does not depend on a reduced expression of w . As the leading term of Y_w with respect to the Bruhat order is h_w , they also form a $Q_{t_1,t_2}(\Lambda)$ -basis $\{Y_w\}_{w \in W}$ of $H_{t_1,t_2}^{Q(\Lambda)}(W)$. We are interested in the transition coefficients $p(w, v)$ and $\tilde{p}(w, v) \in Q_{t_1,t_2}(\Lambda)$ between the two basis $\{Y_w\}_{w \in W}$ and $\{h_w\}_{w \in W}$, i.e.

$$Y_v = \sum_{w \leq v} p(w, v)h_w, \text{ and } h_v = \sum_{w \leq v} \tilde{p}(w, v)Y_w.$$

Take a reduced expression of v e.g. $v = s_{i_1} \cdots s_{i_\ell}$ where $\ell = \ell(v)$ is the length of v (cf. [Hum]). Then Y_v is expressed as follows.

$$Y_v = \prod_{j=1}^{\ell} \left(h_{i_j} + \frac{t_1 + t_2}{E(\beta_j)} \right)$$

where $\beta_j := s_{i_1} \cdots s_{i_{j-1}}(\alpha_{i_j})$ for $j = 1, \dots, \ell$. The set $R(v) := \{\beta_1, \dots, \beta_\ell\} \subset R^+$ is independent of the reduced expression of v . The Yang-Baxter basis defined in [LLT] is normalized as follows.

$$Y_v^{LLT} := \left(\prod_{j=1}^{\ell} \frac{E(\beta_j)}{t_1 + t_2} \right) Y_v = \prod_{j=1}^{\ell} \left(\frac{E(\beta_j)}{t_1 + t_2} h_{i_j} + 1 \right).$$

Remark 2 *The relation to K-theory Schubert calculus is as follows. If we set $t_1 = 0, t_2 = -1$ and replacing α_i by $-\alpha_i$. Then the coefficient of h_w in Y_v^{LLT} is the localization $\psi^w(v)$ at v of the equivariant K-theory Schubert class ψ^w (cf. [LSS]).*

Let w_0 be the longest element in W . Define $Q_{t_1,t_2}(\Lambda)$ -algebra homomorphism $\Omega : H_{t_1,t_2}^{Q(\Lambda)} \rightarrow H_{t_1,t_2}^{Q(\Lambda)}$ by $\Omega(h_w) = h_{w_0 w w_0}$. Let \star be the ring homomorphism on $\mathbb{Z}[\Lambda]$ induced by $\star(e^\lambda) = e^{-\lambda}$ and extend to $Q_{t_1,t_2}(\Lambda)$.

Proposition 2 (Lascoux [Las] Lemma 1.8.1 for type A case) *For $v \in W$,*

$$\Omega(Y_{w_0 v w_0}) = \star[w_0(Y_v)]$$

where W acts only on the coefficients.

Proof: When $\ell(v) > 0$ there exists $s \in S$ such that $v = v's > v'$. Using the induction assumption on v' , we get the formula for v . □

Taking the coefficient of h_w in the above equation, we get

Corollary 1

$$p(w_0 w w_0, w_0 v w_0) = \star[w_0 p(w, v)].$$

2.3 Inner product and orthogonality

Define inner product $(\ , \)^H$ on $H_{t_1, t_2}^{Q(\Lambda)}(W)$ by $(f, g)^H :=$ the coefficient of h_{w_0} in fg^\vee , where $g^\vee = \sum c_w h_{w^{-1}}$ if $g = \sum c_w h_w$. It is easy to see that $(fh_s, g)^H = (f, gh_s)^H$ for $f, g \in H_{t_1, t_2}^{Q(\Lambda)}(W)$ and $s \in S$. There is an involution $\hat{\cdot} : H_{t_1, t_2}^{Q(\Lambda)} \rightarrow H_{t_1, t_2}^{Q(\Lambda)}$ defined by $\hat{h}_i = h_i - (t_1 + t_2), \hat{t}_1 = -t_2, \hat{t}_2 = -t_1$. It is easy to see that $\hat{h}_s h_s = -t_1 t_2$ for $s \in S$.

The following proposition is due to A.Lascoux for the type A case [Las] P.33.

Proposition 3 For all $v, w \in W$,

$$(h_v, \hat{h}_{w_0 w})^H = \delta_{v, w}.$$

Proof: We can use induction on the length $\ell(v)$ of v to prove the equation. □

We have another orthogonality between Y_v and $w_0(Y_{w_0 w})$.

Proposition 4 (Type A case was due to [LLT] Theorem 5.1, [Las] Theorem 1.8.4.)

For all $v, w \in W$,

$$(Y_v, w_0(Y_{w_0 w}))^H = \delta_{v, w}.$$

Proof: We use induction on $\ell(v)$ and use the fact that if $s \in S$ and $u \in W$, then $Y_u h_s = aY_{us} + bY_s$ for some $a, b \in Q_{t_1, t_2}(\Lambda)$. □

2.4 Duality between the transition coefficients

Recall that we have two transition coefficients $p(w, v), \tilde{p}(w, v) \in Q_{t_1, t_2}(\Lambda)$ defined by the following expansions.

$$Y_v = \sum_{w \leq v} p(w, v) h_w$$

$$h_v = \sum_{w \leq v} \tilde{p}(w, v) Y_w$$

Below gives a relation between them.

Theorem 1 (Lascoux [Las] Corollary 1.8.5 for type A case) For $w, v \in W$,

$$\tilde{p}(w, v) = (-1)^{\ell(v) - \ell(w)} p(vw_0, ww_0).$$

Proof: We will calculate $(h_v, w_0(Y_{w_0 w}))^H$ in two ways. As $h_v = \sum_{w \leq v} \tilde{p}(w, v) Y_w$,

$$(h_v, w_0(Y_{w_0 w}))^H = \tilde{p}(w, v)$$

by the orthogonality on Y_v (Proposition 4). On the other hand, as $h_i + \frac{t_1 + t_2}{E(\beta)} = \hat{h}_i - \frac{t_1 + t_2}{E(-\beta)}$ for $\beta \in R$, we can expand Y_v in terms of \hat{h}_w as follows.

$$Y_v = \sum_{w \leq v} (-1)^{\ell(v) - \ell(w)} \star [p(w, v)] \hat{h}_w.$$

So we have

$$w_0(Y_{w_0 w}) = \sum_{w_0 v \leq w_0 w} (-1)^{\ell(v) - \ell(w)} w_0[\star p(w_0 v, w_0 w)] \hat{h}_{w_0 v}.$$

Then using the orthogonality on h_v (Proposition 3) and Corollary 1,

$$(h_v, w_0(Y_{w_0 w}))^H = (-1)^{\ell(v) - \ell(w)} w_0[\star p(w_0 v, w_0 w)] = (-1)^{\ell(v) - \ell(w)} p(vw_0, ww_0).$$

The theorem is proved. □

2.5 Recurrence relations

Here we give some recurrence relations on $p(w, v)$ and $\tilde{p}(w, v)$.

Proposition 5 (left p) For $w \in W$ and $s \in S$, if $sv > v$ then

$$p(w, sv) = \begin{cases} \frac{t_1+t_2}{E(\alpha_s)} s[p(w, v)] - t_1 t_2 s[p(sw, v)] & \text{if } sw > w \\ (t_1 + t_2) \left(\frac{1}{E(\alpha_s)} + 1 \right) s[p(w, v)] + s[p(sw, v)] & \text{if } sw < w. \end{cases}$$

Proof: By the definition we have $Y_{sv} = Y_s s[Y_v]$ from which we can deduce the recurrence formula. □

We note that by this recurrence we can identify $p(w, v)$ as a coefficient of transition between two bases of the space of Iwahori fixed vectors cf. Theorem 3 below.

Proposition 6 (right p) For $w \in W$ and $s \in S$, if $vs > v$ then

$$p(w, vs) = \begin{cases} \frac{t_1+t_2}{vE(\alpha_s)} p(w, v) - t_1 t_2 p(ws, v) & \text{if } ws > w \\ (t_1 + t_2) \left(\frac{1}{vE(\alpha_s)} + 1 \right) p(w, v) + p(ws, v) & \text{if } ws < w. \end{cases}$$

Proof: We can use the equation $Y_{vs} = Y_v v[Y_s]$ and taking the coefficient of h_w , we get the formula. □

Proposition 7 (left \tilde{p}) For $w \in W$ and $s \in S$, if $sv > v$ then

$$\tilde{p}(w, sv) = \begin{cases} -\frac{t_1+t_2}{E(\alpha_s)} \tilde{p}(w, v) + \left(1 + \frac{t_1+t_2}{E(\alpha_s)}\right) \left(1 + \frac{t_1+t_2}{E(-\alpha_s)}\right) s[\tilde{p}(sw, v)] & \text{if } sw > w \\ -\frac{t_1+t_2}{E(\alpha_s)} \tilde{p}(w, v) + s[\tilde{p}(sw, v)] & \text{if } sw < w. \end{cases}$$

Proof: We can prove the recurrence relation using Corollary 2 below. □

Proposition 8 (right \tilde{p}) For $w \in W$ and $s \in S$, if $vs > v$ then

$$\tilde{p}(w, vs) = \begin{cases} -\frac{t_1+t_2}{wE(\alpha_s)} \tilde{p}(w, v) + \left(1 + \frac{t_1+t_2}{wE(\alpha_s)}\right) \left(1 + \frac{t_1+t_2}{wE(-\alpha_s)}\right) \tilde{p}(ws, v) & \text{if } ws > w \\ -\frac{t_1+t_2}{wE(\alpha_s)} \tilde{p}(w, v) + \tilde{p}(ws, v) & \text{if } ws < w. \end{cases}$$

Proof: We can prove the recurrence relation using Corollary 2 below. □

3 Kostant-Kumar's twisted group algebra

Let $Q_{t_1, t_2}^{KK}(W) := Q_{t_1, t_2}(\Lambda) \# \mathbb{Z}[W]$ be the (generic) twisted group algebra of Kostant-Kumar. Its element is of the form $\sum_{w \in W} f_w \delta_w$ for $f_w \in Q_{t_1, t_2}(\Lambda)$ and the product is defined by

$$\left(\sum_{w \in W} f_w \delta_w \right) \left(\sum_{u \in W} g_u \delta_u \right) = \sum_{w, u \in W} f_w w(g_u) \delta_{wu}.$$

Define $y_i \in Q_{t_1, t_2}^{KK}(W)$ ($i = 1, \dots, r$) by

$$y_i := A_i \delta_i + B_i \text{ where } A_i := \frac{t_1 + t_2 e^{-\alpha_i}}{1 - e^{\alpha_i}}, B_i := \frac{t_1 + t_2}{1 - e^{-\alpha_i}}.$$

Proposition 9 *We have the following equations.*

- (1) $(y_i - t_1)(y_i - t_2) = 0$ for $i = 1, \dots, r$.
- (2) $\underbrace{y_i y_j \cdots}_{m_{i,j}} = \underbrace{y_j y_i \cdots}_{m_{i,j}}$, where $m_{i,j}$ is the order of $s_i s_j$.

Proof: These equations can be shown by direct calculations. □

By this proposition we can define $y_w := y_{i_1} \cdots y_{i_\ell}$ for a reduced expression $w = s_{i_1} \cdots s_{i_\ell}$. These $\{y_w\}_{w \in W}$ become a $Q_{t_1, t_2}(\Lambda)$ -basis of $Q_{t_1, t_2}^{KK}(W)$.

Remark 3 *This operator y_i can be seen as a generic Demazure-Lusztig operator. When $t_1 = -1, t_2 = q$, it becomes $y_{s_i}^q$ in Kumar's book[Kum](12.2.E(9)). We can also set A_i which satisfies*

$$A_i A_{-i} = \frac{(t_1 + t_2 e^{-\alpha_i})(t_1 + t_2 e^{\alpha_i})}{(1 - e^{\alpha_i})(1 - e^{-\alpha_i})}.$$

For example, if we set $A_i = \frac{t_1 + t_2 e^{\alpha_i}}{1 - e^{\alpha_i}}$ and $t_1 = q, t_2 = -1$ and replace α_i by $-\alpha_i$, it becomes Lusztig's T_{s_i} [Lul]. If we set $A_i = -\frac{t_1 + t_2 e^{\alpha_i}}{1 - e^{-\alpha_i}}$ and $t_1 = -1, t_2 = v$ and replace α_i by $-\alpha_i$, it becomes \mathcal{T}_i in [BBL].

We can define a $Q_{t_1, t_2}(\Lambda)$ -module isomorphism $\Phi : Q_{t_1, t_2}^{KK}(W) \rightarrow H_{t_1, t_2}^{Q(\Lambda)}(W)$ by $\Phi(y_w) = h_w$.

Let $\Delta_{s_i} := A_i \delta_i$. Define $A(w) := \prod_{\beta \in R(w)} \frac{t_1 + t_2 e^{-\beta}}{1 - e^\beta}$ and $\Delta_w := A(w) \delta_w$. Then it becomes that

$\Delta_{s_{i_1}} \cdots \Delta_{s_{i_\ell}} = A(w) \delta_w = \Delta_w$. In particular, Δ_{s_i} 's satisfy the braid relations. We can show below by induction on length $\ell(w)$.

Theorem 2 *For $w \in W$, we have*

$$\Phi(\Delta_w) = Y_w.$$

Proof: If $w = s_i$, $\Delta_{s_i} = A_i \delta_i = y_i - B_i$. Therefore $\Phi(\Delta_{s_i}) = h_i - B_i = h_i + \frac{t_1 + t_2}{E(\alpha_i)} = Y_{s_i}$. If

$s_i w > w$, by induction hypothesis we can assume $\Phi(\Delta_w) = Y_w = \sum_{u \leq w} p(u, w) h_u$. As Φ is a $Q_{t_1, t_2}(\Lambda)$ -

isomorphism, it follows that $\Delta_w = \sum_{u \leq w} p(u, w) y_u$. Then $\Delta_{s_i w} = \Delta_{s_i} \Delta_w = A_i \delta_i \sum_{u \leq w} p(u, w) y_u =$

$\sum_{u \leq w} s_i [p(u, w)] A_i \delta_i y_u = \sum_{u \leq w} s_i [p(u, w)] (y_i - B_i) y_u = \sum_{u \leq s_i w} p(u, s_i w) y_u$. We used the recurrence relation (Proposition 5) for the last equality. Therefore $\Phi(\Delta_{s_i w}) = \sum_{u \leq s_i w} p(u, s_i w) h_u = Y_{s_i w}$. The theorem is proved. \square

Corollary 2 (Explicit formula for $\tilde{p}(w, v)$)

Let $v = s_{i_1} \cdots s_{i_\ell}$ be a reduced expression. Then we have

$$\tilde{p}(w, v) = \frac{1}{A(w)} \sum_{\epsilon = (\epsilon_1, \dots, \epsilon_\ell) \in \{0, 1\}^\ell, s_{i_1}^{\epsilon_1} \cdots s_{i_\ell}^{\epsilon_\ell} = w} \prod_{j=1}^{\ell} C_j(\epsilon)$$

where for $\epsilon = (\epsilon_1, \dots, \epsilon_\ell) \in \{0, 1\}^\ell$, $C_j(\epsilon) := s_{i_1}^{\epsilon_1} s_{i_2}^{\epsilon_2} \cdots s_{i_{j-1}}^{\epsilon_{j-1}} (\delta_{\epsilon_j, 1} A_{i_j} + \delta_{\epsilon_j, 0} B_{i_j})$.

Proof: Taking the inverse image of the map Φ , the equality $h_v = \sum_{w \leq v} \tilde{p}(w, v) Y_w$ becomes

$$y_v = \sum_{w \leq v} \tilde{p}(w, v) \Delta_w = \sum_{w \leq v} \tilde{p}(w, v) A(w) \delta_w.$$

As $v = s_{i_1} \cdots s_{i_\ell}$ is a reduced expression, $y_v = y_{s_{i_1}} \cdots y_{s_{i_\ell}} = (A_{i_1} \delta_{i_1} + B_{i_1} \delta_e) \cdots (A_{i_\ell} \delta_{i_\ell} + B_{i_\ell} \delta_e)$. By expanding this we get the formula. \square

Remark 4 Using Theorem 1, we also have a closed form for $p(w, v)$. We have another conjectural formula for $p(w, v)$ using λ -chain cf. [Nar].

Example 1 Type A_2 . We use notation $A_{-1} = \star(A_1)$, $B_{-1} = \star(B_1)$, $B_{12} = \frac{t_1 + t_2}{1 - e^{-(\alpha_1 + \alpha_2)}}$.

When $v = s_1 s_2 s_1$, $w = s_1$, then $\epsilon = (1, 0, 0)$, $(0, 0, 1)$ and

$$\tilde{p}(s_1, s_1 s_2 s_1) = (A_1 B_{12} B_{-1} + B_1 B_2 A_1) / A_1 = B_{12} B_{-1} + B_1 B_2 = B_2 B_{12}.$$

When $v = s_1 s_2 s_1$, $w = s_2$, then $\epsilon = (0, 1, 0)$ and

$$\tilde{p}(s_2, s_1 s_2 s_1) = (B_1 A_2 B_{12}) / A_2 = B_1 B_{12}.$$

When $v = s_1 s_2 s_1$, $w = e$, then $\epsilon = (0, 0, 0)$, $(1, 0, 1)$ and

$$\tilde{p}(e, s_1 s_2 s_1) = B_1 B_2 B_1 + A_1 B_{12} A_{-1}.$$

4 Casselman's problem

In his paper [Cas] B. Casselman gave a problem concerning transition coefficients between two bases in the space of Iwahori fixed vectors of a principal series representation of a p -adic group. We relate the problem with the Yang-Baxter basis and give an answer to the problem.

4.1 Principal series representations of p -adic group and Iwahori fixed vector

We follow the notations of M.Reeder [Re1, Re2]. Let G be a connected reductive p -adic group over a non-archimedean local field F . For simplicity we restrict to the case of split semisimple G . Associated to F , there is the ring of integer \mathcal{O} , the prime ideal \mathfrak{p} with a generator ϖ , and the residue field with $q = |\mathcal{O}/\mathfrak{p}|$ elements. Let P be a minimal parabolic subgroup (Borel) of G , and A be the maximal split torus of P so that $A \simeq (F^*)^r$ where r is the rank of G . For an unramified quasi-character τ of A , i.e. a group homomorphism $\tau : A \rightarrow \mathbb{C}^*$ which is trivial on $A_0 = A \cap K$, where $K = G(\mathcal{O})$ is a maximal compact subgroup of G . Let $T = \mathbb{C}^* \otimes X^*(A)$ be the complex torus dual to A , where $X^*(A)$ is the group of rational characters on A , i.e. $X^*(A) = \{\lambda : A \rightarrow F^*, \text{ algebraic group homomorphism}\}$. We have a pairing $\langle, \rangle : A/A_0 \times T \rightarrow \mathbb{C}^*$ given by $\langle a, z \otimes \lambda \rangle = z^{\text{val}(\lambda(a))}$. This gives an identification $T \simeq X^{nr}(A)$ of T with the set of unramified quasi-characters on A (cf. [Bum] Exercise 18,19).

Let $\Delta \subset X^*(A)$ be the set of roots of A in G , Δ^+ be the set of positive roots corresponding to P and $\Sigma \subset \Delta^+$ be the set of simple roots. For a root $\alpha \in \Delta$, we define $e_\alpha \in X^*(T)$ by

$$e_\alpha(\tau) = \langle h_\alpha(\varpi), \tau \rangle$$

for $\tau \in T$ where $h_\alpha : F^* \rightarrow A$ is the one parameter subgroup (coroot) corresponding to α .

Remark 5 As the definition shows, e_α is defined using the coroot $\alpha^* = h_\alpha$. So it should be parametrized by α^* , but for convenience we follow the notation of [Re1]. Later we will identify $e_\alpha(\alpha \in \Delta = R^*)$ with $e^\alpha(\alpha \in R = \Delta^*)$ by the map $*$: $\Delta \rightarrow R$ of root data.

W acts on right of $X^{nr}(A)$ so that $\tau^w(a) = \tau(waw^{-1})$ for $a \in A$, $\tau \in T$ and $w \in W$. The action of W on $X^*(T)$ is given by $(we_\alpha)(\tau) = e_{w\alpha}(\tau) = e_\alpha(\tau^w)$ for $\alpha \in \Delta$, $\tau \in T$ and $w \in W$.

The principal series representation $I(\tau)$ of G associated to an unramified quasicharacter τ of A is defined as follows. As a vector space over \mathbb{C} it consists of locally constant functions on G with values in \mathbb{C} which satisfy the left relative invariance properties with respect to P where τ is extended to P with trivial value on the unipotent radical N of $P = AN$.

$$I(\tau) := \text{Ind}_P^G(\tau) = \{f : G \rightarrow \mathbb{C} \text{ loc. const. function} \mid f(pg) = \tau\delta^{1/2}(p)f(g) \text{ for } \forall p \in P, \forall g \in G\}.$$

Here δ is the modulus of P . The action of G on $I(\tau)$ is defined by right translation, i.e. for $g \in G$ and $f \in I(\tau)$, $(\pi(g)f)(x) = f(xg)$.

Let B be the Iwahori subgroup which is the inverse image $\pi^{-1}(P(\mathbb{F}_q))$ of the Borel subgroup $P(\mathbb{F}_q)$ of $G(\mathbb{F}_q)$ by the projection $\pi : G(\mathcal{O}) \rightarrow G(\mathbb{F}_q)$. Then we define $I(\tau)^B$ to be the space of Iwahori fixed vectors in $I(\tau)$, i.e.

$$I(\tau)^B := \{f \in I(\tau) \mid f(gb) = f(g) \text{ for } \forall b \in B, \forall g \in G\}.$$

This space has a natural basis $\{\varphi_w^\tau\}_{w \in W}$. $\varphi_w^\tau \in I(\tau)^B$ is supported on PwB and satisfies

$$\varphi_w^\tau(pwb) = \tau\delta^{1/2}(p) \text{ for } \forall p \in P, \forall b \in B.$$

4.2 Intertwiner and Casselman's basis

From now on we always assume that τ is regular i.e. the stabilizer $W_\tau = \{w \in W \mid \tau^w = \tau\}$ is trivial. The intertwining operator $\mathcal{A}_w^\tau : I(\tau) \rightarrow I(\tau^w)$ is defined by

$$\mathcal{A}_w^\tau(f)(g) := \int_{N_w} f(wng)dn$$

where $N_w := N \cap w^{-1}N_-w$, with N_- being the unipotent radical of opposite parabolic P_- which corresponds to the negative roots Δ^- . The integral is convergent when $|e_\alpha(\tau)| < 1$ for all $\alpha \in \Delta^+$ such that $w\alpha \in \Delta^-$ (cf. [Bum] Proposition 63), and may be meromorphically continued to all τ . It has the property that for $x, y \in W$ with $\ell(xy) = \ell(x) + \ell(y)$, then

$$\mathcal{A}_y^\tau \mathcal{A}_x^\tau = \mathcal{A}_{xy}^\tau.$$

The Casselman's basis $\{f_w^\tau\}_{w \in W}$ of $I(\tau)^B$ is defined as follows. $f_w^\tau \in I(\tau)^B$ and

$$\mathcal{A}_y^\tau f_w^\tau(1) = \begin{cases} 1 & \text{if } y = w \\ 0 & \text{if } y \neq w. \end{cases}$$

M.Reeder characterizes this using the action of affine Hecke algebra (cf. [Re2] Section 2). The affine Hecke algebra $\mathcal{H} = \mathcal{H}(G, B)$ is the convolution algebra of B bi-invariant locally constant functions on G with values in \mathbb{C} . By the theorem of Iwahori-Matsumoto it can be described by generators and relations. The basis $\{T_w\}_{w \in \tilde{W}_{aff}}$ consists of characteristic functions $T_w := ch_{BwB}$ of double coset BwB . Let \mathcal{H}_W be the Hecke algebra of the finite Weyl group W generated by the simple reflections s_α for simple roots $\alpha \in \Sigma$. As a vector space \mathcal{H} is the tensor product of two subalgebras $\mathcal{H} = \Theta \otimes \mathcal{H}_W$. The subalgebra Θ is commutative and isomorphic to the coordinate ring of the complex torus T with a basis $\{\theta_a \mid a \in A/A_0\}$, where θ_a is defined as follows (cf. [Lu2]). Define $A^- := \{a \in A \mid |\alpha(a)|_F \leq 1 \forall \alpha \in \Sigma\}$. For $a \in A$, choose $a_1, a_2 \in A^-$ such that $a = a_1 a_2^{-1}$. Then $\theta_a = q^{(\ell(a_1) - \ell(a_2))/2} T_{a_1} T_{a_2}^{-1}$ where for $x \in G$, $\ell(x)$ is the length function defined by $q^{\ell(x)} = [BxB : B]$ and $T_x \in \mathcal{H}$ is the characteristic function of BxB .

By Lemma (4.1) of [Re1], there exists a unique $f_w^\tau \in I(\tau)_w \cap I(\tau)^B$ for each $w \in W$ such that

- (1) $f_w^\tau(w) = 1$ and
- (2) $\pi(\theta_a) f_w^\tau = \tau^w(a) f_w^\tau$ for all $a \in A$.

Here $I(\tau)_w := \{f \in I(\tau) \mid \text{support of } f \text{ is contained in } \bigcup_{x \geq w} P_x P\}$.

4.3 Transition coefficients

Let

$$f_w^\tau = \sum_{w \leq v} a_{w,v}(\tau) \varphi_v^\tau$$

and

$$\varphi_w^\tau = \sum_{w \leq v} b_{w,v}(\tau) f_v^\tau.$$

The Casselman's problem is to find an explicit formula for $a_{w,v}(\tau)$ and $b_{w,v}(\tau)$.

To relate the results in Sections 2 and 3 with the Casselman's problem, in this subsection we specialize the parameters $t_1 = -q^{-1}$, $t_2 = 1$ and take tensor product with the complex field \mathbb{C} . For example, the Yang-Baxter basis Y_w will become a $Q_{t_1, t_2}(\Lambda) \otimes \mathbb{C}$ basis in $H_{t_1, t_2}^{Q(\Lambda)}(W)_{\mathbb{C}} = H_{t_1, t_2}^{Q(\Lambda)}(W) \otimes \mathbb{C}$. The generic Demazure-Lusztig operator defined in Section 3 will become

$$y_i := A_i \delta_i + B_i \text{ where } A_i := \frac{-q^{-1} + e^{-\alpha_i}}{1 - e^{\alpha_i}}, B_i := \frac{-q^{-1} + 1}{1 - e^{-\alpha_i}}.$$

Then $(y_i + q^{-1})(y_i - 1) = 0$.

Theorem 3 We identify e^α with e_α (cf. Remark 5). Then,

$$a_{w,v}(\tau) = \tilde{p}(w, v)(\tau)|_{t_1=-q^{-1}, t_2=1}$$

$$b_{w,v}(\tau) = p(w, v)(\tau)|_{t_1=-q^{-1}, t_2=1}.$$

Proof: $b_{w,v}$'s satisfy the same recurrence relation (Proposition 5 with $t_1 = -q^{-1}, t_2 = 1$) as $p(w, v)$'s (cf. [Re2] Proposition (2.2)). The initial condition $b_{w,w} = p(w, w) = 1$ leads to the second equation. The first equation then also holds. Note that the $b_{y,w}$ in [Re2] is our $b_{w,y}$. □

Remark 6 There is also a direct proof that does not use recurrence relation cf. [NN].

Corollary 3 We have a closed formula for $a_{w,v}(\tau)$ and $b_{w,v}(\tau)$ by Corollary 2 and Theorem 1.

Corollary 4 For $v \in W$, we have

$$\sum_{w \leq v} b_{w,v} = \prod_{\beta \in R(v)} \frac{1 - q^{-1}e^\beta}{1 - e^\beta},$$

and

$$\sum_{w \leq v} b_{w,v}(-q^{-1})^{\ell(w)} = \prod_{\beta \in R(v)} \frac{1 - q^{-1}}{1 - e^\beta}.$$

Proof: When $t_1 = -q^{-1}, t_2 = 1$, we can specialize h_i to 1 and we get the first equation from the definition of Y_v , since $1 + \frac{(1-q^{-1})e^\beta}{1-e^\beta} = \frac{1-q^{-1}e^\beta}{1-e^\beta}$. We can also specialize h_i to $-q^{-1}$ and $-q^{-1} + \frac{(1-q^{-1})e^\beta}{1-e^\beta} = \frac{1-q^{-1}}{1-e^\beta}$ gives the second equation. □

Remark 7 The left hand side of the first equation in Corollary 4 is $m(e, v^{-1})$ in [BN]. So this gives another proof of Theorem 1.4 in [BN].

4.4 Whittaker function

M.Reeder [Re2] specified a formula for the Whittaker function $\mathcal{W}_\tau(f_w^\tau)$ and using $b_{w,v}$, he got a formula for $\mathcal{W}_\tau(\varphi_w^\tau)$. For $a \in A$, let $\lambda_a \in X^*(T)$ be

$$\lambda_a(z \otimes \mu) = z^{val(\mu(a))} \text{ for } z \in \mathbb{C}^*, \mu \in X^*(A).$$

Formally the result of M.Reeder [Re2] Corollary (3.2) is written as follows. For $w \in W$ and $a \in A^-$,

$$\mathcal{W}(\varphi_w)(a) = \delta^{1/2}(a) \sum_{w \leq y} b_{w,y} y \left[\lambda_a \prod_{\beta \in R^+ - R(y)} \frac{1 - q^{-1}e^\beta}{1 - e^\beta} \right] \in \mathbb{C}[T].$$

Then using Corollary 3, we have an explicit formula of $\mathcal{W}(\varphi_w)(a)$.

4.5 Relation with Bump-Nakasuji's work

Now we explain the relation between this paper and Bump-Nakasuji [BN]. First of all, the notational conventions are slightly different. Especially in the published [BN] the natural base and intertwiner are differently parametrized. The natural basis ϕ_w in [BN] is our $\varphi_{w^{-1}}$. The intertwiner M_w in [BN] is our $\mathcal{A}_{w^{-1}}$ so that if $\ell(w_1w_2) = \ell(w_1) + \ell(w_2)$, $M_{w_1w_2} = M_{w_1} \circ M_{w_2}$ while $\mathcal{A}_{w_1w_2} = \mathcal{A}_{w_2}\mathcal{A}_{w_1}$.

In the paper [BN], another basis $\{\psi_w\}_{w \in W}$ for the space $I(\tau)^B$ was defined and compared with the Casselman's basis. They defined $\psi_w := \sum_{v \geq w} \varphi_v$ and expand this as $\psi_w = \sum_{v \geq w} m(w, v)f_v$ and conversely $f_w = \sum_{v \geq w} \tilde{m}(w, v)\psi_v$. They observed that the transition coefficients $m(w, v)$ and $\tilde{m}(w, v)$ factor under certain condition. Let $S(w, v) := \{\alpha \in R^+ | w \leq s_\alpha v < v\}$ and $S'(w, v) := \{\alpha \in R^+ | w < s_\alpha w \leq v\}$. Then the statements of the conjectures are as follows.

Conjecture 1 ([BN] Conjecture 1.2) Assume that the root system R is simply-laced. Suppose $w \leq v$ and $|S(w, v)| = \ell(v) - \ell(w)$, then

$$m(w, v) = \prod_{\alpha \in S(w, v)} \frac{1 - q^{-1}z^\alpha}{1 - z^\alpha}.$$

Conjecture 2 ([BN] Conjecture 1.3) Assume that the root system R is simply-laced. Suppose $w \leq v$ and $|S'(w, v)| = \ell(v) - \ell(w)$, then

$$\tilde{m}(w, v) = (-1)^{\ell(v) - \ell(w)} \prod_{\alpha \in S'(w, v)} \frac{1 - q^{-1}z^\alpha}{1 - z^\alpha}.$$

Proposition 10 Conjecture 1.2 and Conjecture 1.3 in [BN] are equivalent.

Proof: We can show $m(w, v) = \sum_{w \leq z \leq v} p(z, v)$ and $\tilde{m}(w, v) = \sum_{w \leq z \leq v} (-1)^{\ell(v) - \ell(z)} \tilde{p}(w, z)$. Then it follows by the Theorem 1 that $\tilde{m}(w, v) = (-1)^{\ell(v) - \ell(w)} m(vw_0, ww_0)$. As $S'(w, v) = S(vw_0, ww_0)$ we get the desired conclusion. \square

Acknowledgements

We would like to thank Shin-ichi Kato for valuable comments on the first version of this paper. We also would like to thank the referees for careful reading and point out the significantly related paper [Che]. This work was supported in part by JSPS Research Fellowship for Young Scientists and by Grant-in-Aid for Scientific Research.

References

[BBL] B. Brubaker, D. Bump and A. Licata, Whittaker functions and Demazure operators, *J. of Number Th.* 146 (2015), 41–68.

[Bum] D. Bump, Hecke Algebras, May 11, 2010. <http://sporadic.stanford.edu/bump/math263/hecke.pdf>

[BN] D. Bump and M. Nakasuji. Casselman's basis of Iwahori vectors and the Bruhat order, *Canadian Journal of Mathematics*, Vol.63, 1238–1253, (2011).

- [Cas] W. Casselman, The unramified principal series of p -adic groups I. The spherical function, *Comp. Math.* 40 (1980), 387–406.
- [Che] I. Cherednik, A unification of Knizhnik-Zamolodchikov and Dunkl operators via affine Hecke algebras, *Invent. Math.* 106 (1991), 411–431.
- [Dem] M. Demazure, Donnees Radicielles, 85–155, SGA3 Exp. 21. Springer Lect. Notes in Math. 153, (1970).
- [Hum] J.E. Humphreys, Reflection Groups and Coxeter Groups, Cambridge studies in adv. math. 29 (1990).
- [Kat] S. Kato, R -matrix arising from affine Hecke algebras and its application to Macdonald's difference operators, *Comm. Math. Phys.* 165 (1994), 533–553.
- [Kum] S. Kumar, Kac-Moody Groups, their Flag Varieties and Representation Theory, Progress in Math. Birkhauser 2002.
- [Las] A. Lascoux. *Polynômes*. <http://phalanstere.univ-mlv.fr/~al/ARTICLES/CoursYGKM.pdf>
- [LSS] T. Lam, A. Schilling and M. Shimozono, K -theory Schubert calculus of the affine Grassmannian, *Comp. Math.* 146 (2010), 811–852.
- [LLT] A. Lascoux, B. Leclerc and J.Y. Thibon, Flag varieties and the Yang-Baxter Equation, *Letters in Math. Phys.*, 40 (1997), 75–90.
- [Lu1] G. Lusztig, Equivariant K -theory and representations of Hecke algebras, *Proc. Amer. Math. Soc.*, 94 (1985), 337–342.
- [Lu2] G. Lusztig, Affine Hecke algebras and their graded version, *J. Amer. Math. Soc.*, 2 (1989), 599–635.
- [NN] M. Nakasuji and H. Naruse, Casselman problem and duality for intertwiner, in preparation.
- [Nar] H. Naruse, Schubert calculus and hook formula, in preparation.
- [Re1] M. Reeder, On certain Iwahori invariants in the unramified principal series, *Pacific J. of Math.* 153 (1992), 313–342.
- [Re2] M. Reeder, p -adic Whittaker functions and vector bundles on flag manifolds, *Comp. Math.* 85 (1993), 9–36.