



HAL
open science

Decomposition of the product of a monomial and a Demazure atom

Anna Ying Pun

► **To cite this version:**

Anna Ying Pun. Decomposition of the product of a monomial and a Demazure atom. 28-th International Conference on Formal Power Series and Algebraic Combinatorics, Simon Fraser University, Jul 2016, Vancouver, Canada. 10.46298/dmtcs.6323 . hal-02168296

HAL Id: hal-02168296

<https://hal.science/hal-02168296>

Submitted on 28 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Decomposition of the product of a monomial and a Demazure atom

Anna Ying Pun[†]

Department of Mathematics, University of Pennsylvania, Philadelphia, 19103, USA

Abstract. We prove that the product of a monomial and a Demazure atom is a positive sum of Demazure atoms combinatorially. This result proves one particular case in a conjecture which provides an approach to a combinatorial proof of Schubert positivity property.

Résumé. Nous montrons combinatoirement que le produit d'un monôme et d'un atome de Demazure est une somme positive d'atomes de Demazure. Ce résultat montre un cas particulier d'une conjecture qui fournit une approche combinatoire de la propriété de positivité de Schubert.

Keywords. generalized Demazure atoms, key polynomials, Schubert positivity, nonsymmetric Macdonald polynomials, skyline filings

1 Introduction


Let \mathbb{N} (or \mathbb{Z}^+) be the set of all positive integers and $\mathbb{Z}_{\geq 0}$ be the set of non-negative integers. Also we denote by S_k the set of all permutations of the set $\{1, \dots, k\}$. For $n \in \mathbb{Z}_{\geq 0}$ and $k \in \mathbb{N}$, we say $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \in (\mathbb{Z}_{\geq 0})^k$ is a weak composition n (denoted as $\alpha \vDash n$) with k parts if $\sum_{i=1}^k \alpha_i = n$ and write $l(\alpha) = k$ to denote the length (the number of parts) of α . Furthermore, if $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_k \geq 0$, we call α a partition of n with k parts and write $\alpha \vdash n$ (usually we denote $l(\alpha) = \max\{i \mid \alpha_i > 0\}$ for α being a partition).

We denote $\bar{\alpha}$ as the *reverse* of α , that is, $\bar{\alpha} = (\alpha_k, \dots, \alpha_1)$. Define ω_α as the permutation of minimal length such that $\omega_\alpha(\alpha) := (\alpha_{\omega_\alpha(1)}, \alpha_{\omega_\alpha(2)}, \dots, \alpha_{\omega_\alpha(k)})$ is a partition. Given two weak compositions α and β , we write $\beta \geq \alpha$ if and only if $\omega_\beta \leq \omega_\alpha$ in the strong Bruhat order.


Let α be a weak composition. The *augmented diagram* of shape α is the figure with $|\alpha| + l(\alpha)$ cells (or boxes) where column i has $\alpha_i + 1$ cells. The bottom row is called the *basement* of the augmented diagram.

For example, if $\alpha = (1, 0, 1, 0, 0, 4, 0, 6, 5)$, then the augmented diagram of α is

[†]Email: punying@sas.upenn.edu



Also we impose an order, called the *reading order*, on the cells of the diagram which reads each row from left to right, starting from the top row and ends in the bottom row. So the order of the above diagram is:



where the number in each cell represents the order of that cell in reading order.

A *filling* of an augmented diagram is an assignment of a positive integer to each cell in the diagram.

Macdonald [Mac96] defined a family of non-symmetric polynomials, called non-symmetric Macdonald polynomials,

$$\{E_\gamma(x_1, \dots, x_n; q, t) \mid \gamma \text{ is a weak composition with } n \text{ parts, } n \in \mathbb{N}\}$$

which shares many properties with the family of symmetric Macdonald polynomials [Mac95]

$$\{P_\lambda(x_1, \dots, x_n; q, t) \mid \lambda \text{ is a partition with } n \text{ parts, } n \in \mathbb{N}\}.$$

Haglund, Haiman and Loehr [HHL08] obtained a combinatorial formula for $E_\gamma(x_1, \dots, x_n; q, t)$ using fillings of augmented diagram of shape γ , called *skyline fillings*, satisfying certain constraints.

Marshall [Mar99] studied the family of non-symmetric Macdonald polynomials using another notation $\hat{E}_\gamma(x_1, \dots, x_n; q, t) := E_{\bar{\gamma}}(x_n, \dots, x_1; \frac{1}{q}, \frac{1}{t})$. In particular, by setting $q = t = 0$ in \hat{E}_γ , one can obtain Demazure atoms (first studied by Lascoux and Schützenberger [LS90]) $\mathcal{A}_\gamma = \hat{E}_\gamma(x_1, \dots, x_n; 0, 0) = E_{\bar{\gamma}}(x_n, \dots, x_1; \infty, \infty)$. Similarly, one can obtain Demazure characters (key polynomials) by setting $q = t = 0$ in E_γ , i.e., $\kappa_\gamma = E_\gamma(x_1, \dots, x_n; 0, 0) = \hat{E}_\gamma(x_n, \dots, x_1; \infty, \infty)$. The set of all Demazure atoms forms a basis for the polynomial ring, as does the set of all key polynomials.

Haglund, Luoto, Mason, Remmel and van Willigenburg [HLMvW11], [HMR13] further studied the combinatorial formulas for Demazure atoms and Demazure characters given by the skyline fillings and obtained results which generalized those for Schurs functions like the Pieri Rule, the Robinson-Schensted-Knuth (RSK) algorithm, and the Littlewood-Richardson (LR) rule.

From now on we only consider fillings whose entries in each column are weakly decreasing from bottom (to top) since those are the fillings related to Demazure atoms and Demazure characters [HHL08].

For any two columns (including the basement cells) i and j with $i < j$, we pick three cells X, Y and Z , where cell X is immediately above cell Y in the ‘taller’ column k , where


$$k = \begin{cases} i & \text{if } \alpha_i \geq \alpha_j, \\ j & \text{if } \alpha_i < \alpha_j, \end{cases}$$

and cell Z from the ‘shorter’ column to form a *triple* (X, Y, Z) in the following way:

$$\begin{cases} \text{Type A triple: cell } Z \text{ is in the same row as cell } X & \text{if } \alpha_i \geq \alpha_j \\ \text{Type B triple: cell } Z \text{ is in the same row as cell } Y & \text{if } \alpha_i < \alpha_j \end{cases}.$$

We say (X, Y, Z) forms a *coinversion triple* if the filling F of the diagram assigns each cell in the triple a positive integer, say $F(X), F(Y), F(Z)$ respectively, in such a way that $F(X) \leq F(Z) \leq F(Y)$. Otherwise we call (X, Y, Z) an *inversion triple*.

Here are some examples of triples:



The first two are type A triples and the last two are type B triples while the second and the third ones are coinversion triples and the first and the last one are inversion triples.

A *semi-standard augmented filling* (SSAF) of an augmented diagram with shape being a weak composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ is a filling satisfying:

1. the basement entries form a permutation σ (in one line notation) of $\{1, \dots, k\}$, i.e., $\sigma \in S_k$;
2. every (Type A or B) triple is an inversion triple.

We denote $SSAF(\sigma, \alpha)$ the set of all SSAF of an augmented diagram of shape $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ with basement entries (from left to right) being $\sigma \in S_k$ (i.e., basement of column i has entry $\sigma(i)$).

The following combinatorial formulas can be found in [HHL08, LS90, Mas09].

1. Let \mathcal{A}_α be the Demazure atom of shape α . Then

$$\mathcal{A}_\alpha = \sum_{F \in SSAF(\epsilon_k, \alpha)} x^F,$$

where $\epsilon_k \in S_k$ is the identity. In particular, if α is a partition, then $\mathcal{A}_\alpha = x^\alpha = \prod_{i=1}^k x_i^{\alpha_i}$ is a monomial.

2. Let $\kappa_{\bar{\alpha}}$ be the key polynomial (Demazure character) of shape α . Then

$$\kappa_{\bar{\alpha}} = \sum_{F \in SSAF(\bar{\epsilon}_k, \alpha)} x^F,$$

where $\bar{\epsilon}_k(i) = k + 1 - i$.

Note that if $\bar{\alpha}$ is a partition, $\kappa_{\bar{\alpha}} = \mathcal{A}_{\bar{\alpha}}$ which is a monomial by 1. The following are some properties among keys and atoms and their products:

3. A key polynomial is a positive sum of Demazure atoms. In fact,

$$\kappa_{\bar{\alpha}} = \sum_{\beta \geq \alpha} \mathcal{A}_{\beta}.$$

4. A key polynomial with a partition shape λ , with $l(\lambda) = k$, is the Schur polynomial s_{λ} , i.e., $\kappa_{\bar{\lambda}} = s_{\lambda}(x_1, \dots, x_k)$.

Let λ, μ be partitions and α, β be weak compositions. Denote $+_{\mathcal{A}}$ and $+_{\kappa}$ as being able to be decomposed into positive sum of atoms and keys respectively. Note that by 3, $+_{\kappa}$ implies $+_{\mathcal{A}}$. Otherwise, we put \times in the cell. For example, a partition (μ) -shaped atom times a key of any shape (α) is key positive and hence we put $+_{\kappa}$ in the corresponding box.

		Atoms		Keys	
		shape	λ	α	λ
Atoms	μ	$+_{\mathcal{A}}$	$+_{\mathcal{A}} \textcircled{1}$	$+_{\mathcal{A}}$	$+_{\kappa} \textcircled{2}$
	β	\times	\times	$+_{\mathcal{A}}$	\times
Keys	μ	\times	\times	$+_{\kappa}$	$+_{\kappa}$
	β	\times	\times	\times	open $\textcircled{3}$

Tab. 1: Decomposition of products of atoms and keys into atoms

The positive results in the table can be found in [HLMvW11], except for the cells marked $\textcircled{1}$, $\textcircled{2}$ and $\textcircled{3}$.

This paper proves $\textcircled{1}$ combinatorially in Section 4 which was open:

Theorem 1 *The product $\mathcal{A}_{\mu} \cdot \mathcal{A}_{\alpha}$ is atom-positive for any partition μ and weak composition α .*

The coefficients in the decomposition into atoms are actually counting the number of ways to insert words arising from SSAF of shape α into a SSAF of shape μ and we will discuss properties of words and how to record different ways of insertion in Section 2 and 3. Also note that the product in the theorem is not key positive. A simple counter example would be just putting μ as the empty partition, that is, with all entries 0.

$\textcircled{2}$ is proved in [Jos03] (the proof involves crystals but does not involve SSAF). Both results $\textcircled{1}$ and $\textcircled{2}$ imply $+_{\mathcal{A}}$ for the $\mathcal{A}_{\mu} \cdot \kappa_{\bar{\alpha}}$ cell. Also we conjecture that we can apply the bijection in the proof in Theorem 6.1 in [HLMvW11] on Theorem 13 in Section 4 to give a tableau-combinatorial proof to $\textcircled{2}$.

As for the product of two keys of arbitrary shapes, that is, the cell marked $\textcircled{3}$, there are examples showing that the product of two key polynomials of arbitrary shapes is not a positive sum of keys. So it remains to check if it is a positive sum of atoms, which is still open. Hence $\textcircled{3}$ gives the following conjecture which we have verified for $l(\alpha), l(\beta) \leq 3$:

Conjecture 1 Let α, β be weak compositions. Then the product of the key polynomials of shape $\bar{\alpha}$ and $\bar{\beta}$ can be written as a positive sum of atoms, i.e.,

$$\kappa_\alpha \cdot \kappa_\beta = \sum_{\gamma \models |\alpha| + |\beta|} c_{\alpha\beta}^\gamma \mathcal{A}_\gamma$$

for some non-negative integers $c_{\alpha\beta}^\gamma$.

It is a classical result in Algebraic Geometry that the product of two Schubert polynomials can be written as a positive sum of Schubert polynomials. A representation theoretic proof is also given recently by using Kraskiewicz-Pragacz modules [Wat14]. However a combinatorial proof of the positivity property of Schubert polynomials has long been open. Since every Schubert polynomial is a positive sum of key polynomials [LS89], the product of two Schubert polynomials is a positive sum of product of two key polynomials and hence by Conjecture 1 is a positive sum of atoms. This provides a possible approach to a combinatorial proof of the positivity property of Schubert polynomials by trying to recombine the atoms into keys and hence into Schubert polynomials.

Since words and insertion are the crucial tools used to prove Theorem 13, we will discuss an algorithm on words in Section 2, then we will relate it to SSAF and recording tableaux in Section 3. Finally we will use the results we obtained in Section 2 and 3 to prove the main theorem that the product of $\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha$ is atom-positive for partition λ and weak composition α in Section 4.

2 Convert a column word to a row word

Definition 1 A *word* is a sequence of positive integers.

Definition 2 Let $a, b, c \in \mathbb{N}$ and u, v be some fixed (can be empty) words. Define twisted Knuth relation \curvearrowright^* by:

1. $ubacv \curvearrowright^* ucavb$ if $c \leq b < a$
2. $uacbv \curvearrowright^* ucabv$ if $c < b \leq a$.

Then we say two words w and w' are twisted Knuth equivalent if w can be transformed to w' by repeated use of 1. and 2. and we write $w \curvearrowright^* w'$.

Definition 3 A word w is a **column word** if it can be broken down into k weakly decreasing subsequences of weakly decreasing lengths

$$w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$$

where $c_1 \geq c_2 \geq \dots \geq c_k > 0$, $c_1, \dots, c_k \in \mathbb{N}$ such that

$$\begin{cases} a_{ij} \geq a_{i,j+1} & \text{for } 1 \leq j < c_i, 1 \leq i \leq k \\ a_{i+1,c_{i+1}-j} > a_{i,c_i-j} & \text{for } 0 \leq j < c_i, 1 \leq i < k \end{cases}$$

Definition 4 A word w is a **row word** if it can be broken down into k strictly increasing subsequences of weakly decreasing lengths

$$w = a_{11} \dots a_{1r_1} | a_{21} \dots a_{2r_2} | \dots | a_{k1} \dots a_{kr_k}$$

where $r_1 \geq r_2 \geq \dots \geq r_k > 0$, $r_1, \dots, r_k \in \mathbb{N}$ such that

$$\begin{cases} a_{ij} < a_{i,j+1} & \text{for } 1 \leq j < r_i, 1 \leq i \leq k \\ a_{i+1,r_{i+1}-j} \leq a_{i,r_i-j} & \text{for } 0 \leq j < r_i, 1 \leq i < k \end{cases}.$$

Given a SSAF, one can get its column word by using the algorithm described in [HLMvW11], while one can get its row word by reading the entries of each row in ascending order, starting from the bottom row to the top row. We call a word a column (row resp.) word because when we insert each subsequence of the word using the insertion in [Mas08], a new column (row resp.) will be created.

Example 1 886531|97643|9764|5|6 is a column word whose corresponding SSAF is:

							6	
							6	4
					3		7	5
					4		7	6
					5		8	9
					6		8	9
1		3						
1	2	3	4	5	6	7	8	9

and the corresponding row word is 13689|589|467|357|46|6.

Lemma 2 Let $u := a_1 a_2 \dots a_n | b$ and $v := a_1 a_2 \dots a_n | b | c$ be two column words, where $c > b > a_n$, $n \in \mathbb{N}$. Then

1. $u \smile^* b' a'_1 \dots a'_n$ where $b' < a'_1$, $a'_1 \geq a'_2 \geq \dots \geq a'_n$, and $b' := a_t$ where $t = \min\{j \mid b > a_j\}$,
2. $v \smile^* b' c' a''_1 a''_2 \dots a''_n$ where b' is defined as in 1., $b' < c' < a''_1$ and $a''_1 \geq a''_2 \geq \dots \geq a''_n$.

Lemma 3 Let $k \geq 2$ be an integer and i_1, \dots, i_k be nonnegative integers. Let

$$w_0 = a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} | a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} | \dots | a_{k1}^{(i_k)} \dots a_{kc_k}^{(i_k)}$$

be a column word. Then there exist $a_{rs}^{(i_r+1)}$ where $1 \leq s \leq c_r$, $1 \leq r < k$ and $a_{k1}^{(i_k+m)}$ where $1 \leq m < k$ satisfying

$$\begin{cases} a_{k-j-1, c_{k-j-1}}^{(i_{k-j-1})} < a_{k1}^{(i_k+j)} < a_{k-j,1}^{(i_{k-j+1})} & \text{for } 1 \leq j < k-1 \quad (k > 3) \\ a_{k1}^{(i_k+k-1)} < a_{11}^{(i_1+1)} \\ a_{1,1}^{(i_1+1)} \dots a_{1,c_1}^{(i_1+1)} | \dots | a_{k-1,1}^{(i_{k-1}+1)} \dots a_{k-1,c_{k-1}}^{(i_{k-1}+1)} | a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)} & \text{is a column word} \end{cases}$$

such that $w_0 \smile^* w_1 \smile^* \dots \smile^* w_{k-1}$, where

$$\begin{cases} w_j := a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{k-j-1,1}^{(i_{k-j-1})} \dots a_{k-j-1, c_{k-j-1}}^{(i_{k-j-1})} a_{k1}^{(i_k+j)} a_{k-j,1}^{(i_{k-j+1})} \dots a_{k-j, c_{k-j}}^{(i_{k-j+1})} \dots \\ \quad \dots a_{k-1,1}^{(i_{k-1}+1)} \dots a_{k-1, c_{k-1}}^{(i_{k-1}+1)} a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)} & \text{for } 1 \leq j < k-1, \\ w_{k-1} := a_{k1}^{i_k+k-1} a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} \dots a_{k-1,1}^{(i_{k-1}+1)} \dots a_{k-1, c_{k-1}}^{(i_{k-1}+1)} a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)}. \end{cases}$$

The proof is basically induction on k .

Lemma 3 shows how to create the first entry of the corresponding row word, leaving the remaining part as a column word. We illustrate the Lemma by an example with $a_{k1}^{(i_k+j)}$ circled for all $0 \leq j \leq k-1$.

Example 2 Let w_0 be the word in Example 1, hence $k = 5$, $c_1 = 6$, $c_2 = 5$, $c_3 = 4$, $c_4 = c_5 = 1$. Set $i_j = 0$ for all j . Then we have

$$\left\{ \begin{array}{l} w_0 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | a_{41}^{(0)} | \textcircled{a_{51}^{(0)}} = 886531|97643|9764|5|\textcircled{6} \\ w_1 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | \textcircled{a_{51}^{(1)}} | a_{41}^{(1)} = 886531|97643|9764|\textcircled{5}|6 \\ w_2 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | \textcircled{a_{51}^{(2)}} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531|97643|\textcircled{4}|9765|6 \\ w_3 = a_{11}^{(0)} \dots a_{16}^{(0)} | \textcircled{a_{51}^{(3)}} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531|\textcircled{3}|97644|9765|6 \\ w_4 = \textcircled{a_{51}^{(4)}} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = \textcircled{1}|886533|97644|9765|6 \end{array} \right. .$$

Lemma 4 Let $w = a_{11} \dots a_{1c_1} \dots a_{k1} \dots a_{kc_k}$ be a column word. Then there exists a sequence $\{b_{ij}\}_{1 \leq j \leq c_i, 1 \leq i \leq k}$ such that $w \rightsquigarrow^* b_{k1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$, where $b_{11} > b_{21} > \dots > b_{k1}$ and $b_{11} \dots b_{1c_1} | b_{22} \dots b_{2c_2} | \dots | b_{k2} \dots b_{kc_k}$ is a column word (with lengths $c_1 > c_2 - 1 \geq \dots \geq c_k - 1$).

The proof is applying Lemma 3 inductively.

Using the notation in Lemma 4, since $w \rightsquigarrow^* b_{k1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$, where $b_{11} > b_{21} > \dots > b_{k1}$, the SSAF with basement being the identity (i.e., $1 \ 2 \ \dots \ n$, for some positive integer n) representing the word w is the same as that representing $b_{k1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$. Denote $F(w)$ as the SSAF created.

Since $b_{k1} b_{k-1,1} \dots b_{11}$ is strictly increasing, by Lemma 15 in [HMR13], they create new cells in ascending reading order (i.e., one after another) and hence is exactly the first entire row of $F(w)$ as the entries are fixed when inserting $b_{k1} b_{k-1,1} \dots b_{11}$ into an empty atom, and there are k columns (as w has k subsequences) and so the first row has length k and hence the row reading word has exactly $b_{k1}, b_{k-1,1}, \dots, b_{11}$ as the first subsequence. That means we can apply Lemma 4 to find the first subsequence of the row reading word of $F(w)$.

Since $b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$ is a column word, we can apply Lemma 4 again and get the second subsequence of the row reading word of $F(w)$, and we can apply Lemma 4 repeatedly on the remaining b_{ij} 's until we get all the subsequences of the row reading word of $F(w)$. As a result, we can convert w into the row reading word of $F(w)$ by applying Lemma 4 repeatedly as described.

3 Convert a Column Recording Tableau to a Row Recording Tableau

This section gives an interpretation of the twisted Knuth equivalence using recording tableaux. We use the insertion in [Mas08] and the generalized Littlewood-Richardson rule in [HLMvW11]. We also use

the notation $U \leftarrow W$ for a SSAF U and a biword $W = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ y_1 & y_2 & \dots & y_n \end{pmatrix}$ for some integer n to denote the pair (U', L) where U' is the SSAF obtained by $U \leftarrow y_1 y_2 \dots y_n$ while L is the recording tableau, i.e., by putting x_i into the cell created when y_i is being inserted. In particular, if $y_1 y_2 \dots y_n$ is a column (row resp.) word, then we call L as a column (row resp.) recording tableau. By abuse of notation, we sometimes refer $U \leftarrow W$ to either U' or L (depending on the context). Note that if we change the basement ϵ_n into the large basement in [HLMvW11], a column recording tableau is the same as an LRS in [HLMvW11].

Lemma 5 *Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & 1 \\ a & b & c \end{pmatrix}$, $a \geq b, c > b$ (i.e., $ab|c$ is a column word.). Let L be the recording tableau of $U \leftarrow W$. Let V be the SSAF representing the word abc and $a'b'c'$ be the row reading word of V (so $a' < b'$ and $c' \leq b'$ and $abc \rightsquigarrow^* a'b'c'$). Consider the biword $\widetilde{W} = \begin{pmatrix} 1 & 1 & 2 \\ a' & b' & c' \end{pmatrix}$ and let \widetilde{L} be the recording tableau of $U \leftarrow \widetilde{W}$. Then L determines \widetilde{L} .*

Lemma 6 *Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \dots & 2 & 1 \\ a_1 & a_2 & \dots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \dots \geq a_k, b > a_k$ (i.e., $a_1 a_2 \dots a_k | b$ is a column word). Let L be the recording tableau of $U \leftarrow W$. If $b > a_1$, then L has reading word $22 \dots 21$, i.e., b creates the last cell in reading order and the insertion of $a_2 \dots a_k$ is independent of the insertion of b .*

Lemma 6 follows by applying Lemma 5 inductively.

Lemma 7 *Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \dots & 2 & 1 \\ a_1 & a_2 & \dots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \dots \geq a_k, b > a_k$ (i.e., $a_1 a_2 \dots a_k | b$ is a column word). Let L be the recording tableau of $U \leftarrow W$. If $i = \min\{j \mid b > a_j\}$, then the cell created by inserting a_m for $m > i$ is not affected by the insertion of b .*

Proof: Since $U \leftarrow a_1 a_2 \dots a_k b = (U \leftarrow a_1 a_2 \dots a_{i-1}) \leftarrow a_i \dots a_k b$. By applying Lemma 6 with the U in the lemma being $U \leftarrow a_1 \dots a_{i-1}$ and result follows. □

Lemma 8 *Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \dots & 2 & 1 \\ a_1 & a_2 & \dots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \dots \geq a_k, b > a_k$ (i.e., $a_1 a_2 \dots a_k | b$ is a column word). Let L be the recording tableau of $U \leftarrow W$. Suppose $b \leq a_{k-1}$. Let \widetilde{L} be the recording tableau of $U \leftarrow \widetilde{W}$ where $\widetilde{W} = \begin{pmatrix} 1 & 1 & 2 & \dots & 2 & 2 \\ a_k & a_1 & a_2 & \dots & a_{k-1} & b \end{pmatrix}$, then L determines \widetilde{L} .*

The above result follows by applying Lemma 5 inductively.

Lemma 9 *Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \dots & 2 & 1 \\ a_1 & a_2 & \dots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \dots \geq a_k, b > a_k$ (i.e., $a_1 a_2 \dots a_k | b$*

is a column word). Let L be the recording tableau of $U \leftarrow W$. Let \tilde{L} be the recording tableau of $U \leftarrow \tilde{W}$ where $\tilde{W} = \begin{pmatrix} 1 & 1 & 2 & \cdots & 2 \\ b' & a'_1 & a'_2 & \cdots & a'_k \end{pmatrix}$, such that $b' = a_i$, where $i = \min\{j \mid b > a_j\}$ and $b' < a'_1, a'_1 \geq a'_2 \geq \cdots \geq a'_k$ (one can verify that $b'a'_1 \dots a'_k$ is indeed the row reading word of the SSAF representing the word $a_1 a_2 \dots a_k b$ by 1. in Lemma 2), then L determines \tilde{L} .

Proof: If $i = k$, then we are done by Lemma 8.

If $i = 1$, then by 1. in Lemma 2, we have $b'a'_1 \dots a'_k = a_1 b a_2 \dots a_k$ and hence by Lemma 6, b would create the last cell in reading order in \tilde{L} when being inserted and hence we know that which two cells are created by a_1 and b in \tilde{L} and hence we know how to label the entries by marking those two cells as 1 and the rest as 2.

Suppose $i < k$ then by 1. in Lemma 2, we have $b'a'_1 \dots a'_k = a_i a_1 \dots a_{i-1} b a_{i+1} \dots a_k$. Now by Lemma 7, we know all a_j , where $j > i$, create the same cells in L and \tilde{L} and since $a_1 \dots a_{i-1} a_i b \curvearrowright^* a_i a_1 \dots a_{i-1} b$, we just need to first apply Lemma 8 on $U \leftarrow a_1 \dots a_i b$ as the insertion recorded by L in the lemma in order to find the first $(i+1)$ cells created in \tilde{L} by the insertion $U \leftarrow a_i a_1 \dots a_{i-1} b$ and then label the rest of cells created by $(U \leftarrow a_i a_1 \dots a_{i-1} b) \leftarrow a_{i+1} \dots a_k$ as 2 and get the entries of \tilde{L} and result follows. \square

Lemma 9 gives a recording tableau interpretation of 1. in Lemma 2 using L and \tilde{L} .

Lemma 10 Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Let $w_0 = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \cdots | a_{k1} \dots a_{kc_k}$ with $k \leq n$ be a column word and using the notation in Lemma 3 while we assume $i_j = 0$ for $1 \leq j \leq c_r, 1 \leq r \leq k$, the recording tableau L of $U \leftarrow W_0$ determines the recording tableau \tilde{L}_m of $U \leftarrow W_m$ for $0 \leq m \leq k-1$, where $L := \tilde{L}_0$ and W_m is a biword with the lower word being w_m and the upper word has entry $k+1-t$ if the lower word entry just below it is $a_{i_j}^{(s)}$ for $s = i_t, i_t + 1$, and $1_{(1)}$ if $m > 0$ and the entry is $a_{k1}^{(i_k+m)}$.

Lemma 10 follows by inductively using Lemma 9.

Lemma 11 Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Let $w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \cdots | a_{k1} \dots a_{kc_k}$ with $k \leq n$ be a column word and using the notation in Lemma 4, the recording tableau L of $U \leftarrow W$ determines the recording tableau \tilde{L} of $U \leftarrow \tilde{W}$, where

$$\begin{cases} W = \begin{pmatrix} k & k & \cdots & k & \cdots & 1 & 1 & \cdots & 1 \\ a_{11} & a_{12} & \cdots & a_{1c_1} & \cdots & a_{k1} & a_{k2} & \cdots & a_{kc_k} \end{pmatrix} \\ \tilde{W} = \begin{pmatrix} 1_{(1)} & 2_{(1)} & \cdots & k_{(1)} & k & k & \cdots & k & \cdots & 1 & 1 & \cdots & 1 \\ b_{k1} & b_{k-1,1} & \cdots & b_{11} & b_{12} & b_{13} & \cdots & b_{1c_1} & \cdots & b_{k2} & b_{k3} & \cdots & b_{kc_k} \end{pmatrix} \end{cases}.$$

We apply Lemma 10 (which is like the tableaux version of Lemma 3) to get the result above.

Lemma 12 Let U be a SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Let $w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \cdots | a_{k1} \dots a_{kc_k}$ with $k \leq n$ be a column word. Let V be the SSAF representing w (i.e., inserting w into an empty atom). Let $\tilde{w} = x_{11} x_{12} \dots x_{r_1} \cdots x_{c_1,1} \dots x_{c_1,r_{c_1}}$ (as V has c_1 rows and k columns and so r_1 is k), where $k = r_1 \geq r_2 \geq \cdots \geq r_{c_1} > 0$ are the row lengths of V from bottom to top, be the row reading word of V . Then the recording tableau L of $U \leftarrow W$ determines the recording tableau R of $U \leftarrow \tilde{W}$, where

$$\left\{ \begin{aligned} W &= \begin{pmatrix} k & k & \dots & k & \dots & 1 & 1 & \dots & 1 \\ a_{11} & a_{12} & \dots & a_{1c_1} & \dots & a_{k1} & a_{k2} & \dots & a_{kc_k} \end{pmatrix} \\ \widetilde{W} &= \begin{pmatrix} 1 & 1 & \dots & 1 & \dots & c_1 & c_1 & \dots & c_1 \\ x_{11} & x_{12} & \dots & x_{1r_1} & \dots & x_{c_1,1} & x_{c_1,2} & \dots & x_{c_1,r_{c_1}} \end{pmatrix} \end{aligned} \right.$$

Proof:

By Lemma 11 and the corresponding word in Lemma 4, given L , we know how to enter all the 1's in R , which are those cells marked $1_{(1)}, 2_{(1)}, \dots, k_{(1)}$ after applying Lemma 11 as the $b_{k1}b_{k-1,1} \dots b_{11} = x_{11}x_{12} \dots x_{1r_1}$ (by the argument after Lemma 4).

We remove the cells from \widetilde{L} in Lemma 11 to create a new L to apply Lemma 11 on, we can get the second row entries (as described in the paragraphs after Lemma 4), and hence we know how to put all the 2's into R . By the same argument, we can fill in all entries in R and hence L determines R . □

4 Decomposition of the product of a monomial and an atom into positive sum of atoms

We prove Theorem 1 mentioned in the Introduction. We rephrase the statement more precisely as follows.

Theorem 13 *Let λ be a partition and α be a weak composition. Let \mathcal{A}_λ and \mathcal{A}_α be atoms of shape λ and α respectively. Then*

$$\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha = \sum_{\beta = |\lambda| + |\alpha|, \lambda \subseteq \beta} c_{\lambda\alpha}^\beta \mathcal{A}_\beta$$

where $c_{\lambda\alpha}^\beta$ is the number of distinct LRS of shape β/λ created by column words whose corresponding SSAF has shape α .

Proof: Since $\mathcal{A}_\lambda = x^\lambda$ (there is exactly one SSAF with shape λ , denoted by U_λ) and $\mathcal{A}_\alpha = \sum_{F \in \text{SSAF}(\epsilon_{l(\alpha)}, \alpha)} x^F$,

we have

$$\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha = \sum_{F \in \text{SSAF}(\epsilon_{l(\alpha)}, \alpha)} x^\lambda x^F$$

To prove the theorem, we only need to check that given a LRS L of shape β/λ created by column words whose corresponding SSAF has shape α , if there is some column word

$$w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$$

and biword

$$W = \begin{pmatrix} k & k & \dots & k & \dots & 1 & 1 & \dots & 1 \\ a_{11} & a_{12} & \dots & a_{1c_1} & \dots & a_{k1} & a_{k2} & \dots & a_{kc_k} \end{pmatrix}$$

such that $U_\lambda \leftarrow W$ creates the same L , then the SSAF corresponding to w also has shape α .

First consider the last cell in reading order among all those containing entry k in L , that means it is the very first entry inserted. Since a_{11} must be inserted in a cell immediately above the cell (including those in basement) containing a_{11} , the position of that cell fixes the value of a_{11} .

Now consider all the cells with k and also the last cell in reading order among all those containing the entry $k - 1$, then these cells are created by $U_\lambda \leftarrow a_{11} \dots a_{1c_1} a_{21}$. By Lemma 12, we know the corresponding row recording tableau and hence we know which two cells are the first two entries being inserted using the corresponding row word. Note that the first row consists of distinct entries and is inserted in ascending order using the row word, then by Lemma 15 in [HMR13], we know the cells are created in ascending reading order (one after another in reading order) and so they must be the cells immediately above those in U_λ , and hence the value inside each of those cells in the SSAF created by inserting the row word into U_λ is exactly the value inside the cell just below it. Hence we know what the first two row entries of the corresponding SSAF of $a_{11} a_{12} \dots a_{1c_1} a_{21}$ are. Since we already know the first row entry, which is the lowest entry of the column corresponding to $a_{11} \dots a_{1c_1}$ is, we now know what the lowest entry of the second column (corresponding to $a_{11} \dots a_{1c_1} a_{21}$ and hence the same for $a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2}$) is.

We can repeat the same process until we get all the last entries of the k columns and hence fix the shape of the SSAF corresponding to w . Since we read those entries just by considering L , this shows that L fixes the shape of the corresponding SSAF of w and result follows. \square

References

- [HHL08] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for nonsymmetric Macdonald polynomials. *Amer. J. Math.*, 130(2):359–383, 2008.
- [HLMvW11] J. Haglund, K. Luoto, S. Mason, and S. van Willigenburg. Refinements of the Littlewood-Richardson rule. *Trans. Amer. Math. Soc.*, 363(3):1665–1686, 2011.
- [HMR13] James Haglund, Sarah Mason, and Jeffrey Remmel. Properties of the nonsymmetric Robinson-Schensted-Knuth algorithm. *J. Algebraic Combin.*, 38(2):285–327, 2013.
- [Jos03] Anthony Joseph. A decomposition theorem for Demazure crystals. *Journal of Algebra*, 265(2):562 – 578, 2003.
- [LS89] A. Lasku and M.-P. Shyuttsenberzhe. Noncommutative Schubert polynomials. *Funktsional. Anal. i Prilozhen.*, 23(3):63–64, 1989.
- [LS90] Alain Lascoux and Marcel-Paul Schützenberger. Keys & standard bases. In *Invariant theory and tableaux (Minneapolis, MN, 1988)*, volume 19 of *IMA Vol. Math. Appl.*, pages 125–144. Springer, New York, 1990.
- [Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.

- [Mac96] I. G. Macdonald. Affine Hecke algebras and orthogonal polynomials. *Astérisque*, (237):Exp. No. 797, 4, 189–207, 1996. Séminaire Bourbaki, Vol. 1994/95.
- [Mar99] Dan Marshall. Symmetric and nonsymmetric Macdonald polynomials. *Ann. Comb.*, 3(2-4):385–415, 1999. On combinatorics and statistical mechanics.
- [Mas08] S. Mason. A decomposition of Schur functions and an analogue of the Robinson-Schensted-Knuth algorithm. *Sém. Lothar. Combin.*, 57:Art. B57e, 24, 2006/08.
- [Mas09] S. Mason. An explicit construction of type A Demazure atoms. *J. Algebraic Combin.*, 29(3):295–313, 2009.
- [Wat14] M. Watanabe. Tensor product of Kraskiewicz and Pragacz’s modules. *ArXiv e-prints*, October 2014.