



HAL
open science

On the dynamics of 3D electrified falling films

Jiao He, Rafael Granero-Belinchón

► **To cite this version:**

Jiao He, Rafael Granero-Belinchón. On the dynamics of 3D electrified falling films. 2019. hal-02168399v1

HAL Id: hal-02168399

<https://hal.science/hal-02168399v1>

Preprint submitted on 28 Jun 2019 (v1), last revised 10 Nov 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE DYNAMICS OF 3D ELECTRIFIED FALLING FILMS

JIAO HE AND RAFAEL GRANERO-BELINCHÓN

ABSTRACT. In this article, we consider a non-local variant of the Kuramoto-Sivashinsky equation in three dimensions (2D interface). Besides showing the global wellposedness of this equation we also obtain some qualitative properties of the solutions. In particular, we prove that the solutions become analytic in the spatial variable for positive time, the existence of a compact global attractor and an upper bound on the number of spatial oscillations of the solutions. We observe that such a bound is particularly interesting due to the chaotic behavior of the solutions.

1. INTRODUCTION

The present work is concerned with the full 3D dynamics of a thin fluid film falling along with a flat inclined plate. Besides gravitational effects, we consider the action of an electric field acting normal to the plate. In particular, for the case where the fluid lies on top of the plate (overlying films), the following equation was derived by Tomlin, Papageorgiou & Pavliotis [27]:

$$(1) \quad \eta_t + \eta\eta_x + (\beta - 1)\eta_{xx} - \eta_{yy} - \gamma\Lambda^3\eta + \Delta^2\eta = 0$$

where $\beta > 0$ is the Reynolds number, $0 \leq \gamma \leq 2$ measures the electric field strength and Λ is a non-local operator corresponding to the electric field effect given on the Fourier variables as

$$\widehat{\Lambda u} = |\xi|\hat{u}(\xi) = (\xi_1^2 + \xi_2^2)^{0.5}\hat{u}(\xi).$$

We observe that the term corresponding to the electric field, $-\gamma\Lambda^3(\eta)$, always has a destabilizing effect, while the term $(\beta - 1)\eta_{xx}$ can be stabilizing or destabilizing depending on the value of the Reynolds number. Namely, for subcritical Reynolds numbers $0 < \beta < 1$, $(\beta - 1)\eta_{xx}$ is a stabilizing term, while for supercritical Reynolds numbers $1 < \beta$, it has a destabilizing effect.

Since falling films have received much attention from many authors, a wide variety of results about their nonlinear stability can be found. In particular, the 2D case (1D interface) was first studied by González & Castellanos [10]. These authors identified a critical electric field strength for sub-critical Reynolds number flows above which instability was found. Later on, Tseluiko & Papageorgiou also considered the 2D case (1D interface). In particular, Tseluiko & Papageorgiou performed a numerical study of the 1D analog of (1) and found attractors for the dynamics for certain values of the physical parameters [30]. The same authors provided analytical bounds on the energy of the solutions and the dimension of the attractor [31] (see also [29] for the case of vertical film flow). Compared with the case of 1D interface, equation (1) generalizes previous works by taking transverse dynamics into consideration.

Equation (1) is similar to the well-known Kuramoto-Sivashinsky equation in one dimension,

$$(KS) \quad \eta_t + \eta\eta_x = -\eta_{xx} - \eta_{xxx}$$

which is a model appearing in several applications. For instance, LaQuey, Mahajan, Rutherford & Tang [18] obtained (KS) as a model of collisional trapped-ion mode in tokamak geometry (see also Cohen, Krommes, Tang & Rosenbluth [3]), Kuramoto & Tsuzuki [17] considered the possible instabilities of a two components reaction-diffusion system and also recovered (KS). Furthermore, Sivashinsky [24] (see also the companion paper by Michelson & Sivashinsky [20]) derived (KS) as a model of the evolution of a disturbed plane flame front. Later on, Sivashinsky & Michelson [25] linked (KS) to the evolution of a film of viscous liquid flowing down a vertical plane. Several equations sharing some similarities were obtained by Topper & Kawahara [28], Lee and Chen [19], Coward & Hall [6], Frenkel & Indireskumar [8] and by James & Wilczek [16] when considering falling fluid films, plasma turbulence and cellular suspensions.

Equation (KS) has rich dynamics. Indeed, applying the Fourier transformation to the linear part of (KS),

$$\partial_t \hat{\eta}(\xi) = (\xi^2 - \xi^4) \hat{\eta}(\xi),$$

it results in the stability of high frequencies ($|\xi| > 1$) and instability of low frequencies ($0 < |\xi| < 1$). Specifically, the term η_{xx} leads to instability at large scales; the dissipative term η_{xxxx} is responsible for damping at small scales. Then we see that for general initial data, the linear problem is unstable and leads to an exponential growth of certain frequencies. When the nonlinear term $\eta\eta_x$ is added, stabilization occurs as this term transfers energy from the long wavelengths to the short wavelengths and balances the exponential growth due to the linear parts. This interaction between the unstable linear parts and a nonlinearity who carries energy between frequencies makes the solution of (KS) to develop chaotic dynamics for certain values of the parameters.

This nonlinear stabilization of the Kuramoto-Sivashinsky equation with L -periodic boundary conditions,

$$\eta(x + L, t) = \eta(x, t), \text{ for all } x \text{ and } t,$$

was considered mathematically by Nicolaenko, Scheurer & Temam in [21] under the hypothesis that the initial data has odd symmetry: $\eta_0(x) = -\eta_0(-x)$. After that, Ilyashenko [14], Collet, Eckmann, Epstein & Stubbe [5] and Goodman [11] found new bounds for the L^2 -norm of the solution of the KS equation without oddness condition for the initial data. The fact that the solutions are uniformly bounded leads us to the question of the optimal bound for the radius of the absorbing set in L^2 for arbitrarily large periods L . In that regards, the known bounds are

$$\limsup_{t \rightarrow \infty} \left(\int_0^L u^2 dx \right)^{0.5} \leq O(L^p)$$

where the original $p = 5/2$ [21] was later improved to $p = 8/5$ [5] and finally to $p = 3/2$ [1]. The global bound has been upgraded recently by Giacomelli & Otto [9], where they proved the bound

$$\limsup_{t \rightarrow \infty} \left(\int_0^L u^2 dx \right)^{0.5} \leq o(L^{1.5}).$$

We observe that the conjectured bound is $O(L^{0.5})$.

The analyticity of solutions is of great interest not only for KS equation, but also for other nonlinear partial differential equations. For instance, we refer the reader to the seminar paper by Foias & Temam [7] where they show that solutions of the Navier-Stokes equations are analytic in time with values in a Gevrey class of functions (in space). This technique has been extended largely to other nonlinear parabolic equations and, in particular, Collet, Eckmann, Epstein & Stubbe [4] addressed the spatial analyticity of

solutions of one-dimensional Kuramoto-Sivashinsky equation. They showed that at large time the solutions are analytic in a strip around the real axis and also gave a rigorous lower bound for its width, i.e. the radius of analyticity is proportional to $L^{-16/25}$. Grujić [13] used a Gevrey class technique to obtain a neighborhood in the global attractor of the set of all stationary solutions in which the radius of analyticity is independent of L . This latter result shed some light on a conjecture in [4] that asks whether there is a $\alpha > 0$, independent of L , such that the solutions of the KS equation are analytic in space in the complex strip $\{x + is, s < \alpha\}$ for sufficiently large time. In higher dimensions, the literature on estimating the radius of analyticity for the Kuramoto-Sivashinsky-type equations is more scarce. For example, we refer to the works by Pinto [22, 23] where, among other properties, the author studied the time analyticity of a variant of the two-dimensional KS equation. More recently, Ioakim & Smyrlis [15] also studied the analyticity properties of solutions of Kuramoto-Sivashinsky type equations and some related systems.

The goal of the present work is to mathematically study the initial value problem for nonlocal two-dimensional Kuramoto-Sivashinsky-type equation with periodic boundary conditions and initial data with zero mean

$$\int_0^L \int_0^L \eta_0(x, y) dx dy = 0.$$

Of course, the zero average condition is propagated by the PDE. We organize this paper as follows. In section 2, we give some notations, definitions and classical results. In section 3, we show the global existence of solutions to initial value problem (1) and in section 4, we prove the existence of an absorbing set in L^2 and in higher Sobolev norms. In section 5, we prove that these solutions are analytic in a strip based on *a priori* estimates in a Gevrey class. Finally, in section 6, we establish a bound for the number of spatial oscillations which are a manifestation of the spatial chaos that this PDE evidences.

1.1. Notation. We will use C to denote a universal constant that can change from one line to another. We will make frequent use of the usual homogeneous Sobolev spaces \dot{H}^s :

$$\dot{H}^s(\mathbb{T}^2) = \{\eta \in L^2(\mathbb{T}^2) : \sum_{\xi \in \mathbb{Z}^2} |\xi|^{2s} |\hat{\eta}(\xi)|^2 < \infty\}.$$

2. RESCALING OF THE EQUATION

We assume that η is L -periodic,

$$\eta(x + L, y) = \eta(x, y), \quad \eta(x, y + L) = \eta(x, y),$$

and define $\mathbb{T}^2 = [0, 2\pi]^2$ and $\lambda = \frac{2\pi}{L}$. We rescale our variables according to

$$\tilde{x} = \lambda x, \quad \tilde{y} = \lambda y, \quad \tilde{\eta} = \lambda^{-1} \eta, \quad \tilde{t} = \lambda^2 t,$$

which gives

$$\lambda^3 \tilde{\eta}_t + \lambda^3 \tilde{\eta} \tilde{\eta}_x + (\beta - 1) \lambda^3 \tilde{\eta}_{xx} - \lambda^3 \tilde{\eta}_{yy} - \gamma \lambda^4 \Lambda^3(\tilde{\eta}) + \lambda^5 \Delta^2 \tilde{\eta} = 0.$$

Then we obtain

$$(2) \quad \tilde{\eta}_t + \tilde{\eta} \tilde{\eta}_x + (\beta - 1) \tilde{\eta}_{xx} - \tilde{\eta}_{yy} - \gamma \lambda \Lambda^3(\tilde{\eta}) + \lambda^2 \Delta^2 \tilde{\eta} = 0.$$

Denoting $\delta = \gamma \lambda$ and $\epsilon = \lambda^2$, we can equivalently consider the following initial-value problem

$$(3) \quad \eta_t + \eta \eta_x + (\beta - 1) \eta_{xx} - \eta_{yy} - \delta \Lambda^3(\eta) + \epsilon \Delta^2 \eta = 0, \quad (x, y) \in \mathbb{T}^2, t > 0$$

with initial data

$$\eta(x, y, 0) = \eta_0(x, y), \quad (x, y) \in \mathbb{T}^2.$$

In what follows, we will drop the tilde notation.

3. GLOBAL EXISTENCE OF STRONG SOLUTIONS

In this section, we will state the global well-posedness result of the initial-value problem (3):

Theorem 1. *If $\eta_0 \in H^2(\mathbb{T}^2)$, then for every $0 < T < \infty$ the initial value problem (3) has a unique solution*

$$\eta \in C([0, T]; H^2(\mathbb{T}^2)) \cap L^2(0, T; H^4(\mathbb{T}^2)).$$

Proof. Step 1 : L^2 estimate. We multiply (3) by η and integrate by parts to obtain

$$\frac{1}{2} \frac{d}{dt} \|\eta\|_{L^2}^2 = (\beta - 1) \|\eta_x\|_{L^2}^2 - \|\eta_y\|_{L^2}^2 + \delta \int_{\mathbb{T}^2} \Lambda^3(\eta) \eta - \epsilon \|\Delta \eta\|_{L^2}^2.$$

Using the inequality

$$(\beta - 1) |\xi_1|^2 - |\xi_2|^2 + \delta |\xi|^3 - \frac{\epsilon}{2} |\xi|^4 \leq \frac{\delta^2}{2\epsilon} + (\beta - 2),$$

together with the Fourier transform, we find that

$$\frac{d}{dt} \|\eta\|_{L^2}^2 \leq -\epsilon \|\Delta \eta\|_{L^2}^2 + \left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) \|\eta\|_{L^2}^2.$$

An application of Gronwall's inequality lead us to

$$\|\eta\|_{L^2}^2 + \epsilon \int_0^t \exp \left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) (t - s) \right) \|\Delta \eta\|_{L^2}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp \left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) t \right).$$

Hence,

$$\|\eta\|_{L^2}^2 + \epsilon \int_0^t \|\eta\|_{H^2}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp \left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) t \right)$$

Step 2 : H^1 estimate. Now we multiply (3) by $-\Delta \eta$ and integrate by parts to obtain that

$$\frac{1}{2} \frac{d}{dt} \|\eta\|_{H^1}^2 = (\beta - 1) \|\eta_x\|_{H^1}^2 - \|\eta_y\|_{H^1}^2 + \delta \|\Lambda^{\frac{3}{2}} \eta\|_{H^1}^2 - \epsilon \|\Delta \eta\|_{H^1}^2 + \|\eta\|_{L^\infty} \|\eta\|_{H^1} \|\Delta \eta\|_{L^2}.$$

Using the same method as in step 1, we get

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta\|_{H^1}^2 &\leq -\frac{3\epsilon}{4} \|\Delta \eta\|_{H^1}^2 + \left(\frac{\delta^2}{\epsilon} + \beta - 2 \right) \|\eta\|_{H^1}^2 + \frac{1}{2\epsilon} \|\eta\|_{L^\infty}^2 \|\eta\|_{H^1}^2 + \frac{\epsilon}{2} \|\Delta \eta\|_{L^2}^2 \\ &\leq -\frac{\epsilon}{4} \|\Delta \eta\|_{H^1}^2 + \left(\frac{\delta^2}{\epsilon} + \beta - 2 + \frac{1}{2\epsilon} \|\eta\|_{L^\infty}^2 \right) \|\eta\|_{H^1}^2, \end{aligned}$$

which implies

$$\frac{d}{dt} \|\eta\|_{H^1}^2 \leq -\frac{\epsilon}{2} \|\Delta \eta\|_{H^1}^2 + \left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 + \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2 \right) \|\eta\|_{H^1}^2.$$

Using Gronwall's inequality, we find that

$$\begin{aligned} \|\eta\|_{H^1}^2 + \frac{\epsilon}{2} \int_0^t \exp \left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 \right) (t - s) + \int_s^t \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2 \right) \|\Delta \eta\|_{H^1}^2 ds \\ \leq \|\eta_0\|_{H^1}^2 \exp \left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 \right) t + \int_0^t \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2 \right). \end{aligned}$$

From step 1, we already have that

$$\epsilon \int_0^t \|\eta\|_{H^2}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp\left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4\right)t\right).$$

Using the Sobolev embedding, we get that

$$\epsilon \int_0^t \|\eta\|_{L^\infty}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp\left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4\right)t\right).$$

Inserting this into the inequality above, we obtain

$$\begin{aligned} \|\eta\|_{H^1}^2 + \frac{\epsilon}{2} \int_0^t \exp\left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4\right)(t-s) + \int_s^t \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2\right) \|\Delta\eta\|_{H^1}^2 ds \\ \leq \|\eta_0\|_{H^1}^2 \exp\left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4\right)t + \frac{1}{\epsilon^2} \|\eta_0\|_{L^2}^2 \exp\left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4\right)t\right)\right) \\ \leq \|\eta_0\|_{H^1}^2 \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^1})t)). \end{aligned}$$

Hence, we conclude that

$$\begin{aligned} \|\eta\|_{H^1}^2 + \frac{\epsilon}{2} \int_0^t \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^1})(t-s))) \|\Delta\eta\|_{H^1}^2 ds \\ \leq \|\eta_0\|_{H^1}^2 \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^1})t)). \end{aligned}$$

In particular,

$$\|\eta\|_{H^1}^2 + \frac{\epsilon}{2} \int_0^t \|\Delta\eta\|_{H^1}^2 ds \leq \|\eta_0\|_{H^1}^2 \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^1})t)).$$

Step 3 : H^2 estimate. We multiply (3) by $\Delta^2\eta$ and integrate by parts to obtain that

$$\frac{1}{2} \frac{d}{dt} \|\eta\|_{H^2}^2 = (\beta - 1) \|\eta_x\|_{H^2}^2 - \|\eta_y\|_{H^2}^2 + \delta \|\Lambda^{\frac{3}{2}}\|_{H^2}^2 - \epsilon \|\Delta\eta\|_{H^2}^2 + \|\eta\|_{L^\infty} \|\eta\|_{H^1} \|\Delta^2\eta\|_{L^2}.$$

Using the same method as in step 1 and step 2, we can obtain

$$\begin{aligned} \frac{d}{dt} \|\eta\|_{H^2}^2 &\leq -\frac{\epsilon}{2} \|\Delta\eta\|_{H^2}^2 + \left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 + \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2\right) \|\eta\|_{H^1}^2 \\ &\leq -\frac{\epsilon}{2} \|\Delta\eta\|_{H^2}^2 + \left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 + \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2\right) \|\eta\|_{H^2}^2, \end{aligned}$$

Using Gronwall's inequality again, we obtain that

$$\begin{aligned} \|\eta\|_{H^2}^2 + \frac{\epsilon}{2} \int_0^t \exp\left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4\right)(t-s) + \int_s^t \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2\right) \|\Delta\eta\|_{H^2}^2 ds \\ \leq \|\eta_0\|_{H^2}^2 \exp\left(\left(\frac{2\delta^2}{\epsilon} + 2\beta - 4\right)t + \int_0^t \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2\right). \end{aligned}$$

Then, the Sobolev embedding implies that

$$\begin{aligned} \|\eta\|_{H^2}^2 + \frac{\epsilon}{2} \int_0^t \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^2})(t-s))) \|\Delta\eta\|_{H^2}^2 ds \\ \leq \|\eta_0\|_{H^2}^2 \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^2})t)). \end{aligned}$$

Finally, we have

$$\|\eta\|_{H^2}^2 + \frac{\epsilon}{2} \int_0^t \|\Delta\eta\|_{H^2}^2 ds \leq \|\eta_0\|_{H^2}^2 \exp(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{H^2})t)).$$

Step 4 : Existence of solution We consider a positive, symmetric mollifier $J_{\epsilon'}$ (such as the periodic heat kernel), to approximate the initial value problem (3) by the regularized problem

$$\begin{aligned} \partial_t \eta_{\epsilon'} + J_{\epsilon'} * \frac{\partial_x \left((J_{\epsilon'} * \eta_{\epsilon'})^2 \right)}{2} \\ = J_{\epsilon'} * \left((1 - \beta) \partial_{xx} (J_{\epsilon'} * \eta_{\epsilon'}) + \partial_{yy} (J_{\epsilon'} * \eta_{\epsilon'}) + \delta \Lambda^3 (J_{\epsilon'} * \eta_{\epsilon'}) - \epsilon \Delta^2 (J_{\epsilon'} * \eta_{\epsilon'}) \right), \end{aligned}$$

with initial data

$$\eta_{\epsilon'}(0) = J_{\epsilon'} * \eta_0.$$

By the Picard's theorem, these sequence of regularized problems have a unique solution $\eta_{\epsilon'} = C^1([0, T_{\epsilon'}], H^2(\mathbb{T}^2))$. Moreover, these problems verify the same energy estimates as in steps 1-3 and, as a consequence, we can take $T = T(\eta_0)$ independent of ϵ' . Passing to the limit we conclude the existence of at least one solution in

$$\eta \in L^\infty(0, T; H^2) \cap L^2(0, T; H^4)$$

and since we have *a priori* estimates, these solutions exist for arbitrary long time T .

Step 5 : Uniqueness. We can prove the uniqueness of the solutions by contradiction, i.e. assuming that there exists two solutions of the problem (3), η_1 and η_2 , corresponding to the same initial data η_0 . We denote their difference by $\underline{\eta}$. Then we have

$$(4) \quad \underline{\eta}_t + \frac{1}{2} (\eta_1^2 - \eta_2^2)_x + (\beta - 1) \underline{\eta}_{xx} - \underline{\eta}_{yy} - \delta \Lambda^3(\underline{\eta}) + \epsilon \Delta^2 \underline{\eta} = 0.$$

As the proof of L^2 estimate in step 1, we multiply (4) by $\underline{\eta}$ and integrate by parts:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\underline{\eta}\|_{L^2}^2 &= (\beta - 1) \|\underline{\eta}_x\|_{L^2}^2 - \|\underline{\eta}_y\|_{L^2}^2 + \delta \|\Lambda^{\frac{3}{2}}(\underline{\eta})\|_{L^2}^2 - \epsilon \|\Delta \underline{\eta}\|_{L^2}^2 - \int_{\mathbb{T}^2} \frac{1}{2} (\eta_1^2 - \eta_2^2)_x \underline{\eta} \\ &\leq \left(\frac{\delta^2}{3\epsilon} + \beta - 2 \right) \|\underline{\eta}\|_{L^2}^2 - \frac{\epsilon}{4} \|\Delta \underline{\eta}\|_{L^2}^2 + \frac{1}{2} \|\eta_1 + \eta_2\|_{L^\infty} \|\underline{\eta}\|_{H^1} \|\underline{\eta}\|_{L^2} \\ &\leq \left(\frac{\delta^2}{3\epsilon} + \beta - 2 \right) \|\underline{\eta}\|_{L^2}^2 - \frac{\epsilon}{4} \|\Delta \underline{\eta}\|_{L^2}^2 + \frac{\epsilon}{4} \|\underline{\eta}\|_{H^1}^2 + \frac{1}{4\epsilon} \|\eta_1 + \eta_2\|_{L^\infty}^2 \|\underline{\eta}\|_{L^2}^2 \\ &\leq \left(\frac{\delta^2}{3\epsilon} + \beta - 2 + \frac{1}{4\epsilon} \|\eta_1 + \eta_2\|_{L^\infty}^2 \right) \|\underline{\eta}\|_{L^2}^2 \\ &\leq \left(\frac{\delta^2}{3\epsilon} + \beta - 2 + \frac{1}{2\epsilon} (\|\eta_1\|_{L^\infty}^2 + \|\eta_2\|_{L^\infty}^2) \right) \|\underline{\eta}\|_{L^2}^2 \end{aligned}$$

Using Gronwall's inequality, we have that

$$\|\underline{\eta}\|_{L^2}^2 \leq \|\underline{\eta}_0\|_{L^2}^2 \exp \left(\left(\frac{2\delta^2}{3\epsilon} + 2\beta - 4 \right) t + \frac{1}{\epsilon} \int_0^t \|\eta_1\|_{L^\infty}^2 + \|\eta_2\|_{L^\infty}^2 \right).$$

From step 1, we already have that

$$\epsilon \int_0^t \|\eta_1\|_{H^2}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp \left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) t \right),$$

and

$$\epsilon \int_0^t \|\eta_2\|_{H^2}^2 ds \leq \|\eta_0\|_{L^2}^2 \exp \left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4 \right) t \right).$$

Thus, the uniqueness of solution follows from the inequality

$$\|\underline{\eta}\|_{L^2}^2 \leq \|\underline{\eta}_0\|_{L^2}^2 \exp \left(\exp(C(\delta, \epsilon, \beta, \|\eta_0\|_{L^2})t) \right).$$

Step 6 : Endpoint continuity in time. To conclude the endpoint continuity, we can perform a standard argument using the parabolic gain of regularity $L^2(0, T; H^4)$. Indeed, we can take $0 < \sigma \ll 1$ as small as desired and there exists a $0 < \sigma' < \sigma$ such that $u(\sigma') \in H^4$. Repeating the same argument as before, we find a solution

$$\eta_\sigma \in L^\infty(\sigma', T; H^4) \cap C([\sigma', T], H^2).$$

Because of the uniqueness of solution we obtain the continuity of the original solution

$$\eta \in C((0, T], H^2).$$

Finally, the continuity at the origin is a consequence of the energy estimates. \square

4. LARGE TIME DYNAMICS

The goal of this section is to prove uniform boundedness of solutions $\eta \in L^\infty([0, \infty); L^2(\mathbb{T}^2))$. In other words, we establish the existence of an absorbing ball in L^2 by collecting global bounds showing the dissipative character of the equation. We start by proving the following two Gagliardo-Nirenberg inequalities

Lemma 1. *For smooth enough periodic functions we have that the following two inequalities hold true*

$$\begin{aligned} \|\nabla\eta\|_{L^4(\mathbb{T}^2)}^2 &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)}, \\ \|\Delta\eta^2\|_{L^2(\mathbb{T}^2)} &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)}. \end{aligned}$$

Proof. We start proving the first inequality:

$$\begin{aligned} \|\nabla\eta\|_{L^4(\mathbb{T}^2)}^4 &= \int_{\mathbb{T}^2} (\nabla\eta \cdot \nabla\eta)^2 \\ &= - \int_{\mathbb{T}^2} \eta \nabla \cdot (\nabla\eta |\nabla\eta|^2) \\ &= - \int_{\mathbb{T}^2} \eta \Delta\eta |\nabla\eta|^2 + \eta \nabla\eta \cdot \nabla |\nabla\eta|^2 \\ &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)}\|\nabla\eta\|_{L^2(\mathbb{T}^2)}^2 \\ &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)}\|\nabla\eta\|_{L^4(\mathbb{T}^2)}^2. \end{aligned}$$

Therefore, we conclude our result by noticing that

$$\begin{aligned} \|\Delta\eta^2\|_{L^2(\mathbb{T}^2)} &\leq C\|\eta\Delta\eta + |\nabla\eta|^2\|_{L^2(\mathbb{T}^2)} \\ &\leq C\|\eta\Delta\eta\|_{L^2(\mathbb{T}^2)} + \|\nabla\eta\|_{L^2(\mathbb{T}^2)}^2 \\ &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)} + \|\nabla\eta\|_{L^4(\mathbb{T}^2)}^2 \\ &\leq C\|\eta\|_{L^\infty(\mathbb{T}^2)}\|\Delta\eta\|_{L^2(\mathbb{T}^2)}. \end{aligned}$$

\square

Remark 1. *We observe that the previous constants C can be computed explicitly.*

We now require the following lemma which is a 2D version of [21] (see also [12]).

Lemma 2. *Let $(x_0, y_0) \in \mathbb{T}^2$, then for any $M > 0$ large, $\eta \in C^\infty(\mathbb{T}^2)$ with $\eta(x_0, y_0) = 0$, there exists a smooth periodic function $\varphi_M^{x_0, y_0} \in C^\infty(\mathbb{T}^2)$, such that the following inequality holds:*

$$\left| \int_{\mathbb{T}^2} \varphi_M^{x_0, y_0}(x, y) \eta^2(x, y) dx dy \right| \leq \frac{C}{M} \|\Delta\eta\|_{L^2(\mathbb{T}^2)} \|\eta\|_{L^\infty(\mathbb{T}^2)}.$$

Proof. We first define

$$\varphi_M^{x_0, y_0}(x, y) = \sum_{|\xi| \leq M} e^{-i\xi_1(x-x_0)} e^{-i\xi_2(y-y_0)}, \quad \xi = (\xi_1, \xi_2).$$

From Plancherel theorem, we have that

$$\begin{aligned} \int_{\mathbb{T}^2} \varphi_M^{x_0, y_0}(x, y) \eta^2(x, y) dx dy &= \sum_{|\xi| \leq M} \int_{\mathbb{T}^2} \eta^2(x, y) e^{-i\xi_1(x-x_0)} e^{-i\xi_2(y-y_0)} dx dy \\ &= \sum_{|\xi| \leq M} \int_{\mathbb{T}^2} \eta^2(x+x_0, y+y_0) e^{-i\xi_1 x} e^{-i\xi_2 y} dx dy \\ &= \sum_{|\xi| \leq M} \int_{\mathbb{T}^2} g(x, y) e^{-i\xi_1 x - i\xi_2 y} dx dy \\ &= 2\pi \sum_{|\xi| \leq M} \widehat{g}(\xi_1, \xi_2), \end{aligned}$$

where $g(x, y) = \eta^2(x+x_0, y+y_0)$. By the condition $\eta(x_0, y_0) = 0$, we immediately have $g(0, 0) = 0$. In the form of Fourier variables, we have that

$$\sum_{|\xi|} \widehat{g}(\xi_1, \xi_2) = 0.$$

Splitting the left-hand side of the equality above into high-frequency part and low-frequency part, we have that

$$\sum_{|\xi| \leq M} \widehat{g}(\xi_1, \xi_2) = - \sum_{|\xi| > M} \widehat{g}(\xi_1, \xi_2).$$

Then, we bound

$$\begin{aligned} \left| \int_{\mathbb{T}^2} \varphi_M^{x_0, y_0}(x, y) \eta^2(x, y) dx dy \right| &\leq C \left| \sum_{|\xi| \leq M} \widehat{g}(\xi_1, \xi_2) \right| \\ &\leq C \left| \sum_{|\xi| > M} \widehat{g}(\xi_1, \xi_2) \right| \\ &\leq C \left| \sum_{|\xi| > M} |\xi|^2 \widehat{g}(\xi_1, \xi_2) \frac{1}{|\xi|^2} \right| \\ &\leq C \left(\sum_{|\xi| > M} (|\xi|^2 \widehat{g}(\xi_1, \xi_2))^2 \right)^{\frac{1}{2}} \left(\sum_{|\xi| > M} \frac{1}{|\xi|^4} \right)^{\frac{1}{2}} \\ &\leq C \|\Delta g\|_{L^2(\mathbb{T}^2)} \left(\sum_{|\xi| > M} \frac{1}{|\xi|^4} \right)^{\frac{1}{2}}. \end{aligned}$$

We use polar coordinates to estimate $\sum_{|\xi| > M} \frac{1}{|\xi|^4}$:

$$\sum_{|\xi| > M} \frac{1}{|\xi|^4} = \sum_{\sqrt{\xi_1^2 + \xi_2^2} > M} \frac{1}{(\xi_1^2 + \xi_2^2)^2} \leq \int_0^{2\pi} \int_M^\infty \frac{1}{r^3} dr d\theta \leq \int_0^{2\pi} \frac{1}{2M^2} d\theta \leq \frac{\pi}{M^2}.$$

Using the second estimate in Lemma 1, we conclude that

$$\begin{aligned} \left| \int_{\mathbb{T}^2} \varphi_M^{x_0, y_0}(x, y) \eta^2(x, y) dx dy \right| &\leq \frac{C}{M} \|\Delta g\|_{L^2(\mathbb{T}^2)} \\ &\leq \frac{C}{M} \|\Delta \eta^2\|_{L^2(\mathbb{T}^2)} \\ &\leq \frac{C}{M} \|\eta\|_{L^\infty(\mathbb{T}^2)} \|\Delta \eta\|_{L^2(\mathbb{T}^2)}. \end{aligned}$$

□

The rest of this section is devoted to prove that the solutions of problem (3) remain uniformly bounded in L^2 . The following background flow method was first used by Nicolaenko, Scheurer, & Temam [21] and then improved by Collet, Eckmann, Epstein & Stubbe [5], Goodman [11] and Bronski & Gambill [1]. In terms of the rigorous results, a global bound on the solution is given by the following theorem :

Theorem 2. *Let $\eta_0 \in H^2(\mathbb{T}^2)$ be the zero mean initial data. Then the solution η of the initial-value problem (3) satisfies*

$$(5) \quad \limsup_{t \rightarrow \infty} \|\eta(t)\|_{L^2(\mathbb{T}^2)} \leq R_{\epsilon, \delta}.$$

Proof. The proof is based on the construction of a Lyapunov functional, $\mathcal{F}(t)$, such that

$$\frac{d}{dt} \mathcal{F}(t) \leq 0,$$

if

$$\mathcal{F}(t) \geq R_{\epsilon, \delta},$$

i.e. implying the existence of an absorbing set in L^2 . We first let ϕ be a smooth, 2π -periodic function, which we will choose later. Then, we multiply equation (3) by $\eta - \phi$, and integrate by parts:

$$\int \int ((\eta - \phi)_t + \eta \eta_x + (\beta - 1) \eta_{xx} - \eta_{yy} - \delta \Lambda^3(\eta) + \epsilon \Delta^2 \eta) (\eta - \phi) = 0,$$

thus,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta - \phi\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{T}^2} \phi_x \eta^2 - (\beta - 1) \|\eta_x\|_{L^2}^2 + \|\eta_y\|_{L^2}^2 + \epsilon \|\Delta \eta\|_{L^2}^2 \\ - \int_{\mathbb{T}^2} (\beta - 1) \eta_{xx} \phi + \int_{\mathbb{T}^2} \eta_{yy} \phi - \delta \int_{\mathbb{T}^2} \Lambda^3 \eta (\eta - \phi) - \int_{\mathbb{T}^2} \epsilon \Delta \eta \Delta \phi = 0. \end{aligned}$$

For the term corresponding to the nonlocal operator Λ , we note that, by the Plancherel theorem,

$$\int_{\mathbb{T}^2} \Lambda \eta \phi = \int_{\mathbb{T}^2} |\xi| \hat{\eta} \bar{\hat{\phi}} = \int_{\mathbb{T}^2} \hat{\eta} \overline{|\xi| \hat{\phi}} = \int_{\mathbb{T}^2} \eta \Lambda \phi,$$

and, with the help of Young's inequality,

$$\delta \int_{\mathbb{T}^2} \Lambda^3 \eta \phi = \delta \int_{\mathbb{T}^2} \eta \Lambda^3 \phi \leq \frac{\delta}{2} \|\eta\|_{L^2}^2 + \frac{\delta}{2} \|\phi\|_{H^3}^2.$$

Then we have that,

$$\begin{aligned}
(6) \quad \frac{1}{2} \frac{d}{dt} \|\eta - \phi\|_{L^2}^2 &= (\beta - 1) \|\eta_x\|_{L^2}^2 - \|\eta_y\|_{L^2}^2 - \epsilon \|\Delta \eta\|_{L^2}^2 - \int_{\mathbb{T}^2} \frac{\phi_x}{2} \eta^2 \\
&\quad - \int_{\mathbb{T}^2} (\beta - 1) \eta_x \phi_x + \int_{\mathbb{T}^2} \eta_y \phi_y + \delta \int_{\mathbb{T}^2} \Lambda^3 \eta (\eta - \phi) + \int_{\mathbb{T}^2} \epsilon \Delta \eta \Delta \phi \\
&\leq \frac{3}{2} (\beta - 1) \|\eta_x\|_{L^2}^2 - \frac{1}{2} \|\eta_y\|_{L^2}^2 - \frac{2\epsilon}{3} \|\Delta \eta\|_{L^2}^2 - \int_{\mathbb{T}^2} \frac{\phi_x}{2} \eta^2 + \delta \int_{\mathbb{T}^2} \Lambda^3 \eta \eta \\
&\quad + \frac{\delta}{2} \|\eta\|_{L^2}^2 + \frac{\delta}{2} \|\phi\|_{H^3}^2 + \frac{\beta - 1}{2} \|\phi_x\|_{L^2}^2 + \frac{1}{2} \|\phi_y\|_{L^2}^2 + \frac{3}{4\epsilon} \|\epsilon \Delta \phi\|_{L^2}^2.
\end{aligned}$$

Now, we define the following function

$$f(|\xi|) = f(z) := \frac{\delta}{2} + \left(\frac{3\beta}{2} - 2\right) z^2 + \delta z^3 - \frac{2\epsilon}{3} z^4, \quad z \geq 0.$$

Observe that $f(z)$ has at most three zeros. Since $\epsilon > 0$, we find that $f(z)$ is bounded and has global maximum at point $z_1 = \frac{9\delta - \sqrt{81\delta^2 + 288\epsilon\beta - 384\epsilon}}{16\epsilon}$. Then we have the following inequality

$$\frac{\delta}{2} + \frac{3}{2} (\beta - 1) |\xi_1|^2 - \frac{1}{2} |\xi_2|^2 + \delta |\xi|^3 - \frac{2\epsilon}{3} |\xi|^4 \leq f(z_1) := C(\beta, \delta, \epsilon).$$

Inserting this relation into (6), we obtain that

$$(7) \quad \frac{1}{2} \frac{d}{dt} \|\eta - \phi\|_{L^2(\mathbb{T}^2)}^2 = -\frac{\epsilon}{3} \|\Delta \eta\|_{L^2(\mathbb{T}^2)}^2 - \|\eta\|_{L^2}^2 + \int_{\mathbb{T}^2} \left(\lambda - \frac{\phi_x}{2}\right) \eta^2 + F(\phi),$$

where

$$\lambda = C(\beta, \delta, \epsilon) + 1,$$

and

$$F(\phi) = \frac{\beta - 1}{2} \|\phi_x\|_{L^2}^2 + \frac{1}{2} \|\phi_y\|_{L^2}^2 + \frac{3}{4\epsilon} \|\epsilon \Delta \phi\|_{L^2}^2 + \frac{\delta}{2} \|\phi\|_{H^3}^2.$$

For the moment (we will get rid of this assumption later), we assume that $\eta(0, 0, t) = 0$ and define $\varphi_M^{0,0}(x, y)$ as in the proof of lemma 2 and $\phi(x, y)$ such that

$$(8) \quad \lambda - \frac{\phi_x(x, y)}{2} = \lambda \varphi_M^{0,0}(x, y).$$

We consider the functional

$$\mathcal{F}(t) = \|\eta - \phi\|_{L^2(\mathbb{T}^2)}^2.$$

Inserting this into (7) and using lemma 2, we obtain

$$\frac{1}{2} \frac{d}{dt} \|\eta - \phi\|_{L^2(\mathbb{T}^2)}^2 \leq -\frac{\epsilon}{3} \|\Delta \eta\|_{L^2(\mathbb{T}^2)}^2 - \|\eta\|_{L^2}^2 + \lambda \frac{c}{M} \|\Delta \eta\|_{L^2(\mathbb{T}^2)}^2 + F(\phi)$$

Then we take $M = M(\beta, \delta, \epsilon)$ such that

$$\lambda \frac{c}{M} = (C(\beta, \delta, \epsilon) + 1) \frac{c}{M} \leq \frac{\epsilon}{3},$$

and we get

$$(9) \quad \frac{1}{2} \frac{d}{dt} \|\eta - \phi\|_{L^2(\mathbb{T}^2)}^2 \leq -\|\eta - \phi\|_{L^2(\mathbb{T}^2)}^2 + \|\phi\|_{L^2}^2 + F(\phi),$$

or, equivalently,

$$\frac{1}{2} \frac{d}{dt} \mathcal{F} \leq -\mathcal{F} + \|\phi\|_{L^2}^2 + F(\phi),$$

which allows us to conclude the uniform boundedness of \mathcal{F} . Thus, if $1 \ll \|\eta(t)\|_{L^2}$ and $\eta(0, 0, t) = 0$, we have that

$$\begin{aligned} \|\eta\|_{L^2(\mathbb{T}^2)} &\leq \|\eta - \phi\|_{L^2(\mathbb{T}^2)} + \|\phi\|_{L^2(\mathbb{T}^2)} \\ &\leq \left(\|\eta_0 - \phi\|_{L^2(\mathbb{T}^2)}^2 + \|\phi\|_{L^2}^2 + F(\phi) \right)^{\frac{1}{2}} e^{-4t} + 2\|\phi\|_{L^2} + F(\phi)^{\frac{1}{2}} \\ &\leq \left(\|\eta_0\|_{L^2} + 2\|\phi\|_{L^2} + \frac{\beta}{2}\|\phi\|_{H^1} + \frac{3}{4\epsilon}\|\phi\|_{H^2} + \frac{\delta}{2}\|\phi\|_{H^3} \right) e^{-4t} \\ &\quad + 2\|\phi\|_{L^2} + \frac{\beta}{2}\|\phi\|_{H^1} + \frac{3}{4\epsilon}\|\phi\|_{H^2} + \frac{\delta}{2}\|\phi\|_{H^3} \\ &:= R_{\epsilon, \delta}. \end{aligned}$$

Now we want to get rid of the assumption $\eta(0, 0, t)$. According to the mean-zero assumption, for every t^* ,

$$\int_{\mathbb{T}^2} \eta(x, y, t) dx dy = 0, \forall t > 0,$$

and we are able to find at least one point $(x_0(t^*), y_0(t^*))$ such that

$$\eta(x_0(t^*), y_0(t^*), t^*) = 0.$$

We define $\varphi_M^{x_0(t^*), y_0(t^*)}(x, y)$ as in the proof of lemma 2 and $\phi_{t^*}(x, y)$ such that

$$(10) \quad \lambda - \frac{(\phi_{t^*})_x(x, y)}{2} = \lambda \varphi_M^{x_0(t^*), y_0(t^*)}(x, y).$$

Now, following [11], we introduce the following set of functions

$$\mathcal{S} = \{\tilde{\phi} : \tilde{\phi}(x, y) = \phi(x + r_x, y + r_y) \mid -\pi \leq r_x, r_y \leq \pi\},$$

where ϕ is defined by (8) and the functional

$$\mathcal{F}(t) = \text{dist}(\eta(t), \mathcal{S})^2 = \inf_{\tilde{\phi}} \|\eta(t) - \tilde{\phi}\|_{L^2(\mathbb{T}^2)}^2.$$

We observe that \mathcal{S} is contained in a (large enough) ball in L^2 . We also note that the sequence of $\phi_{t^*} \in \mathcal{S}$ for all t^* . Then, we want to prove that, at this time instant t^* , we have that

$$\left. \frac{d}{dt} \|\eta - \phi_{t^*}\|_{L^2} \right|_{t=t^*} < 0.$$

In particular, this implies that η can not leave a ball around \mathcal{S} without being immediately attracted back again.

Then, we assume that, at a certain t^* , we have that $\mathcal{F}(t^*) \geq R_{\epsilon, \delta} \gg 1$. We chose $(x(t^*), y(t^*))$ and define ϕ_{t^*} . As a consequence of the definition of the Lyapunov functional, we have that

$$\|\eta(t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 \geq \mathcal{F}(t^*) \geq R_{\epsilon, \delta} \gg 1.$$

We can obtain the inequality (analog to (9))

$$(11) \quad \frac{1}{2} \frac{d}{dt} \|\eta(t + t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 \leq -\|\eta(t + t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 + \|\phi_{t^*}\|_{L^2}^2 + F(\phi_{t^*}).$$

It follows that

$$\left. \frac{d}{dt} \|\eta(t + t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 \right|_{t=0} \leq 0, \text{ if } \|\eta(t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)} \geq R_{\epsilon, \delta} \gg 1,$$

where $R_{\epsilon, \delta}$ is independent of t^* . Therefore,

$$\mathcal{F}(t) \leq \|\eta(t + t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 \leq \|\eta(t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2$$

if $\|\eta(t^*) - \tilde{\phi}_{t^*}\|_{L^2(\mathbb{T}^2)}^2 \geq R_{\epsilon, \delta}$. Thus, $\mathcal{F}(t)$ is a bounded function of time. \square

Similarly as we have obtained that there exists an absorbing set in L^2 , we can conclude the existence of an absorbing set in higher Sobolev norms.

Theorem 3. *Let $\eta_0 \in H^2(\mathbb{T}^2)$ be the zero mean initial data. Then the solution η of the initial-value problem (3) satisfies*

$$\limsup_{t \rightarrow \infty} \|\eta(t)\|_{H^2(\mathbb{T}^2)} \leq R'_{\epsilon, \delta}.$$

Proof. Recalling the existence of an absorbing set in the L^2 -norm and the regularity results in Theorem 1, so the proof is straightforward by using a bootstrap argument.

We first show that there exists an absorbing set in the H^1 -norm. Inequality (5) implies that for a $T > 0$ large enough, we have

$$\|\eta(t)\|_{L^2(\mathbb{T}^2)} \leq R_{\epsilon, \delta} + 1, \quad \forall t > T.$$

Combining this inequality with the L^2 energy estimate in the proof of Theorem 1, we obtain that

$$\|\eta\|_{L^\infty([0, T]; L^2)}^2 \leq \|\eta_0\|_{L^2}^2 \exp\left(\left(\frac{\delta^2}{\epsilon} + 2\beta - 4\right)T\right),$$

which results in that there exists a constant depending on initial data, δ and ϵ such that

$$\max_{0 \leq t < \infty} \|\eta(t)\|_{L^2(\mathbb{T}^2)}^2 \leq C(\|\eta_0\|_{L^2}, \epsilon, \delta).$$

We multiply (3) by $-\Delta\eta$ and integrate by parts to obtain that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta\|_{H^1}^2 &\leq (\beta - 1) \|\eta_x\|_{H^1}^2 - \|\eta_y\|_{H^1}^2 + \delta \|\Lambda^{\frac{3}{2}}\|_{H^1}^2 - \epsilon \|\Delta\eta\|_{H^1}^2 + \|\eta\|_{L^4}^2 \|\Delta\eta_x\|_{L^2} \\ &\leq c_\epsilon \|\eta\|_{L^4}^4 - \frac{\epsilon}{2} \|\eta\|_{H^3}^2 + (\beta - 1) \|\eta_x\|_{H^1}^2 - \|\eta_y\|_{H^1}^2 + \delta \|\Lambda^{\frac{3}{2}}\|_{H^1}^2 \\ &\leq C_{\epsilon, \delta, \eta_0} \|\eta\|_{H^1}^2 - \frac{\epsilon}{2} \|\eta\|_{H^3}^2 + (\beta - 1) \|\eta_x\|_{H^1}^2 - \|\eta_y\|_{H^1}^2 + \delta \|\Lambda^{\frac{3}{2}}\|_{H^1}^2, \\ &\leq C_{\epsilon, \delta, \eta_0} \|\eta\|_{H^1}^2 - \frac{\epsilon}{4} \|\eta\|_{H^3}^2 + C_{\epsilon, \delta} \|\eta\|_{L^2}^2 \\ &\leq -\frac{\epsilon}{8} \|\eta\|_{H^3}^2 + C_{\epsilon, \delta, \eta_0} \\ &\leq -\frac{\epsilon}{16} \|\eta\|_{H^1}^2 - \frac{\epsilon}{16} \|\eta\|_{H^3}^2 + C_{\epsilon, \delta, \eta_0}. \end{aligned}$$

where we used the Plancherel Theorem, the Poincaré inequality and the Sobolev inequality

$$\|\eta\|_{L^4} \leq C \|\eta\|_{H^{1/2}} \leq C \|\eta\|_{L^2}^{1/2} \|\eta\|_{H^1}^{1/2}.$$

It follows that

$$\frac{d}{dt} \|\eta\|_{H^1}^2 + \frac{\epsilon}{8} \|\eta\|_{H^1}^2 \leq C(\|\eta_0\|_{L^2}, \epsilon, \delta).$$

Using the Gronwall inequality, we immediately obtain the uniform bound

$$\|\eta(t)\|_{H^1}^2 \leq C(\|\eta_0\|_{H^1}, \epsilon, \delta).$$

Recall that the H^2 energy estimate is

$$\frac{d}{dt} \|\eta\|_{H^2}^2 + \frac{\epsilon}{2} \|\eta\|_{H^4}^2 \leq \left(\frac{2\delta^2}{\epsilon} + 2\beta - 4 + \frac{1}{\epsilon} \|\eta\|_{L^\infty}^2 \right) \|\eta\|_{H^1}^2,$$

so we can mimic the previous proof to obtain that there exists an absorbing set in H^2 . The proof is completed. \square

5. ANALYTICITY

The aim of this section is to show instant analyticity for the solutions of (3). We shall prove that the solutions of (3) are analytic in a strip. In order to do this, we use the method developed by Collet, Eckmann, Epstein & Stubbe in [4] (see also [7]). Roughly speaking, our proof is based on *a priori* estimates for functions in certain Gevrey class.

Given a function $\sigma(t)$, we consider the weighted exponential operators

$$e^{\sigma(t)\Lambda}\eta = \sum_{\xi \in \mathbb{T}^2} \hat{\eta}(\xi) e^{\sigma(t)|\xi|} e^{i\xi \cdot x}$$

for functions in the space

$$G := \left\{ \eta \in L^2(\mathbb{T}^2) : \sum_{\xi \in \mathbb{Z}^2} e^{2\sigma(t)|\xi|} |\hat{\eta}(\xi)|^2 < \infty \right\}.$$

We observe that the functions in G are analytic. We also define the inner product and norm on this Hilbert space by

$$\langle \mu, \eta \rangle_{\sigma(t)} = \int_{\mathbb{T}^2} e^{\sigma(t)\Lambda} \mu e^{\sigma(t)\Lambda} \eta dx = 4\pi^2 \sum_{\ell \in \mathbb{Z}^2} e^{2\sigma(t)|\ell|} \widehat{\mu(\ell)} \overline{\widehat{\eta(\ell)}},$$

$$\|\eta\|_{\sigma(t)}^2 = \|e^{\sigma(t)\Lambda} \eta\|_{L^2}^2.$$

With these previous definitions, we can state the main result of this section.

Theorem 4. *Let η_0 be given in $H^2(\mathbb{T}^2)$. Then, there exists T_0 depending on $\eta_0, \epsilon, \beta, \delta$ such that the solution of (3) satisfies*

$$\|e^{\sigma(t)\Lambda} \eta(t)\|_{L^2}^2 \leq 1 + 2C_{\epsilon, \delta, \eta_0}^2, \quad \forall t \geq 0$$

where $\sigma(t) = \min\{\tanh(t), \tanh(\frac{T_0}{2})\}$. In particular, it becomes analytic for $t > 0$.

Before proving theorem 4, we first state some auxiliary lemmas:

Lemma 3. *For every $b > a \geq 0$,*

$$\|\Lambda^{\frac{a}{2}} \eta\|_{\sigma(t)}^2 \leq \|\Lambda^{\frac{b}{2}} \eta\|_{\sigma(t)}^{\frac{2a}{b}} \|\eta\|_{\sigma(t)}^{2 - \frac{2a}{b}}.$$

Proof.

$$\begin{aligned}
\|\Lambda^{\frac{a}{2}}\eta\|_{\sigma(t)}^2 &= \|e^{\sigma(t)\Lambda}\Lambda^{\frac{a}{2}}\eta\|_{L^2}^2 \\
&= \sum_{\xi \in \mathbb{Z}^2} e^{2\sigma(t)|\xi|} |\xi|^a |\hat{\eta}(\xi)|^2 \\
&= \sum_{\xi \in \mathbb{Z}^2} e^{\sigma(t)|\xi|^{\frac{2a}{b}}} |\xi|^a |\hat{\eta}(\xi)|^{\frac{2a}{b}} e^{\sigma(t)|\xi|(2-\frac{2a}{b})} |\hat{\eta}(\xi)|^{2-\frac{2a}{b}} \\
&\leq \left(\sum_{\xi \in \mathbb{Z}^2} \left(e^{\sigma(t)|\xi|^{\frac{2a}{b}}} |\xi|^a |\hat{\eta}(\xi)|^{\frac{2a}{b}} \right)^{\frac{b}{a}} \right)^{\frac{a}{b}} \left(\sum_{\xi \in \mathbb{Z}^2} \left(e^{\sigma(t)|\xi|(2-\frac{2a}{b})} |\hat{\eta}(\xi)|^{2-\frac{2a}{b}} \right)^{\frac{b}{b-a}} \right)^{\frac{b-a}{b}} \\
&\leq \left(\sum_{\xi \in \mathbb{Z}^2} e^{2\sigma(t)|\xi|} |\xi|^b |\hat{\eta}(\xi)|^2 \right)^{\frac{a}{b}} \left(\sum_{\xi \in \mathbb{Z}^2} e^{2\sigma(t)|\xi|} |\hat{\eta}(\xi)|^2 \right)^{\frac{b-a}{b}} \\
&\leq \|\Lambda^{\frac{b}{2}}\eta\|_{\sigma(t)}^{\frac{2a}{b}} \|\eta\|_{\sigma(t)}^{2-\frac{2a}{b}}.
\end{aligned}$$

□

And an auxiliary lemma estimating the nonlinear term:

Lemma 4. $|\langle \eta\eta_x, \eta \rangle_{\sigma(t)}| \leq c \|\Lambda\eta\|_{\sigma(t)} \|\Lambda^{\frac{1}{2}}\eta\|_{\sigma(t)}^2.$

Proof. We first denote $\eta^* = e^{\sigma(t)\Lambda}\eta$, then

$$\widehat{\eta^*(j)} = e^{\sigma(t)|j|} \hat{\eta}(j).$$

By the definition of Fourier series, we have

$$\eta = \sum_{j \in \mathbb{Z}^2} \hat{\eta}(j) e^{ij \cdot x}, \quad e^{\sigma(t)\Lambda}\eta = \sum_{j \in \mathbb{Z}^2} \hat{\eta}(j) e^{ij \cdot x} e^{\sigma(t)|j|}.$$

In fact,

$$\begin{aligned}
\langle \eta\eta_x, \eta \rangle_{\sigma(t)} &= (2\pi)^2 i \sum_{\ell \in \mathbb{Z}^2} \sum_{j \in \mathbb{Z}^2} (\hat{\eta}(j) \hat{\eta}(\ell - j) j_1 \overline{\hat{\eta}(\ell)}) e^{2\sigma(t)|\ell|} \\
&= (2\pi)^2 i \sum_{\ell \in \mathbb{Z}^2} \sum_{j \in \mathbb{Z}^2} (\widehat{\eta^*}(j) \widehat{\eta^*}(\ell - j)) j_1 \overline{\widehat{\eta^*}(\ell)} e^{\sigma(t)(|\ell| - |j| - |\ell - j|)}.
\end{aligned}$$

Since $|\ell| \leq |j| + |\ell - j|$ leads to $|\ell| - |j| - |\ell - j| \leq 0$, we have that

$$e^{\sigma(t)(|\ell| - |j| - |\ell - j|)} \leq 1.$$

Moreover,

$$\begin{aligned}
|\langle \eta\eta_x, \eta \rangle_{\sigma(t)}| &\leq (2\pi)^2 \left| i \sum_{\ell \in \mathbb{Z}^2} \sum_{j \in \mathbb{Z}^2} (\widehat{\eta^*}(j) \widehat{\eta^*}(\ell - j)) j_1 \overline{\widehat{\eta^*}(\ell)} e^{\sigma(t)(|\ell| - |j| - |\ell - j|)} \right| \\
&\leq (2\pi)^2 \sum_{\ell \in \mathbb{Z}^2} \sum_{j \in \mathbb{Z}^2} |\widehat{\eta^*}(j)| |\widehat{\eta^*}(\ell - j)| |j| |\overline{\widehat{\eta^*}(\ell)}| \\
&= (2\pi)^2 \sum_{k+j=\ell} |\widehat{\eta^*}(j)| |j| |\widehat{\eta^*}(k)| |\overline{\widehat{\eta^*}(\ell)}| \\
&= \int_{\mathbb{T}^2} \phi(x) \theta(x) \overline{\psi(x)} dx
\end{aligned}$$

where

$$\phi(x) = \sum_{j \in \mathbb{Z}^2} |\widehat{\eta}^*(j)| |j| e^{ijx}, \quad \theta(x) = \sum_{k \in \mathbb{Z}^2} |\widehat{\eta}^*(k)| e^{ikx}, \quad \psi(x) = \sum_{\ell \in \mathbb{Z}^2} |\widehat{\eta}^*(\ell)| e^{i\ell x}.$$

Notice that $|\widehat{\phi}(j)| = |j| |\widehat{\eta}^*(j)|$ and $|\widehat{\theta}(k)| = |\widehat{\eta}^*(k)|$. We now bound the last integral by the Hölder inequality,

$$\begin{aligned} \int_{\mathbb{T}^2} \phi(x) \theta(x) \bar{\psi}(x) dx &\leq \|\phi(x)\|_{L^2} \|\theta(x)\|_{L^4} \|\psi(x)\|_{L^4} \\ &\leq c \|\phi(x)\|_{L^2} \|\theta(x)\|_{H^{\frac{1}{2}}}^2 \\ &= c \left(\sum_{j \in \mathbb{Z}^2} |\widehat{\phi}(j)|^2 \right)^{\frac{1}{2}} \sum_{j \in \mathbb{Z}^2} \left(|k|^{\frac{1}{2}} |\widehat{\theta}(k)| \right)^2 \\ &= c \left(\sum_{j \in \mathbb{Z}^2} |j|^2 |\widehat{\eta}^*(j)|^2 \right)^{\frac{1}{2}} \sum_{j \in \mathbb{Z}^2} \left(|k|^{\frac{1}{2}} |\widehat{\eta}^*(k)| \right)^2 \\ &= c \|\Lambda \eta^*\|_{L^2} \|\Lambda^{\frac{1}{2}} \eta^*\|_{L^2}^2 \\ &\leq c \|\Lambda^{\frac{1}{2}} \eta\|_{\sigma(t)}^2 \|\Lambda \eta\|_{\sigma(t)} \end{aligned}$$

where we used the Sobolev embedding $H^{\frac{1}{2}}(\mathbb{T}^2) \hookrightarrow L^4(\mathbb{T}^2)$ and the Plancherel theorem in the computation above. \square

We begin now the proof of theorem 4:

Proof of Theorem 4. We first take inner product of (3) with $\eta(t)$ in Gevrey class G ,

$$(12) \quad \langle \eta_t, \eta \rangle_{\sigma(t)} + \langle \eta \eta_x, \eta \rangle_{\sigma(t)} + \langle (\beta - 1) \eta_{xx} - \eta_{yy} - \delta \Lambda^3(\eta) + \epsilon \Delta^2 \eta, \eta \rangle_{\sigma(t)} = 0$$

Note that

$$\frac{1}{2} \frac{d}{dt} \langle \eta, \eta \rangle_{\sigma(t)} = \sigma'(t) \Lambda \langle \eta, \eta \rangle_{\sigma(t)} + \langle \eta_t, \eta \rangle_{\sigma(t)}$$

then we have

$$\langle \eta_t, \eta \rangle_{\sigma(t)} = \frac{1}{2} \frac{d}{dt} \langle \eta, \eta \rangle_{\sigma(t)} - \sigma'(t) \Lambda \langle \eta, \eta \rangle_{\sigma(t)}.$$

Substituting this into (12),

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \langle \eta, \eta \rangle_{\sigma(t)} &= \sigma'(t) \langle \Lambda \eta, \eta \rangle_{\sigma(t)} - \langle (\beta - 1) \eta_{xx} - \eta_{yy} - \delta \Lambda^3(\eta) + \epsilon \Delta^2 \eta, \eta \rangle_{\sigma(t)} - \langle \eta \eta_x, \eta \rangle_{\sigma(t)} \\ &\leq \sigma'(t) \|\Lambda^{\frac{1}{2}} \eta\|_{\sigma(t)}^2 + \beta \|\Lambda \eta\|_{\sigma(t)}^2 + \delta \|\Lambda^{\frac{3}{2}} \eta\|_{\sigma(t)}^2 - \epsilon \|\Lambda^2 \eta\|_{\sigma(t)}^2 + c \|\Lambda \eta\|_{\sigma(t)} \|\Lambda^{\frac{1}{2}} \eta\|_{\sigma(t)}^2 \\ &\leq \sigma'(t) \|\Lambda^2 \eta\|_{\sigma(t)}^{\frac{1}{2}} \|\eta\|_{\sigma(t)}^{\frac{3}{2}} + \beta \|\Lambda^2 \eta\|_{\sigma(t)} \|\eta\|_{\sigma(t)} + \delta \|\Lambda^2 \eta\|_{\sigma(t)}^{\frac{3}{2}} \|\eta\|_{\sigma(t)}^{\frac{1}{2}} \\ &\quad - \epsilon \|\Lambda^2 \eta\|_{\sigma(t)}^2 + c \|\Lambda^2 \eta\|_{\sigma(t)} \|\eta\|_{\sigma(t)}^2. \end{aligned}$$

By the Young inequality, we have

$$(13) \quad \frac{d}{dt} \|\eta\|_{\sigma(t)}^2 \leq \left(D_1 (\sigma'(t))^{\frac{4}{3}} + D_2 + D_3 \right) \|\eta\|_{\sigma(t)}^2 + D_4 \|\eta\|_{\sigma(t)}^4$$

where $D_1 = (\frac{3}{2\epsilon})^{1/3}$, $D_2 = \frac{9}{4\epsilon} \beta^2$, $D_3 = (\frac{27}{4\epsilon})^3 \delta^4$, $D_4 = \frac{9}{4\epsilon} c^2$.

By the definition of $\sigma(t) = \min\{\tanh(t), \tanh(\frac{T_0}{2})\}$, we have

$$\sigma'(t) \leq 1.$$

Inserting this into (13), we obtain

$$\frac{d}{dt} \left(1 + \|\eta\|_{\sigma(t)}^2\right) \leq K \left(1 + \|\eta\|_{\sigma(t)}^2\right)^2$$

with

$$K = D_1 + D_2 + D_3 + D_4.$$

Define $y_1(t) = 1 + \|\eta\|_{\sigma(t)}^2$, it turns out to deal with the following ordinary differential inequality

$$\begin{cases} y_1'(t) \leq K y_1(t)^2 \\ y_1(0) = 1 + \|\eta_0\|_{L^2}^2. \end{cases}$$

After solving this ODI, we obtain

$$\|\eta(t)\|_{\sigma(t)}^2 \leq 1 + 2\|\eta_0\|_{L^2}^2 \text{ for } t \in [0, T_0],$$

where

$$T_0 = \frac{1}{2K(1 + \|\eta_0\|_{L^2}^2)}.$$

Moreover, according to theorem 1 and theorem 2, we know that the solution is unique, global in time and stays in a ball of Radius $R_{\epsilon, \delta}$ once it has entered it, that is to say, $\|\eta(t)\|_{L^2}$ remains bounded for all time,

$$\limsup_{t \rightarrow \infty} \|\eta(t)\|_{L^2(\mathbb{T}^2)} \leq R_{\epsilon, \delta}.$$

So up to now, we already prove local analyticity of $\eta(t)$

$$(14) \quad \|\eta(t)\|_{\sigma(t)}^2 \leq 1 + 2C_{\epsilon, \delta, \eta_0}^2 \text{ for } t \in [0, T_0].$$

In order to obtain global analyticity, we follow the previous idea and repeat the argument above starting at $\frac{T_0}{2}$. We consider time $t \in [\frac{T_0}{2}, \frac{3T_0}{2}]$, and let $y_2(t) = 1 + \|\eta(t)\|_{\sigma(t - \frac{T_0}{2})}^2$, so $y_2(\frac{T_0}{2}) = 1 + \|\eta(\frac{T_0}{2})\|_{L^2}^2$. Thus, solving the following ordinary differential inequality

$$\begin{cases} y_2'(t) \leq K y_2(t)^2 \\ y_2\left(\frac{T_0}{2}\right) = 1 + \left\| \eta\left(\frac{T_0}{2}\right) \right\|_{L^2}^2, \end{cases}$$

we have that,

$$\|\eta(t)\|_{\sigma(t - \frac{T_0}{2})}^2 \leq 1 + 2 \left\| \eta\left(\frac{T_0}{2}\right) \right\|_{L^2}^2 \leq 1 + 2C_{\epsilon, \delta, \eta_0}^2,$$

for time $t \in [\frac{T_0}{2}, \frac{3T_0}{2}]$.

By the definition of $\sigma(t)$ and observe that $\tanh(t)$ is strictly increasing, we know that $\sigma(t)$ remains being a constant after time $\frac{T_0}{2}$, and this constant is $\sigma(\frac{T_0}{2}) = \tanh(\frac{T_0}{2})$, so we choose $t = T_0$ in the inequality above, then

$$\|\eta(t)\|_{\sigma(\frac{T_0}{2})}^2 \leq 1 + 2C_{\epsilon, \delta, \eta_0}^2 \text{ for } t \in [T_0, \frac{3T_0}{2}].$$

We mimic this argument by adding $\frac{T_0}{2}$ each time, so we obtain that $\eta(t)$ is analytic in the time intervals $[\frac{3T_0}{2}, 2T_0], [2T_0, \frac{5T_0}{2}] \dots$ Recalling local analyticity (14), we finally obtain

$$\|\eta(t)\|_{\sigma(t)}^2 \leq 1 + 2C_{\epsilon, \delta, \eta_0}^2 \text{ for any } t \geq 0.$$

This completes our proof. \square

Remark 2. *It is of interest to point out that the global analyticity of the solution which we show here is better than the result in [12], where the argument given by the authors can be extended to prove the global analyticity of solutions of the Kuramoto-Sivashinsky equation, outside a set of time instants with zero measure.*

6. EXISTENCE OF ATTRACTOR AND THE NUMBER OF PEAKS

In this section, we are interested in the existence of the attractor and its properties. By applying the Theorem 1.1 in [26], we can prove that the initial value problem (3) possesses a compact global attractor in $H^2(\mathbb{T}^2)$. First, we denote by $S(t)$ the solution operator, where $S(t)\eta_0 = \eta(x, y, t)$.

Definition 1. *The solution operator $S(t)\eta_0 = \eta(x, y, t)$ defines a compact semiflow in H^2 , if for every initial data $\eta_0 \in H^2$, the following four statements holds:*

- (i) $S(0)\eta_0 = \eta_0$;
- (ii) $S(t+s)\eta_0 = S(t)S(s)\eta_0$, for all t, s ;
- (iii) For every $t > 0$,

$$S(t)(\cdot) : H^2 \rightarrow H^2$$

is continuous;

- (iv) *There exists $T^* > 0$ such that $S(T^*)$ is a compact operator, i.e. for every bounded set $B \subset H^2$, $S(T^*)B \subset H^2$ is a compact set.*

Definition 2. *An attractor $\mathcal{A} \subset H^2$ is a set that satisfies the following properties:*

- (i) *\mathcal{A} is an invariant set, i.e., $S(t)\mathcal{A} = \mathcal{A}, \forall t \geq 0$;*
- (ii) *there is an open neighborhood $\mathcal{B} \subset H^2$ of \mathcal{A} such that, for every η_0 in \mathcal{B} , $S(t)\eta_0$ converges to \mathcal{A} as $t \rightarrow \infty$:*

$$\text{dist}(S(t)\eta_0, \mathcal{A}) \rightarrow 0 \text{ as } t \rightarrow \infty.$$

We shall need the following lemma:

Lemma 5. *Let $\eta_0 \in H^2(\mathbb{T}^2)$ be the initial data. Then $S(\cdot)\eta_0 \in \mathcal{C}([0, T]; H^2(\mathbb{T}^2))$ defines a compact semiflow in $H^2(\mathbb{T}^2)$.*

Proof. In order to show that $S(t)\eta_0 = \eta(\cdot, \cdot, t)$ defines a compact semiflow, we must verify (i)-(iv) in definition 1. If we fix a t_0 , the continuity of $S(t_0)(\cdot)$ from H^2 to H^2 is strightforward by energy estimates. Then, as in Theorem 3, we have the existence of an absorbing set in H^2 -norm, so there exists T^* such that

$$\|\eta(t)\|_{H^2} \leq R'_{\epsilon, \delta}(\eta_0, \delta, \epsilon), \forall t \geq T^* .$$

Since (i) and (ii) are obvious, we conclude our proof by invoking the analyticity of solutions. \square

Theorem 5. *The system (3) has a maximal, connected, compact attractor in the space $H^2(\mathbb{T}^2)$.*

Proof. By applying Theorem 1.1 in [26] and Lemma 5. \square

The rest of this section is devoted to studying a particular feature of the chaotic behavior of (3), namely, the number of spatial oscillations. We shall need the lemma proved by Grujić in [13], which gives us an effective method to study the number of peaks (see also [12, 2]). We cannot use directly the method in [13], mainly because Lemma 8.1 in [13] is quite suitable to bound the number of peaks in one space dimension, but not appropriate for our two-dimensional model.

We first let $\eta(x, y) = f_y(x)$, where $f_y(x)$ depends on both x and y . Then our original problem (3) deduces to

$$(15) \quad \partial_t f_y + f_y \partial_x f_y + (\beta - 1) \partial_x^2 f_y + F_y(x) = 0,$$

where $F_y(x) = -\eta_{yy}(x, y) - \delta \Lambda^3(\eta)(x, y) + \epsilon \Delta^2 \eta(x, y)$ can be seen as a forcing term.

In the previous section, we have shown that $\eta(x, y, t)$ is analytic in a growing complex strip

$$\mathbb{S}_{\sigma(t)} = \{(x, y) + i(\tilde{x}, \tilde{y}) : (x, y) \in \mathbb{T}^2, |(\tilde{x}, \tilde{y})| < \sigma(t)\}.$$

Recalling that $\eta(x, y, t) = f_y(x, t)$, then $f_y(x, t)$ is analytic in

$$\mathbb{S}'_{\sigma(t)} = \{x + i\tilde{x} : x \in \mathbb{T}, |\tilde{x}| < \sigma(t)\}.$$

Now we use the following Lemma (from [13]):

Lemma 6. *Fix y in problem (15). Let $\sigma > 0$, and let $f_y(x)$ be analytic in the neighbourhood of $\{z = x + i\tilde{x} : |\Im z| \leq \sigma\}$ and be 2π -periodic in x -direction. Then, for any $\mu > 0$, $\mathbb{T} = I_\mu \cup R_\mu$, where I_μ is a union of at most $\lceil \frac{4\pi}{\sigma} \rceil$ intervals open in \mathbb{T} , and*

- $|\partial_x f_y(x)| \leq \mu$, for all $x \in I_\mu$;
- $\text{card}\{x \in R_\mu : \partial_x f_y(x) = 0\} \leq \frac{2}{\log 2} \frac{2\pi}{\sigma} \log \left(\frac{\max_{|\Im z| \leq \sigma} |\partial_x f_y(z)|}{\mu} \right)$.

With the help of the lemma above, we have our main result.

Theorem 6. *Let η be a solution of system (3) for initial data $\eta_0 \in H^2(\mathbb{T}^2)$ and define T_0 as in Theorem 4. Then, $\mathbb{T} = I \cup R$, where I is a union of at most $\lceil \frac{4\pi}{\tanh(\frac{T_0}{2})} \rceil$ open intervals in \mathbb{T} and the following estimates hold for $t \geq \frac{T_0}{2}$,*

$$|\partial_x \eta(x, y, t)| \leq 1, \text{ for all } x \in I, y \in \mathbb{T}$$

and

$$\text{card}\{x \in R : |\nabla \eta(x, y, t)| = 0\} \leq \frac{4\pi}{\log 2} \frac{\log C_{\epsilon, \delta, \eta_0}}{\tanh\left(\frac{T_0}{2}\right)}.$$

where $C_{\epsilon, \delta, \eta_0}$ is a constant depending on ϵ, δ, η_0 .

Proof. From the results of Theorem 4 and Theorem 5, we know that the system has an attractor and the solution η is analytic at least in the strip of width $\sigma(t)$.

Now, we can apply Lemma 6 with $\mu = 1$ and bound

$$\begin{aligned} \text{card}\{x \in R : |\nabla \eta(x, y, t)| = 0\} &\leq \text{card}\{x \in R : \partial_x \eta = 0\} \\ &\leq \frac{2}{\log 2} \frac{2\pi}{\sigma} \log \left(\frac{\max_{|\Im z| \leq \sigma} |\partial_x \eta(z, y, t)|}{\mu} \right) \\ &\leq \frac{4\pi}{\log 2} \frac{1}{\sigma} \log \left(C \|e^{\sigma(t)\Lambda} \eta\|_{L^2} \right). \end{aligned}$$

In the last inequality above, we used the following estimate

$$\begin{aligned}
\|\partial_x \eta(z, y, t)\|_{L^\infty(|\Im z| \leq \frac{\sigma}{2})} &= \|\partial_x \eta(x + i\tilde{x}, y, t)\|_{L^\infty(|\Im z| \leq \frac{\sigma}{2})} \\
&\leq \left\| \sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} |\xi_1| \widehat{\eta}(\xi_1, \xi_2, t) e^{i(x\xi_1 + y\xi_2)} e^{-\tilde{x}\xi_1} \right\|_{L^\infty(|\Im z| \leq \frac{\sigma}{2})} \\
&\leq \left\| \sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} |\xi| |\widehat{\eta}(\xi_1, \xi_2, t)| e^{|\tilde{x}||\xi|} \right\|_{L^\infty(|\Im z| \leq \frac{\sigma}{2})} \\
&\leq \sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} |\xi| |\widehat{\eta}(\xi_1, \xi_2, t)| e^{\frac{\sigma(t)}{2}|\xi|} \\
&\leq \sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} |\xi|^3 |\widehat{\eta}(\xi_1, \xi_2, t)| e^{\frac{\sigma(t)}{2}|\xi|} \frac{1}{|\xi|^2} \\
&\leq \sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} |\widehat{\eta}(\xi_1, \xi_2, t)| e^{\sigma(t)|\xi|} \frac{1}{|\xi|^2} \\
&\leq \|e^{\sigma(t)\Lambda} \eta\|_{L^2} \left(\sum_{(\xi_1, \xi_2) \in \mathbb{Z}^2} \frac{1}{|\xi|^4} \right)^{\frac{1}{2}} \\
&\leq C \|e^{\sigma(t)\Lambda} \eta\|_{L^2}
\end{aligned}$$

where we used the fact that

$$|\xi|^3 e^{\frac{\sigma(t)}{2}|\xi|} \leq C e^{\sigma(t)|\xi|}.$$

Since η has global analyticity

$$\|e^{\sigma(t)\Lambda} \eta(t)\|_{L^2} \leq 1 + 2C_{\epsilon, \delta, \eta_0}, \quad \forall t \geq 0,$$

we can conclude that for $t \geq \frac{T_0}{2}$

$$\text{card}\{x \in R : |\nabla \eta(x, y, t)| = 0\} \leq \frac{4\pi}{\log 2} \frac{\log C_{\epsilon, \delta, \eta_0}}{\tanh\left(\frac{T_0}{2}\right)}.$$

where $C_{R_{\epsilon, \delta, \eta_0}}$ depends on $R_{\epsilon, \delta}$. □

Theorem 6 gives us a bound of the number of wild spatial oscillations of the solution, then the following corollary is a direct result of it.

Corollary 1. *Let η be a solution corresponding to the initial data $\eta_0 \in H^2(\mathbb{T}^2)$, then for $t \geq \frac{T_0}{2}$, the number of peaks for η can be bounded as*

$$\text{card}\{\text{peaks for } \eta\} \leq \frac{4\pi}{\log 2} \frac{\log C_{\epsilon, \delta, \eta_0}}{\tanh\left(\frac{T_0}{2}\right)}.$$

where $C_{\epsilon, \delta, \eta_0}$ depends on ϵ, δ, η_0 and T_0 is defined as before.

ACKNOWLEDGMENT

R.GB was supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). J.H has been partially funded by the ANR project Dyficoliti ANR-13-BS01-0003-01.

REFERENCES

- [1] J. C. Bronski and T. N. Gambill. Uncertainty estimates and L^2 bounds for the Kuramoto-Sivashinsky equation. *Nonlinearity*, 19(9):2023, 2006.
- [2] J. Burczak and R. Granero-Belinchón. On a generalized doubly parabolic Keller–Segel system in one spatial dimension. *Mathematical Models and Methods in Applied Sciences*, 26(01):111–160, 2016.
- [3] B. I. Cohen, J. Krommes, W. Tang, and M. Rosenbluth. Non-linear saturation of the dissipative trapped-ion mode by mode coupling. *Nuclear fusion*, 16(6):971, 1976.
- [4] P. Collet, J.-P. Eckmann, H. Epstein, and J. Stubbe. Analyticity for the Kuramoto-Sivashinsky equation. *Physica D: Nonlinear Phenomena*, 67(4):321–326, 1993.
- [5] P. Collet, J.-P. Eckmann, H. Epstein, and J. Stubbe. A global attracting set for the Kuramoto-Sivashinsky equation. *Communications in mathematical physics*, 152(1):203–214, 1993.
- [6] A. V. Coward and P. Hall. On the nonlinear interfacial instability of rotating core-annular flow. *Theoretical and Computational Fluid Dynamics*, 5(6):269–289, 1993.
- [7] C. Foias and R. Temam. Gevrey class regularity for the solutions of the Navier-Stokes equations. *Journal of Functional Analysis*, 87(2):359–369, 1989.
- [8] A. Frenkel and K. Indreshkumar. Wavy film flows down an inclined plane: perturbation theory and general evolution equation for the film thickness. *Physical Review E*, 60(4):4143, 1999.
- [9] L. Giacomelli and F. Otto. New bounds for the Kuramoto-Sivashinsky equation. *Communications on pure and applied mathematics*, 58(3):297–318, 2005.
- [10] A. Gonzalez and A. Castellanos. Nonlinear electrohydrodynamic waves on films falling down an inclined plane. *Physical Review E*, 53(4):3573, 1996.
- [11] J. Goodman. Stability of the Kuramoto-Sivashinsky and related systems. *Communications on Pure and Applied Mathematics*, 47(3):293–306, 1994.
- [12] R. Granero-Belinchón and J. K. Hunter. On a nonlocal analog of the Kuramoto-Sivashinsky equation. *Nonlinearity*, 28(4):1103, 2015.
- [13] Z. Grujić. Spatial analyticity on the global attractor for the Kuramoto-Sivashinsky equation. *Journal of Dynamics and Differential Equations*, 12(1):217–228, 2000.
- [14] J. S. Il’Yashenko. Global analysis of the phase portrait for the Kuramoto-Sivashinsky equation. *Journal of Dynamics and Differential Equations*, 4(4):585–615, 1992.
- [15] X. Ioakim and Y.-S. Smyrlis. Analyticity for Kuramoto-Sivashinsky type equations and related systems. *Procedia IUTAM*, 11:69–80, 2014.
- [16] M. James and M. Wilczek. Vortex dynamics and lagrangian statistics in a model for active turbulence. *The European Physical Journal E*, 41(2):21, 2018.
- [17] Y. Kuramoto and T. Tsuzuki. Persistent propagation of concentration waves in dissipative media far from thermal equilibrium. *Progress of theoretical physics*, 55(2):356–369, 1976.
- [18] R. E. LaQuey, S. Mahajan, P. Rutherford, and W. Tang. Nonlinear saturation of the trapped-ion mode. *Physical Review Letters*, 34(7):391, 1975.
- [19] Y. Lee and H. Chen. Nonlinear dynamical models of plasma turbulence. *Physica Scripta*, 1982(T2A):41, 1982.
- [20] D. Michelson and G. Sivashinsky. Nonlinear analysis of hydrodynamic instability in laminar flames—II. numerical experiments. *Acta Astronautica*, 4(11-12):1207–1221, 1977.
- [21] B. Nicolaenko, B. Scheurer, and R. Temam. Some global dynamical properties of the Kuramoto-Sivashinsky equations: nonlinear stability and attractors. *Physica D: Nonlinear Phenomena*, 16(2):155–183, 1985.
- [22] F. C. Pinto. Nonlinear stability and dynamical properties for a Kuramoto-Sivashinsky equation in space dimension two. *Discrete & Continuous Dynamical Systems-A*, 5(1):117–136, 1999.
- [23] F. C. Pinto. Analyticity and Gevrey class regularity for a Kuramoto-Sivashinsky equation in space dimension two. *Applied mathematics letters*, 14(2):253–260, 2001.
- [24] G. Sivashinsky. Nonlinear analysis of hydrodynamic instability in laminar flames—I. derivation of basic equations. *Acta astronautica*, 4(11-12):1177–1206, 1977.
- [25] G. Sivashinsky and D. Michelson. On irregular wavy flow of a liquid film down a vertical plane. *Progress of theoretical physics*, 63(6):2112–2114, 1980.
- [26] R. Temam. *Infinite-dimensional dynamical systems in mechanics and physics*, volume 68. Applied Mathematical Sciences. Springer-Verlag, 1997.
- [27] R. Tomlin, D. Papageorgiou, and G. Pavliotis. Three-dimensional wave evolution on electrified falling films. *Journal of Fluid Mechanics*, 822:54–79, 2017.

- [28] J. Topper and T. Kawahara. Approximate equations for long nonlinear waves on a viscous fluid. *Journal of the Physical society of Japan*, 44(2):663–666, 1978.
- [29] D. Tseluiko and D. Papageorgiou. Dynamics of an electrostatically modified Kuramoto-Sivashinsky–Korteweg–de Vries equation arising in falling film flows. *Physical Review E*, 82(1):016322, 2010.
- [30] D. Tseluiko and D. T. Papageorgiou. A global attracting set for nonlocal Kuramoto-Sivashinsky equations arising in interfacial electrohydrodynamics. *European Journal of Applied Mathematics*, 17(6):677–703, 2006.
- [31] D. Tseluiko and D. T. Papageorgiou. Wave evolution on electrified falling films. *Journal of Fluid Mechanics*, 556:361–386, 2006.

J. He: Université de Lyon, Université Lyon 1 – CNRS UMR 5208 Institut Camille Jordan – 43 bd. du 11 Novembre 1918 – Villeurbanne Cedex F-69622, France.

Email: jiao.he@math.univ-lyon1.fr

R. Granero-Belinchón: Departamento de Matemáticas, Estadística y Computación, Universidad de Cantabria. Avda. Los Castros s/n, Santander, Spain.

Email: rafael.granero@unican.es