



Genetic diversity and population structure of *Saccharomyces cerevisiae* strains isolated from traditional alcoholic beverages of Côte d'Ivoire

Youan Charles Tra Bi, Tiemele Laurent Simon Amoikon, Clémentine A. Kouakou, Jacques Noemie, Marine Lucas, Cecile Grondin, Jean-Luc Legras, Florent K. N'Guessan, Theodore N. Djeni, Marcellin K. Djè, et al.

► To cite this version:

Youan Charles Tra Bi, Tiemele Laurent Simon Amoikon, Clémentine A. Kouakou, Jacques Noemie, Marine Lucas, et al.. Genetic diversity and population structure of *Saccharomyces cerevisiae* strains isolated from traditional alcoholic beverages of Côte d'Ivoire. *International Journal of Food Microbiology*, 2019, 297, pp.1-10. 10.1016/j.ijfoodmicro.2019.03.001 . hal-02168620

HAL Id: hal-02168620

<https://hal.science/hal-02168620>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Genetic diversity and population structure of *Saccharomyces cerevisiae* strains isolated from traditional alcoholic beverages of Côte d'Ivoire

Charles Y. Tra Bi^{a,b,†*}, Tiémélé L.S. Amoikon^{a,b,†}, Clémentine A. Kouakou^a, Jacques Noémie^{b,‡}, Marine Lucas^{b,¶}, Cécile Grondin^b, Jean-Luc Legras^c, Florent K. N'guessan^a, Théodore N. Djéni^a, Marcellin K. Djè^a, Serge Casaregola^b

^a Laboratoire de Biotechnologie et Microbiologie des Aliments, Unité de Formation et de Recherche en Sciences et Technologie des Aliments (UFR-STA), Université Nanguigou, 02 BP 801 Abidjan 02, Côte d'Ivoire

^b Micalis Institute, INRA, AgroParisTech, CIRM-Levures, Université Paris-Saclay, 78350 Jouy-en-Josas, France

^c SPO, Université Montpellier INRA, Montpellier SupAgro, , F-34060, Montpellier, France

[#] Present address: Bioger, INRA/AgroParisTech, 78850, Thiverval-Grignon, France

[¶] These authors have contributed equally to this work.

*Corresponding author, phone: +225 09 96 97 48; +225 06 18 80 99, E-mail: charlestrabiyouan@yahoo.fr

Abstract

In order to assess the genetic diversity and population structure of indigenous *S. cerevisiae* from Côte d'Ivoire, a total of 170 strains were isolated from four traditional alcoholic beverages through nine regions. Microsatellite analysis performed at 12 loci revealed that strains of palm oil and raffia wine were genetically related, unlike those of tchapalo and ron wine which formed two distinct and homogeneous clusters. Furthermore, *S. cerevisiae* isolates of traditional beverages from Côte d'Ivoire appear to be differentiated according to the ecological niche. Indeed, a much higher heterozygosity was observed for isolates from ron wine and tchapalo beer, whereas isolates from palm oil wine and raffia wine were clearly inbred. In comparison with the European, North American, Asian and others West African populations, Ivorian population was well defined, although most of these strains were admixed. Among these strains, only isolates from raffia wine appeared to have alleles in common to all populations.

Keywords: Microsatellite, population, *Saccharomyces cerevisiae*, diversity, traditional beverage, genotyping.

1 Introduction

Saccharomyces cerevisiae is one of the main actor involved in various biotechnology processes. These last decades, several studies have been focused on genetic diversity and biogeographical distribution of diverse *S. cerevisiae* strains. Molecular techniques used for this purpose among many others are pulsed field gel electrophoresis (PFGE) (Vezinhet *et al.*, 1990), restriction analysis of the mitochondrial DNA (mtDNA-RFLP) (Aigle *et al.*, 1984; Querol *et al.*, 1992), amplified fragment length polymorphism (AFLP) (Azumi and Goto-Yamamoto., 2001), amplification of interdelta regions by PCR (Ness *et al.*, 1993; Legras and Karst 2003) and microsatellite markers (Techera *et al.*, 2001; Legras *et al.*, 2005). Microsatellite analyses have offered significant advances in understanding distribution and dynamics of *S. cerevisiae* population from different terroirs. Thus, Viel *et al.* (2017) have detected specificities of *S. cerevisiae* at vineyard scale from the North-East of Italy, suggesting a geographic differentiation according to the winemaking region with a widespread dissemination of *S. cerevisiae* industrial strains that was very high in the areas where the native strains abundance was low. Likewise, microsatellite analyses on *S. cerevisiae* populations isolated from African traditional fermented foods allowed to highlight the presence of an African phylogenetic branch separate from European (bread, wine, beer, cheese) and Asian strains (sake) (Legras *et al.*, 2007; Ezeronye and Legras, 2009; Tapsoba *et al.*, 2015), which was confirmed with recent genomic data (Liti *et al.*, 2009; Peter *et al.*, 2018). This suggests that African traditional fermented foods could be a reservoir of genetically different strains of *S. cerevisiae* to be explored for applications such as food, bioethanol production, biopreservation, biocontrol, etc. Most of *S. cerevisiae* commonly used in biotechnology have been isolated from natural niches and have undergone various domestication events that have resulted in the selection of strains adapted to human activities (technological strains, etc.). Although several studies have been performed on *S. cerevisiae* isolated from some natural niches (Naumov *et al.*, 1998; Sniegowski *et al.*, 2002; Lorca *et al.*,

2018), it seems important to assess the structure, the genetic diversity and the distribution of native *S. cerevisiae* obtained from African, and Ivoirian niches particularly, which has been very little studied so far. The characterization of these unexplored ecological niches may provide new genotypes with beneficial properties for biotechnological applications. The aim of this work was firstly to assess the genetic diversity of *S. cerevisiae* strains present in traditional beverages from Côte d'Ivoire compared to other African beverages and then characterize the population structure of the isolates of this species. For this purpose, this study focused on the yeast populations isolated from four fermented beverages obtained with different vegetal material. These beverages were tchapalo beer produced from sorghum (*Sorghum bicolor*) and palm oil, raffia and ron wines respectively obtained from *Elaeis guineensis*, *Raphia hookeri* and *Borassus aethiopum* sap fermentation.

2 Material and Methods

2.1 Sample collection and yeast isolation

Samples of palm wine (raffia, palm oil and ron wines) were collected during ten sampling campaigns in different regions of Côte d'Ivoire. Sampling areas for raffia wine were located at Grand-Lahou, Alepé, Adzopé and Abengourou; those for palm oil wine were located at Bingerville, Bonoua, Grand-Lahou, Alépé and Attinguié while those for ron wine were located at Toumodi. Sorghum beer or tchapalo samples were collected during four sampling campaigns at Abidjan and Bingerville. These sampling areas were located in south-eastern and middle of Côte d'Ivoire as indicated by the sample map (Figure 1). Distance between the sampling areas and the city of Abidjan as well as their geographical coordinates are shown in the Table 1. Local producers in 26 different sites were contacted and sampling was carried out from September 2014 to January 2015 and from August 2016 to September 2016. Sampling was twice a day (morning and evening) done at weekly intervals, to collect samples of 9 to 15hr fermentation duration. Samples were collected into pre-sterilized 250 ml Plexiglas containers, and immediately immersed in an isothermal box containing dry ice, and brought to the laboratory within 3 h.

Total titratable acidity (TTA), pH, and total soluble sugars (TSS) for each sample were determined as previously described (Tra Bi *et al.*, 2016). For yeast isolation, each beverage sample (1 mL) was directly diluted in tenfold series in buffered peptone water (BIO-RAD, France) and aliquots (0.1 mL) were plated in duplicate on Yeast Extract Peptone Dextrose (YPD) agar. After incubation at 30°C for 3-5 days, ten yeast colonies were randomly selected from each sample and purified by streaking two times on YPD agar. *Saccharomyces* yeasts

were presumptively isolated by cultivation of purified colonies on lysine agar medium (Sigma-Aldrich, France). Thus, a total of 262 isolates were selected for molecular identification and maintained at -80°C in YPD broth containing 20% (v/v) glycerol (Merck, France).

Variance analysis (ANOVA) and Tukey HSD tests (Honestly Significant Difference) with R statistical software v3.4.1 (R Core Team, 2017) were used to compare the physicochemical and microbiological characteristics of each beverage taken from the different sampling areas. Differences were considered significant for values of $P < 0.05$.

2.2 Molecular identification of *Saccharomyces* species

The species level identity of *Saccharomyces* strains was checked initially by non-transcribed spacer PCR-Restriction Fragment Length Polymorphism (NTS2 PCR-RFLP) analysis using *AluI* and *BanI* enzymes in comparison with reference strains. From strains showing *S. cerevisiae* NTS2 PCR-RFLP profile (Nguyen and Gaillardin, 1997), thirty representative strains were selected for D1/D2 region sequencing (Kurtzman and Robnett, 1998). The D1/D2 region of 26S rRNA gene was amplified with the primer pair NL-1 (5'-GCATATCAATAAGCGGAGGAAAAG-3') and NL-4 (5'-GGTCCGTGTTTCAAGACGG-3') (Kurtzman and Robnett, 1998). Reactions were performed in an automatic thermal cycler (Gene Amp PCR System 2700, Thermo Fisher Scientific) under the following conditions: initial denaturation at 94°C for 4 min followed by 30 cycles of denaturation at 94°C for 30 s, primer annealing at 54°C for 40 s and DNA extension at 72°C for 1 min 30 s. A final extension was completed at 72°C for 7 min. Amplicons were visualized on 0.8% agarose gel. Finally, 20 µL of PCR products were sequenced on both strands with the primers used for the PCR amplification (Eurofins, Germany). Assembly of both strands sequences was performed using the phred/phrap/consed suite (Gordon *et al.*, 1998). Identification queries were fulfilled by a BLAST search of the National Center for Biotechnology Information database (NCBI, Bethesda, USA) and the YeastIP databases (<http://genome.jouy.inra.fr/yeastip/>).

2.3 Microsatellite analysis

The genomic DNA extraction of isolates was performed following the procedure described by Hoffman and Winston (1987), based on a mechanical grinding of cells in presence of detergent, followed by several steps of protein and RNA removal from a 24 hour yeast culture in YPD broth. Microsatellite loci analysis was performed according to Legras *et al.* (2007). Amplification was performed at 12 loci combined in two multiplex of 6 loci (Table S1 in the

supplemental material), and the size of fluorescent amplicons was measured on an ABI3100 (Applied Biosystems) capillary electrophoresis device, using Gene Scan HD400Rox size standard (Applied Biosystems). These genotypes were compared to data obtained in previous work on strains from other origins.

2.4 Population analysis

The evaluation of relationships between *S. cerevisiae* strains was carried out from the genetic relative dissimilarity distance which reflects the number of alleles differing between two individuals, and calculated with the library POPPR (v2.5.0) under R statistical software v3.4.1 (R Core Team, 2017). Trees were obtained from distance matrices with R library ape v5.0, and drawn using R library ggtree v1.10.5 (Yu *et al.*, 2017). In order to assess genetic structure in *S. cerevisiae* populations, two different approaches were used. The first one was Bayesian clustering analysis implemented with InStruct 2.3.4 software (Gao *et al.*, 2007) to identify the most likely number of genetic clusters or ‘populations’ (denoted by K) and assign individuals to their most likely cluster. In order to deduce ancestry on that set of strains, 8 runs were performed with Instruct for values from K = 2 to 17, and 8 additional runs for values from K = 14 to 16. The most likely K value was identified as the value giving the lowest mean value of Deviance Information Criterion (DIC) of the different runs. The results of the 16 runs of the best value of K were combined with CLUMPAK (Kopelman *et al.*, 2015) according to the LargeK Greedy method. The second was a multivariate method based on discriminant analysis in principal components (DAPC) (Jombart *et al.*, 2010) and implemented with the R Adegenet package v2.1.0 (Jombart, 2008) in R v3.4.1 (R Core Team, 2017). DAPC identified clusters of genetically related organisms by partitioning genetic variability into clusters that maximize between-group and minimize within-group differentiation. This approach was an ideal clustering algorithm for datasets that does not followed Hardy–Weinberg equilibrium. Analysis was performed taking into account alpha score function to choose the optimal number of principal components for the analysis of dataset (López-Uribe *et al.*, 2016) due to the changes of the individual membership probability according to the number of PCA axes retained. Population analyses were performed using the hierfstat v0.04-22 R package (Goudet, 2005) for Nei (1973) estimates of effect of subpopulations compared to the total population (Fst). Population differentiation according to sampling area was tested with OBSTRUCT (Gayevskiy *et al.*, 2014). The inbreeding coefficient of an individual relative to the subpopulation (Fis) was estimated with hierfstat. Average selfing rates were estimated as $1 - o$, where o the outcrossing rate is $o = (1 - F_{is}) / (1 + F_{is})$

3 Results

3.1 Characteristics of beverages

The global characteristics of the four different beverages from which strains were isolated in this study were presented in Table 2. It appeared from this investigation that these four beverages had in general a similar acidic pH and a Total Titratable Acidity (TTA) varying between 0.47 and 1.35. The soluble sugars content varied from one beverage to another with the highest content in tchapalo. The yeast load observed in the Ivorian beverages sampled was generally of the same order. Average values were 1.1×10^8 CFU/mL in ron wine, 1.7×10^7 CFU/mL in tchapalo beer, 1.4×10^7 CFU/mL in palm oil wine and 1.3×10^7 CFU/mL in raffia wine (Table 2).

3.2 Assessment of *Saccharomyces cerevisiae* diversity

A total of 170 isolates from Côte d'Ivoire traditional beverages (Table S2 in the supplemental material) were identified as belonging to the species *S. cerevisiae* by using molecular methods. Indeed, the RFLP of NTS2 PCR products showed 2 bands for *AluI* (1500 and 200 pb) and 3 bands for *BanI* (660, 450 and 220 pb) typical to *S. cerevisiae* strains (Nguyen and Gaillardin, 1997). Then, the sequencing of the D1/D2 domains of the 26S rDNA confirmed the identity of some strains tested as *S. cerevisiae* with a little sequence divergence of 1 to 4 bp for 16 strains and 100% identity for 14 strains (Table S3 in the supplemental material). Sequence divergence of 3 to 4 bp was already observed from *S. cerevisiae* strains isolated from tchapalo (N'Guessan *et al.*, 2011). The 170 isolates derived from four ecological niches sampled at 26 different sites from 9 regions (south-eastern and middle) of Côte d'Ivoire. Among them, 58 have been isolated from ron wine, 64 from raffia wine, 29 from palm oil wine and 19 from tchapalo beer. These isolates were subjected to a microsatellite analysis which allowed to generate genotypic data. In order to compare the populations between the different type of beverages, Fst statistics between isolates from tchapalo beer, palm oil, raffia and ron wines were calculated with these data. Population comparisons indicated significant differences between all populations ($P < 0.01$). Fst data showed weak values ranging from 0-0.107 (Table 3a), indicating a little genetic differentiation between the significantly different beverage isolates. The smallest differentiation was observed between palm oil wine and raffia wine strains, and in contrast, tchapalo was the most differentiated population from other populations. Ron wine population presented a moderate differentiation from tchapalo and raffia wine.

Impact of geographical origin on *S. cerevisiae* diversity was studied for isolates from palm oil wine, raffia wine and tchapalo beer. Ron wine with only one sampling area was not considered for this study. The Nei pairwise F_{st} distances matrix for the two sampling areas of tchapalo beer (Abidjan and Bingerville) showed low F_{st} values and non-significant differentiation. In contrast the four sampling areas of raffia wine (Grand-Lahou, Alépé, Adzopé and Abengourou) and most of the five sampling areas of palm oil wine showed low F_{st} values and significant differentiation between sites (Table 3). Furthermore, pairwise F_{st} statistic values for all sampling areas demonstrated no correlation to geographical distance, suggesting that diversity of *S. cerevisiae* was not related to geographical origin. Although isolates of palm oil and raffia wines were genetically related, *S. cerevisiae* isolates from Côte d'Ivoire traditional beverages appeared to be differentiated according to the ecological niche. The F_{is} values evaluated for each population were 0.603 and 0.544, respectively, for individuals from palm oil wine and raffia wine, 0.234 and 0.138 for individuals from ron wine and tchapalo beer that contained strains not corresponding to one specific fermentation (Table 4). Such high values of F_{is} which indicate a low outcrossing rate (and a high mean inbreeding rate) as well as the high observed homozygosity of strains isolated from palm oil wine and raffia wine (26 and 25 % of observed sites are heterozygotes), suggest that palm oil wine and raffia wine are inbred. In contrast, the low F_{is} values which indicate a high outcrossing rate (and a low mean selfing rate) as well as the high observed heterozygosity (52 and 76 % of observed sites are heterozygotes) for isolates from ron wine and tchapalo beer suggest more outbred populations.

3.3 Genetic relationship among strains

The phylogenetic relationships among Ivorian strains of *S. cerevisiae* were highlighted by a phylogenetic tree (Figure 2a). This dendrogram showed that strains of tchapalo (in green) and ron wine (in blue) formed two distinct and homogeneous clusters, unlike those of palm oil (in brown) and raffia wines (in red). Strains of the latter two groups appeared mixed in the remaining clusters.

To further detect the specificities of these Ivorian *S. cerevisiae* populations in comparison to populations from other African countries, and in comparison to well characterized populations such as those from wine, sake, and bread, we built a tree including the 170 Ivorian strains and 214 strains from multiple African countries or other origins (Figure 2b) analysed in a previous work (Tapsoba *et al.*, 2015). This consensus neighbour-joining tree was obtained from a pairwise individual distance matrix calculated with relative dissimilarity distance. It showed

several clusters of different genetic origin closely related to geographic origin. Thus, there was a cluster formed from European strains: wine and bread strains. Five clusters of strains originated from African substrates, one of which consists of tchapalo and millet beer (different shades of green) and the four others of different types of palm wine. The last two clusters encompassed Asian and North American strains. Ivorian strains are mainly encountered as sub groups inside the main clusters. In contrast to sorghum beer strains, ron wine strains from Côte d'Ivoire are found in two main clusters, one close to Burkina Faso ron wine strains, and in a second different cluster. A similar situation can be observed for raffia wine strains: one group of strains is detected close to Nigerian raffia wine strains, whereas the second group is different.

3.4 Population structure

Population structure was inferred with the Bayesian clustering software InStruct which does not assume Hardy-Weinberg equilibrium, but instead assigns individuals proportional membership coefficients in different populations based on inbreeding rates, which is well suited for *S. cerevisiae* population analysis (Gao *et al.*, 2007; Martiniuk *et al.*, 2016). Deviance information criterion indicated the most likely ancestral population number to be $K = 15$.

Each population was labelled by the geographic location and/or substrate from which the strains were isolated (Figure 3). The 15 populations included 11 African beverage populations among them, five originated from Côte d'Ivoire, two from European wine and bread populations, one from the Asian sake population and one from the North American oak population. The InStruct inferences revealed that most strains were assigned to a single ancestry cluster although admixtures were observed. About 8 out of 18 clusters were composed of individuals with one ancestral cluster membership coefficients higher than 0.80. The 10 other clusters (bread, sake from Japan, oak from North-America, laboratory, Bertram palm from Malaysia, various origins from Africa, *Hyphaene thebaica* from Djibouti, raffia, palm oil and ron wines from Côte d'Ivoire) appeared as admixtures. A small number of individuals within ron (*Borassus aethiopum*) wine strains had alleles common with those of ron strains from Burkina Faso, raffia and palm oil wines from Côte d'Ivoire. Strains of palm oil wine have some alleles in common with strains from Djibouti, Ghana (bili), Burkina Faso (dolo), other strains from Côte d'Ivoire (tchapalo and millet beer) and European wine strains. Strains of raffia wine appeared to have alleles in common with all populations, but in small proportion (Table S2 in the supplemental material).

In order to improve the detection of population structure of this study, a discriminant principal component analysis (Jombart *et al.*, 2010) based on 80 % of the global variation obtained from the 50 first PCA axis was conducted (Figure 4). The first two axes enabled differentiation of sorghum and millet beers strains (different shades of pink in the lower part of Figure 4a) from Burkina Faso ron wines strains (shade of brown), Côte d'Ivoire (coral) and miscellaneous African strains (magenta) in the upper part of Figure 4a. Genetic differentiation between African strains following some of their ecological niche was observed in the second axis: strains from the sap of *Arecaceae* family plants of the (palm oil palm, raffia palm, *Borassus* palm...) in the upper quadrant of Figure 4a were distinguished from those *Poaceae* family (millet and sorghum) in the lower part. The cluster strains corresponding to Ivorian tchapalo beer isolates were colocalized with Burkina Faso dolo beer isolates and Côte d'Ivoire millet beer strains. As for axes 3 and 4 (Figure 4b), they highlighted the differentiation between North-America populations and Ghana cocoa strains from that of the rest of the world, and the axis 4 differentiate Nigeria palm strains (and to a lower extent sake strains) from other strains. The two plots (Figure 4) showed that palm oil, raffia and ron wines from Côte d'Ivoire were also clustered but these clusters overlapped with clusters strains from Djibouti palm, Nigeria raffia wine, Burkina Faso ron wine and African miscellaneous origins. This clustering was also in agreement with the clustering obtained at $K = 15$ for InStruct. Obstruct tests were performed to evaluate if the population structure revealed in Bayesian ancestry profiles obtained from Instruct was correlated with the geography and confirmed of Nei pairwise F_{st} distance. For this analyse, only palm oil and raffia wines isolates, which had more than two sampling areas were considered. Obstruct analysis and canonical discriminant analysis clearly showed that the differences between isolates for most sampling areas of palm oil wine (Attinguié, Grand-Lahou, Alépé, Bingerville and Bonoua) were significant (Figure 5, Table S4a in the supplemental material). Strains isolated from Bonoua and Grand-Lahou were very different from the others. For strains from sampling areas of raffia wine, the differences between populations were also significant with the isolates from Alépé and Grand-Lahou and very different from the others isolates (Figure 5, Table S4b in the supplemental material).

4 Discussion

This study was centered on the genetic diversity and population structure of indigenous *S. cerevisiae* from Cote d'Ivoire, as well as the factors that may explain the structure of the resulting populations. For this purpose, four traditional alcoholic beverages (raffia wine, palm oil wine, ron wine and tchapalo beer) collected in nine regions of Côte d'Ivoire were studied.

The pH of these beverages, which fluctuated between 3 and 5, were similar to those reported by several authors in palm wines after the first day of tapping (Amoa-Awua *et al.*, 2007; Karamoko *et al.*, 2012) and in tchapalo beer produced by traditional brewers (Aka *et al.*, 2008). These ranges of pH could be a major advantage for the bio-preservation of these beverages by protecting them from the development of some pathogenic flora (Aka *et al.*, 2008; Yao *et al.*, 2009). The Total Titratable Acidity (TTA) and Total Soluble Sugars (TSS) contents were variable from one beverage to another, while the yeast average load was around 10^7 to 10^8 CFU /mL in the four beverages. The low TSS contents observed in beverages of this study could be related to the high yeast load, but also to the rest of the microbiota of these beverages. TTA rates observed in beverages could be due to the production of a variety of organic acids by yeasts and bacteria (N'guessan *et al.*, 2008; Karamoko *et al.*, 2012). Yeast loads found in this study could probably be due to the sum of various physical, chemical, and biotic factors such as temperature, humidity, presence of nutrients, such as sugars (Santiago-Urbina *et al.*, 2015). Similar yeast loads were reported in traditional fermented beverages produced in Africa (Aka *et al.*, 2008; Stringini *et al.*, 2009; Karamoko *et al.*, 2012). Diversity analysis indicated that ecological niche had a major influence on *S. cerevisiae* population's native of Côte d'Ivoire. Fis values of palm oil and raffia wines strains (0.603 and 0.544) revealed an excess of homozygote individuals which is very likely due to high inbreeding within each population rather than to population differentiation (Albertin *et al.*, 2014). Schuller and Casal (2007) also observed a significant excess of homozygotes in their study on three vineyards located in the northwest of Portugal. These authors explained this fact by asexual reproduction with some cycles of homothallic self-mating (genome renewal). Moreover, these authors proposed as an alternative explanation that a mitotic recombination or gene conversion during asexual reproduction may generate such a low of heterozygosity. The high degree of homozygosity would point to the existence of genetically isolated clonal subpopulations of *S. cerevisiae* strains with distinct genetic constitution. However, the microsatellite loci analyzed here represent only a tiny fraction of the genome and provide an indication of the mean heterozygosity. The heterozygosity detected at these microsatellite loci was found in good agreement with genomic data (i.e. Legras *et al.*, 2007; 2018) but other loci in the genome including those with technological relevance can present at different heterozygosity status.

Evaluation of genetic diversity of Ivorian strains only revealed that isolates sharing the same micro-environment would be genetically similar to each other, regardless of their geographical origin, implying that ecological origin is the main factor responsible for the

close phylogenetic relationship between *S. cerevisiae* strains. Impact of geographical origin on genetic diversity evaluated with Obstruct showed a difference between most of strains population according to sampling area, which was in accordance with Nei Fst pairwise distance. However these genetic distances remained low. Phylogenetic tree, Instruct and Discriminant Analysis of Principal Components (DAPC), considering only African strains, presented that subgroups appeared to be formed by differentiation according to ecological origin. One of the possible explanations is that at a relative close geographical distance, insects like bees, wasps, and fruit flies, as well as birds, which are known to be vectors for yeasts, could have homogenized these yeast populations. Also, Human activities (sharing seasonal staff and equipment during the harvest and fermentation periods) could influence the yeast population structure and promote dispersal (Börlin *et al.*, 2016). Although strains of palm oil wine, raffia wine and ron wine were isolated in Côte d'Ivoire, they have been associated to the same cluster containing the strains of other African countries which derived also of plant sap from the *Arecaceae* family (*Elaeis guineensis*, *Raphia hookeri*, *Borassus aethiopum*, *Borassus akeassii* and *Hyphaene thebaica*). The clustering of strains from different traditional beers, notably tchapalo in Côte d'Ivoire, dolo in Burkina Faso and bili bili in Ghana in the same cluster would reflect the history of the inoculum used. In Côte d'Ivoire, sorghum beer was initially produced by populations living in the North and North-East of the country (Djè *et al.*, 2008; N'guessan *et al.*, 2010). Inocula found in Southern and Eastern Côte d'Ivoire may have been introduced by traditional producers from North and North East of Côte d'Ivoire and those of Burkina Faso, Mali and Ghana. An exception to this distribution of strains according to their geographical origin was the genetic affiliation of strains from Asia (Japanese sake, Malaysian palm wine), and from North America (oak) of some Ivorian strains, a result in agreement with those of Liti *et al.* (2009). Strains from palm oil, raffia wines and tchapalo beer had common alleles with those of European wine, suggesting that these strains could have a hybrid origin as reported by Ezeronye and Legras (2009). This origin could always be explained by the fact that the European colonization of Africa would have led to the introduction of *S. cerevisiae* in the same way, as cachaça strains in Brazil (Barbosa *et al.*, 2018). This particular ecological situation, where African and European wine or bread species are present in the same environment, may have provided the opportunity for cross-breeding (Legras *et al.*, 2007). Our results confirm that the beverage production areas represent a reservoir of indigenous strains with a particular genotypic profile, selected by interactions between yeasts and their environment, as suggested by Martínez *et al.* (2007).

The presence of few European wine and bread strains alleles, in low proportions among Ivorian strains would have provided them with some genes of interest, expressing phenotypic characteristics which could be interesting to exploit in fermentation processes. However, a thorough analysis of the biotechnological properties of these strains will be necessary to confirm this idea.

Acknowledgment

This work was financially supported by the International Foundation for Science, Sweden (E/5560-1) and the National Institute for Agricultural Research (INRA, France). We thank Dr. Didier Montet who help for the project design, Christelle Louis-Mondésir for expert technical assistance and Tionhonkélé Drissa Soro for design of sampling map.

Reference

- Aigle, M., Erbs, D., Moll, M., 1984. Some molecular structures in the genome of lager brewing yeasts. *J. Am. Soc. Brew. Chem.* 42, 1-7. <https://doi.org/10.1094/ASBCJ-42-0001>
- Aka, S., Djeni, N.T., N'guessan, K.F., Yao, K.C., Djè, K.M., 2008. Variabilité des propriétés physico-chimiques et dénombrement de la flore fermentaire du tchapalo, une bière traditionnelle de sorgho en Côte d'Ivoire. *Afr. Sci.* 4, 274-286.
- Albertin, W., Panfili, A., Miot-Sertier, C., Goulielmakis, A., Delcamp, A., Salin, F., Lonvaud-Funel, A., Curtin, C., Masneuf-Pomarède, I., 2014. Development of microsatellite markers for the rapid and reliable genotyping of *Brettanomyces bruxellensis* at strain level. *Food Microbiol.* 42, 188-195. <https://doi.org/10.1016/j.fm.2014.03.012>
- Amoa-awua, W.K., Sampson, E., Tano-Debrah, K., 2007. Growth of yeasts, lactic and acetic bacteria in palm wine during tapping and fermentation from felled oil palm (*Elaeis guineensis*) in Ghana. *J. Appl. Microbiol.* 102, 599-606. <https://doi.org/10.1111/j.1365-2672.2006.03074.x>.
- Azumi, M., Goto-Yamamoto, N., 2001. AFLP analysis of type strains and laboratory and industrial strains of *Saccharomyces sensu stricto* and its application to phenetic clustering. *Yeast*. 18, 1145–1154. <https://doi.org/10.1002/yea.767>.
- Barbosa, R., Pontes, A., Santos, R.O., Montandon, G.G., de Ponzzes-Gomes, C.M., Morais, P.B., Gonçalves, P., Rosa, C.A., Sampaio, J.P., 2018. Multiple rounds of artificial

selection promote microbe secondary domestication – the case of cachaça yeasts. *Genome Biol Evol.* evy132, <https://doi.org/10.1093/gbe/evy132>.

Börlin, M, Venet, P, Claisse, O, Salin, F, Legras, J-L, Masneuf-Pomarede, I., 2016. Cellar-associated *Saccharomyces cerevisiae* population structure revealed high level diversity and perennial persistence at Sauternes wine estates. *Appl Environ Microbiol* 82, 2909–2918. <https://doi.org/10.1128/AEM.03627-15>.

Cockerham, C.C., 1969. Variance of gene frequencies. *Evolution.* 23, 72-84. <https://doi.org/10.2307/2406485>

Djè, K.M., N’guessan, K.F., Djeni, N.T., Dadié, A.T., 2008. Biochemical changes during alcoholic fermentation in the production of tchapalo, a traditional sorghum beer. *Int. J. Food Eng.* 4, 1-15. <https://doi.org/10.2202/1556-3758.1408>

Ezeronye, O.U., Legras, J.L., 2009. Genetic analysis of *Saccharomyces cerevisiae* strains isolated from palm wine in eastern Nigeria. Comparison with other African strains. *J. Appl. Microbiol.* 106, 1569-1578. <https://doi.org/10.1111/j.13652672.2008.04118.x>

Gao, H., Williamson S., Bustamante C.D., 2007. A Markov chain Monte Carlo approach for joint inference of population structure and inbreeding rates from multilocus genotype data. *Genetics*, 176, 1635-1651. <https://doi.org/10.1534/genetics.107.072371>

Gayevskiy, V., Klaere, S., Knight, S., Goddard, M.R., 2014. ObStruct: A Method to Objectively Analyse Factors Driving Population Structure Using Bayesian Ancestry Profiles. *PLoS ONE.* 9 (1), e85196. <https://doi.org/10.1371/journal.pone.0085196>

Gordon, D., Abajian, C., Green, P., 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* 8, 195-202.

Goudet, J., 2005. Hierfstat, a package for R to compute and test hierarchical F-statistics. *Mol Ecol Notes.* 5, 184-186. <https://doi.org/10.1111/j.1471-8286.2004.00828.x>

Hoffman, C.S., Winston, F., 1987. A ten-minute preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene*, 57, 267-272. [http://dx.doi.org/10.1016/0378-1119\(87\)90131-4](http://dx.doi.org/10.1016/0378-1119(87)90131-4)

Jombart, T., 2008. Adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24, 1403-1405. <https://doi.org/10.1093/bioinformatics/btn129>

Jombart, T., Devillard, S., Balloux, F., 2010. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genomics.* 11, 94. <https://doi.org/10.1186/1471-2156-11-94>.

Karamoko, D., Djeni, N.T., N’guessan, K.F., Bouatenin, K.M.J-P., Djè, K.M., 2012. The biochemical and microbiological quality of palm wine samples produced at different

periods during tapping and changes which occurred during their storage. *Food Control*. 26, 504-511. <https://doi.org/10.1016/j.foodcont.2012.02.018>

Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., Mayrose, I., 2015. Clumpak: a program for identifying clustering modes and packaging population structure inferences across. *Mol. Ecol. Resour.* 15, 1179-1191. <https://doi.org/10.1111/17550998.12387>

Kurtzman, C.P., Robnett, C.J., 1998. Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie Van Leeuwenhoek*. 73, 331–371. <https://doi.org/10.1023/A:1001761008817>

Legras, J-L., Karst F., 2003. Optimisation of Inter-Deltas analysis for *Saccharomyces cerevisiae* strain characterisation. *FEMS Microbiol. Lett.* 221, 249-255. [https://doi.org/10.1016/S0378-1097\(03\)00205-2](https://doi.org/10.1016/S0378-1097(03)00205-2)

Legras, J-L., Ruh, O., Merdinoglu, D., Karst, F., 2005. Selection of hypervariable microsatellite loci for the characterization of *Saccharomyces cerevisiae* strains. *Int. J. Food. Microbiol.* 102, 73-83. <https://doi.org/10.1016/j.ijfoodmicro.2004.12.007>

Legras, J-L., Merdinoglu, D., Cornuet, J-M., Karst, F., 2007. Bread, beer and wine: *Saccharomyces cerevisiae* diversity reflects human history. *Mol. Ecol.* 16, 2091-2102. <https://doi.org/10.1111/j.1365-294X.2007.03266.x>

Legras, J-L., Galeote, V., Camarasa, C., Blondin, B., Dequin, S. 2017. Ecology, Diversity and Applications of *Saccharomyces* Yeasts in Food and Beverages. In: Satyanarayana T., Kunze G. (eds) *Yeast Diversity in Human Welfare*. Springer, Singapore. https://doi.org/10.1007/978-981-10-2621-8_12

Legras, J-L., Galeote, V., Bigey, F., Camarasa, C., Marsit, S., Nidelet, T., Sanchez, I., Couloux, A., Guy, J., Franco-Duarte, R., Marcet-Houben, M., Gabaldon, T., Schuller, D., Sampaio, J.P., Dequin, S., 2018. Adaptation of *S. cerevisiae* to Fermented Food Environments Reveals Remarkable Genome Plasticity and the Footprints of Domestication. *Mol. Biol. Evol.* 35, 1712-1727. <https://doi.org/10.1093/molbev/msy066>

Liti, G., Carter, D.M., Moses, A.M., Warringer, J., Parts, L., James, S.A., Davey, R.P., Roberts, I.N., Burt, A., Koufopanou, V., Tsai, I.J., Bergman, C.M., Bensasson, D., O'Kelly, M.J., Van Oudenaarden, A., Barton, D.B., Bailes, E., Nguyen, A.N., Jones, M., Quail, M.A., Goodhead, I., Sims, S., Smith, F., Blomberg, A., Durbin, R., Louis, E.J., 2009. Population genomics of domestic and wild yeasts. *Nature*. 458, 337-341. <https://doi.org/10.1038/nature07743>

- López-Urbe, M.M., Cane, J.H., Minckley, R.L., Danforth, B.N., 2016. Crop domestication facilitated rapid geographical expansion of a specialist pollinator, the squash bee *Peponapis pruinosa*. *Proc. Biol. Scis.* 283, 20160443. <https://doi.org/10.1098/rspb.2016.0443>
- Lorca, G., Uribe, S., Godoy, L., Ganga, M.A., Martinez Fernandez, C.A., Martinez Fernandez, C. A., 2018. Screening of native *S. cerevisiae* strains in the production of Pajarete wine: a tradition of Atacama Region, Chile. *J. Wine. Res.* 29, 130-142. <https://doi.org/10.1080/09571264.2018.1465900>
- Martínez, C., Cosgaya, P., Vásquez, C., Gac, S., Ganga, A., 2007. High degree of correlation between molecular polymorphism and geographic origin of wine yeast strains. *J. Appl. Microbiol.* 103, 2185-2195. <https://doi.org/10.1111/j.13652672.2007.03493.x>
- Martiniuk, J.T., Pacheco, B., Russel, G., Tong, S., Backstrom, I., Measday, V., 2016. Impact of commercial strain use on *Saccharomyces cerevisiae* population structure and dynamics in pinot noir vineyards and spontaneous fermentations of a Canadian Winery. *PLoS One.* 11, e0160259. <https://doi.org/10.1371/journal.pone.0160259>
- N'guessan, F.K., Djeni, N.T., Dadié, A.T., Djè K.M, 2008. Influence of alcoholic fermentation temperature on sorghum beer quality. *Food.* 2, 119-123.
- N'guessan, F.K., N'Dri, D.Y., Camara, F., Djè K.M., 2010. *Saccharomyces cerevisiae* and *Candida tropicalis* as starters cultures for the alcoholic fermentation of tchapalo, a traditional sorghum beer. *World J. Microbiol. Biotechnol.* 26, 693-699. <https://doi.org/10.1007/s11274-009-0224-y>
- N'guessan, F.K., Brou, K., Jacques, N., Casaregola, S., Djè, K.M., 2011. Identification of yeasts during alcoholic fermentation of tchapalo, a traditional sorghum beer from Côte d'Ivoire. *Antonie van Leeuwenhoek.* 99, 855-864. <https://doi.org/10.1007/s10482-011-9560-7>.
- Naumov, G.I., Naumova, E.S., Sniegowski, P.D., 1998. *Saccharomyces paradoxus* and *Saccharomyces cerevisiae* are associated with exudates of North American oaks. *Can. J. Microbiol.* 44, 1045-1050.
- Nei, M., 1973. Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. U.S.A.* 70, 3321-3323.
- Ness, F., Lavallée, F., Dubourdieu, D., Aigle, M., Dulau, L., 1993. Identification of yeast strains using the polymerase chain reaction. *J. Sci. Food Agric.* 62, 89-94. <https://doi.org/10.1002/jsfa.2740620113>

- Nguyen, H.-V., Gaillardin, C., 1997. Two subgroups within the *Saccharomyces bayanus* species evidenced by PCR amplification and restriction polymorphism of the nontranscribed spacer 2 in the ribosomal DNA unit. *Syst. Appl. Microbiol.* 20, 286-294. [https://doi.org/10.1016/S0723-2020\(97\)80075-6](https://doi.org/10.1016/S0723-2020(97)80075-6)
- Peter, J., De Chiara, M., Friedrich, A., Yue, J.X., Pflieger, D., Bergström, A., Sigwalt, A., Barre, B., Freil, K., Llored, A., Cruaud, C., Labadie, K., Aury, J.M., Istace, B., Lebrigand, K., Barbry, P., Engelen, S., Lemainque, A., Wincker, P., Liti, G., Schacherer, J., 2018. Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature*. 556, 339-344. <https://doi.org/10.1038/s41586-018-0030-5>
- Querol, A., Barrio, E., Huerta, T., Ramon, D., 1992. Molecular monitoring of wine fermentations conducted by active dry yeast strains. *Appl. Environ. Microbiol.* 58, 2948-2953.
- Santiago-Urbina, J.A., Arias-García, J.A, Ruiz-Terán, F., 2015. Yeast species associated with spontaneous fermentation of taberna, a traditional palm wine from the southeast of Mexico. *Ann. Microbiol.* 65, 287-296. <https://doi.org/10.1007/s13213-014-0861-8>
- Schuller, D., Casal, M., 2007. The genetic structure of fermentative vineyard-associated *Saccharomyces cerevisiae* populations revealed by microsatellite analysis. *Antonie Van Leeuwenhoek*. 91, 137-150. <https://doi.org/10.1007/s10482-006-9104-8>
- Sniegowski, P.D., Dombrowski, P.G., Fingerman, E., 2002. *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* coexist in a natural woodland site in North America and display different levels of reproductive isolation from European conspecifics. *FEMS Yeast Res.* 1, 299-306. <https://doi.org/10.1111/j.1567-1364.2002.tb00048.x>
- Stringini, M., Comitini, F., Taccari, M., Ciani, M., 2009. Yeast diversity during tapping and fermentation of palm wine from Cameroon. *Food Microbiol.* 26, 415-420. <https://doi.org/10.1016/j.fm.2009.02.006>
- Tapsoba, F., Legras, J-L., Savadogo, A., Dequin, S., Traore A.S., 2015. Diversity of *Saccharomyces cerevisiae* strains isolated from *Borassus akeassii* palm wines from Burkina Faso in comparison to other African beverages. *Int. J. Food Microbiol.* 15, 30065-30069. <https://doi.org/10.1016/j.ijfoodmicro.2015.07.010>
- Techera, A.G., Jubany, S., Carrau, F.M., Gaggero, C., 2001. Differentiation of industrial wine yeast strains using microsatellite markers. *Lett. Appl. Microbiol.* 33, 71-75. <https://doi.org/10.1046/j.1472-765X.2001.00946.x>
- Tra Bi, C.Y., N'guessan, F.K., Kouakou, C.A., Noémie, J., Casaregola, S., Djé, M.K, 2016. Identification of yeasts isolated from raffia wine (*Raphia hookeri*) produced in Côte

d'Ivoire and genotyping of *Saccharomyces cerevisiae* strains by PCR inter-delta.
World J. Microbiol. Biotechnol. 32, 125-134. <https://doi.org/10.1007/s11274-0162095-3>

Vezinhet, F., Blondin, B., Hallet, J.N., 1990. Chromosomal DNA patterns and mitochondrial DNA polymorphisms as tools for identification of enological strains of *Saccharomyces cerevisiae*. Appl. Microbiol. Biotechnol. 32, 568-571. <https://doi.org/10.1007/BF00173729>

Viel, A., Legras, J.L., Nadai, C., Carlot, M., Lombardi, A., Crespan, M., Migliaro, D., Giacomini, A., Corich, V., 2017. The geographic distribution of *Saccharomyces cerevisiae* isolates within three Italian neighboring winemaking regions reveals strong differences in yeast abundance, genetic diversity and industrial strain dissemination. Front Microbiol. 8, 1595. <https://doi.org/10.3389/fmicb.2017.01595>

Yao, A., Egounlety, M., Kouamé, L.P., Thonart, P., 2009. Les bactéries lactiques dans les aliments ou boissons amylacés et fermentés de l'Afrique de l'Ouest: leur utilisation actuelle. Ann. Med. Vet. 153, 54-65.

Yu, J., Dossa K., Wang, L., Zhang, Y., Wei, X., Liao, B., Zhang, X., 2017. PMDBase: a database for studying microsatellite DNA and marker development in plants. Nucleic. Acids Res. D1046–D1053. <https://doi.org/10.1093/nar/gkw906>




Figure 1: Map of Côte d'Ivoire showing the areas from where samples were collected.




Figure 2a: Phylogenetic tree among Ivorian *S. cerevisiae* strains. The tree is drawn according to neighbour-joining method from a pairwise individual distance matrix calculated with relative dissimilarity distance between 170 strains and based on the polymorphism at 12 loci.




Figure 2b: Phylogenetic tree among *S. cerevisiae* strains of diverse geographical origins and substrates. The tree is drawn according to the neighbour-joining method from a pairwise individual distance matrix calculated with relative dissimilarity distance between the 284 strains (from 18 different ecological niches) and based on the polymorphism at 12 loci.





Figure 3: Inferred population structure using InStruct program on the 384 *S. cerevisiae* strains. Deviance information criterion indicated the most likely population number to be $K = 15$. *S. cerevisiae* populations are delimited vertically according to their ecological origin. Colours represent the different alleles that constitute each population.

(a)



(b)





Figure 4: Discriminant principal component analysis (DPCA) (a): Plot showing representation of the different clusters according to axes 1 and 2; (b) plot showing representation of the different clusters according to axes 3 and 4. Clusters are shown by different colours and inertia ellipses, while dots represent individuals.

(a)



(b)




Figure 5: Canonical discriminant analysis of the ancestry obtained for palm oil wine (a) and raffia wine (b) isolates from Côte d’Ivoire with OBSTRUCT, after removal of strains from other countries. The outer blue ellipsoid labelled group reflects the variation of the group means around the grand mean while the red circle reflects the pooled within-group dispersion and covariation. The black points indicate predefined populations while numbers at the arrows indicate inferred populations.

Table 1: Geographical information of the sampling areas and sample codes

Sampling areas	Geographical coordinates	Distance between the sampling area and the city of Abidjan (Km)	Beverage collected	Code
Abidjan	$1 : 5^{\circ}18.5796' \text{ N}$ $L : 4^{\circ}0.7596' \text{ W}$	0	Tchapalo beer	$\frac{\text{TLM/ Ab/}}{\text{TT14/ TPA/}}$
Bingerville	$1 : 5^{\circ}21.3486' \text{ N}$	18	Palm oil wine	BingerVP
	$L : 3^{\circ}53.1222' \text{ W}$		Tchapalo beer	Binger-T
Attinguié	$1 : 5^{\circ}29'40'' \text{ N}$ $L : 4^{\circ}03'06'' \text{ W}$	22	Palm oil wine	AttgiéP-Vp
Alépé	$1 : 5^{\circ}29'46'' \text{ N}$	45	Palm oil wine	PA
	$L : 3^{\circ}39'49'' \text{ W}$		Raffia wine	AR
Bonoua	$1 : 5^{\circ}16.3482' \text{ N}$ $L : 3^{\circ}35.775' \text{ W}$	59	Palm oil wine	Bon Vp
Adzopé	$1 : 6^{\circ}06'24'' \text{ N}$ $L : 3^{\circ}51'42'' \text{ W}$	104	Raffia wine	Adz-R
Grand-Lahou	$1 : 5^{\circ} 8' 7 \text{ N}$	152	Palm oil wine	PG
	$L : 5^{\circ} 1' 26 ' \text{ W}$		Raffia wine	GR
Toumodi	$1 : 6^{\circ}33.4794' \text{ N}$ $L : 5^{\circ}1.0614' \text{ W}$	198	Ron wine	TR
Abengourou	$1 : 6^{\circ}43'46'' \text{ N}$ $L : 3^{\circ}29'47'' \text{ W}$	210	Raffia wine	Abeng R

Table 2: Physico-chemical characteristics and yeast count of alcoholic beverages of the study

Beverages		pH	Total titratable acidity (%)	Total soluble sugars (°Brix)	Yeast count (CFU/mL)
Palm oil wine	Average	$3.73 \pm 0.09^*$	$0.80 \pm 0.05^*$	$5.41 \pm 0.77^*$	$1.4 \pm 0.1 \times 10^{7*}$
	Minimum	3.45 ± 0.04	0.75 ± 0.07	3.72 ± 1.13	$9.7 \pm 0.5 \times 10^6$
	Maximum	4.30 ± 0.19	0.87 ± 0.03	6.65 ± 1.41	$1.7 \pm 0.2 \times 10^7$
Raffia wine	Average	$3.80 \pm 0.12^*$	$0.81 \pm 0.10^*$	$4.35 \pm 0.79^\dagger$	$1.3 \pm 0.2 \times 10^{7*}$
	Minimum	3.21 ± 0.26	0.47 ± 0.04	3.00 ± 0.30	$9.5 \pm 2.3 \times 10^6$
	Maximum	4.35 ± 0.01	1.35 ± 0.07	7.00 ± 0.40	$4.8 \pm 1.8 \times 10^7$
Ron wine	Average	$3.72 \pm 0.12^*$	$0.68 \pm 0.09^\dagger$	$4.88 \pm 0.40^\dagger$	$1.1 \pm 0.3 \times 10^{8\dagger}$
	Minimum	3.54 ± 0.08	0.48 ± 0.11	4.03 ± 0.25	$1.9 \pm 0.7 \times 10^7$
	Maximum	3.91 ± 0.21	1.00 ± 0.14	5.80 ± 0.30	$2.6 \pm 0.6 \times 10^8$
Tchapalo beer	Average	$3.46 \pm 0.13^*$	$0.93 \pm 0.03^\ddagger$	$7.70 \pm 1.04^\ddagger$	$1.7 \pm 0.1 \times 10^{7*}$
	Minimum	3.38 ± 0.1	0.92 ± 0.02	5.81 ± 0.57	$1.2 \pm 0.1 \times 10^7$
	Maximum	3.5 ± 0.14	0.94 ± 0.04	10.87 ± 2.0	$2.4 \pm 0.1 \times 10^7$

The values expressed are the average of three measurements. On the same column, mean values with the same symbol are not significantly different (Tukey HSD test range at $P < 0.05$).

Table 3a: Pairwise Fst distance matrix between populations isolated from different beverages. Fst is calculated according to Nei (1973). Fst values are given in the lower matrix and upper matrix indicates P-value estimated from 1000 permutations of the genotypes dataset.

	Palm oil wine	Raffia wine	Ron wine	Tchapalo beer
Palm oil wine	0.000	0.008	0.001	0.001
Raffia wine	0.021	0.000	0.001	0.001
Ron wine	0.061	0.069	0.000	0.001
Tchapalo beer	0.107	0.074	0.073	0.000

Table 3b: Pairwise Fst distance matrix between sampling areas. Fst is calculated according to Nei (1973). Fst values are given in the lower matrix and upper matrix indicates P-value estimated from 1000 permutations of the genotypes dataset.

Palm oil wine					
	Attingué	Grand-Lahou	Alépé	Bingerville	Bonoua
Number of individuals	6	7	7	5	4
Attingué	0.000	0.022	0.035	0.077	0.002
Grand-Lahou	0.066	0.000	0.001	0.033	0.001
Alépé	0.045	0.070	0.000	0.008	0.001
Bingerville	0.093	0.092	0.084	0.000	0.001
Bonoua	0.070	0.064	0.073	0.118	0.000

Raffia wine				
	Grand-Lahou	Alépé	Adzopé	Abengourou
Number of individuals	16	35	6	7
Grand-Lahou	0.000	0.003	0.008	0.044
Alépé	0.011	0.000	0.019	0.031
Adzopé	0.049	0.022	0.000	0.051
Abengourou	0.054	0.026	0.061	0.000

Tchapalo beer				
---------------	--	--	--	--

	Abidjan	Bingerville
Number of individuals	17	2
Abidjan	0.000	0.198
Bingerville	0.022	0.000

Values are significantly different for P-value < 0.05.

Table 4: Fis value per population and corresponding estimate of selfing rate

	Ecological niche			
	Raffia wine	Palm oil wine	Tchapalo beer	Ron wine
Number of individuals	64	29	19	58
Fis	0.603	0.544	0.138	0.214
Outcrossing estimate ($o=(1-Fis)/(1+Fis)$)	0.25	0.30	0.76	0.65
Selfing estimate ($=1-o = 2Fis/(1+Fis)$)	0.75	0.70	0.24	0.35