



Testing Linear Regressions by StatsModel Library of Python for Oceanological Data Interpretation

Polina Lemenkova

► To cite this version:

Polina Lemenkova. Testing Linear Regressions by StatsModel Library of Python for Oceanological Data Interpretation. Aquatic Sciences and Engineering, 2019, 34 (2), pp.51-60. 10.26650/ASE2019547010 . hal-02168755

HAL Id: hal-02168755

<https://hal.science/hal-02168755>

Submitted on 29 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Testing Linear Regressions by StatsModel Library of Python for Oceanological Data Interpretation

Polina Lemenkova^{1*} 

Cite this article as: Lemenkova, P. (2019). Testing Linear Regressions by StatsModel Library of Python for Oceanological Data Interpretation. *Aquatic Sciences and Engineering*, 34(2), 51-60.

ABSTRACT

The study area is focused on the Mariana Trench, west Pacific Ocean. The research aim is to investigate correlation between various factors, such as bathymetric depths, geomorphic shape, geographic location on four tectonic plates of the sampling points along the trench, and their influence on the geologic sediment thickness. Technically, the advantages of applying Python programming language for oceanographic data sets were tested. The methodological approaches include GIS data collecting, data analysis, statistical modelling, plotting and visualizing. Statistical methods include several algorithms that were tested: 1) weighted least square linear regression between geological variables, 2) autocorrelation; 3) design matrix, 4) ordinary least square regression, 5) quantile regression. The spatial and statistical analysis of the correlation of these factors aimed at the understanding, which geological and geodetic factors affect the distribution of the steepness and shape of the trench. Following factors were analysed: geology (sediment thickness), geographic location of the trench on four tectonics plates: Philippines, Pacific, Mariana and Caroline and bathymetry along the profiles: maximal and mean, minimal values, as well as the statistical calculations of the 1st and 3rd quantiles. The study revealed correlations between the sediment thickness and distinct variations of the trench geomorphology and sampling locations across various segments along the crescent of the trench.

Keywords: Programming language, Python, Statistical analysis, Pacific Ocean, Hadal trenches, Mariana Trench, oceanology, marine geology

ORCID IDs of the authors:
P.L. 0000-0002-5759-1089

¹Ocean University of China,
College of Marine Geo-sciences.
238 Songling Road, Laoshan,
266100, Qingdao, Shandong
Province, People's Republic of
China

Submitted:
30.03.2019

Revision Requested
22.05.2019

Last Revision Received
13.06.2019

Accepted:
13.06.2019

Online published:
26.06.2019

Correspondence:
Polina Lemenkova
E-mail:
lemenkovapolina@stu.ouc.edu.cn
pauline.lemenkova@gmail.com

©Copyright 2019 by Aquatic
Sciences and Engineering
Available online at
<https://dergipark.org.tr/ase>

INTRODUCTION

Multiple approaches and GIS methods have been used so far to model ocean seafloor, the most unreachable part of the Earth. These include echo sounding (Smith, & Sandwell, 1997), CTD (conductivity-temperature-depth profiler) technique (Taira et al., 2005), acoustic methods, continual profiling with single-beam systems and bottom coverage capability, multi-beam swath-mapping systems (Dierssen, & Theberge, 2014), classic approaches of the GIS mapping and other tools of geoinformatics (Fujie et al., 2006), remote sensing images analysis, navigation charts and data modelling using schematic cross-sections of the subduction

zones (e.g. Schellart, 2008), statistical modelling using R and packages, e.g. dplyr, ggplot2, PMCMR, car (Reid et al., 2018). Of all these, statistical modelling of the oceanological data sets by means of R and Python programming languages is the most cost-effective for investigating hadal trench geomorphology.

Various studies have been reported on the geologic variations of the Mariana Trench involving uneven distribution of various geomorphic phenomena across the seafloor (e.g., Michibayashi et al., 2007; Grand et al., 1997). Amongst these, the questions of how the trench shape is varies and what are the factors affective its geomorphology are the most chal-


Regional studies of the marine geology of the trenches across the Pacific Ocean (e.g., Bello-González et al., 2018; Boston et al., 2017), modelling and predictions made upon analysis of the geophysical settings of various trench, produced by these investigators were instrumental in understanding current issues of the marine geological studies. The concepts of these reports on seafloor spreading, tectonic slab subduction, continental drift, and plate tectonics in the Pacific Ocean were analysed in the current research.

The study area is located in the Mariana Trench, west Pacific Ocean, where the deepest place of the Earth is recorded (Theberge, 2008).

First, during the geospatial part of the research, the data were collected from the Quantum GIS project as vector layers. The attribute tables contained numerical data on bathymetry, geology, tectonic plates and geometric features of the Mariana Trench in its various segments of the geographic location: north-west, centre, south-west.

METHODOLOGY

The variables of geologic interest were stored in the table consisting of 18 rows where numeric information describes geology, bathymetry, geodesy and tectonics of the Mariana Trench. To fit most of the models covered by StatsModels Python library, the design or regressor matrix was created using existing approaches (Everitt, 2002; Box, & Tiao, 1992; Millman & Aivazis, 2011). The



```

1 from __future__ import print_function
2 %matplotlib inline
3 import numpy as np
4 import pandas as pd
5 import statsmodels.api as sm
6 from patsy import dmatrices
7 import os
8 os.chdir('/Users/pauline/Documents/Python')
9 df = pd.read_csv('Tab-Morph.csv')
10 df = df.dropna()
11 df[-10:]
12 y, X = dmatrices('profile ~ sedim_thick + igneous_volc + slope_angle',
13                 data=df, return_type='dataframe')
14 y[:7]
15 X[:7]

```

	Intercept	sedim_thick	igneous_volc	slope_angle
0	1.0	132.0	112.0	25.0
1	1.0	103.0	71.0	32.0
2	1.0	96.0	0.0	51.0
3	1.0	109.0	0.0	64.0
4	1.0	127.0	3.0	52.0
5	1.0	135.0	5.0	70.0
6	1.0	142.0	0.0	55.0

OLS Regression Results					
Dep. Variable:	profile	R-squared:		0.457	
Model:	OLS	Adj. R-squared:		0.379	
Method:	Least Squares	F-statistic:		5.881	
Date:	Sun, 24 Mar 2019	Prob (F-statistic):		0.00445	
Time:	19:36:32	Log-Likelihood:		-77.241	
No. Observations:	25	AIC:		162.5	
Df Residuals:	21	BIC:		167.4	
Df Model:	3				
Covariance Type:	nonrobust				
	coef	std err	t	P> t	[0.025 0.975]
Intercept	17.9305	6.362	2.819	0.010	4.701 31.160
sedim_thick	-0.1320	0.048	-2.743	0.012	-0.232 -0.032
igneous_volc	0.0445	0.016	2.819	0.010	0.012 0.077
slope_angle	0.1328	0.115	1.151	0.263	-0.107 0.373
Omnibus:		1.669	Durbin-Watson:		0.654
Prob(Omnibus):		0.434	Jarque-Bera (JB):		1.095
Skew:		-0.198	Prob(JB):		0.578
Kurtosis:		2.054	Cond. No.		884.

Warnings:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

```

Intercept    17.930487
sedim_thick  -0.131967
igneous_volc  0.044532
slope_angle   0.132841
dtype: float64

```

Table 1. Python code for design matrix 'dmatrices' (left) and OLS computation by StatsModels of Python (right), Mariana Trench data frame.

first is a matrix of endogenous variables of sediment thickness, which show the response or geological regressand on changed environmental conditions: geographic location, depth or tectonic plate.

The design matrix (Table 1, left) shows the results of the first six lines representing values of explanatory variables in a set of geological attributes, Mariana Trench.

The computing of the Ordinary Least Square (OLS) was based on the formula (1):

$$b_i = \frac{\sum_{j=1}^n (x_j - \bar{x})(y_j - \bar{y})}{\sum_{j=1}^n (x_j - \bar{x})^2} \quad (1)$$

Where,

n is the sample size;

x is a constant and a scalar regressor;

y is a random regressor, sampled together with x ;

h is the number of lags being tested;

Each row of the calculated OLS coefficient estimates (Table 1, right) shows an individual bathymetric profile with the successive columns corresponding to the geologic and oceanographic variables and their specific values across the profiles.

Quantile statistics (QQ)

The used algorithm is very straightforward with a selected function of qqplot() by StatsModels to perform this task. The QQ regression is a common abbreviation for 'quantile by quantile' statistical plot. The plot shows (Figure 2) one quantile against another across various geological parameters (from left to right): A) Sediment thickness; B) Slope angle degrees; C) Pacific Plate; D); Philippine Plate E) Mariana Plate; F) Distribution of samples of igneous volcanic areas.

Technically, the plotting was performed using following code of Python for each corresponding plot:

```

ax1=plt = qqplot(df.sedim_thick, line='q',
ax=ax1, fit=True,
linewidth=.5, alpha=.5, markerfacecolor='#00a497',
markeredgecolor='grey', )

```

The QQ statistics calculation has been based on the following formula (2) after Ljung, & Box (1978):

$$Q = n(n+2) \sum_{k=1}^n \frac{I_k^2}{(n-k)} \quad (2)$$

Where,

n is the sample size;

ρ is the sample autocorrelation at lag k , and

h is the number of lags being tested.

The comparison of all the six subplots enables to analyse the form of their shape against a straight line. The quantiles are bathymetric sample observations with geologic attribute values placed in the ascending order. The QQ statistics are used over the pool of the sampling data to study their distribution. A QQ statistic is a visual representation of the quantiles of a standard normal distribution of the geological data set across the Mariana Trench, showing their variation in space.

Weighted Least Squares

A Weighted Least Squares (WLS) for the geological variables are shown on Figure 3. The approach of a weighted least squares is a standard approach in regression analysis to approximate the solution of overdetermined systems which is the case for the complex marine geological systems. The least squares algorithms has two sub-types: linear or ordinary least squares and nonlinear least squares. In the scope of this research, only the linear least squares were tested: ordinary least squares and weight-



Figure 2. Plotted QQ statistics for the data distribution: A) Sediment thickness; B) Slope angle degrees; C) Pacific Plate; D); Philippine Plate E) Mariana Plate; F) Volcanic spots.

ed least square which is a generalization of the first one. In this case, the off-diagonal entries of the correlation matrix of the geological residuals are null the variances of the observations, yet unequal along the covariance matrix.

A WLS is a statistical approach representing a special case of the generalized least squares.

The calculation is based on the principle of the Gauss–Newton algorithm that solves a non-linear least squares problem by modifying a Newton’s method for finding a minimum of a function. The computation was based on the general approach of existing equation (after Björck, 1996):

$$J^T W J \Delta \beta = J^T W \Delta y \quad (3)$$

Where

W is a diagonal when the observational errors are uncorrelated and the weight matrix;

$J(t)$ is a transposed Jacobian matrix;

β are unbiased estimators as linear column vectors, the entries of the Jacobian matrix;

y is a vector of the response values.

The calculations of the WLS for the data set (Table 2) were done according to the reported procedures (Seabold & Perktold, 2010; Strutz, T. (2016) by Python code snippet:

```
# Step-1.
mod_wls = sm.WLS(y, X, weights=1./(w ** 2))
```

```
res_wls = mod_wls.fit()
print(res_wls.summary())
# Step-2.
res_ols = sm.OLS(y, X).fit()
print(res_ols.params)
print(res_wls.params)
# Step-3.
se = np.vstack([[res_wls.bse], [res_ols.bse],
                [res_ols.HC0_se], [res_ols.HC1_se], [res_ols.HC2_se], [res_ols.HC3_se]])
se = np.round(se,4)
colnames = ['x1', 'const']
rownames = ['WLS', 'OLS', 'OLS_HC0', 'OLS_HC1', 'OLS_HC3', 'OLS_HC3']
tabl = SimpleTable(se, colnames, rownames, txt_fmt=default_txt_fmt)
print(tabl)
```

Quantile regressions

Quantile regression shows (Figure 4) the estimated conditional median and other quantiles of the response geological variables. Thus the upper two rows of the plot show (Figure 4, A, B, C, D) data distribution across tectonic plates: Pacific Plate, Philippine Plate, Mariana Plate and Caroline Plate. The lower row of the plot (Figure 4, E, F) shows data distribution for the cumulative sediment thickness and slope angle degree by profiles.




Figure 3. Weighted Least Squares plotted for data distribution: A) Pacific Plate, B) Philippine Plate, C) Mariana Plate, D) Sediment thickness, E) Depths (max); F) Slope angle degree.

The quantile regressions were plotted (Figure 4) using Python code by StatsModel:

```
# Step-1. Least Absolute Deviation
mod = smf.quantreg('profile ~ slope_angle', data)
res = mod.fit(q=.5)
print(res.summary())
# Step-2. Placing the quantile regression
```

```
results in a Pandas DataFrame, and the OLS
results in a dictionary
quantiles = np.arange(.05, .96, .1)
def fit_model(q):
    res = mod.fit(q=q)
    return [q, res.params['Intercept'], res.
            params['slope_angle']] + res.conf_
int().loc['slope_angle'].tolist()
```



```
models = [fit_model(x) for x in quantiles]
models = pd.DataFrame(models, columns=['q', 'a', 'b', 'lb', 'ub'])
ols = smf.ols('profile ~ slope_angle', data).fit()
ols_ci = ols.conf_int().loc['slope_angle'].tolist()
```

```
ols = dict(a = ols.params['Intercept'],
           b = ols.params['slope_angle'],
           lb = ols_ci[0],
           ub = ols_ci[1])
print(models)
print(ols)
```




Figure 4. Quantile regression plotted for sediment thickness (m) versus geologic parameters: A) Pacific Plate, B) Philippine Plate, C) Mariana Plate, D) Caroline Plate, E) Cumulative sediment thickness and F) Slope angle degree by profiles.

Essentially, quantile regression is another approach of the linear regression tested in the current research. The quantitative results of the quantile regression are shown on Table 3, with respect to the relevant plots shown on Figure 4 (corresponding to Figure 4: A, B, C, D, E, F).

Dynamic regression model: State Autoregressive Moving Average (SARIMA)

The methodology of the dynamic regression model is based on the StatsModels embedded algorithm (Seabold & Perktold, 2010).

The abbreviation of SARIMA is the Space AutoRegressive Moving Average model, initially developed by Ansley & Kohn, (1985). The concept of the application of the SARIMA time series estimation and post-estimation lies in the following. The changes in the variables that are taken as time lags in classic dynamic regression models are applied towards bathymetric profiles from 1 to 25. In this way, unlike in time series case when the SARIMA fits the univariate models with time-dependent disturbances, this case applies space-dependent disturbances, crossing the sampling selection from 1 to 25 (X axe).

Because the model includes both dependent and independent variables, the selected type of SARIMA was SARIMAX (see the Python code snippet below). The first group consists in changing variables that is geologic settings and bathymetry (depths). The

second group (independent variables) is presented by the profiles lags that cross the Mariana Trench with the distance between each of 100 km and the length of 1000 km. This cross-section profiles are taken as independent variables. Therefore, the dependant variables differ spatially in different segments of the trench.

The model (Figure 5) fits univariate model of the geomorphic structure of the trench by independent values of the distribution of the bathymetric observation by profiles with dependent disturbances of depths.

Fitting the model was done using the Python snippet:

```
mod = sm.tsa.statespace.SARIMAX(data['sedim_thick'], trend='c', order=(1,1,1))
res = mod.fit(dis=False)
print(res.summary())
fig = sm.graphics.tsa.plot_pacf(data.iloc[1:]
['Ddf.geology'], lags=25, ax=axes[1])
```

The algorithm fits a model where the disturbances follow a linear specification of the bathymetric distribution across the trench. The dependent and independent geological variables vary by profiles (Figure 5). Plotting was done using subplot function of Matplotlib:

```
fig, axes = plt.subplots(2, 2, figsize=(15,8))
```




Figure 5. Plotted SARIMAX statistics for the bathymetry: A) Sediment thickness index; B) Data distribution index, Philippine Plate; C) Autocorrelation; D); Partial Autocorrelation.

Script-040-Sarimax Stata.py - /Users/pauline/Documents/Python/Script-040-Sarimax Stata.py (3.7.2)*

```
# Step-1. Import data
os.chdir('/Users/pauline/Documents/Python')
data = pd.read_csv('Tab-Morph.csv')
data.index = data.profile

# Step-2. Fitting the model by sediment thickness
mod = sm.tsa.statespace.SARIMAX(data['sedim_thick'], trend='c', order=(1,1,1))
res = mod.fit(disp=False)
print(res.summary())

# Step-3. Dataset
data['sedim_thick'] = np.log(data['sedim_thick'])
data['D.sedim_thick'] = data['plate_phill'].diff()

# Step-4. Graph data
fig, axes = plt.subplots(1, 2, figsize=(15,4))
axes[0].plot(data.index_mpl_repr(), data['sedim_thick'], '-')
axes[0].set_xlabel('Bathymetric profiles, 1:25', fontsize=10)
axes[0].set_title('Mariana Trench: Sediment thickness index, \nby bathymetric profiles')
axes[0].annotate('A', xy=(0.95, .92), xycoords='axes fraction', fontsize=12,
                bbox=dict(boxstyle='round', pad=0.3, fc='w', edgecolor='grey', linewidth=1, alpha=0.9))

# Step-5. Log difference
axes[1].plot(data.index_mpl_repr(), data['D.sedim_thick'], '-')
axes[1].hlines(0, data.index[0], data.index[-1], 'r')
axes[1].set_xlabel('Bathymetric profiles, 1:25', fontsize=10)
axes[1].set_title('Mariana Trench: Data distribution index on \nPhilippine Plate, difference of logs')
axes[1].annotate('B', xy=(0.95, .92), xycoords='axes fraction', fontsize=12,
                bbox=dict(boxstyle='round', pad=0.3, fc='w', edgecolor='grey', linewidth=1, alpha=0.9))

# Step-6. Graph data
fig, axes = plt.subplots(1, 2, figsize=(15,4))
fig = sm.graphics.tsa.plot_acf(data.iloc[1:]['D.sedim_thick'], lags=21, ax=axes[0])
axes[0].annotate('C', xy=(0.95, .92), xycoords='axes fraction', fontsize=12,
                bbox=dict(boxstyle='round', pad=0.3, fc='w', edgecolor='grey', linewidth=1, alpha=0.9))
fig = sm.graphics.tsa.plot_pacf(data.iloc[1:]['D.sedim_thick'], lags=21, ax=axes[1])
axes[1].annotate('D', xy=(0.95, .92), xycoords='axes fraction', fontsize=12,
                bbox=dict(boxstyle='round', pad=0.3, fc='w', edgecolor='grey', linewidth=1, alpha=0.9))

plt.show()
```

Statespace Model Results

Dep. Variable:

sedim_thick

No. Observations:

25

Model:

SARIMAX(1, 1, 1)

Log Likelihood

-108.677

Date:

Fri, 29 Mar 2019

AIC

225.354

Time:

10:49:01

BIC

230.066

Sample:

0

HQIC

226.604

Covariance Type:

opg

coef

std err

z

P>|z|

[0.025

0.975]

intercept

-1.8019

2.957

-0.609

0.542

-7.597

3.993

ar.L1

0.2696

0.390

0.692

0.489

-0.494

1.033

ma.L1

-0.6914

0.414

-1.669

0.095

-1.503

0.121

sigma2

495.4968

197.897

2.504

0.012

107.625

883.369

Ljung-Box (Q):

18.06

Jarque-Bera (JB):

4.25

Prob(Q):

0.75

Prob(JB):

0.12

Heteroskedasticity (H):

0.38

Skew:

-0.73

Prob(H) (two-sided):

0.19

Kurtosis:

4.46

Warnings:

[1] Covariance matrix calculated using the outer product of gradients (complex-step).

>>>

Ln: 28

Col: 29

Table 4. Python programming code for used SARIMAX algorithm (left) and Statespace Model Results (right). Tested case in Mariana Trench tectonics: sediment thickness across Philippine Plate.

RESULTS AND DISCUSSIONS

The conducted statistical data modelling and several types of the regression analysis were aimed to compare the variance in the geological data sets of the Mariana Trench explained by the complex interplay of the geomorphic, geological and oceanological attributes of the data and the bathymetric factors of the location of several segments of the trench in various parts of the Pacific Ocean. Using StatsModels library of Python, in particular several linear models of the correlation between various factors were computed, analysed and explained by the groups of variables.

Tested environmental variables of the Mariana Trench include four main geological factors (location on the tectonic plates, slope steepness degree, sediment thickness of the layer and bathymetric depth and submarine volcanism) and attributes of the 25 cross-section bathymetric profiles (mean values, maximal depths, median values and two quartile sub-division of the data sets) and the shared variances between environment and attributes. Shared variance arises due to correlations between the factor of sediment thickness and slope angle degree, e.g. because the attributes of the sediment accumulation are influenced by the canyon shape apart from the directly or indirectly depending on the oceanological conditions of the submarine currents.

The graphical output shows normal data distribution as demonstrated by (Figure 2), where each QQ profile has a given value for data distribution by quantiles across the trench profiles. The distribution of the geologic residuals is shown on Figure 3 showing particularly data correlation for several cases: frequency of the data distribution by Pacific Plate, Philippine Plate, Mariana Plate, sediment thickness, and range of the bathymetric depths taken for the maximal values, and finally, geomorphological shape as slope angle degree. The results shown on Table 2 represent the computed numerical values of the previous graph (Figure 3).

The results on the Quantile regression (Figure 4) show the conditional median of the response geologic variable given changing

bathymetric values with movement southwards across the Mariana Trench. Thus, the upper plots shows (Figure 4, A, B, C) data distribution across tectonic plates: Pacific Plate, Philippine Plate, Mariana Plate and Caroline Plate. The lower plot (Figure 4, D, E, F) shows data distribution for the Caroline Plate and cumulative sediment thickness and slope angle degree by profiles.

The numerical explanation of the Figure 4 with corresponding sub-plots is presented in Table 3. The Figure 5 shows dynamic regression model using Python function embedded in StatsModel: State Autoregressive Moving Average. Finally, Table 4 shows the Python code that was used to perform the procedure of SARIMAX and the resulting output table. The model shows autocorrelation of the data by bathymetric profiles. The results demonstrated a correlation between the geological variables and geospatial location of the samplings across Mariana Trench, which proves the interplay between multiple factors affecting its geomorphology.

CONCLUSIONS

While the usage of the traditional methods of geoinformatics and spatial analysis is, beyond doubts, strongly recommended for any research in geoscience, there is another powerful tool for the geospatial data processing other than GIS, sometimes overlooked or skipped by the geographers: a data modelling by use of Python or R programming languages. Python, an open source free programming language is highly suitable for the statistical analysis in geoscience research, since it has a powerful statistical and math libraries, e.g. StatsModel, highly effective for scientific computing and used in the current research. The functionality of Python language and StatsModel, tested in this work, is proved to be highly effective for the statistical analysis of the geo-marine sets.

The proposed approach of the Python based statistical analysis enables accurate and efficient computation and modelling of the large data sets in marine geology and oceanology. A challenge in the evaluation of geological big data sets (that is, several thou-

sand of observation points as in this case, 12.590 samples) concerns the difficulty in manual identifying a correct algorithms in the computations and data distribution analysis.

The necessity to apply a precise machine learning algorithms is recently increased in geographic sciences with respect to the importance of choosing an effective method for data visualization and computation. The solutions to the mentioned above problems are provided by Pandas data frames: for example, optimizing structure of the data, selecting the correct parts from the whole data frame (columns, rows in the data arrays) and plotting. Based on the presented results, the application of Python programming language is strongly recommended in geoscience research as an addition to the traditional GIS methods.

Conflict of Interests: The author declares no conflict of interest.

Funding: This research was funded by the China Scholarship Council (CSC), State Oceanic Administration (SOA), Marine Scholarship of China, Grant Nr. 2016SOA002, Beijing, People's Republic of China.

REFERENCES

- Ansley, C. F. & R. J. Kohn. (1985). Estimation, filtering, and smoothing in state space models with incompletely specified initial conditions. *Annals of Statistics* 13, 1286–1316. [\[CrossRef\]](#)
- Bello-González, J. P., Contreras-Reyes, E. & Arriagada, C. (2018). Predicted path for hotspot tracks off South America since Paleocene times: Tectonic implications of ridge-trench collision along the Andean margin. *Gondwana Research*, 64, 216–234. [\[CrossRef\]](#)
- Boston, B., Moore, G. F., Nakamura, Y. & Kodaira, S. (2017). Forearc slope deformation above the Japan Trench megathrust: Implications for subduction erosion. *Earth and Planetary Science Letters*, 462, 26–34. [\[CrossRef\]](#)
- Björck, A. (1996). Numerical methods for least squares problems. SIAM, Philadelphia. ISBN 0-89871-360-9. [\[CrossRef\]](#)
- Box, G. E. P.; Tiao, G. C. (1992). *Bayesian Inference in Statistical Analysis*. New York: John Wiley and Sons. ISBN 0-471-57428-7. (Section 8.1.1). [\[CrossRef\]](#)
- Dierssen, H. M. & Theberge Jr. A. E. (2014). Bathymetry: History of Seafloor Mapping. *Encyclopedia of Natural Resources*, Taylor & Francis. [\[CrossRef\]](#)
- Fujie, G., Ito, A., Kodaira, S., Takahashi, N., & Kaneda, Y. (2006). Confirming sharp bending of the Pacific plate in the northern Japan trench subduction zone by applying a traveltimes mapping method. *Physics of the Earth and Planetary Interiors*, 157, 72–85. [\[CrossRef\]](#)
- Grand, S. P., Hilst, R. D. van der, & Widiyantoro, S. (1997). Global Seismic Tomography: A Snapshot of Convection in the Earth. *GSA Today*, 7(4), 2–7.
- Lemenkova, P. (2019). Processing Oceanographic Data by Python Libraries NumPy, SciPy And Pandas. *Aquatic Research*, 2(2), 73-91. [\[CrossRef\]](#)
- Ljung, G. M. & Box, G. E. P. (1978). On a Measure of a Lack of Fit in Time Series Models. *Biometrika*. 65 (2): 297–303. [\[CrossRef\]](#)
- Michibayashi, K., Tasaka, M., Ohara, Y., Ishii, T., Okamoto, A., & Fryer, P. (2007). Variable microstructure of peridotite samples from the southern Mariana Trench: Evidence of a complex tectonic evolution. *Tectonophysics*, 444, 111–118. [\[CrossRef\]](#)
- Millman, K. J. & Aivazis, M. (2011). Python for Scientists and Engineers, *Computing in Science & Engineering*, 13, 9-12. [\[CrossRef\]](#)
- Oliphant, T. (2015). *Guide to NumPy* (2 ed.). CreateSpace. ISBN 978-1517300074.
- Oliphant, T. E. (2007). *Python for scientific computing*. Computing in Science & Engineering 9(3), 10-20. [\[CrossRef\]](#)
- Reid, W. D. K., Cuomo, N. J., & Jamieson, A. J. (2018). Geographic and bathymetric comparisons of trace metal concentrations (Cd, Cu, Fe, Mn, and Zn) in deep-sea lysianassoid amphipods from abyssal and hadal depths across the Pacific Ocean. *Deep-Sea Research Part I*, 138, 11–21. [\[CrossRef\]](#)
- Schellart, W. P. (2008). Subduction zone trench migration: Slab driven or overriding-plate-driven? *Physics of the Earth and Planetary Interiors*, 170, 73–88. [\[CrossRef\]](#)
- Seabold, S. & Perktold, J. (2010). Statsmodels: Econometric and statistical modeling with python. *Proceedings of the 9th Python in Science Conference*.
- Smith, W. H. F., & Sandwell, D. T. (1997). Global Sea Floor Topography from Satellite Altimetry and Ship Depth Soundings. *Science*, 277, 1956–1962. [\[CrossRef\]](#)
- Strutz, T. (2016). *Data Fitting and Uncertainty (A practical introduction to weighted least squares and beyond)*. Springer Vieweg. ISBN 978-3-658-11455-8.
- Taira, K., Yanagimoto, D., & Kitagawa, S. (2005). Deep CTD Casts in the Challenger Deep, Mariana Trench. *Journal of Oceanography*, 61, 447–454. [\[CrossRef\]](#)
- Theberge, A. (2008). Thirty years of discovering the Mariana Trench. *Hydro International*, 12, 38–39.
- Timm, N. H. (2007). *Applied Multivariate Analysis*. Springer Science & Business Media, 695 p. ISBN: 978-0-387-95347-2. [\[CrossRef\]](#)