



HAL
open science

Complete Genome Sequences of Five *Acinetobacter baumannii* Phages from Abidjan, Côte d'Ivoire

Christiane Esoh, Jean-Philippe Vernadet, Gilles Vergnaud, Adama Coulibaly, Adèle Kakou-N'Douba, Assanvo S.-P. N'Guetta, Gregory Resch, Christine Pourcel

► **To cite this version:**

Christiane Esoh, Jean-Philippe Vernadet, Gilles Vergnaud, Adama Coulibaly, Adèle Kakou-N'Douba, et al.. Complete Genome Sequences of Five *Acinetobacter baumannii* Phages from Abidjan, Côte d'Ivoire. *Microbiology Resource Announcements*, 2019, 8 (1), <10.1128/MRA.01358-18>. <hal-02170278>

HAL Id: hal-02170278

<https://hal.science/hal-02170278v1>

Submitted on 10 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Complete Genome Sequences of Five *Acinetobacter baumannii* Phages from Abidjan, Côte d'Ivoire

Christiane Essoh,^a Jean-Philippe Vernadet,^b Gilles Vergnaud,^b Adama Coulibaly,^a Adèle Kakou-N'Douba,^c Assanvo S.-P. N'Guetta,^d Gregory Resch,^e Christine Pourcel^b

^aDepartment of Biochemistry-Genetic, School of Biological Sciences, Université Peleforo Gon Coulibaly, Korhogo, Côte d'Ivoire

^bInstitute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Université Paris-Sud, Université Paris-Saclay, Paris, France

^cLaboratory of Bacteriology-Virology, Department of Microbiology, School of Medical Sciences, Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire

^dLaboratory of Genetic, School of Biosciences, Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire

^eDepartment of Fundamental Microbiology, University of Lausanne, Lausanne, Switzerland

ABSTRACT Five bacteriophages of *Acinetobacter baumannii* were isolated from sewage water in Abidjan, Côte d'Ivoire. Phages Aci01-1, Aci02-2, and Aci05 belong to an unclassified genus of the *Myoviridae* family, with double-stranded DNA (dsDNA) genomes, whereas Aci07 and Aci08 belong to the *Fri1virus* genus of the *Podoviridae* family of phages.

Using two clinical strains of *Acinetobacter baumannii*, B09 and Abidjan-46-62, we isolated five lytic bacteriophages from hospital sewage water in Abidjan, Côte d'Ivoire. Five milliliters of water was clarified by centrifugation at $2,500 \times g$ for 20 min and then filtrated through a 0.45- μm filter. Two hundred microliters of sample was then incubated for 20 h in 2 ml LB medium with 20 μl of an overnight bacterial culture, and after centrifugation, the supernatant was spotted onto bacteria in soft LB agar. Phages recovered in lysis zones were amplified as previously described (1). Phages vB_AbaM_B09_Aci01-1 (short name, Aci01-1), vB_AbaM_B09_Aci02-2 (Aci02-2), and vB_AbaM_B09_Aci05 (Aci05) formed small clear plaques, whereas vB_AbaP_46-62_Aci07 (Aci07) and vB_AbaP_B09_Aci08 (Aci08) produced clear plaques with a large halo (Fig. 1A and B). The morphology of phage virions was determined using transmission electron microscopy (Fig. 1C and D). Phage suspensions were adsorbed on a grid coated with carbon (EMS, Hatfield, PA, USA) and then washed with distilled water, followed by staining with 1% uranyl acetate (Sigma, St. Louis, MO, USA) in H_2O for 1 min. Electron micrographs (EM) were taken with a Philips CM100 transmission electron microscope (Thermo Fisher Scientific, Hillsboro, USA) at an acceleration voltage of 80 kV. EM for Aci01-1, Aci02-2, and Aci05 showed a $75 \pm 4\text{-nm}$ head and a $140 \pm 2\text{-nm}$ nonflexible contractile tail, which are characteristic of *Myoviridae*, and EM for Aci07 and Aci08 showed a $60 \pm 2\text{-nm}$ icosahedral capsid with a short tail that is characteristic of *Podoviridae*. For genome purification, bacteriophages were lysed in lysis buffer (10 mM Tris [pH 7.8]; 10 mM EDTA; 10 mM NaCl; 0.5% SDS [wt/vol]) and treated with proteinase K at $50 \mu\text{g ml}^{-1}$ for 2 h at 50°C , followed by one phenol and one chloroform extraction and ethanol precipitation. Each phage genome was sequenced separately in an Illumina MiSeq 300-bp paired-end run with a 900-bp insert library, producing 2.7 to 5.5 million reads (800 to 1,700 Mb), including a variable proportion of reads derived from contaminating bacterial host DNA. Quality control was performed with FastQC version 0.11.5, and reads were trimmed with GeneiousR11 (Biomatters Ltd., Auckland, New Zealand) with default trim parameters. Assembly was performed on 10% of the reads using the Geneious assembler with the medium sensitivity/fast option to a single linear contig. The total amount of reads was then mapped to the contig to produce a

Citation Essoh C, Vernadet J-P, Vergnaud G, Coulibaly A, Kakou-N'Douba A, N'Guetta AS-P, Resch G, Pourcel C. 2019. Complete genome sequences of five *Acinetobacter baumannii* phages from Abidjan, Côte d'Ivoire. *Microbiol Resour Announc* 8:e01358-18. <https://doi.org/10.1128/MRA.01358-18>.

Editor David A. Baltrus, University of Arizona

Copyright © 2019 Essoh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christine Pourcel, christine.pourcel@u-psud.fr.

Received 8 October 2018

Accepted 19 November 2018

Published 3 January 2019




FIG 1 Plaque morphology of phage Aci05 (0.5 mm) (A) and phage Aci07 (2 mm) (B). EM of phage Aci05 (C) and phage Aci07 (D). Bar, 100 nm.

high-quality sequence at a mean coverage of 1,000 \times or more. The presence of direct terminal repeats (DTRs) which induced a higher read coverage allowed for identification of the phage genome ends. Automatic annotation was performed with Rapid Annotations using Subsystems Technology (RAST) (2).

The Aci01-1, Aci02-2, and Aci05 genomes were 103,628 bp, 104,354 bp, and 102,789 bp long with DTRs of 1,184 bp, 1,198 bp, and 1,151 bp, respectively. A comparison was performed by aligning the genomes using MAFFT version 7 (3) as embedded in GeneiousR11. The average nucleotide identity (ANI) among the Aci01-1, Aci02-2, and Aci05 genomes was higher than 90% compared with less than 65% toward phage vB_AbaM_phiAbaA1 from Poland (GenBank accession number [KJ628499](#)) and phage Acibel004 from Belgium (GenBank accession number [KJ473422](#)) (4), two presently unclassified myoviruses (5). The genome size and architecture and the presence of a DTR and tRNAs suggest that these phages may be members of the *Felixounavirinae* subfamily (1, 6).

The Aci07 and Aci08 genomes were 42,330 bp and 42,067 bp long with 397-bp and 410-bp DTRs, respectively. As estimated by eye, the overall gene organization of Aci07 and Aci08 was that of ApiP_P1 (GenBank accession number [MF033350](#)) and AS12 (GenBank accession number [KY268295](#)), respectively, belonging to the genus *Fris1virus* of the *Autographinivirinae* subfamily (7). The ANI among these phages as determined by MAFFT alignment varied from 77% to 86%.

Data availability. The genome sequences of the five phages are available at GenBank under accession numbers [MH800198](#) (Aci01-1), [MH800199](#) (Aci02-2), [MH746814](#) (Aci05), [MH800200](#) (Aci07), and [MH763831](#) (Aci08). The raw read archives have been deposited in the European Nucleotide Archive (ENA) under BioProject accession number [PRJEB28456](#).

ACKNOWLEDGMENTS

This research was funded by grant Resisphage, ANR-13-ASTRID-0011-01, from Direction Générale de l'Armement (DGA) and Agence Nationale de la Recherche (ANR, France). This work was also supported by Program d'Appui Stratégique à la Recherche Scientifique (PASRES) of Côte d'Ivoire. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

This work has benefited from the facilities and expertise of the molecular biology platform of TEFOR and I2BC and of the high-throughput sequencing platform of I2BC. We are thankful to Aurélie Marchet for excellent technical assistance and to Jean Daraspe for his help with the electron microscopy.

REFERENCES

1. Eshoh C, Latino L, Midoux C, Blouin Y, Loukou G, Nguetta SP, Lathro S, Cablanmian A, Kouassi AK, Vergnaud G, Pourcel C. 2015. Investigation of a large collection of *Pseudomonas aeruginosa* bacteriophages collected from a single environmental source in Abidjan, Cote d'Ivoire. *PLoS One* 10:e0130548. <https://doi.org/10.1371/journal.pone.0130548>.
2. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
3. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780. <https://doi.org/10.1093/molbev/mst010>.
4. Merabishvili M, Vandenheuvel D, Kropinski AM, Mast J, De Vos D, Verbeke G, Noben JP, Lavigne R, Vanechoutte M, Pirnay JP. 2014. Characterization of newly isolated lytic bacteriophages active against *Acinetobacter baumannii*. *PLoS One* 9:e104853. <https://doi.org/10.1371/journal.pone.0104853>.
5. Turner D, Ackermann HW, Kropinski AM, Lavigne R, Sutton JM, Reynolds DM. 2017. Comparative analysis of 37 *Acinetobacter* bacteriophages. *Viruses* 10:5. <https://doi.org/10.3390/v10010005>.
6. Henry M, Bobay LM, Chevallereau A, Sausseureau E, Ceysens PJ, Debarbieux L. 2015. The search for therapeutic bacteriophages uncovers one new subfamily and two new genera of *Pseudomonas*-infecting *Myoviridae*. *PLoS One* 10:e0117163. <https://doi.org/10.1371/journal.pone.0117163>.
7. Popova AV, Lavyshe DG, Klimuk EI, Edelstein MV, Bogun AG, Shneider MM, Goncharov AE, Leonov SV, Severinov KV. 2017. Novel FriI-like viruses infecting *Acinetobacter baumannii*—vB_AbaP_AS11 and vB_AbaP_AS12—characterization, comparative genomic analysis, and host-recognition strategy. *Viruses* 9:188. <https://doi.org/10.3390/v9070188>.