



# Easy installation of 1,2,3-triazoles or iodo-1,2,3-triazoles onto indole-fused oxazinones via CuAAC-based MCR in the presence of 18-crown-6

A Mayooufi, M Romdhani-Younes, Y Carcenac, Jérôme Thibonnet

## ► To cite this version:

A Mayooufi, M Romdhani-Younes, Y Carcenac, Jérôme Thibonnet. Easy installation of 1,2,3-triazoles or iodo-1,2,3-triazoles onto indole-fused oxazinones via CuAAC-based MCR in the presence of 18-crown-6. Synthetic Communications, In press, <10.1080/00397911.2019.1611857>. <hal-02172442>

**HAL Id: hal-02172442**

**<https://hal.science/hal-02172442v1>**

Submitted on 3 Jul 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Easy installation of 1,2,3-triazoles or iodo-1,2,3-triazoles onto indole-fused oxazinones *via* CuAAC-based MCR in the presence of 18-crown-6


A. Mayooufi<sup>a,b</sup>, M. Romdhani-Younes<sup>b</sup>, Y. Carcenac<sup>a</sup>, J. Thibonnet<sup>a</sup>

<sup>a</sup>Laboratoire Synthèse et Isolement de Molécules BioActives (SIMBA), EA7502, Université de Tours, Faculté des Sciences et Techniques, Parc de Grandmont, 32 av. Monge, 37200 Tours, France

<sup>b</sup>Laboratoire de Chimie Organique Structurale et Macromoléculaire, Département de Chimie, Faculté des Sciences de Tunis, Campus Universitaire El-Manar, Rue Béchir Salem Belkheiria, 2092 Tunis, Tunisie

## Abstract

An efficient protocol was developed to prepare indole-fused oxazinones using silver nitrate. The latter substrates were subjected to multicomponent reactions in the presence of 18-crown-6, which afforded diverse new heterocycles based on an indole-fused oxazinone-1,2,3-triazole scaffold.




## Key words

click chemistry, iodolactonization, multicomponent reactions, azides, iodo-triazoles

## Introduction


Polycyclic indole scaffolds constitute the structural core of numerous important classes of molecules that display an extensive range of biological and pharmacological properties.<sup>[1]</sup> For example HKI 0231A and HKI 0231B have shown the ability to inhibit 3 $\alpha$ -hydroxysteroid dehydrogenase, which is a key enzyme in the inflammatory process.<sup>[2]</sup> More complex polycyclic indole alkaloids, such as vincristine and vinblastine (antitumor agents),<sup>[3]</sup> have been widely investigated.<sup>[4]</sup> Triazole-bearing indoles (Figure 1) have also shown wide-ranging biological activities, such as compound **A** (anti-adipogenic and anti-dyslipidemic activity),<sup>[5]</sup> molecule **B** (selective inhibitors and inducers of bacterial biofilms),<sup>[6]</sup> compound **C** (antimicrobial activity),<sup>[7]</sup> and compound **D** (antifungal agent).<sup>[8]</sup> In addition, indole-fused oxazinone derivatives are present in a large number of bioactive compounds and some representative members of this family are also depicted in Figure 1. For example, compound **E**,<sup>[9]</sup> compound **F**,<sup>[10]</sup> and compound **G**,<sup>[11]</sup> show anticancer properties. Etodolac **H** is used for

the treatment of arthritis,<sup>[12]</sup> and oxazinoindolones **I** showed anti-inflammatory activity.<sup>[13]</sup> The aforementioned findings (Figure 1) gained our attention because of their great importance in biological sciences, and because these structures have rarely been described in the literature.



**Figure 1.** Selected biologically active indole-fused oxazinones and indoles incorporating the 1,2,3-triazole moiety.


We envisioned that combining a triazole moiety with the indole fused oxazinones scaffold may lead to potentially bioactive molecules. In the present work, we describe an easy way to install 1,2,3-triazole and iodo-triazole moieties onto indole-fused oxazinones through an efficient CuAAC-based multicomponent reaction in the presence of 18-crown-6 (Figure 2).



**Figure 2.** Retrosynthetic strategy for the sequential synthesis of indole-fused oxazinones incorporating the 1,2,3-triazole moiety.

## Results and discussion

To achieve our goals, the appropriate cyclization precursors **3a-b** were obtained in two steps starting from ester indoles **1a-b** (Scheme 1). After *N*-alkylation of the latter with allyl bromide, the resulting indole was saponified to **3a-b** in good yields.




**Scheme 1.** Preparation of acids **3a-b**.

For the iodocyclization reaction, other workers<sup>14</sup> reported that the use of I<sub>2</sub> and NaHCO<sub>3</sub> at 70 °C afforded the products **4a** in 84% yield while using NIS and lutidine gave the desired compound **4a** in 81% yield (Table 1, entries 1 and 2). For our part, we wished to re-examine this iodocyclization reaction. Firstly, we tested the efficacy of different sources of electrophilic iodine, such as ICl and NIS instead of I<sub>2</sub> without changing other factors (Table 1, entries 3 and 4), but we observed a decrease in the yield of the desired product **4a**. We then evaluated the effect of different bases on the iodine-mediated electrophilic cyclization of **3a** (Table 1, entries 5-8). The results showed that sodium hydride was the most effective base at room temperature (Table 1, entry 8). To optimize the reaction conditions, we tested the addition of AgNO<sub>3</sub>, and this effectively improved the conversion to reach 99% of the desired product (Table 1, entry 9). An increase in the quantity of the latter additive from 0.5 to 1 equiv reduced the reaction time from 2 h to 30 min, but the yield was not improved (Table 1, entries 10 and 11). Compound **4b** was also obtained in good yield (95%) under the same conditions. This protocol displayed

higher yields and lower reaction times than that previously reported by Joseph and co-workers.<sup>[14]</sup>

**Table 1.** Optimization of Iodolactonization.




Entry	Electrophile (equiv)	Temp (°C)	Base (equiv)	Solvent (ratio)	Additif (equiv)	Yield <sup>a</sup> (%)
1	I <sub>2</sub> (3)	70	NaHCO <sub>3</sub> (3)	CHCl <sub>3</sub> /H <sub>2</sub> O (1/3)	-	84
2	NIS (2.3)	-20	NaHCO <sub>3</sub> (3)	CH <sub>2</sub> Cl <sub>2</sub>	2,6-lutidine	81
3	ICl (3)	70	NaHCO <sub>3</sub> (3)	CH <sub>2</sub> Cl <sub>2</sub>	-	54
4	NBS (3)	70	NaHCO <sub>3</sub> (3)	CH <sub>2</sub> Cl <sub>2</sub>	-	- <sup>b</sup>
5	I <sub>2</sub> (3)	70	K <sub>2</sub> CO <sub>3</sub> (3)	CHCl <sub>3</sub> /H <sub>2</sub> O (1/3)	-	80
6	I <sub>2</sub> (3)	70	Na <sub>2</sub> CO <sub>3</sub> (3)	CHCl <sub>3</sub> /H <sub>2</sub> O (1/3)	-	80
7	I <sub>2</sub> (3)	70	KOH (3)	CHCl <sub>3</sub> /H <sub>2</sub> O (1/3)	-	80
8	I <sub>2</sub> (3)	25	NaH (2)	CH <sub>2</sub> Cl <sub>2</sub>	-	88
9	I <sub>2</sub> (3)	25	NaH (2)	CH <sub>2</sub> Cl <sub>2</sub>	AgNO <sub>3</sub> (0.5)	99
10	I <sub>2</sub> (3)	25	NaH (2)	CH <sub>2</sub> Cl <sub>2</sub>	AgNO <sub>3</sub> (1)	96 <sup>c</sup>
11	I <sub>2</sub> (3)	25	NaH (2)	CH <sub>2</sub> Cl <sub>2</sub>	AgNO <sub>3</sub> (1.5)	96

<sup>a</sup> Isolated yields after column chromatography.

<sup>b</sup> Starting material, no bromolactonization was observed.


<sup>c</sup> 30 min time of reaction.

Next, we examined the reactivity of the di-iodinated indole **4a** *via* CuAAC-based MCR in order to obtain the desired triazole compound **6a**. Various one-pot multicomponent syntheses of 1,2,3-triazoles starting from organic halides have been reported in the literature. We first examined the reaction of di-iodinated indole **4a** according to a literature procedure for an analogous transformation.<sup>[15]</sup> In our case, the starting di-iodinated indole **4a** was reacted with sodium azide and phenyl acetylene at 50 °C in EtOH/H<sub>2</sub>O for 15 h, in the presence of copper iodide. Under these conditions, no desired product **6a** was formed. Although we tested various conditions by changing the solvents, copper catalysts, base and temperature used for this one-pot two-step sequence, no successful combination was found.




**Scheme 2.** Synthetic routes towards triazole-bearing oxazinoindolones.

To access the desired compound **6a**, a two-step strategy for the click reaction was ~~therefore~~ adopted (Scheme 2). Di-iodinated precursor **4a** was subjected to nucleophilic substitution to introduce the azide group under conditions similar to those already employed for related substrates.<sup>[16]</sup> Preliminary experiments were carried out using sodium azide in MeOH/H<sub>2</sub>O at room temperature. Only a complex mixture of products was obtained, so different reaction conditions were tested. Initially, we screened the solvent used for the substitution of **4a** to form the azide **5a**, and found that the use of aprotic solvents DMF, acetone and MeCN gave the desired product in only low yield (30%). An increase in the quantity of NaN<sub>3</sub> from 3 to 6 equivalents did not improve the yield of the reaction, and in all cases the isolated yield of the desired product **5a** did not exceed 40%. We also found that increasing the temperature from 25 °C to 55 °C gave only a mixture of products and no desired product **5a** was isolated. However, the use of phase transfer agents ~~as additives~~ proved to be crucial to improve the outcome of the reaction (Scheme 3).<sup>[17]</sup> We found that the best phase transfer agent was 18-crown-6 rather than TBAI or TBAF. Further investigations showed that increasing the amount of additive reduced the reaction time but did not give better yields. Significantly, the protocol was also tolerant of the di-iodinated precursor **3b** to furnish the product **4b** in 88% yield.



**Scheme 3.** Optimization of the reaction between indole-fused oxazinones **4a-b** and sodium azide.

This success in the synthesis of **5a-b** led us to investigate cycloaddition (CuAAC) reactions of the azide for building systems bearing a 1,2,3-triazole moiety. Traditional approaches for the synthesis of 1,2,3-triazoles starting from an azide precursor have been reported in the literature.<sup>[18]</sup> A first attempt with CuSO<sub>4</sub>·5H<sub>2</sub>O (10 mol%), sodium ascorbate (20 mol%) and phenylacetylene (1.5 equiv) in toluene/H<sub>2</sub>O (3:1) at room temperature provided the corresponding product with a moderate yield (53%). After testing different solvents, we found that by using chlorinated solvents, such as CH<sub>2</sub>Cl<sub>2</sub> or CHCl<sub>3</sub>, the products **6a-b** could be obtained in higher yields (89% and 82% respectively) (Scheme 4).




**Scheme 4.** 1,3-Dipolar cycloaddition between **5a-b** and phenylacetylene.

~~In a continuation of~~ In keeping with our efforts to examine the reactivity of **5a**, we attempted to synthesize 5-iodotriazole **7** through a simple one-pot cycloaddition/iodination reaction. Initially, we tested this reaction under conditions previously defined by our group (Table 2, entry 1).<sup>[19]</sup> This protocol gave the desired product **7** in 55% yield and with a 27/73 ratio of **6a/7**, respectively. To increase the selectivity of the reaction, we examined a range of solvents, catalysts, and reaction times. The use of I<sub>2</sub> or NBS as an electrophilic trapping reagent did not give better results than with ICl (Table 2, entries 2 and 3). Next, we screened other organic solvents (Table 2, entry 4 and 5). The use of chlorinated solvents, such as CH<sub>2</sub>Cl<sub>2</sub> or CHCl<sub>3</sub>, improved the yield of this one pot reaction but always gave a mixture of **7** and the mono-iodinated compound **6a** (85:15). The addition of catalysts, such as NaI or LiI was unsuccessfully tested (Table 2, entries 6 and 7). We therefore turned our attention to the reaction duration, and

it appeared that extended reaction times increased the yield of the side product **6a** (Table 2, entries 8-11). The best result in terms of yield and selectivity was obtained when the reaction time was limited to 48-50 h (Table 2, entries 8 and 9).

**Table 2.** Optimization of one-pot cycloaddition/iodation reaction.



Entry	Electrophile (equiv)	Solvent	Time0 (h)	Ratio (7/6a) <sup>a</sup>	Yield of 7 (%) <sup>b</sup>
1	ICl (1.2)	THF	60	(73/27)	55
2	NBS (1.2)	THF	60	-	- <sup>c</sup>
3	I <sub>2</sub> (1.2)	THF	60	(71/29)	46
4	ICl (1.2)	CHCl <sub>3</sub>	60	(85/15)	74
5	ICl (1.2)	CH <sub>2</sub> Cl <sub>2</sub>	60	(88/12)	70
6	ICl (1.2))	CHCl <sub>3</sub>	60	(88/12)	72 <sup>d</sup>
7	ICl (1.2)	CHCl <sub>3</sub>	60	(88/12)	74 <sup>e</sup>
8	ICl (1.2)	CHCl <sub>3</sub>	50	(90/10)	77
9	ICl (1.2)	CHCl <sub>3</sub>	48	(93/7)	76
10	ICl (1.2)	CHCl <sub>3</sub>	40	(95/5)	60
11	ICl (1.2)	CHCl <sub>3</sub>	72	(66/34)	53

<sup>a</sup> Ratios were measured *via* <sup>1</sup>H NMR.

<sup>b</sup> Isolated yields.

<sup>c</sup> Degradation.


<sup>d</sup> Addition of 20 mol% of LiI.

<sup>e</sup> Addition of 20 mol% of NaI.

Multicomponent reactions are chemical processes which represent an economically useful way to synthesize a large number of potentially bioactive molecules by combining several reaction steps in a single operation. For example, various MCR syntheses of 1,2,3-triazoles have been published in the past decade starting from organic halides, sodium azide, and terminal alkynes. Our preliminary results showed that 1,2,3-triazoles could be obtained in very good yields only in the presence of 18-crown-6 starting from the corresponding **4a** through the classical two-step route.<sup>[20]</sup> With this knowledge in hand, we re-attempted the one-pot process for the two previously defined steps to prepare the substrate **6a-b** from the organic halides **4a-b**. Pleasingly, this time the multicomponent reaction worked well in the presence of 18-crown-6, and the




desired 1,2,3-triazoles **6a-b** were obtained in 90 and 88% yield respectively. Using the optimized conditions, we then focused on exploring the scope of this reaction with various alkynes and sodium azide. The corresponding indole-fused oxazinone-1, 2, 3-triazole scaffolds **6a-l** were isolated in good to excellent yields. As shown in Scheme 5, both electron-withdrawing and electron-donating groups can be successfully incorporated in the alkyne component, without substantially altering the one-pot process' efficiency.



**Scheme 5.** Synthesis of triazole-bearing oxazinoindolones through a CuAAC one-pot procedure.

We have also shown that iodo-triazole **7** could efficiently be obtained *via* a simple one-pot cycloaddition/iodination reaction starting from azido-indole-fused oxazinones. We supposed that indole-fused oxazinone **4a** could be converted into iodotriazole **7** in the presence of 18-crown-6 *via* a MCR process (Scheme 6). In this context, indole-fused oxazinone **4a**, phenylacetylene, sodium azide and ICl were reacted in the presence of copper iodide, 18-crown-6 and Et<sub>3</sub>N as base. Under these conditions, the desired iodo-triazole **7** was obtained in 76% yield as a major product but mixed with 5% of **6a**. It was found that 18-crown-6 is indispensable for this multicomponent click reaction. This one-pot reaction led to the isolation of the desired iodo-triazole **7** with a better yield than in the two-step sequence (70%).



**Scheme 6.** One-pot azidation/cycloaddition/iodination reaction.

## Conclusions

In summary, we have developed a new protocol to prepare indole-fused oxazinones. We have also demonstrated that these gave access to a novel series of heterocyclic derivatives based on triazole and indole-fused oxazinones, prepared using CuAAC based multicomponent reactions. Further investigations concerning the resulting products **6a-i** are ongoing in our laboratory, and an evaluation of their biological activity is underway at the Lilly platform.

## Experimental

### *General procedure for the synthesis of ethyl 1-Allyl-5-bromo-1H-indole-2-carboxylate (2b)*

In a two-necked round bottom flask, NaH 60 % (44 mg, 1.1 mmol, 1.1 equiv.) was dissolved in DMF (6 mL). Compound **1b** (1 mmol, 1 equiv.) in DMF (3 mL) was added dropwise at 0 °C under argon. The mixture was stirred for 30 min at room temperature. A solution of allyl

bromide (145.2 mg, 1.2 mmol, 1.2 equiv.) in DMF (3 mL) was added dropwise at 0 °C. After 4 h of stirring at room temperature, the mixture was hydrolyzed by aqueous NH<sub>4</sub>Cl (20 mL), extracted by diethyl ether (6 × 10 mL), and the organic phases were washed with brine (3 × 10 mL), dried over MgSO<sub>4</sub> and concentrated under vacuum.

#### *General procedure for the synthesis of indole-2-carboxylic acids 3*

In a round bottom flask, NaOH 12% (240 mg, 6 mmol, 3 equiv.) was slowly added to a solution of indole ester **2a-b** (2 mmol, 1 equiv.) in ethanol (6 mL). The mixture was stirred at 45 °C for 3 h, cooled to 0 °C and acidified with HCl (1 M) to obtain pH = 1. The resulting precipitate was filtered and washed with hexane.

#### *General procedure for the synthesis of iodoindole-fused oxazinones 4*

In a round bottom flask containing carboxylic acid **3** (1.11 mmol) and 25 mL of CH<sub>2</sub>Cl<sub>2</sub> were added sodium hydride (88 mg, 2.22 mmol), iodine (845 mg, 3.33 mmol) and silver nitrate (94 mg, 0.55 mmol). The mixture was stirred overnight at room temperature. The medium was hydrolyzed by saturated solution of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (30 mL) then extracted by dichloromethane (3 × 25 mL). The organic phase was washed with NaCl (3 × 10 mL), dried over MgSO<sub>4</sub> and concentrated under vacuum to give the products **4a-b**. Spectroscopic data of known compound **4a** are identical to those reported previously.<sup>[14]</sup>

#### *General procedure for the synthesis of azides 5*

A mixture of the appropriate **4a-b** (0.76 mmol), sodium azide (148 mg, 2.28 mmol), 18-crown-6 (100 mg, 0.38 mmol) and acetone (20 mL) was stirred at room temperature until completion of the reaction (TLC). The solvents were evaporated under reduced pressure, and the mixture was then poured into water, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3×40 mL), dried (MgSO<sub>4</sub>), and concentrated in vacuum. The crude product was purified by silica gel chromatography using PE/EtOAc (1:4) mixture as eluent.

#### *General procedure for the synthesis of triazoles 6*

**Starting from Azide 5:**

Finely powdered  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (10 mol%) and sodium ascorbate (20 mol%) were slowly added to a stirred solution of **5** (0.8 mmol) and appropriate terminal alkyne (1.2 mmol) in  $\text{H}_2\text{O}/\text{CHCl}_3$  (3:1 = 22.5 mL/7.5 mL) at 0-10 °C. The mixture was then warmed to room temperature and stirred until completion (TLC). The mixture was filtered, concentrated and diluted with water (30 mL). The aqueous layer was extracted with  $\text{CHCl}_3$  (3 x 20 mL). The combined organic layers were washed with  $\text{H}_2\text{O}$  and then with brine, dried over  $\text{MgSO}_4$  and concentrated in vacuum. The crude products were purified by flash chromatography on silica gel ( $\text{EtOAc}/\text{PE}$  = 4:1) to afford the pure triazoles **6a** and **6c-l**.

#### **Starting from bis-iodine 4:**

Sodium azide (148 mg, 2.28 mmol) and 18-crown-6 (100 mg, 0.38 mmol) were added to a solution of compound **4** (0.76 mmol) in  $\text{H}_2\text{O}/\text{CHCl}_3$  (3:1 = 22.5 mL/7.5 mL), and the suspension was stirred for 30 min. The mixture was then degassed at 0°C and terminal alkyne (4.0 mmol), sodium ascorbate (641 mg, 4.0 mmol) and  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (76 mg, 0.4 mmol) were successively added. The mixture was stirred overnight. The reaction was quenched by the addition of an aqueous saturated  $\text{NH}_4\text{Cl}$  solution. After being stirred for further 15 min, the mixture was filtered through a Celite pad. The aqueous phase was extracted with  $\text{EtOAc}$  (3 x 20 mL) and the combined organic layers were washed with brine, dried over  $\text{MgSO}_4$  and concentrated under reduced pressure. The crude material was then purified by flash chromatography on silica gel ( $\text{EtOAc}/\text{PE}$  = 4:1) giving compounds **6a** and **6b**.

#### *General procedure for the synthesis of triazole 7*

##### **Starting from Azide 5:**

A mixture of azide **5a** (1 mmol), terminal alkyne (1.3 mmol),  $\text{Et}_3\text{N}$  (1.2 mmol),  $\text{CHCl}_3$  (100 mL),  $\text{ICl}$  (195 mg, 1.2 mmol), and  $\text{CuI}$  (190 mg, 1 mmol) was stirred at room temperature under an argon atmosphere for 48 h. The solvent was removed by distillation under reduced pressure and the crude residue was purified by flash chromatography over silica gel ( $\text{CHCl}_3/\text{MeOH}$  = 88:12) to afford the pure desired triazole product **7**.

##### **Starting from bis-iodine 4:**

Compound **4a-b** (0.76 mmol) was dissolved in  $\text{CHCl}_3$  (20 mL). Sodium azide (148 mg, 2.28 mmol) and 18-crown-6 (100 mg, 0.38 mmol) were then added to the solution and the suspension

was stirred for 30 min. The mixture was then degassed at 0 °C and terminal alkyne (4.0 mmol), Et<sub>3</sub>N (125 µL, 0.92 mmol), CuI (145 mg, 0.76 mmol) and ICl (148 mg, 0.91 mmol) were successively added. The mixture was stirred at room temperature under an argon atmosphere for 60 h. The reaction mixture was quenched by the addition of an aqueous saturated NH<sub>4</sub>Cl solution and stirred for further 15 min. The mixture was filtered through a pad of Celite. The aqueous phase was extracted with EtOAc (3 x 20 mL) and the combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The crude product was then purified by flash chromatography on silica gel (EtOAc/PE = 4:1).

## Notes

The authors declare no competing financial interest.

## ORCID

Jérôme Thibonnet ID <http://orcid.org/0000-0003-3817-210X>

## References

- [1] (a) Verpoorte, R. Antimicrobially Active Alkaloids. *Alkaloids* **1998**, 397–433. (b) McKay, M. J.; Carroll, A. R.; Quinn, R. J.; Hooper, J. N. A. 1,2-Bis(1*H*-indol-3-yl)ethane-1,2-dione, an Indole Alkaloid from the Marine Sponge *Smenospongia* sp. *J. Nat. Prod.* **2002**, 65, 595–597. doi:10.1021/np010347v
- [2] Kleinwachter, P.; Schlegel, B.; Groth, I.; Hartl, A.; Grafe, U. New Inhibitors of 3  $\alpha$  - Hydroxysteroid Dehydrogenase, 0231A and 0231B from *Streptomyces* sp. HKI 0231. *J. Antibiotics* **2001**, 54, 510–512. doi:10.7164/antibiotics.54.510
- [3] Fernández-Pérez, F.; Almagro, L.; Pedreño, M.A.; GómezRos, L.V. Synergistic and cytotoxic action of indole alkaloids produced from elicited cell cultures of *Catharanthus roseus*. *Pharm. Biol.* **2013**, 51, 304–310. doi:10.3109/13880209.2012.722646

- [4] Kaushik, N. K.; Kaushik, N.; Attri, P.; Kumar, N.; Kim, C. H.; Verma, A. K.; Choi, E. H. Biomedical importance of indoles. *Molecules* **2013**, *18*, 6620–6662. doi:10.3390/molecules18066620
- [5] Rajan, S.; Puri, S.; Kumar, D.; Babu, M. H.; Shankar, K.; Varshney, S.; Srivastava, A.; Gupta, A.; Reddy, M. S.; Gaikad, A. N. Novel indole and triazole based hybrid molecules exhibit potent anti-adipogenic and antidyslipidemic activity by activating Wnt3a/ $\beta$ -catenin pathway. *Eur. J. Med. Chem.* **2018**, *143*, 1345–1360. doi:10.1016/j.ejmech.2017.10.034
- [6] Minvielle, M. J.; Bunders, C. A.; Melander, C. Indole–triazoleconjugates are selective inhibitors and inducers of bacterial biofilms. *Med. Chem. Commun.* **2013**, *4*, 916–919. doi:10.1039/C3MD00064H
- [7] Pawar, K.; Yadav, A.; Prasher, P.; Mishra, S.; Singh, B.; Singh, P.; Komath, S. S. Identification of an indole–triazole–amino acid conjugate as a highly effective antifungal agent. *Med. Chem. Commun.* **2015**, *6*, 1352–1359. doi:10.1039/C5MD00156K
- [8] Reddy, B. S.; Reddy, B. S.; Rao, C.; Subha, M. C. S. Synthesis of some new 1,2,3-triazole derivatives of indole with potential antimicrobial activity. *World J. Pharm. Pharm. Sci.* **2015**, *4*, 1084–1090.
- [9] Naveen, P.; Nikhila, G.; Udayakumar, D.; Pulla, V. K.; Perumal, Y.; Dharmarajan, S. Synthesis of novel 5-[(1,2,3-triazol-4-yl)methyl]-1-methyl-3Hpyridazino[4,5-b]indol-4-one by click reaction and exploration anticancer activity. *Med. Chem. Res.* **2016**, *25*, 135–148. doi:10.1007/s00044-015-1473-y
- [10] D'Ambra, T. E.; Estep, K. G.; Bell, M. R.; Eissenstat, M. A.; Josef, K. A.; Ward, S. J.; Haycock, D. A.; Baizman, E. R.; Casiano, F. M.; Beglin, N. C.; Chippari, S. M.; Grego, J. D.; Kullnig, R. K.; Daley, G. T. Conformationally restrained analogs of pravadoline: nanomolar potent, enantioselective, (aminoalkyl)indole agonists of the cannabinoid receptor. *J. Med. Chem.* **1992**, *35*, 124–135. doi: 10.1021/jm00079a016.
- [11] Ayral-Kaloustian, S.; Zhang, N.; Venkatesan, A. M.; Mansour, T. S.; Nguyen, T. H.; Anderson, J. T. U.S. Pat. Appl. Publ. **2009**, US 20090192147 A1 20090730.
- [12](a) Radwan, M. A. A.; Ragab, E. A.; Sabry, N. M.; El-Shenawy, S. M. Synthesis and biological evaluation of new 3-substituted indole derivatives as potential anti-inflammatory and analgesic agents. *Bioorg. Med. Chem.* **2007**, *15*, 3832–3841. doi:

10.1016/j.bmc.2007.03.024. (b) Sharma, V.; Kumar, P.; Pathak, D. Biological importance of the indole nucleus in recent years: A comprehensive review. *J. Heterocycl. Chem.* **2010**, *47*, 491–502. doi:10.1002/jhet.349

[13] Inaba, S.; Ishizumi, K.; Akatsu, M.; Kume, R.; Mori, K.; Yamamoto, H. Jpn. Tokkyo Koho, **1974**, JP 49004238 B 19740131.

[14](a) Putey, A.; Fournet, G.; Joseph, B. General and Easy Access to 11-Substituted 4-Hydroxy-2,3,4,5-tetrahydro[1,4]diazepino[1,2-*a*]indol-1-one Derivatives. *Synlett* **2006**, 2755–2758. doi: 10.1055/s-2006-950249. (b) Putey, A.; Popowycz, F.; Joseph, B. A Non-Classical Route to 2,3-Diiodoindoles from Indole-2-carboxylic Acids. *Synlett* **2007**, 419–422. doi:10.1055/s-2007-967953

[15](a) Kumar, D.; Patel, G.; Reddy, V. B. Greener and Expedient Synthesis of 1,4-Disubstituted 1,2,3-Triazoles from Terminal Acetylenes and in situ Generated  $\alpha$ -Azido Ketones. *Synlett* **2009**, 399–402. doi:10.1055/s-0028-1087556. (b) Alonso, F.; Moglie, Y.; Radivoy, G.; Yus, M. Copper-Catalysed Multicomponent Click Synthesis of 5-Alkynyl 1,2,3-Triazoles under Ambient Conditions. *Synlett* **2012**, 2179–2182. doi:10.1055/s-0031-1290445. (c) Mukherjee, N.; Ahammed, S.; Bhadra, S.; Ranu, B. C. Solvent-free one-pot synthesis of 1,2,3-triazole derivatives by the ‘Click’ reaction of alkyl halides or aryl boronic acids, sodium azide and terminal alkynes over a Cu/Al<sub>2</sub>O<sub>3</sub> surface under ball-milling. *Green Chem.* **2013**, *15*, 389–397. doi:10.1039/C2GC36521A. (d) Feldman, A. K.; Colasson, B.; Fokin, V. V. One-Pot Synthesis of 1,4-Disubstituted 1,2,3-Triazoles from In Situ Generated Azides. *Org. Lett.* **2004**, *6*, 3897–3899. doi:10.1021/ol048859z. (e) Safa, K. D.; Mousazadeh, H. Synthesis of novel 1,4-disubstituted 1,2,3-triazoles bearing organosilicon-sulfur groups via the click reaction sonocatalyzed by LaCu<sub>x</sub>Mn<sub>1-x</sub>O<sub>3</sub>. *Synth. Commun.* **2016**, *46*, 1595–1604. doi:10.1080/00397911.2016.1217339. (f) Guo, S.; Zhou, Y.; Dai, B.; Huo, C.; Liu, C.; Zhao, Y. CuI/Et<sub>2</sub>NH-Catalyzed One-Pot Highly Efficient Synthesis of 1,4-Disubstituted 1,2,3-Triazoles in Green Solvent Glycerol. *Synthesis* **2018**, 2191–2199. doi:10.1055/s-0036-1591557.

[16](a) Feula, A.; Dhillon, S. S.; Byravan, R.; Sangha, M.; Ebanks, R.; Salih, M. A. H.; Spencer, N.; Male, L.; Magyary, I.; Deng, W.-P.; Müller, F.; Fossey, J. S. Synthesis of azetidines and pyrrolidines *via* iodocyclisation of homoallyl amines and exploration of activity in a zebrafish embryo assay. *Org. Biomol. Chem.* **2013**, *11*, 5083–5093. doi:10.1039/C3OB41007B. (b) Yi, M.; Gu, P.; Kang, X.-Y.; Sun, J.; Li, R.; Li, X.-Q. Total

synthesis of ( $\pm$ )-antofine. *Tetrahedron Lett.* **2014**, 55, 105–107. doi:10.1016/j.tetlet.2013.10.124.

- [17] Patonay, T.; Juhász-Tóth, E.; Bényei, A. Base-Induced Coupling of  $\alpha$ -Azido Ketones with Aldehydes—An Easy and Efficient Route to Trifunctionalized Synthons 2-Azido-3-hydroxy Ketones, 2-Acylaziridines, and 2-Acylspiroaziridines. *A. Eur. J. Org. Chem.* **2002**, 285–295. doi:10.1002/1099-0690(20021)2002:2<285::AID-EJOC285>3.0.CO;2-J.
- [18](a) Meldal, M.; Tornøe, C. W. *Chem. Rev.* **2008**, 108, 2952–3015. doi:10.1021/cr0783479. (b) Hein, J. E.; Fokin, V. V. *Chem. Soc. Rev.* **2010**, 39, 1302–1315. doi:10.1039/B904091A. (c) Hassan, S.; Müller, T. J. J. Multicomponent Syntheses based upon Copper-Catalyzed Alkyne-Azide Cycloaddition. *Adv. Synth. Catal.* **2015**, 357, 617–666. doi:10.1002/adsc.201400904. (d) Baltus, C. B.; Jorda, R.; Marot, C.; Berk, K.; Bazgier, V.; Krystof, V.; Prié, G.; Viaud-Massuard, M. C. Synthesis, biological evaluation and molecular modeling of a novel series of 7-azaindole based tri-heterocyclic compounds as potent CDK2/Cyclin E inhibitors. *Eur. J. Org. Chem.* **2016**, 701–719. doi:10.1016/j.ejmech.2015.12.023. (e) Camp, C.; Dorbes, S.; Picard, C.; Benoist, E. Efficient and tunable synthesis of new polydentate bifunctional chelating agents using click chemistry. *Tetrahedron Lett.* **2008**, 49, 1979–1983. doi:10.1016/j.tetlet.2008.01.086. (f) Mishra, K. B.; Tiwari, V. K. Click Chemistry Inspired Synthesis of Morpholine-Fused Triazoles. *J. Org. Chem.* **2014**, 79, 5752–5762. doi:10.1021/jo500890w. (g) Shamim, A.; Vasconcelos, S. N. S.; De Oliveira, I. M.; Reis, J. S.; Pimenta, D. C.; Zukerman-Schpector, J.; Stefani, H. A. Synthesis of Glycosyl Azides and Their Applications Using CuAAC Click Chemistry to Generate Bis- and Tris(triazolyl)glycosyl Derivatives. *Synthesis* **2017**, 5183–5196. doi:10.1055/s-0036-1589090. (h) Martinelli, M.; Milcent, T.; Onger, S.; Crousse, B. Synthesis of new triazole-based trifluoromethyl scaffolds. *Beilstein J. Org. Chem.* **2008**, 4 (19). doi:10.3762/bjoc.4.19.
- [19] Carcenac, Y.; David-Quillot, F.; Abarbri, M.; Duchêne, A.; Thibonnet, J. Palladium-Catalyzed Cross-Coupling of 1,4-Disubstituted 5-Iodo-1,2,3-triazoles with Organotin Reagents. *Synthesis* **2013**, 633–638. doi:10.1055/s-0032-1318112.
- [20] Mayooufi, A.; Romdhani-Younes, M.; Thibonnet, J. *SynOpen* **2018**, 2, 298–305. doi:10.1055/s-0037-1610399.