



Cardiolipin-Based Lipopolyplex Platform for the Delivery of Diverse Nucleic Acids into Gram-Negative Bacteria

Federico Perche, Tony Le Gall, Tristan Montier, Chantal Pichon, Jean-Marc Malinge

► To cite this version:

Federico Perche, Tony Le Gall, Tristan Montier, Chantal Pichon, Jean-Marc Malinge. Cardiolipin-Based Lipopolyplex Platform for the Delivery of Diverse Nucleic Acids into Gram-Negative Bacteria. *Pharmaceuticals*, 2019, 12 (2), pp.81. 10.3390/ph12020081 . hal-02172973

HAL Id: hal-02172973

<https://hal.science/hal-02172973>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Article

Cardiolipin-Based Lipopolyplex Platform for the Delivery of Diverse Nucleic Acids into Gram-Negative Bacteria

Federico Perche ¹, Tony Le Gall ² , Tristan Montier ² , Chantal Pichon ¹ and Jean-Marc Malinge ^{1,*}

¹ Centre de Biophysique Moléculaire, UPR4301 CNRS, Rue Charles Sadron, CEDEX 02, 45071 Orléans, France; federico.perche@cnrs-orleans.fr (F.P.); chantal.pichon@cnrs.fr (C.P.)

² Unité INSERM 1078, Faculté de Médecine, Université de Bretagne Occidentale, Université Européenne de Bretagne, 22 avenue Camille Desmoulins, CEDEX 3, 29238 Brest, France; tony.legall@univ-brest.fr (T.L.G.); tristan.montier@univ-brest.fr (T.M.)

* Correspondence: jean-marc.malinge@cnrs.fr; Tel.: +33-238-255-544

Received: 30 April 2019; Accepted: 24 May 2019; Published: 28 May 2019



Abstract: Antibiotic resistance is a growing public health concern. Because only a few novel classes of antibiotics have been developed in the last 40 years, such as the class of oxazolidinones, new antibacterial strategies are urgently needed (Coates, A.R. et al., 2011). Nucleic acid-based antibiotics are a new type of antimicrobials. However, free nucleic acids cannot spontaneously cross the bacterial cell wall and membrane; consequently, their intracellular delivery into bacteria needs to be assisted. Here, we introduce an original lipopolyplex system named liposome polymer nucleic acid (LPN), capable of versatile nucleic acid delivery into bacteria. We characterized LPN formed with significant therapeutic nucleic acids: 11 nt antisense single-stranded (ss) DNA and double-stranded (ds) DNA of 15 and 95 base pairs (bp), 9 kbp plasmid DNA (pDNA), and 1000 nt ssRNA. All these complexes were efficiently internalized by two different bacterial species, i.e., *Escherichia coli* and *Pseudomonas aeruginosa*, as shown by flow cytometry. Consistent with intracellular delivery, LPN prepared with an antisense oligonucleotide and directed against an essential gene, induced specific and important bacterial growth inhibition likely leading to a bactericidal effect. Our findings indicate that LPN is a versatile platform for efficient delivery of diverse nucleic acids into Gram-negative bacteria.

Keywords: bacterial delivery; oligonucleotide; RNA; pDNA; ssDNA; dsDNA

1. Introduction

Antibiotic resistance is becoming a threat to global human health both in high- and low-income countries, resulting in a growing number of infections and increased mortality, especially due to Gram-negative bacteria [1–3]. The concept of using small nucleic acids as therapeutics to fight antibiotic resistant bacteria is very promising. Different types of strategies can be developed depending on the nature, form, and length of the nucleic acid to be delivered. Chemically synthesized ss- and dsDNA oligonucleotides can be used in order to target either mRNA (antisense strategy) [4] or transcription factor proteins (decoy strategy) [5], with the aim of specifically inhibiting critical bacterial gene expression. Aptamers and CRISPR-Cas nucleases have also recently emerged as potential antibacterial strategies [6–8]. Moreover, functional studies on non-coding bacterial small RNAs have recently highlighted their role in the regulation of genes associated with multiresistance [9] and opened the way to exploit ssRNA molecules as potential antimicrobial molecules [10]. RNA can be delivered or produced in bacteria by pDNA used as expression vector [11].

Because the cell wall of bacteria strongly impedes the penetration of any nucleic acids into the bacterial cytoplasm, a single versatile system suitable for efficient delivery of different types of nucleic acid would be particularly useful. To our knowledge, such a universal platform has not yet been reported.

Indeed, various delivery systems have been used, until now, to condense, protect, and carry nucleic acids into bacteria. Antisense oligodeoxynucleotides (ASOs) were mainly conjugated to cell-penetrating peptides (CPP) [4,12–14]. However, the CPP-dependent approach showed several limitations, including the requirement of designing derivatives when resistance to delivery arises from mutations occurring in bacterial transporter proteins [15]. Other strategies involve formulations with anionic liposomes for delivery of ASO [16] and DNA decoy oligonucleotide [17]. A more recent strategy uses lipopolyplexes with ASO being first complexed with polyethylenimine (PEI) before encapsulation of the PEI/ASO polyplex into anionic liposomes, yielding efficient delivery into Gram-negative bacteria [18–21].

Our group has developed a ternary delivery system named liposome polymer nucleic acid (LPN) for mRNA vaccine delivery [22–25]. LPN is relatively simple to prepare and stable in physiological fluids. We used them for the delivery of pDNA [26], mRNA [23–25], and siRNA [27] into mammalian cells. The two-step complexation protocol of LPN preparation results in the formation of cationic polymer/nucleic acid complexes (polyplexes) of similar size and charge—with different types of nucleic acids—before encapsulation into anionic liposomes, and is required to avoid inactivation of polyplexes in serum [28]. We mixed cationic polyplexes with anionic liposomes aiming to produce near-neutral nanoparticles for further in vivo antibacterial applications, as such nanoparticles are expected to show prolonged blood circulation and therefore higher delivery activity [29–31]. Indeed, neutral nanoparticles show negligible protein adsorption and decreased complement activation as compared to positive and negative ones [29,30,32].

Whereas the delivery of nucleic acids into bacteria has been developed with one formulation per nucleic acid type limiting the development of antibacterial nucleic acid therapeutics, we herein designed a novel LPN platform for the delivery of nucleic acids of diverse size (from 11 nt to 9514 bp) and nature (DNA or RNA) into bacteria.

We tested our LPN strategy on two model species of infectious bacteria, i.e., *Escherichia coli* and *Pseudomonas aeruginosa*. Most of antibiotic resistant infections worldwide are mainly due to *E. coli* [2]. *E. coli* is a Gram-negative bacteria responsible for causing severe foodborne infections which can lead to life-threatening hemolytic uremic syndrome [2]. Moreover, *E. coli* strains resistant to conventional antibiotics have an increasing prevalence in hospital-acquired infections [33,34], making *E. coli* a suitable model for nucleic acid delivery. *P. aeruginosa* is another serious cause of human healthcare-associated infections, with a high prevalence of multidrug-resistant strains and chronic infection in cystic fibrosis patients [35,36].

The present study reports the delivery of various forms of potential nucleic acid therapeutics into bacteria using novel LPNs as shown by flow cytometry. Moreover, antibacterial activity data support the therapeutic potential of LPN–ASO.

2. Results

2.1. Preparation and Characterization of LPN

Lipopolyplexes prepared with anionic liposomes composed of egg phosphatidylcholine (EPC)/dimyristoylphosphatidylglycerol (DMPG)/distearoyl-glycero-phosphoethanolamine-polyethyleneglycol (DSPE-PEG) liposomes encapsulating PEI polyplexes have been reported to deliver ASOs into bacteria [18–21]. Instead of using EPC/DMPG/DSPE-PEG liposomes to encapsulate the PEI polyplexes, we used a more simple mix of two components i.e., distearoyl-glycero-phosphocholine (DSPC) with the negatively charged phospholipid cardiolipin (CL) [37,38]. This choice was driven by the fact that CL interacts with the membrane of Gram-negative bacteria [39].

LPNs are especially suited for the versatile delivery of nucleic acids in eukaryotic cells [22]. Nucleic acids are first complexed with the established cationic polymer 25 kDa PEI, which has already been used in clinical trials [40], into polyplexes before mixing with anionic liposomes to form LPN (Figure 1).




Figure 1. Two-step preparation of liposome polymer nucleic acid (LPN).

Polyplexes size increased along with nucleic acid molecular weight: from 100–120 nm for 11nt ss ASO, 15 and 95 bp ds DNA to 166 nm for 1000 nt RNA and 214 nm for 9514 bp pDNA (Table 1). Entrapping the polyplexes increased the liposome diameter from 100 nm for liposome alone to 140–180 nm for LPN (Table 2) and resulted in near-neutral zeta potentials, physico-chemical features previously observed [24,41,42].

Table 1. Size and zeta potential of polyethylenimine (PEI) polyplexes.

Nucleic Acid	Size (nm)	Zeta Potential (mV)
11 nt ssASO	100 ± 4	8 ± 1.5
15 bp dsDNA	120 ± 8	10 ± 3
95 bp dsDNA	100 ± 7	6.2 ± 1.7
9 kbp pDNA	214 ± 18	9 ± 4
1000 nt RNA	166 ± 10	13 ± 2

Table 2. Size and zeta potential of LPN.

Nucleic Acid	Size (nm)	PDI	Zeta Potential (mV)
11 nt ssASO	140 ± 8	0.20 ± 0.02	6.1 ± 4.7
15 bp dsDNA	150 ± 12	0.26 ± 0.04	5 ± 3
95 bp dsDNA	157 ± 9	0.24 ± 0.04	9.7 ± 4.7
9 kbp pDNA	174 ± 10	0.40 ± 0.08	0.5 ± 0.4
1000 nt RNA	156 ± 15	0.27 ± 0.03	7.4 ± 3.7

PDI: polydispersity index.

For DNA complexes, the size of LPN increased along with the molecular weight: 140/8 nm for ssASO DNA, 157/9 nm for 95 bp DNA oligonucleotide and 174/10 nm for 9514 bp pDNA. The size of RNA LPN was 156/15 nm, a value between that of 95 bp DNA and pDNA. Concerning the charge, all LPNs had a neutral to slightly positive charge (0.5–7 mV) (Table 2). Gel retardation electrophoresis confirmed the condensation of nucleic acids with no detectable free nucleic acid for ssASO DNA, 95 bp dsDNA, pDNA, and ssRNA (Figure 2A).

The morphology of LPN was determined by electron microscopy (Figure 2B) and was found in accordance with previous studies, with a dense polyplex surrounded by a liposome [42,43].




Figure 2. Characterization of LPN: (a) Gel retardation of nucleic acids either free or LPN-complexed. (b) Morphology of LPN–ASO complexes imaged by electron microscopy, scale bar represents 100 nm.

2.2. Delivery

Next, we evaluated nucleic acid delivery into bacteria using flow cytometry. In order to exclusively consider the intracellular signal of the FITC-labeled nucleic acids, extracellular fluorescence of the labeled nucleic acids into bacteria was quenched with trypan blue [23,44,45]. LPNs were able to deliver the different nucleic acids, irrespective of the size; indeed, ssASO DNA, 15 bp and 95 bp dsDNA, and 9514 bp pDNA were delivered into nearly 50% of *E. coli* and *P. aeruginosa* cells, with ASO delivery into cells having the highest efficacy (Figure 3). In both bacterial species, the percentage of transfected cells was lower with RNA, i.e., 15% and 25% in *E. coli* and *P. aeruginosa*, respectively.




Figure 3. Nucleic acid delivery into bacteria. Delivery of LPN prepared with FITC-labeled nucleic acids was evaluated by flow cytometry after 2 h incubation at 37 °C with bacteria.

2.3. Antibacterial Activity

To evaluate the efficiency of LPN in delivering ASO endowed with effect on bacteria growth inhibition, we delivered ASO targeting the acyl carrier protein (AcpP) mRNA whose sequence is well conserved between *E. coli* and *P. aeruginosa* species, with such mRNA targeting yielding effective growth inhibition as previously observed [14,46–50]. Delivery of 11 nt ssASO targeting AcpP mRNA reduced significantly the growth of *E. coli* (Figure 4A) and *P. aeruginosa* (Figure 4B). Such growth delay inhibition lasted at least 20 h (not shown). Growth decrease was no longer observed when LPNs were prepared with a single-stranded control ASO, confirming the specificity of AcpP mRNA targeting. In our experimental conditions, we also checked that cell viability was expectedly reduced when ampicillin was present in the growth medium.




Figure 4. Antibacterial activity of LPN. The growth of *E. coli* (a) and *P. aeruginosa* (b) after treatment with LPN–ASO complexes (1 μ M ASO) or ampicillin (AMP, 150 μ g/mL) was monitored for 5 h at 37 °C.

2.4. Eukaryotic Safety

Next, we evaluated the toxicity of LPN–ASO towards mammalian cells. We observed that LPN–ASO was not toxic towards these eukaryotic cells with no loss in cell viability at the dose used on bacteria (Figure 5). Between 1 and 4 μ M ASO, we have observed that the absence of toxicity was

noteworthy here that these antibacterial effects were safe, specific and relevant, since they could be obtained for doses (of both the cargo and its vehicle) that were found to be fully safe for eukaryotic cells.

We observed that the uptake of 1000 nt RNA by both bacterial species was less important as compared to ASO and the different DNA species, although the extent of complexation of all nucleic acids considered was identical. It is plausible that unlike ssASO, which contains phosphorothioate backbone modifications for resistance to exonuclease activity, chemically unmodified ssRNA was partly digested outside and/or once inside bacteria, resulting in lower intracellular fluorescence. Both dsDNA duplex of 15 and 95 bp (which possessed three phosphorothioate linkages at each DNA end) showed identical delivery efficacy. This observation is of great interest in the search for efficient decoys as the length of consensus sequence recognized by bacterial targeted transcription factors is variable, and depends on the neighboring nucleotide sequence. In addition, the formulation reported herein could also be used to deliver circular DNA ds duplex (DNA minicircle), designed recently to trap several transcription factors for increased decoy strategy efficacy [58].

4. Materials and Methods

4.1. Reagents

All reagents were purchased from Sigma-Aldrich (St. Quentin Fallavier, France) unless otherwise stated. DSPC (1,2-distearoyl-*sn*-glycero-3-phosphocholine), DMPG (1,2-dimyristoyl-*sn*-glycero-3-phospho-(1'-rac-glycerol), CL (cardiolipin), NBD-PE (1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine-*N*-(7-nitro-2-1,3-benzoxadiazol-4-yl) (ammonium salt), Rhod-PE (1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine-*N*-(lissamine rhodamine B sulfonyl), were from Avanti Polar Lipids (Alabaster, AL, USA). Chemical structures of polymer and lipids used are provided in Figure A1.

4.2. Bacteria

E. coli DH5 α (ATCC[®] 53868) was obtained from Thermo Fisher (Les Ulis, France). *P. aeruginosa* (ATCC[®] 10145) was obtained from the Pasteur Institute (Paris, France).

4.3. Nucleic Acids

ASOs with a phosphorothioate backbone were synthesized by Eurogentec (Angers, France). The sequence of the two ASOs was EC AcpP: C*T*T*C*G*A*T*A*G*T*G against *E. coli* [15] and PA AcpP: C*T*T*C*G*A*T*G*G*T*G against *P. aeruginosa* [59], which are complementary to the acyl carrier protein mRNA gene (bases 6–16 of the mRNA coding region) [46]. A control mismatch designated as control ASO (TCTCAGATGGT) was also used [15]. The pTG11033 pDNA (9514 bp) was a gift from Transgène S.A., Strasbourg, France [23]. Luciferase mRNA (1000 nt) was produced by in vitro transcription as in [23]. Nucleic acids were labeled with FITC (Fluorescein Isothiocyanate) using a commercial kit (Label IT[®] Nucleic Acid Labeling Kits, Mirus Bio, Madison, WI, USA).

4.4. Liposome Preparation

Liposomes were prepared by film hydration as in [60]. Liposomes contained 42% CL and 58% DSPC (molar percentages). Chloroform solutions of lipids were evaporated in a rotary evaporator to form lipid films. The latter were hydrated with 10 mM pH 7.4 HEPES at a final lipid concentration of 5.4 mM. Liposomes were then sonicated for 15 min at 20 °C at 37 kHz using an ultrasonic bath (Fischer Bioblock Scientific, Illkirch, France).

4.5. Preparation of Lipopolyplexes

Lipopolyplexes were prepared according to a procedure modified from [23,25]. Instead of mixing polyplexes with a dried lipid film followed by sonication and extrusion, we mixed polyplexes with liposomes in HEPES solution. Briefly, to prepare lipopolyplexes, nucleic acids were mixed with branched PEI of 25kDa in HEPES 10 mM pH 7.4 at a weight ratio (PEI/nucleic acid) of 6 and at room

temperature for 20 min, to allow for polyplex formation. Note that PEI solution was added to nucleic acid solution (and not the opposite), as the order of addition is important for LPN formation [61]. Liposomes were then added to polyplexes using 2 μ L of liposome suspension per μ g of nucleic acid.

4.6. LPN Characterizations

Size and zeta potential of LPNs were determined by DLS using an SZ-100 nanoparticle analyzer (Horiba, Longjumeau, France). LPNs were diluted in 10 mM pH 7.4 HEPES 40 mM NaCl and diameter and zeta potential were measured at 25 °C. Complexation of nucleic acids was validated by gel shift assays: 2 μ g of free or LPN-complexed nucleic acids were run on 1% agarose-formaldehyde gels containing ethidium bromide. Gels were imaged using a Gene Flash imager (Syngene, Cambridge, UK). Transmission electronic microscopy samples were prepared according to the technique of negative staining using uranyl acetate. To this end, 5 μ L of LPN solution in HEPES buffer was deposited on a carbon-coated copper grid for 5 min, and then adsorbed with filter paper. Then, 5 μ L of uranyl acetate 2% in endonuclease-free water was deposited on the grid for 10 s and then adsorbed. Samples were dried at room temperature for 20 min before TEM observation. LPN structure was analyzed using a Philips CM20/STEM electron microscope operating at 50 kV (Centre de Microscopie Electronique, Université d'Orléans, France).

4.7. Flow Cytometry

Cell suspensions were analyzed with a FORTESSA X20 flow cytometer (Beckton Dickinson, Franklin Lakes, NJ, USA). The extracellular fluorescence of the NBD-PE lipid present in the liposomes or the FITC conjugated to the nucleic acid was quenched using trypan blue as in [23,44,45]. The cell-associated fluorescence was then measured with a flow cytometer (FACSsort; Becton Dickinson, Franklin Lakes, New Jersey) with $\lambda_{\text{ex}} = 488$ nm; $\lambda_{\text{em}} = 530 \pm 30$ nm. The fluorescence intensity was expressed as the mean fluorescence intensity of 10,000 events.

4.8. Bacterial Growth Assay

Single-colony overnight cultures were diluted to 5 10^6 cfu/mL in LB broth. Aliquots (100 μ L) of this culture were transferred to 96-well plates. Aliquots (100 μ L) of treatments in LB broth were then added to the wells at a final ASO concentration of 1 μ M. Plates were grown at 37 °C with shaking, and bacterial growth was monitored by absorbance measurements at 600 nm every 15 min using a Victor I spectrophotometer (1420 Multilabel Counter Victor, Wallac, PerkinElmer, Courtaboeuf, France). As a control for growth inhibition, bacteria were treated with ampicillin at 150 μ g/mL.

4.9. Cytotoxicity Towards Eukaryotic Cells

DC2.4 murine immortalized DC cells were a gift from Kenneth L. Rock [62], and were grown at 37 °C in a humidified atmosphere containing 5% CO₂ in RPMI1640 medium supplemented with 10% heat-inactivated fetal bovine serum, 100 units/mL of penicillin and 100 μ g/mL of streptomycin (Fischer Bioblock, Illkirch, France). Cells were mycoplasma-free as evidenced by MycoAlert Mycoplasma Detection Kit (Lonza, Levallois Perret, France). Cytotoxicity was evaluated performing an MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay as in [23].

5. Conclusions

Our results indicate that LPN is a safe multipotent system efficient to deliver a wide range of nucleic acids into Gram-negative bacteria. Indeed, we have delivered DNA and RNA molecules of varying size and nature: RNA (1000 nt), pDNA (9514 bp), linear dsDNA (15 bp and 95 bp), and ASO (11 nt). This broad spectrum activity supports the potential of the LPN platform to treat bacterial infection by delivery of specific ASO or small RNA in different bacterial species.

Author Contributions: Conceptualization: F.P. and J.M.M.; methodology: F.P. and T.L.G.; validation and formal analysis: all authors; writing: F.P., J.M.M., T.L.G., C.P.; funding acquisition: F.P., J.M.M.

Funding: This research was funded by CNRS and the University of Orléans.

Acknowledgments: We are grateful to Bertrand Castaing for providing access to the Nano S zetasizer and to Geoffrey Casas for helpful discussions. We also thank Audrey Sauldubois from the Centre de Microscopie Electronique platform (Université d'Orléans) for TEM studies.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A




Figure A1. Chemical structures of polymer and lipids. PEI (polyethylenimine), DSPC (1,2-distearoyl-sn-glycero-3-phosphocholine), NBD-PE (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(7-nitro-2,1,3-benzoxadiazol-4-yl) (ammonium salt)).

References

- Coates, A.R.; Halls, G.; Hu, Y. Novel classes of antibiotics or more of the same? *Br. J. Pharmacol.* **2011**, *163*, 184–194. [[CrossRef](#)]
- Nambiar, S.; Laessig, K.; Toerner, J.; Farley, J.; Cox, E. Antibacterial drug development: Challenges, recent developments, and future considerations. *Clin. Pharmacol. Ther.* **2014**, *96*, 147–149. [[CrossRef](#)] [[PubMed](#)]
- Organization, W.H. *Antimicrobial Resistance: Global Report on Surveillance*; World Health Organization: Geneva, Switzerland, 2014.
- Xue, X.-Y.; Mao, X.-G.; Zhou, Y.; Chen, Z.; Hu, Y.; Hou, Z.; Li, M.-K.; Meng, J.-R.; Luo, X.-X. Advances in the delivery of antisense oligonucleotides for combating bacterial infectious diseases. *Nanomed. Nanotechnol. Biol. Med.* **2018**, *14*, 745–758. [[CrossRef](#)] [[PubMed](#)]
- McArthur, M.; Bibb, M.J. Manipulating and understanding antibiotic production in *Streptomyces coelicolor* A3 (2) with decoy oligonucleotides. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 1020–1025. [[CrossRef](#)]
- Bikard, D.; Barrangou, R. Using CRISPR-Cas systems as antimicrobials. *Curr. Opin. Microbiol.* **2017**, *37*, 155–160. [[CrossRef](#)] [[PubMed](#)]
- Hong, K.L.; Sooter, L.J. Single-stranded DNA aptamers against pathogens and toxins: Identification and biosensing applications. *BioMed Res. Int.* **2015**, *2015*. [[CrossRef](#)]
- Davydova, A.; Vorobjeva, M.; Pyshnyi, D.; Altman, S.; Vlassov, V.; Venyaminova, A. Aptamers against pathogenic microorganisms. *Crit. Rev. Microbiol.* **2016**, *42*, 847–865. [[CrossRef](#)]
- Law, C.O.K.; Huang, C.; Pan, Q.; Lee, J.; Hao, Q.; Chan, T.-F.; Lo, N.W.S.; Ang, I.L.; Koon, A.; Ip, M. A Small RNA Transforms the Multidrug Resistance of *Pseudomonas aeruginosa* to Drug Susceptibility. *Mol. Ther.-Nucleic Acids* **2019**, *16*, 218–228. [[CrossRef](#)]

10. Chan, H.; Ho, J.; Liu, X.; Zhang, L.; Wong, S.H.; Chan, M.T.; Wu, W.K. Potential and use of bacterial small RNAs to combat drug resistance: A systematic review. *Infect. Drug Res.* **2017**, *10*, 521. [[CrossRef](#)] [[PubMed](#)]
11. Nakashima, N.; Tamura, T. Gene silencing in *Escherichia coli* using antisense RNA s expressed from doxycycline-inducible vectors. *Lett. Appl. Microbiol.* **2013**, *56*, 436–442. [[CrossRef](#)]
12. Santos, R.S.; Figueiredo, C.; Azevedo, N.F.; Braeckmans, K.; De Smedt, S.C. Nanomaterials and molecular transporters to overcome the bacterial envelope barrier: Towards advanced delivery of antibiotics. *Adv. Drug Deliv. Rev.* **2018**, *136*, 28–48. [[CrossRef](#)] [[PubMed](#)]
13. Abushahba, M.F.; Mohammad, H.; Thangamani, S.; Hussein, A.A.; Seleem, M.N. Impact of different cell penetrating peptides on the efficacy of antisense therapeutics for targeting intracellular pathogens. *Sci. Rep.* **2016**, *6*, 20832. [[CrossRef](#)]
14. Daly, S.M.; Sturge, C.R.; Marshall-Batty, K.R.; Felder-Scott, C.F.; Jain, R.; Geller, B.L.; Greenberg, D.E. Antisense Inhibitors Retain Activity in Pulmonary Models of *Burkholderia* Infection. *ACS infect. Dis.* **2018**, *4*, 806–814. [[CrossRef](#)]
15. Puckett, S.E.; Reese, K.A.; Mitev, G.M.; Mullen, V.; Johnson, R.C.; Pomraning, K.R.; Mellbye, B.L.; Tilley, L.D.; Iversen, P.L.; Freitag, M. Bacterial resistance to antisense peptide phosphorodiamidate morpholino oligomers. *Antimicrob. Agents Chemother.* **2012**, *56*, 6147–6153. [[CrossRef](#)] [[PubMed](#)]
16. Fillion, P.; Desjardins, A.; Sayasith, K.; Lagacé, J. Encapsulation of DNA in negatively charged liposomes and inhibition of bacterial gene expression with fluid liposome-encapsulated antisense oligonucleotides. *Biochim. Biophys. Acta (BBA)-Biomembr.* **2001**, *1515*, 44–54. [[CrossRef](#)]
17. González-Paredes, A.; Sitia, L.; Ruyra, A.; Morris, C.J.; Wheeler, G.N.; McArthur, M.; Gasco, P. Solid lipid nanoparticles for the delivery of anti-microbial oligonucleotides. *Eur. J. Pharm. Biopharm.* **2019**, *134*, 166–177. [[CrossRef](#)] [[PubMed](#)]
18. Meng, J.; Bai, H.; Jia, M.; Ma, X.; Hou, Z.; Xue, X.; Zhou, Y.; Luo, X. Restoration of antibiotic susceptibility in fluoroquinolone-resistant *Escherichia coli* by targeting *acrB* with antisense phosphorothioate oligonucleotide encapsulated in novel anion liposome. *J. Antibiot.* **2012**, *65*, 129. [[CrossRef](#)] [[PubMed](#)]
19. Meng, J.; He, G.; Wang, H.; Jia, M.; Ma, X.; Da, F.; Wang, N.; Hou, Z.; Xue, X.; Li, M. Reversion of antibiotic resistance by inhibiting *mecA* in clinical methicillin-resistant *Staphylococci* by antisense phosphorothioate oligonucleotide. *J. Antibiot.* **2015**, *68*, 158. [[CrossRef](#)]
20. Meng, J.; Wang, H.; Hou, Z.; Chen, T.; Fu, J.; Ma, X.; He, G.; Xue, X.; Jia, M.; Luo, X. Novel anion liposome-encapsulated antisense oligonucleotide restores susceptibility of methicillin-resistant *Staphylococcus aureus* and rescues mice from lethal sepsis by targeting *mecA*. *Antimicrob. Agents Chemother.* **2009**, *53*, 2871–2878. [[CrossRef](#)]
21. Wang, H.; Meng, J.; Jia, M.; Ma, X.; He, G.; Yu, J.; Wang, R.; Bai, H.; Hou, Z.; Luo, X. *oprM* as a new target for reversion of multidrug resistance in *Pseudomonas aeruginosa* by antisense phosphorothioate oligodeoxynucleotides. *FEMS Immunol. Med. Microbiol.* **2010**, *60*, 275–282. [[CrossRef](#)]
22. Midoux, P.; Pichon, C. Lipid-based mRNA vaccine delivery systems. *Expert Rev. Vaccines* **2015**, *14*, 221–234. [[CrossRef](#)]
23. Perche, F.; Benvegna, T.; Berchel, M.; Lebegue, L.; Pichon, C.; Jaffrès, P.-A.; Midoux, P. Enhancement of dendritic cells transfection in vivo and of vaccination against B16F10 melanoma with mannosylated histidylated lipopolyplexes loaded with tumor antigen messenger RNA. *Nanomed. Nanotechnol. Biol. Med.* **2011**, *7*, 445–453. [[CrossRef](#)] [[PubMed](#)]
24. Le Moignic, A.; Malard, V.; Benvegna, T.; Lemiègre, L.; Berchel, M.; Jaffrès, P.-A.; Baillou, C.; Delost, M.; Macedo, R.; Rochefort, J.; et al. Preclinical evaluation of mRNA trimannosylated lipopolyplexes as therapeutic cancer vaccines targeting dendritic cells. *J. Control. Release* **2018**, *278*, 110–121. [[CrossRef](#)] [[PubMed](#)]
25. Van der Jeught, K.; De Koker, S.; Bialkowski, L.; Heirman, C.; Tjok Joe, P.; Perche, F.; Maenhout, S.; Bevers, S.; Broos, K.; Deswarte, K.; et al. Dendritic Cell Targeting mRNA Lipopolyplexes Combine Strong Antitumor T-Cell Immunity with Improved Inflammatory Safety. *ACS Nano* **2018**, *12*, 9815–9829. [[CrossRef](#)]
26. Perche, F.; Lambert, O.; Berchel, M.; Jaffrès, P.-A.; Pichon, C.; Midoux, P. Gene transfer by histidylated lipopolyplexes: A dehydration method allowing preservation of their physicochemical parameters and transfection efficiency. *Int. J. Pharm.* **2012**, *423*, 144–150. [[CrossRef](#)] [[PubMed](#)]

27. Gonçalves, C.; Berchel, M.; Gosselin, M.-P.; Malard, V.; Cheradame, H.; Jaffres, P.-A.; Guégan, P.; Pichon, C.; Midoux, P. Lipopolyplexes comprising imidazole/imidazolium lipophosphoramidate, histidinylated polyethyleneimine and siRNA as efficient formulation for siRNA transfection. *Int. J. Pharm.* **2014**, *460*, 264–272. [[CrossRef](#)]
28. Godbey, W.; Wu, K.K.; Mikos, A.G. Poly (ethylenimine) and its role in gene delivery. *J. Control. Release* **1999**, *60*, 149–160. [[CrossRef](#)]
29. Ranneh, A.H.; Takemoto, H.; Sakuma, S.; Awaad, A.; Nomoto, T.; Mochida, Y.; Matsui, M.; Tomoda, K.; Naito, M.; Nishiyama, N. An Ethylenediamine-based Switch to Render the Polyzwitterion Cationic at Tumorous pH for Effective Tumor Accumulation of Coated Nanomaterials. *Angew. Chem.* **2018**, *130*, 5151–5155. [[CrossRef](#)]
30. Chen, L.; Simpson, J.D.; Fuchs, A.V.; Rolfe, B.E.; Thurecht, K.J. Effects of surface charge of hyperbranched polymers on cytotoxicity, dynamic cellular uptake and localization, hemotoxicity, and pharmacokinetics in mice. *Mol. Pharm.* **2017**, *14*, 4485–4497. [[CrossRef](#)]
31. Meng, H.; Leong, W.; Leong, K.W.; Chen, C.; Zhao, Y. Walking the line: The fate of nanomaterials at biological barriers. *Biomaterials* **2018**, *174*, 41–53. [[CrossRef](#)]
32. Chonn, A.; Cullis, P.; Devine, D. The role of surface charge in the activation of the classical and alternative pathways of complement by liposomes. *J. Immunol.* **1991**, *146*, 4234–4241.
33. Fasugba, O.; Gardner, A.; Mitchell, B.G.; Mnatzaganian, G. Ciprofloxacin resistance in community-and hospital-acquired *Escherichia coli* urinary tract infections: A systematic review and meta-analysis of observational studies. *BMC Infect. Dis.* **2015**, *15*, 545. [[CrossRef](#)]
34. Zervos, M.J.; Hershberger, E.; Nicolau, D.P.; Ritchie, D.J.; Blackner, L.K.; Coyle, E.A.; Donnelly, A.J.; Eckel, S.F.; Eng, R.H.; Hiltz, A. Relationship between fluoroquinolone use and changes in susceptibility to fluoroquinolones of selected pathogens in 10 United States teaching hospitals, 1991–2000. *Clin. Infect. Dis.* **2003**, *37*, 1643–1648. [[CrossRef](#)] [[PubMed](#)]
35. Atkin, S.D.; Abid, S.; Foster, M.; Bose, M.; Keller, A.; Hollaway, R.; Sader, H.S.; Greenberg, D.E.; Finklea, J.D.; Castanheira, M. Multidrug-resistant *Pseudomonas aeruginosa* from sputum of patients with cystic fibrosis demonstrates a high rate of susceptibility to ceftazidime–avibactam. *Infect. Drug Resist.* **2018**, *11*, 1499. [[CrossRef](#)] [[PubMed](#)]
36. Centres for Disease Control and Prevention. *Antibiotic Resistance Threats in the United States, 2013*; Centres for Disease Control and Prevention, US Department of Health and Human Services: Atlanta, GA, USA, 2013.
37. Paradies, G.; Paradies, V.; De Benedictis, V.; Ruggiero, F.M.; Petrosillo, G. Functional role of cardiolipin in mitochondrial bioenergetics. *Biochim. Biophys. Acta (BBA)-Bioenerget.* **2014**, *1837*, 408–417. [[CrossRef](#)] [[PubMed](#)]
38. Sosunov, V.; Mischenko, V.; Eruslanov, B.; Svetoch, E.; Shakina, Y.; Stern, N.; Majorov, K.; Sorokoumova, G.; Selishcheva, A.; Apt, A. Antimycobacterial activity of bacteriocins and their complexes with liposomes. *J. Antimicrob. Chemother.* **2007**, *59*, 919–925. [[CrossRef](#)]
39. Lin, T.-Y.; Weibel, D.B. Organization and function of anionic phospholipids in bacteria. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 4255–4267. [[CrossRef](#)] [[PubMed](#)]
40. Lungwitz, U.; Breunig, M.; Blunk, T.; Göpferich, A. Polyethylenimine-based non-viral gene delivery systems. *Eur. J. Pharm. Biopharm.* **2005**, *60*, 247–266. [[CrossRef](#)] [[PubMed](#)]
41. Pinnapireddy, S.R.; Duse, L.; Strehlow, B.; Schäfer, J.; Bakowsky, U. Composite liposome-PEI/nucleic acid lipopolyplexes for safe and efficient gene delivery and gene knockdown. *Colloids Surf. B Biointerfaces* **2017**, *158*, 93–101. [[CrossRef](#)]
42. Wang, L.-L.; Feng, C.-L.; Zheng, W.-S.; Huang, S.; Zhang, W.-X.; Wu, H.-N.; Zhan, Y.; Han, Y.-X.; Wu, S.; Jiang, J.-D. Tumor-selective lipopolyplex encapsulated small active RNA hampers colorectal cancer growth in vitro and in orthotopic murine. *Biomaterials* **2017**, *141*, 13–28. [[CrossRef](#)] [[PubMed](#)]
43. Persano, S.; Guevara, M.L.; Li, Z.; Mai, J.; Ferrari, M.; Pompa, P.P.; Shen, H. Lipopolyplex potentiates anti-tumor immunity of mRNA-based vaccination. *Biomaterials* **2017**, *125*, 81–89. [[CrossRef](#)]
44. Benincasa, M.; Barrière, Q.; Runti, G.; Pierre, O.; Bourge, M.; Scocchi, M.; Mergaert, P. Single cell flow cytometry assay for peptide uptake by bacteria. *Bio-Protocol* **2016**, *6*, e2038. [[CrossRef](#)]
45. Nuutila, J.; Lilius, E.M. Flow cytometric quantitative determination of ingestion by phagocytes needs the distinguishing of overlapping populations of binding and ingesting cells. *Cytom. Part A* **2005**, *65*, 93–102. [[CrossRef](#)]

46. Tilley, L.D.; Hine, O.S.; Kellogg, J.A.; Hassinger, J.N.; Weller, D.D.; Iversen, P.L.; Geller, B.L. Gene-specific effects of antisense phosphorodiamidate morpholino oligomer-peptide conjugates on *Escherichia coli* and *Salmonella enterica* serovar typhimurium in pure culture and in tissue culture. *Antimicrob. Agents Chemother.* **2006**, *50*, 2789–2796. [[CrossRef](#)] [[PubMed](#)]
47. Geller, B.L.; Deere, J.; Tilley, L.; Iversen, P.L. Antisense phosphorodiamidate morpholino oligomer inhibits viability of *Escherichia coli* in pure culture and in mouse peritonitis. *J. Antimicrob. Chemother.* **2005**, *55*, 983–988. [[CrossRef](#)] [[PubMed](#)]
48. Ma, J.-C.; Wu, Y.-Q.; Cao, D.; Zhang, W.-B.; Wang, H.-H. Only acyl carrier protein 1 (AcpP1) functions in *Pseudomonas aeruginosa* fatty acid synthesis. *Front. Microbiol.* **2017**, *8*, 2186. [[CrossRef](#)] [[PubMed](#)]
49. Nikravesh, A.; Dryselius, R.; Faridani, O.R.; Goh, S.; Sadeghizadeh, M.; Behmanesh, M.; Ganyu, A.; Klok, E.J.; Zain, R.; Good, L. Antisense PNA accumulates in *Escherichia coli* and mediates a long post-antibiotic effect. *Mol. Ther.* **2007**, *15*, 1537–1542. [[CrossRef](#)] [[PubMed](#)]
50. Sousa, S.A.; Ramos, C.G.; Almeida, F.; Meirinhos-Soares, L.; Wopperer, J.; Schwager, S.; Eberl, L.; Leitao, J.H. Burkholderia cenocepacia J2315 acyl carrier protein: A potential target for antimicrobials' development? *Microb. Pathog.* **2008**, *45*, 331–336. [[CrossRef](#)]
51. McArthur, M. Transcription Factor Decoys for the Treatment and Prevention of Infections Caused by Bacteria Including *Clostridium Difficile*. U.S. Patent App. 13/802,103, 18 September 2014.
52. McArthur, M. Transcription Factor Decoys. European Patent EP2552455B1, 28 December 2016.
53. Renner, L.D.; Weibel, D.B. Cardiolipin microdomains localize to negatively curved regions of *Escherichia coli* membranes. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 6264–6269. [[CrossRef](#)]
54. Hegarty, J.P.; Krzeminski, J.; Sharma, A.K.; Guzman-Villanueva, D.; Weissig, V.; Stewart Sr, D.B. Bolaamphiphile-based nanocomplex delivery of phosphorothioate gapmer antisense oligonucleotides as a treatment for *Clostridium difficile*. *Int. J. Nanomed.* **2016**, *11*, 3607. [[CrossRef](#)]
55. Marín-Menéndez, A.; Montis, C.; Díaz-Calvo, T.; Carta, D.; Hatzixanthis, K.; Morris, C.J.; McArthur, M.; Berti, D. Antimicrobial nanoplexes meet model bacterial membranes: The key role of cardiolipin. *Sci. Rep.* **2017**, *7*, 41242. [[CrossRef](#)] [[PubMed](#)]
56. Traglia, G.M.; Sala, C.D.; Fuxman Bass, J.I.; Soler-Bistué, A.J.; Zorreguieta, A.; Ramírez, M.S.; Tolmasky, M.E. Internalization of locked nucleic acids/DNA hybrid oligomers into *Escherichia coli*. *BioRes. Open Access* **2012**, *1*, 260–263. [[CrossRef](#)]
57. Setyawati, M.I.; Kutty, R.V.; Tay, C.Y.; Yuan, X.; Xie, J.; Leong, D.T. Novel theranostic DNA nanoscaffolds for the simultaneous detection and killing of *Escherichia coli* and *Staphylococcus aureus*. *ACS Appl. Mater. Interfaces* **2014**, *6*, 21822–21831. [[CrossRef](#)] [[PubMed](#)]
58. Thibault, T.; Degrouard, J.; Baril, P.; Pichon, C.; Midoux, P.; Malinge, J.-M. Production of DNA minicircles less than 250 base pairs through a novel concentrated DNA circularization assay enabling minicircle design with NF- κ B inhibition activity. *Nucleic Acids Res.* **2016**, *45*, e26. [[CrossRef](#)]
59. Ghosal, A.; Nielsen, P.E. Potent antibacterial antisense peptide–peptide nucleic acid conjugates against *Pseudomonas aeruginosa*. *Nucleic Acid Ther.* **2012**, *22*, 323–334. [[CrossRef](#)]
60. Apte, A.; Koren, E.; Koshkaryev, A.; Torchilin, V.P. Doxorubicin in TAT peptide-modified multifunctional immunoliposomes demonstrates increased activity against both drug-sensitive and drug-resistant ovarian cancer models. *Cancer Biol. Ther.* **2014**, *15*, 69–80. [[CrossRef](#)] [[PubMed](#)]
61. Tros de Ilarduya, C.; García, L.; Düzgünes, N. Liposomes and lipopolymeric carriers for gene delivery. *J. Microencapsul.* **2010**, *27*, 602–608. [[CrossRef](#)]
62. Shen, Z.; Reznikoff, G.; Dranoff, G.; Rock, K.L. Cloned dendritic cells can present exogenous antigens on both MHC class I and class II molecules. *J. Immunol.* **1997**, *158*, 2723–2730.

