



**HAL**  
open science

## Consequences of analgesics use in early pregnancy: Results of tests on mice

Radu E. Iacob, Daniela Iacob, Radu Dumitru Moleriu, Delia Mirela Tit, Simona Bungau, Pavel Otrisal, Selim Aleya, Claudia Judea-Pusta, Gabriela Cioca, Ovidiu Gabriel Bratu, et al.

### ► To cite this version:

Radu E. Iacob, Daniela Iacob, Radu Dumitru Moleriu, Delia Mirela Tit, Simona Bungau, et al.. Consequences of analgesics use in early pregnancy: Results of tests on mice. *Science of the Total Environment*, 2019, 10.1016/j.scitotenv.2019.07.212 . hal-02184585

**HAL Id: hal-02184585**

**<https://hal.science/hal-02184585v1>**

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Consequences of Analgesics Use in Early Pregnancy: Results of tests on mice**

2  
3 Radu E. Iacob<sup>1</sup>, Daniela Iacob<sup>2</sup>, Radu Dumitru Moleriu<sup>3</sup>, Delia Mirela Tit<sup>4</sup>, Simona Bungau<sup>4</sup>,  
4 Pavel Otrisal<sup>5</sup>, Selim Aleya<sup>6</sup>, Claudia Judea-Pusta<sup>7</sup>, Gabriela Cioca<sup>8</sup>, Ovidiu Gabriel Bratu<sup>9</sup>,  
5 Lotfi Aleya<sup>10\*</sup>, Izabella Petre<sup>11</sup>

6  
7 <sup>1</sup>Department of Pediatric Surgery, Victor Babeş University of Medicine and Pharmacy Timisoara,  
8 Romania radueiacob@yahoo.com

9 <sup>2</sup>Department of Neonatology, Victor Babeş University of Medicine and Pharmacy Timisoara, Romania

10 <sup>3</sup> Department of Mathematics, Faculty of Mathematics and Computer Science, West University of  
11 Timisoara, Timisoara, Romania danielariacob@yahoo.com

12 <sup>4</sup>Department of Pharmacy, Faculty of Medicine and Pharmacy, University of Oradea, Oradea 410028,  
13 Romania; mirela\_tit@yahoo.com\_(D.M.T.), simonabungau@gmail.com\_(S.B.)

14 <sup>5</sup>Nuclear, Biological and Chemical Defence Institute, University of Defence, Vítá Nejedlého, 682 01  
15 Vyškov, Czech Republic pavel.otrisal@unob.cz (P.O.)

16 <sup>6</sup>FacultyofMedicine,Besançon,Chrono-EnvironnementLaboratory,UMRCNRS6249,France<sup>[1]</sup><sub>[SEP]</sub>

17 <sup>7</sup>Department of Morphological Disciplines, Faculty of Medicine and Pharmacy, University of Oradea,  
18 Oradea 410028, Romania claupustaml@yahoo.com (C.J.-P.)

19 <sup>8</sup>Faculty of Medicine, “Lucian Blaga” University of Sibiu, Sibiu, Romania gabriela.cioca@ulbsibiu.ro  
20 (G.C.)

21 <sup>9</sup>Clinical Department 3, University of Medicine and Pharmacy “Carol Davila”, Bucharest, Romania;  
22 ovi78doc@yahoo.com\_(O.G.B)

23 <sup>10</sup>Laboratoire Chrono-environnement CNRS 6249, Université de Franche-Comté, Besançon, France;

24 <sup>11</sup>Department of Obstetrics and Gynecology, “Victor Babeş” University of Medicine and Pharmacy  
25 Timisoara, Romania dr.petreizabella@yahoo.com

26

27 \*Corresponding author: lotfi.aleya@univ-fcomte.fr

28 Phone: + 33 (0) 3 81 66 66 66

29 Fax: + 33 (0) 3 81 66 57 97

30

31 **Abstract**

32 Self-medication during pregnancy continues to be an issue in developing countries due to  
33 poor medical education. The most commonly used drugs are analgesics, mainly  
34 acetaminophen (paracetamol, APAP) and, to a lesser extent, ketoprofen (KPF). The aim of the  
35 study was to establish whether there are consequences of accidental use of these two drugs  
36 during early embryogenesis. The experimental study was performed on 30 pregnant white  
37 mice, divided into three groups: a first group to which APAP was administered, a second  
38 group to which KPF was administered, and group 3 as a control group. At delivery, the baby  
39 mice were examined, and after their first parturition, they were taken into study and examined  
40 according to the established protocol. Macroscopic and microscopic examinations of the liver  
41 and kidney were performed; liver and renal changes were recorded. Regarding the fertility,  
42 the number of fetuses born to mothers that were administered APAP and KPF corresponded  
43 to the normal values recorded in this species. Microscopic changes that were found in the  
44 kidney were as follows: APAP group – necrosis of the urinary tube, vascular congestions and  
45 vascular disorders; KPF group – ectasia, especially in the medullary cavity. Microscopic  
46 hepatic changes showed in the APAP group – vascular congestions, vascular disorders and  
47 hemosiderin deposits in the Kupffer cells; in the KPF group were found – discrete vascular  
48 disorders consisting in sinusoidal capillary ectasia and vascular congestions, as well as the  
49 presence of lymphocyte conglomerates. The aforementioned lesions indicate hepatic and renal  
50 distress with variable degrees of severity, but they appear to be reversible (the longer the time  
51 from the maximum effect of the drug the lower its toxicity).

52

53 **Key words:** paracetamol; ketoprofen; mice embryogenesis; microscopic changes; kidney;  
54 liver.

55

56

## 57 **1. Introduction**

58 Despite the development and diversification of contraception methods and particularly of  
59 prenatal diagnosis of congenital malformations, each year there are still a large number of  
60 preterm and full-term newborns who are born with these conditions, due to the high  
61 proportion of cases of social difficulties and the absence of pregnancy checkups.

62 Meanwhile, the tendency toward self-medication with analgesics is a problem worldwide.  
63 Paracetamol (APAP), under various dosage forms and trade names, is one of the products  
64 most used, even during the pregnancy period. To a lesser extent ketoprofen (KPF) is also  
65 used by pregnant women (Andrade et al., 2004; Bayene and Beza, 2018). APAP use is so  
66 widespread among the population that studies are showing its ubiquitous presence in human  
67 urine (Modick et al., 2014).

68 The occasional self-medication by pregnant women, when they are not aware of their  
69 pregnancy, is an additional aspect of the above situation. Consequences may be more severe if  
70 self-medication occurs during embryogenesis, in cases when pregnancy has not yet been  
71 identified. These cases may be referred to as accidental use of such substances by the  
72 pregnant women.

73 Although many studies confirm the effects of long-term exposure of embryos to these drugs  
74 (Adjei, et al., 2008; Feldkamp et al., 2010; Kristensen et al., 2011; Rebordosa et al., 2008),  
75 as do many experimental studies on mice and rats (Burdan et al., 2001; Kristensen et al.,  
76 2011), data establishing the correlations between short-term occasional use of these  
77 substances and the emergence of abnormalities are still insufficient.

78 Most problems occur with so-called teratogenic drugs that cross the placenta and enter the  
79 placental fetal circulation. Sometimes the effects of drugs on the fetus are known only after  
80 many years. Thalidomide, for example, used in Europe in the 1960s, resulted in numerous  
81 serious birth defects, such as missing or incompletely formed limbs (Gilbert-Barnes, 2010;

82 Vargesson, 2015). It had been put on the market at the end of the 1950s and was used for a  
83 considerable period of time, until it was banned and withdrawn from the market in the UK, at  
84 the end of 1961, and in 1962 almost worldwide (Gilbert-Barnes, 2010).

85 For medicines / drugs, the list of agents shown to be potentially teratogenic is surprisingly  
86 small (Mitchell et al., 2011). The term “teratogen” is attributed to any environmental factor  
87 that causes permanent abnormalities in the structure, function or development of the embryo  
88 or fetus (Gilbert-Barnes, 2010). In the first two weeks of pregnancy (the pre-embryonic  
89 stage), the embryo appears to be resistant to the action of teratogenic factors, but a significant  
90 exposure can be lethal. Fetal malformations are caused by certain medications, narcotics,  
91 chemicals (Abdel-Daim et al., 2018 a and b), maternal conditions or diseases (Stoicescu et al.,  
92 2017; Tataru et al., 2019), including infections, that can affect the normal intrauterine  
93 development of the fetus. A connection between the dose of medication intake and its effect  
94 has been found for animals or humans, a greater intake during pregnancy determining more  
95 severe phenotypic effects on the fetus (Frias and Gilbert Barnes, 2008).

96 During the first three to eight weeks, there is a period of maximum susceptibility to the action  
97 of teratogenic agents because all the major organs are formed now (organogenesis). Most fetal  
98 malformations occur in this stage of pregnancy (although less than 1% of total congenital  
99 malformations are caused by human teratogenic drugs) (Mazzu-Nascimento et al., 2017).

100 The induction of fetal malformations by these agents depends on the maternal and paternal  
101 genetic predisposition, the stage of fetal development at the time of exposure, the route of  
102 administration and the duration of exposure to the teratogenic agent (Brent, 2001; Gilbert-  
103 Barnes, 2010). It is crucial to develop medical education programs targeting all social  
104 categories, which inform on the possibilities of family planning and the importance of proper  
105 dispensation of all medications during pregnancy and of avoiding self-medication (Bayene  
106 and Beza, 2018). Such programs would help reduce the number of unwanted pregnancies,  
107 and—if medications are dispensed properly during pregnancy—exposure to a number of

108 teratogenic factors would be avoided, thereby leading to a significant decrease in the  
109 percentage of serious birth defects/malformations.

110 The purpose of this experimental study was to identify the effects of the maximum tolerated  
111 dose of certain drugs during embryogenesis. It was intended to establish a correlation between  
112 acetaminophen or ketoprofen and disorders in the first part of pregnancy, during  
113 embryogenesis. Our research, which was performed on female mice during pregnancy,  
114 examined the possibility of congenital disorders.

115

## 116 **2. Materials and methods**

### 117 **2.1. Study design**

118 Experiments were performed on white mice for the following reasons: they have a high  
119 reproduction potential, thereby ensuring the number of animals needed; they are easy to  
120 handle; they can easily be put into homogenous groups (by weight, age); they do not require  
121 large spaces or special care conditions; and they are resistant to infections and stress from  
122 experiments. Thirty pregnant white mice were selected and divided into three groups: group A  
123 – to which APAP was administered, group B – to which KPF was administered, and group C  
124 – the control group.

### 125 **2.2. Ethical Statement**

126 This research was performed in compliance with the European Convention principles for the  
127 protection of vertebrate animals used in experimental and other scientific purposes, adopted in  
128 1986, in Strasbourg (ECPVAEOSP, 1986), to the 2010/63/EU Directive of the European  
129 Parliament and of the European Council on the protection of animals used for scientific  
130 purposes, adopted on September 22, 2010, and in accordance with laboratory good practices.

131 The mice were kept under optimal microclimate conditions, locked in cages with feeders and  
132 waterers, in the same room. The study was approved by the Ethical Local Commission of the

133 Pius Brânzeu Center for Laparoscopic Surgery and Microsurgery of the Victor Babeș  
134 University of Medicine and Pharmacy Timisoara (reference number 680/12.04.2006).

### 135 **2.3. Technique of medical substances administration**

136 After weighing, all animals were caught by the tail, pulled out of the scale box and placed on  
137 a smooth drape. The animals were then covered with another drape, held firmly but delicately,  
138 and marked on the tail with a permanent marker: blue for APAP and red for KPF. The  
139 posterior girdle was lifted to expose the hypogastric region, where the drug dose (calculated  
140 based on the weight obtained by scaling and presented in **Table 1**), was injected  
141 intraperitoneally by means of a fine-needle syringe; the procedure was repeated strictly on  
142 each animal, for three consecutive days. The substances were administrated to the mice  
143 between the fifth and seventh gestation days. This time interval was chosen because nidation  
144 during embryogenesis in mice occurs 4.5 days after conception, corresponding to the early  
145 second week of human embryogenesis, when nidation occurs. Day 7 of embryogenesis in  
146 mice matches the middle of the third week in human embryogenesis (Day 17), when the  
147 delayed menstrual cycle may lead the woman to perform a pregnancy test and to stop self-  
148 medication in the event of positive diagnosis (Hill, 2019). The three days of the mouse  
149 embryogenesis during which drugs were administered (5<sup>th</sup> to 7<sup>th</sup> days) are equivalent to 10  
150 days of human embryogenesis (7<sup>th</sup> to 17<sup>th</sup> days) (Hill, 2018; Theiler Staging Criteria, 2019).

151 The mice were observed until the end of gestation and kept under optimal microclimate  
152 conditions. After delivery, the baby mice were examined, to assess the possible consequences  
153 of drug administration on fertility. After the first reproductive cycle, the mice were euthanized  
154 and macroscopic and microscopic examinations of the liver and kidney performed.

155 Macroscopic changes of liver and kidneys were assessed by comparing them to the aspect of  
156 organs harvested from the control group, focusing on their appearance upon inspection,  
157 preservation of shape and surface uniformity, sizes, and consistency at palpation. After the

158 macroscopic examination of the harvested organs, microscopic examination using the  
159 hematoxylin-eosin (H.E.) staining method was performed.

#### 160 **2.4. Statistical analysis**

161 To gather the database, Microsoft Excel (version 2013) was used, and for the statistical  
162 analysis two different programs were available: SPSS (version 17) for the statistical  
163 significance and Microsoft Excel for the figures and the descriptive statistics. A normality test  
164 was applied, in our case a Shapiro–Wilk test (because our study group consisted of less than  
165 50 subjects) to see if our data were normally distributed. In order to describe the database, the  
166 central tendency and the dispersion parameters were calculated. For the statistical  
167 significance, an ANOVA test was used.

168

### 169 **3. Results**

170 With regard to the *fertility* of the females exposed to intrauterine action of the drugs, no  
171 changes were recorded in comparison with the control group. The number of fetuses born to  
172 mothers who were administered APAP and KPF corresponded to the normal values recorded  
173 in this species, 6-10 per litter apparently being normal. Their growth and development were  
174 monitored up to the sexual maturity and the birth of the first generation of babies (normal  
175 growth as regards the number and morphology of animals).

176 Upon *morphological examination* of the subjects, no differences were recorded in comparison  
177 with the control group. Macroscopic examination of liver and kidneys indicated no  
178 abnormalities in the animals exposed to intrauterine action of the drug in comparison with the  
179 control group.

#### 180 **Microscopic examination of the organs**

##### 181 *APAP group*

182 Microscopic changes in the kidney (after administration of the drug) were found in the form  
183 of necrosis of the urinary tube, desquamation of epithelium, vascular congestion and vascular



184 disorders. At the same time, microscopic hepatic changes showed vascular congestion,  
185 vascular disorders and hemosiderin deposits in the Kupffer cells.

186 *KPF group*

187 The microscopic renal changes after administration of KPF were vascular disorders consisting  
188 of ectasia, especially in the medullary cavity. In the liver, discrete vascular disorders were  
189 observed, consisting in sinusoidal capillary ectasia and vascular congestion, as was the  
190 presence of lymphocyte conglomerates. Such lesions indicate hepatic and renal distress in the  
191 animals undergoing intrauterine exposure to APAP and KPF during embryogenesis. Lesions  
192 were severe to variable degrees but appear to be reversible. The histopathological changes  
193 described were also observed in overdose situations in animals such as dogs and cats which  
194 for instance developed transient renal and / or hepatic impairment. In addition, the changes  
195 assessed through biopsy puncture were less pronounced since the period between  
196 administration and puncture was higher. Basically, if biopsy puncture was made at peak effect  
197 of the active substance, the changes would more pronounced: the longer the time from the  
198 maximum effect of the drug the lower its toxicity. The most representative aspects of the  
199 lesions are presented in **Figures 1 and 2**.

200 The main observation that can be extracted from the descriptive statistical analysis is that the  
201 mean weight of our subjects is almost the same regardless of whether they received either  
202 drug (APAP or KPF) or no drug. The mean values for each mouse over the three tested days,  
203 for weight and for the drug doses, were obtained (**Table 2**).

204 Later, the Shapiro–Wilk normality test was applied. The  $p > 0.05$  value assures the use of the  
205 ANOVA test for establishing the statistical significance between the tested groups. The main  
206 part of the analysis consisted in applying the ANOVA Two Factor Without Replication test in  
207 order to see if there were any differences among the weight or the drug doses over the three  
208 tested days, or if there were any differences among the ten tested mice for either drug.

209 Therefore, four different ANOVA Two Factor Without Replication tests were performed, and  
210 their results are presented in **Tables 3** (for APAP) and **4** (for KPF).

211 In the first test, which compared the mice weight over the three days, considerably significant  
212 differences ( $p < 0.001$ ) were obtained in the following two scenarios: when the differences  
213 among the ten mice were examined and when the differences between the studied days were  
214 examined. In the second test, concerning the APAP dose, extremely significant differences  
215 were obtained for both scenarios: when the differences among the subjects were tested and  
216 when the differences among the studied days were tested. The same observation holds true for  
217 the KPF mice group, under the same circumstances, regarding the  $p < 0.001$  value (**Tables 3**  
218 and **4**).

219 A final hypothesis was tested at the end of this study. We wanted to see if any differences  
220 could be found in the weights of the mice used in this experimental study, in all three groups  
221 (mice with APAP, with KPF and drug-free). Therefore, an average value for weight over the  
222 three days studied was calculated. Application of the ANOVA Single Factor test showed that  
223 there were no statistically significant differences ( $p = 0.883$ , which is  $> 0.05$ ) among the  
224 weights of the mice in all tested groups. In short, the results showed that the drugs did not  
225 influence weight dynamics (**Figure 3**).

226

#### 227 **4. Discussion**

228 Some studies have reported side effects on fetuses from APAP use during pregnancy, as seen  
229 by the presence of congenital malformations clinically diagnosed at birth (Feldkamp et al.,  
230 2010). Thielea et al. (2013) published one of the most complex analyses on the short- and  
231 long-term effects of APAP use during pregnancy. Data in the field also includes information  
232 about the possible onset of gastroschisis. A retrospective study which included 206 cases of  
233 gastroschisis reported an increased risk of this malformation in the event of APAP use during  
234 pregnancy (odds ratio = 1.5, 95% confidence interval: 1.1, 2.2) (Werler et al., 2002). Other

235 authors have reported an increased odds ratio for neural tube defects (Lynberg et al., 1004)  
236 and an increased odds ratio for congenital cryptorchidism, especially in the second trimester,  
237 and reduced testosterone production in *ex vivo* fetal rat testes (Kristensen et al., 2011; Jensen  
238 et al., 2010). In a study published in 2014 and conducted on 64,322 live-born children and  
239 mothers enrolled in the Danish National Birth Cohort during 1996-2002, Liew et al. (2014)  
240 show that prenatal exposure to APAP action increases the risk of behavioral disorders or  
241 hyperkinetic syndrome.

242 Studies regarding the effects of KPF use during pregnancy are fewer in number. A study on  
243 drug prescriptions during pregnancy reports on KPF administration during the last trimester of  
244 pregnancy (Garcia et al., 2000). Other data concluded on the consequence of KPF use during  
245 pregnancy on a group of 11 newborns from mothers who experienced adverse effects to this  
246 drug. Ten of them had renal dysfunction and three deaths were recorded (Llanas et al., 1996).  
247 A case of a newborn with kidney failure following KPF administration to the mother during  
248 the last four days of pregnancy was also presented (1991).

249 In this study, administration of APAP or KPF for a short period of time, between the 5<sup>th</sup> to 7<sup>th</sup>  
250 days of gestation period in mice, does not influence pregnancy evolution. Mice resulting from  
251 such pregnancies have normal morphological and behavioral development in comparison with  
252 the control group, and the fertility of these animals has the values characteristic to the species.  
253 Macroscopic examination of liver and kidneys of animals exposed to the action of these  
254 substances during embryogenesis did not show pathological aspects.

255 Our research did not find any congenital malformations, behavioral disorders or hyperkinetic  
256 syndrome, cryptorchidism or affected fertility in mice which, in the early pregnancy period,  
257 were exposed, for a short time, to one of these drugs (APAP or KPF). However, this paper  
258 mentions some hepatic and renal microscopic changes noted in standard histologic  
259 examinations, but with no vital risk. But these changes call for more in-depth studies  
260 regarding the accidental use of APAP and/or KPF during the first period of pregnancy. Thus,

261 there is a clear need for complex educational programs dedicated to reducing self-medication  
262 among the general population, especially among fertile or pregnant women (in order to  
263 drastically reduce the phenomenon of self-medication in the case of undiagnosed pregnancies)  
264 (Bungau et al., 2015). Reducing self-medication is essential along with taking into account the  
265 threats caused to the environment by increased drug consumption, especially those commonly  
266 in use such as APAP and KPF. For example, the disposal of expired or unused drugs into  
267 waste or landfilled waste exacerbates their concentration in the environment (Cycońet al.,  
268 2016). Since paracetamol is among the most frequently used nonprescription drugs to reduce  
269 fever and relieve pain it is often found in surface water, sewage treatment plants (Taschina et  
270 al., 2017) and drinking water.

271 APAP concentrations in aquatic environments increased over the last years, since they are not  
272 completely removed in wastewater treatment plants, and their residues may be released  
273 through effluents into the aquatic environment (Al-Kaf et al., 2017). The continuous dumping  
274 of non-steroidal anti-inflammatory drugs is of concern since it is practically impossible to  
275 assess their concentration in the environment along with the potential negative impacts on  
276 living organisms in the future (Cycońet al., 2016).

277

## 278 **5. Conclusions**

279 Our study revealed that administration of APAP or KPF drugs does not influence the weight  
280 dynamics of mice during pregnancy; they developed normally, and no differences are  
281 observed between mice administered APAP or KPF and other healthy mice. Changes /  
282 abnormalities that occur can be seen only at the level of the kidney and liver and are visible  
283 just in a histological examination. At present, we cannot determine whether these  
284 abnormalities (caused by developing tissue damage) are due to the direct *in utero* toxicity of  
285 the analgesic, its metabolites or other factors. We believe that further studies are needed to  
286 determine the mechanisms leading to these lesions.

287

288 **Acknowledgements**

289 We would like to thank the entire team of the Veterinary Anatomy Department of Banat  
290 University of Agricultural Science and Veterinary Medicine of Timisoara, for their support  
291 during this study. We express our appreciation to the editor, Dr. Damia Borcelo, and to the  
292 anonymous reviewers for helping to improve our paper.

293

294 **Conflicts of interest:** The authors declare that there was no conflict of interest.

295 **Contribution:** All authors had equal contribution to this paper.

296 **Funding:** This research received no external funding.

297 **References**

298 Abdel-Daim M.M., Abo-El-Sooud K., Aleya L., Bungau S.G., Najda A., Saluja R., 2018a.

299 Alleviation of Drugs and Chemicals Toxicity: Biomedical Value of Antioxidants. *Oxid.*  
300 *Med. Cell. Longev.*, ID 6276438, 2018. <https://doi.org/10.1155/2018/6276438>

301 Abdel-Daim, M.M., Zakhary, N.I., Aleya, L., Bungau, S.G., Bohara, R.A., Siddiqi, N.J.,

302 2018b. Aging, metabolic and degenerative disorders: Biomedical Value of  
303 Antioxidants. *Oxid. Med. Cell. Longev.*, 2018, ID 2098123.  
304 <https://doi.org/10.1155/2018/2098123>

305 Adjei, A.A., Gaedigk, A., Simon, S.D., Weinshilboum, R.M., Leeder, J.S., 2008.

306 Interindividual variability in acetaminophen sulfation by human fetal liver: implications  
307 for pharmacogenetic investigations of drug-induced birth defects. *Birth Defects Res. A*  
308 *Clin. Mol. Teratol.* 82(3), 155-165.

309 Al-Kaf, A.G., Naji, K.M., Abdullah, Q.Y.M., Edrees, W.H.A., 2017. *Chronicles of*  
310 *Pharmaceutical Science*, 1(6), 341-355.

311 Andrade, S.E., Gurwitz, J.H., Davis, R.L., Chan, K.A., 2004. Prescription drug use in  
312 pregnancy. *Am. J. Obstet. Gynecol.*, 191, 398-407.

- 313 Beyene, K.G., Beza, S.W., 2018. Self-medication practice and associated factors among  
314 pregnant women in Addis Ababa, Ethiopia. *Trop Med Health.* 46: 10.  
315 <https://doi.org/10.1186/s41182-018-0091-z>
- 316 Brent, R.L., 2001. The cause and prevention of human birth defects: what have we learned in  
317 the past 50 years? *Congenit Anom.*, 41(1), 3–21.
- 318 Bungau, S., Bungau, C., Tit, D.M., Pallag, A., 2015. The influence of specialized academic  
319 studies on self-medication. *Rev. Rom. Bioet.* 13(1), 153-158.
- 320 Burdan, F., Siezienievska, Z., Kis, G., Blicharski, T., 2001. Embryofetotoxicity of  
321 acetaminophen (paracetamol) in experimental in vivo model. *Ann. Univ. Mariae Curie*  
322 *Sklodowska Med.*, 56, 89-94.
- 323 Cycoń, M, Borymski, S, Żołnierczyk, B, Piotrowska-Seget, Z., 2016. Variable Effects of  
324 Non-steroidal Anti-inflammatory Drugs (NSAIDs) on Selected Biochemical Processes  
325 Mediated by Soil Microorganisms. *Front. Microbiol.* 7:1969.  
326 [doi:10.3389/fmicb.2016.01969](https://doi.org/10.3389/fmicb.2016.01969)
- 327 Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010  
328 on the protection of animals used for scientific purposes, 2010. Available: [http://eur-](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32010L0063)  
329 [lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32010L0063](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32010L0063). Accessed 2018  
330 March 21.
- 331 European Convention for the Protection of Vertebrate Animals used for Experimental and  
332 Other Scientific Purposes (ECPVAEOSP) European Convention for the Protection of  
333 Vertebrate Animals used for Experimental and Other Scientific Purposes - Council of  
334 Europe, 1986. Available: <http://conventions.coe.int/treaty/en/treaties/html/123.htm>.  
335 Accessed 2018 March 21.
- 336 Feldkamp, M.L., Meyer, R.E., Krikov, S., Botto, L.D., 2010. Acetaminophen use in  
337 pregnancy and risk of birth defects: findings from the National Birth Defects Prevention  
338 Study. *Obstet. Gynecol.*, 115, 109-115.

- 339 Frias, J.L., Gilbert-Barness, E., 2008. Human teratogens: current controversies. *Advanc*  
340 *Pediat.*, 55, 171–211.
- 341 Garcia, I., Beyens, M.N., Gauchoux, R., Guy, C., Ollagnier, M., 2000. Drug prescription for  
342 pregnant women in the department of the Loire. *Therapie*, 55(5), 605-611.
- 343 Gilbert-Barness, E., 2010. Teratogenic Causes of Malformations. *Ann Clin Lab Sci.*, 40(2),  
344 99-114
- 345 Gouyon, J.B., Petion, A.M., Sandre, D., Sgro, C., Bavoux, F., 1991. Neonatal kidney  
346 insufficiency and intrauterine exposure to ketoprofen. *Arch. Fr. Pediatr.*, 48(5), 347-  
347 348.
- 348 Hill, M.A., 2019, Embryology Carnegie Stage Comparison. Retrieved  
349 from [https://embryology.med.unsw.edu.au/embryology/index.php/Carnegie\\_Stage\\_Co](https://embryology.med.unsw.edu.au/embryology/index.php/Carnegie_Stage_Comparison)  
350 [mparison](https://embryology.med.unsw.edu.au/embryology/index.php/Carnegie_Stage_Comparison)
- 351 Jensen, M.S., Rebordosa, C., Thulstrup, A.M., Toft, G., Sørensen, H.T., Bonde, J.P.,  
352 Henriksen, T.B., Olseng, J., 2010. Maternal use of acetaminophen, ibuprofen, and  
353 acetylsalicylic acid during pregnancy and risk of cryptorchidism. *Epidemiology*, 21,  
354 779-785.
- 355 Kristensen, D.M., Hass, U., Lesne, L., Lottrup, G., Jacobsen, P.R., Desdoits-Lethimonier, C.,  
356 Boberg, J., Petersen, J.H., Toppari, J., Jensen, T.K., Brunak, S., Skakkebaek, N.E.,  
357 Nellemann, C., Main, K.M., Jegou, B., Leffers, H., 2011. Intrauterine exposure to mild  
358 analgesics is a risk factor for development of male reproductive disorders in human and  
359 rat. *Hum. Reprod.*, 26, 235-244.
- 360 Liew, Z., Ritz, B., Rebordosa, C., Lee, P.C., Olsen, J., 2014. Acetaminophen use during  
361 pregnancy, behavioral problems, and hyperkinetic disorders. *JAMA Pediatr.*, 168(4),  
362 313-320.

- 363 Llanas, B., Cavert, M.H., Apere, H., Demarquez, J.L., 1996, Adverse effects of ketoprofen  
364 after intrauterine exposure. Value of plasma determination. Arch. Pediatr., 3(3), 248-  
365 253.
- 366 Lynberg, M.C., Khoury, M.J., Lu, X., Cocian, T., 1994. Maternal flu, fever, and the risk of  
367 neural tube defects: a population-based case-control study. Am. J. Epidemiol., 140,  
368 244-255.
- 369 Mazzu-Nascimento, T., Melo, D.G., Morbioli, G.G., Carrilho, E., Vianna, F.S.L., Silva, A.A.,  
370 Schuler-Faccini, L., 2017. Teratogens: a public health issue - a Brazilian  
371 overview. Genet. Mol. Biol. 40(2), 387-397. doi: 10.1590/1678-4685-GMB-2016-0179.
- 372 Mitchell, A.A., Gilboa, S.M., Werler, M.M., Kelley, K.E., Louik, C., Hernández-Díaz, S.,  
373 2011. National Birth Defects Prevention Study. Medication use during pregnancy, with  
374 particular focus on prescription drugs: 1976-2008. Am. J. Obstet. Gynecol. 205(1),  
375 51.e1-8. doi: 10.1016/j.ajog.2011.02.029.
- 376 Modick, H., Weiss, T., Dierkes, G., Bruning, T., Koch, H.M., 2014. Ubiquitous presence of  
377 paracetamol in human urine: sources and implications. Reproduction. 147(4), 105-117.
- 378 Rebordosa, C., Kogevinas, M., Sorensen, H.T, Olsen, J., 2008. Pre-natal exposure to  
379 paracetamol and risk of wheezing and asthma in children: a birth cohort study. Int. J.  
380 Epidemiol., 37, 583-590.
- 381 Stoicescu, M., Bungau, S., Tit, D.M., Mutiu, G., Purza, L., Iovan, C., Pop, O.L., 2017.  
382 Carcinogenic uterine risk of repeated abortions: hormonal receptors tumoral expression.  
383 Rom. J. Morphol. Embriol., 58(4), 1429-1434.
- 384 Taschina M., Copolovici D.M., Bungau S., Lupitu A.I., Copolovici L., Iovan C., 2017. The Influence  
385 of Residual Acetaminophen on Beans (*Phaseolus Vulgaris* L.) Plants Secondary Metabolites,  
386 Farmacia, 65(5), 709-713.
- 387 Tataru, A.-L., Furau, G., Afilon, J., Ionescu, C., Dimitriu, M., Bratu, O.G., Tit, D.M.,  
388 Bungau, S., Furau, C., 2019. The situation of cervical cancers in the context of female



389 genital cancers clustering and burden of disease in Arad County, Romania. *J. Clin.*  
390 *Med.*, 8(1), 96. <https://doi.org/10.3390/jcm8010096>

391 Theiler Staging Criteria for Mouse Embryo Development  
392 <http://www.emouseatlas.org/emap/ema/home.html> (accessed on 24 March. 2019)

393 Thielea, K., Kesslerb, T., Arcka, P., Erhardt, A., Tiegs, G., 2013. Acetaminophen and  
394 pregnancy: short- and long-term consequences for mother and child. *J. Reprod.*  
395 *Immunol.*, 97, 128-139.

396 Vargesson, N., 2015. Thalidomide-induced teratogenesis: History and mechanisms. *Birth*  
397 *Defects Res C Embryo Today*. 105(2), 140–156.

398 Werler, M.M., Sheehan, J.E., Mitchell, A.A., 2002. Maternal medication use and risks of  
399 gastroschisis and small intestinal atresia. *Am. J. Epidemiol.*, 155, 26-31.

#### 400 **Figure legends**

401  
402 **Figure 1.** (a) Urinary tube necrosis with epithelium desquamation, section in HE staining of  
403 kidney, detailed image, 40x; (b) Erythrocyte agglutinations indicating vascular congestion,  
404 section in HE staining of kidney, detailed image, 40x; (c) Vascular disorders consisting of  
405 vascular congestions, section in HE staining of liver, detailed image, 40x; (d) Congestions,  
406 hemosiderin deposits in the Kupffer cells, section in HE staining of liver, detailed image, 40x.  
407

408 **Figure 2.** (a) Vascular disorders consisting in ectasia, section in HE staining of kidney,  
409 detailed image, 40x; (b) Vascular disorders consisting in regional ectasia of the sinusoidal  
410 capillaries and passive vascular congestions, lymphocyte conglomerate, section in HE  
411 staining liver, detailed image, 40x.

412

413 **Figure 3.** The weight distribution within the 3 groups





Figure 1 (a); (b); (c); (d).



(a)

(b)

Figure 2 (a); (b).




Figure 3.

Table 1. APAP and KPF doses/animal weight/day (50 mg/kg/day and 5 mg/kg/day respectively)

Animal No.	Day	Weight (g)	Dose (mg)	Weight (g)	Dose (mg)
		APAP		KPF	
1	I	28	1.4	27	0.135
	II	29	1.45	29	0.145
	III	32	1.6	31	0.155
2	I	34	1.7	36	0.18
	II	33	1.65	37	0.185
	III	37	1.85	38	0.19
3	I	29	1.45	29	0.145
	II	33	1.65	31	0.155
	III	36	1.8	34	0.17
4	I	35	1.75	33	0.165
	II	36	1.8	36	0.18
	III	39	1.95	38	0.19
5	I	34	1.7	34	0.17
	II	34	1.7	35	0.175
	III	36	1.8	38	0.19
6	I	37	1.85	36	0.18
	II	38	1.9	38	0.19
	III	39	1.95	41	0.205
7	I	32	1.6	32	0.16
	II	35	1.75	35	0.175
	III	37	1.85	37	0.185
8	I	29	1.45	29	0.145
	II	33	1.65	33	0.165
	III	35	1.75	35	0.175
9	I	32	1.6	32	0.16
	II	35	1.75	35	0.175
	III	37	1.85	37	0.185
10	I	29	1.45	29	0.145
	II	32	1.6	32	0.16
	III	34	1.7	34	0.17

Table 2. The main tendency and dispersion parameters for the mean values (the average values computed for the 3 tested days) for the mice weight and APAP/KPF doses

Statistics	Mean Weight Group 1	Mean APAP dose	Mean Weight Group 2	Mean KPF dose
Mean	33.97	1.70	34.23	17.83
Standard Error	0.77	0.04	0.87	0.41
Median	34.67	1.73	34.67	18.20
Mode	34.67	1.73	35.67	18.20
Standard Deviation	2.44	0.12	2.75	1.28
Sample Variance	5.96	0.01	7.56	1.64
Kurtosis	-0.05	-0.05	0.16	-0.05
Skewness	-0.11	-0.11	-0.51	-0.11
Range	8.33	0.42	9.33	4.38
Minimum	29.67	1.48	29.00	15.58
Maximum	38	1.90	38.33	19.95
Sum	339.67	16.98	342.33	178.33
Count	10	10	10	10



Table 3. The ANOVA Two-Factor Without Replication test applied on the APAP injected mice group for weight and for drug dose.

Source of variation	SS	df	MS	F	F crit.
Weight					
The analysis on mice	160.9667	9	17.88519	16.82578	2.456281149
The analysis on the 3 studied days	92.86667	2	46.43333	43.68293	3.554557146
Error	19.13333	18	1.062963		
Total	272.9667	29			
Drug dose					
The analysis on mice	0.402417	9	0.044713	16.82578	2.456281
The analysis on the 3 studied days	0.232167	2	0.116083	43.68293	3.554557
Error	0.047833	18	0.002657		
Total	0.682417	29			

Table 4. The ANOVA Two-Factor Without Replication test applied on the KPF injected mice group for weight and for drug dose.




Source of variation	SS	df	MS	F	F crit.
Weight					
The analysis on mice	223.6333	9	24.84815	59.90179	2.456281
The analysis on the 3 studied days	105.8667	2	52.93333	127.6071	3.554557
Error	7.466667	18	0.414815		
Total	336.9667	29			
Drug dose					
The analysis on mice	0.005591	9	0.000621	59.90179	2.456281
The analysis on the 3 studied days	0.002647	2	0.001323	127.6071	3.554557
Error	0.000187	18	1.04E-05		
Total	0.008424	29			







30 pregnant white female mice

3 groups



Fertility  
Morphological examination  
Microscopic examination of the organs



No congenital malformations, behavioral disorders or impaired fertility in mice (to APAP or KPF influence) were found. Some liver and kidney microscopic changes have been described in the standard histological examination but without vital risk.