



**HAL**  
open science

## Half-space theorems for the Allen-Cahn equation and related problems

Francois Hamel, Yong Liu, Pieralberto Sicbaldi, Kelei Wang, Juncheng Wei

► **To cite this version:**

Francois Hamel, Yong Liu, Pieralberto Sicbaldi, Kelei Wang, Juncheng Wei. Half-space theorems for the Allen-Cahn equation and related problems. 2019. hal-02184621v1

**HAL Id: hal-02184621**

**<https://hal.science/hal-02184621v1>**

Preprint submitted on 16 Jul 2019 (v1), last revised 27 Jul 2019 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# HALF-SPACE THEOREMS FOR THE ALLEN-CAHN EQUATION AND RELATED PROBLEMS

FRANÇOIS HAMEL, YONG LIU, PIERALBERTO SICBALDI, KELEI WANG,  
AND JUNCHENG WEI

ABSTRACT. In this paper we obtain rigidity results for a bounded non-constant entire solution  $u$  of the Allen-Cahn equation in  $\mathbb{R}^n$ , whose level set  $\{u = 0\}$  is contained in a half-space. If  $n \leq 3$  we prove that the solution must be one-dimensional. In dimension  $n \geq 4$ , we prove that either the solution is one-dimensional or stays below a one-dimensional solution and converges to it after suitable translations. Some generalizations to one phase free boundary problems are also obtained.

## 1. INTRODUCTION AND MAIN RESULTS

We are interested in rigidity results for bounded entire solutions of the Allen-Cahn equation

$$(1) \quad -\Delta u = u - u^3, \quad x = (x_1, \dots, x_n) \in \mathbb{R}^n.$$

In the simplest case  $n = 1$ , equation (1) reduces to an ODE and has a heteroclinic solution

$$H(x) = \tanh\left(\frac{x}{\sqrt{2}}\right).$$

Phase plane analysis tells us that up to a translation,  $H$  is the unique monotone increasing solution in  $\mathbb{R}$ . The one-dimensional solution  $H$  actually plays an important role in the theory of Allen-Cahn equation in general dimensions. Indeed, De Giorgi [10] conjectured that for  $n \leq 8$ , if a bounded solution  $u$  to (1) is strictly monotone in one direction, say  $x_n$ , then it must be one-dimensional, which then means that  $u$  is identically equal to  $H(x \cdot e + a)$  for some unit vector  $e$ , with  $e_n > 0$ , and some real number  $a$ . This conjecture has been proved to be true for  $n = 2$  (Berestycki, Caffarelli and Nirenberg [4], Ghoussoub and Gui [18]),  $n = 3$  (Ambrosio and Cabré [3]), and for  $4 \leq n \leq 8$  (Savin [26]) under the additional limiting condition

$$\lim_{x_n \rightarrow \pm\infty} u(x', x_n) = \pm 1$$

pointwise in  $x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}$ . This condition implies that the level sets  $\{x \in \mathbb{R}^n : u(x) = \mu\}$ , for every  $\mu \in (-1, 1)$ , of the function  $u$  are entire graphs with respect to the first  $n - 1$  variables. On the other hand, for  $n \geq 9$ , Del Pino, Kowalczyk and the fifth author [13] constructed monotone solutions which are not one-dimensional, showing that the condition  $n \leq 8$  in the De Giorgi conjecture cannot be relaxed.

The De Giorgi conjecture can be regarded as a rigidity result for the Allen-Cahn equation. The second rigidity result we would like to mention here is about the classification of the solutions which are global minimizers of the associated energy functional. Savin [26]

proved that global minimizers are one-dimensional up to dimensions  $n \leq 7$ . The second, fourth and fifth authors constructed in [21] counterexamples in dimension 8, i.e. global minimizers which are not one-dimensional. For related rigidity results for the solutions to the Allen-Cahn equation, we refer to [2, 7, 15, 16] and the references therein.

In this paper, we are interested in rigidity results for the Allen-Cahn equation when the zero level set

$$\Gamma_0 := \{u = 0\} = \{x \in \mathbb{R}^n : u(x) = 0\}$$

of the solution is contained in a half-space, say  $\{x_n > 0\} := \{x \in \mathbb{R}^n : x_n > 0\}$  up to translation and rotation. However, we point out that we make no assumption on the monotonicity of  $u$  in a direction nor on its stability or minimizing properties.

Our first result is the following half-space rigidity result:

**Theorem 1.** (Weak half-space theorem) *Let  $n \leq 3$  and  $u$  be a bounded non-constant solution of (1). Suppose that the zero level set  $\{u = 0\}$  is contained in  $\{x_n > 0\}$ . Then  $u$  is one-dimensional. More precisely, there exists  $a \in \mathbb{R}$  such that either  $u(x) = H(-x_n + a)$  for all  $x \in \mathbb{R}^n$ , or  $u(x) = H(x_n + a)$  for all  $x \in \mathbb{R}^n$ .*

Note that, in any dimension  $n \geq 1$ , if  $u$  is a bounded solution of (1), then necessarily

$$-1 \leq u \leq 1 \quad \text{in } \mathbb{R}^n$$

from the maximum principle. Furthermore, if  $\{u = 0\}$  is empty, then by constructing suitable comparison functions as in the first part of the proof of Theorem 3 (see Section 2 below), we can show that  $u \equiv \pm 1$  in  $\mathbb{R}^n$ . Therefore, if  $u$  is bounded and non-constant, then  $\{u = 0\} \neq \emptyset$  and  $-1 < u < 1$  in  $\mathbb{R}^n$  from the strong maximum principle. Moreover, if  $u \geq 0$  (resp.  $u \leq 0$ ) in  $\mathbb{R}^n$  and  $\{u = 0\} \neq \emptyset$ , then  $u \equiv 0$  in  $\mathbb{R}^n$  from the strong maximum principle. Hence we can assume without loss of generality that the three sets  $\{u = 0\}$ ,  $\{u > 0\}$  and  $\{u < 0\}$  are not empty.

We point out that here it is not assumed that the nodal set  $\{u = 0\}$  is a graph, that is, the sets  $\{u > 0\}$  or  $\{u < 0\}$  are not assumed to be epigraphs. For rigidity results in the epigraph case we refer to [14, 17] and the references therein.

As an application of Theorem 1, using the classification result of stable solutions of the Allen-Cahn equation in the plane, we get the following *strong half-space theorem*:

**Corollary 2.** (Strong half-space theorem) *Suppose  $n = 2$ . Let  $u_1 < u_2$  be two bounded non-constant solutions of (1) in  $\mathbb{R}^2$ . Then  $u_1$  and  $u_2$  are one-dimensional, namely there exist a unit vector  $e$  and some real numbers  $a < b$  such that  $u_1(x) = H(x \cdot e + a)$  and  $u_2(x) = H(x \cdot e + b)$  for all  $x \in \mathbb{R}^2$ .*

We will also generalize Theorem 1 to a free boundary problem. We refer to Section 4 for the precise statement and its proof.

Our results are inspired by analogous results in the minimal surface theory. A half-space theorem for minimal surfaces in  $\mathbb{R}^3$  was proved by Hoffman and Meeks [20]. It states that connected, proper, minimal surfaces in  $\mathbb{R}^3$  are necessarily planes. A version of a half-space theorem for minimal surfaces with bounded Gaussian curvature is proved in [27]. The half-space theorem plays an important role in the understanding of the structure of minimal

spaces, and there is a vast literature on this subject. It is used in the proof of the local removal singularity theorem [23], it is also used to study the properness of minimal surfaces (see, for instance, [24]). In [9], Colding and Minicozzi proved that the plane is the only complete embedded minimal disk in  $\mathbb{R}^3$ , by establishing a chord-arc bound and applying Hoffman-Meeks half-space theorem.

We remark that the half-space theorem is not true for minimal hypersurfaces in  $\mathbb{R}^n$  with  $n \geq 4$ . For example the higher dimensional catenoid provides a counterexample. However, for the Allen-Cahn equation, this question is still open in higher dimensions. In view of the construction of solutions concentrated on higher dimensional catenoid [1], we turn to believe that the half-space theorem of the Allen-Cahn equation should be true also for all  $n \geq 4$ . Intuitively, for solutions of the Allen-Cahn equation there are strong interactions between different ends, while this is not the case for minimal surfaces.

The proof of the half-space theorem for minimal surfaces uses sweeping principle. It appears that this idea does not work for the Allen-Cahn case, although we can prove partial results along this direction. Our main result for solutions of the Allen-Cahn equation in arbitrary dimension with zero level set contained in a half-space is the following:

**Theorem 3.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of the Allen-Cahn equation (1) in  $\mathbb{R}^n$ . If  $u < 0$  in the half-space  $\{x_n < 0\}$ , then there exists  $a \in \mathbb{R}$  such that*

$$u(x) \leq H(x_n + a)$$

for all  $x \in \mathbb{R}^n$ , and either

- (1)  $u(x) = H(x_n + a)$  for all  $x \in \mathbb{R}^n$ , or
- (2)  $u(x) < H(x_n + a)$  for all  $x \in \mathbb{R}^n$  and there exists a sequence  $(y_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times \{0\}$  such that  $|y_k| \rightarrow +\infty$  as  $k \rightarrow +\infty$ , and the functions  $u(\cdot + y_k)$  converge in  $C_{loc}^2(\mathbb{R}^n)$  to the function  $x \mapsto H(x_n + a)$  as  $k \rightarrow +\infty$ .

In Theorem 3 and throughout the paper,  $|\cdot|$  denotes the Euclidean norm in  $\mathbb{R}^n$ . We also use the following notations:  $B_R(x) = \{y \in \mathbb{R}^n : |y - x| < R\}$  and  $B_R = B_R(0)$  for  $R > 0$  and  $x \in \mathbb{R}^n$ , and we denote

$$\text{dist}(x, E) = \inf_{y \in E} |x - y|$$

for  $x \in \mathbb{R}^n$  and  $E \subset \mathbb{R}^n$ .

We complete the introduction by listing some corollaries following from Theorem 3.

**Corollary 4.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of the Allen-Cahn equation (1) in  $\mathbb{R}^n$ . Then there does not exist a non-degenerate cone containing  $\{u = 0\}$ .*

**Corollary 5.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of the Allen-Cahn equation (1) in  $\mathbb{R}^n$ . If  $\{u = 0\}$  is contained in a slab  $\{x \in \mathbb{R}^n : |x \cdot e| < A\}$  for some unit vector  $e$  and some real number  $A > 0$ , then there exists  $a \in \mathbb{R}$  such that either  $u(x) = H(-x \cdot e + a)$  or  $u(x) = H(x \cdot e + a)$ , for all  $x \in \mathbb{R}^n$ .*

Corollary 5 is similar to Theorem 2.1 by Farina [14]. Actually, the same conclusion was obtained in [14] under the additional assumption that both sets  $\{u > 0\}$  and  $\{u < 0\}$  are unbounded with respect to the direction  $e$ . We actually show in the proof of Corollary 5 at

the end of Section 2 that this last property is necessarily satisfied under the assumptions of Corollary 5, thanks to Theorem 3.

We remark also that Corollary 5 points out the difference between minimal surfaces and level sets of Allen-Cahn solutions from the point of view of half-space theorems: for  $n \geq 4$  the minimal catenoid is contained in a slab, while bounded entire solutions of the Allen-Cahn equation cannot have the zero level set contained in a slab. In particular, the zero level set of the solutions obtained in [1] is not contained in a slab, although such level set approaches the minimal catenoid in a compact region.

**Corollary 6.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of the Allen-Cahn equation (1) in  $\mathbb{R}^n$ . If there exists a closed half-space  $E$  such that  $\{u = 0\} \subset E$  and  $\{u = 0\} \cap \partial E \neq \emptyset$ , then there is  $a \in \mathbb{R}$  such that  $u(x) = H(x \cdot e + a)$  for all  $x \in \mathbb{R}^n$ , where  $e$  is a unit vector orthogonal to  $\partial E$ .*

It should be interesting to link such kind of half-space results to the De Giorgi conjecture. In this sense, an open question is to understand if Theorem 1 can be true in all dimensions, at least with the hypothesis of the monotonicity of the solution in one direction. Note, in particular, that the zero level sets of the  $x_n$ -monotone and non-planar solutions of (1) in  $\mathbb{R}^n$  with  $n \geq 9$ , constructed by Del Pino, Kowalczyk and the fifth author in [13], are not included in any half-space.

**Remark 7.** Theorem 1 and Corollary 4 and 5 do not hold good if the zero level set  $\{u = 0\}$  is replaced by another level set  $\{u = \mu\}$  with  $\mu \neq 0$ . Similarly, Theorem 3 does not hold good either if one assumes that  $u < \mu$  in  $\{x_n < 0\}$  for some  $\mu > 0$ . Indeed, equation (1) admits solutions  $u_L$  which vanish on  $(L\mathbb{Z})^n$  with  $L > \pi\sqrt{n}$ , are  $2L$ -periodic with respect to each variable  $x_i$ , which satisfy  $\max_{\mathbb{R}^n} |u_L| \rightarrow 0$  as  $L \xrightarrow{\geq} \pi\sqrt{n}$ , and which are not one-dimensional!

**Outline of the paper.** Section 2 is devoted to the proof of Theorem 3 and its corollaries. The proof of Theorem 3 is itself used in the proof of Theorem 1 and Corollary 2 done in Section 3. Lastly, Section 4 is concerned with a half-space theorem for a related free boundary problem.

## 2. HALF-SPACE THEOREMS IN GENERAL DIMENSION: PROOF OF THEOREM 3 AND ITS COROLLARIES

The half-space theorem of minimal hypersurfaces in  $\mathbb{R}^n$  is not true when  $n \geq 4$ , because the higher dimensional catenoids lie in a half-space. For the Allen-Cahn equation (1), we still do not know whether there is version of the half-space theorem in dimension  $n \geq 4$ . We have obtained partial classification results in this direction, based on the maximum principle.

We start this section with a general property holding in any dimension  $n \geq 1$ . This can essentially be found in [5, Lemmas 3.2 and 3.3] and [14, Lemma 2.3].

**Proposition 8.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of (1) in any dimension  $n \geq 1$ . Then  $\{u = 0\} \neq \emptyset$  and  $|u(x)| \rightarrow 1$  as  $\text{dist}(x, \{u = 0\}) \rightarrow +\infty$ .*

*Proof.* The proof is standard, we briefly sketch it for the sake of completeness. We recall from the introduction that  $u$  necessarily satisfies  $-1 < u < 1$  in  $\mathbb{R}^n$  and  $\{u = 0\} \neq \emptyset$ . Consider any  $R > 0$  and  $x \in \mathbb{R}^n$  such that

$$\text{dist}(x, \{u = 0\}) > R$$

(hence,  $\overline{B(x, R)} \cap \{u = 0\} = \emptyset$ ).

Let  $\lambda_R$  be the principal eigenvalue of  $-\Delta$  in the ball  $B_R$  with Dirichlet boundary condition, that is, there is a function  $\varphi_R \in C^2(\overline{B_R})$  such that  $-\Delta\varphi_R = \lambda_R\varphi_R$  in  $\overline{B_R}$ ,  $\varphi_R = 0$  on  $\partial B_R$ , and  $\varphi_R > 0$  in  $B_R$ . From the classical radial symmetry result [19], the function  $\varphi_R$  is radially symmetric and decreasing with respect to the distance from the origin. Therefore, up to multiplication by a positive constant, one can assume without loss of generality that  $\varphi_R(0) = 1 = \max_{\overline{B_R}} \varphi_R$ . Notice also that  $\lambda_R = \lambda_1/R^2$ . Since the conclusion is concerned with the limit as  $\text{dist}(x, \{u = 0\}) \rightarrow +\infty$ , one can assume without loss of generality that  $R > 0$  is large enough so that  $0 < \lambda_R < 1$ .

In the closed ball  $\overline{B(x, R)}$ , the continuous function  $u$  does not vanish. Up to changing  $u$  into  $-u$ , let us assume without loss of generality that  $u > 0$  in  $\overline{B(x, R)}$ . There exists then  $\varepsilon_0$  such that  $\varepsilon\varphi_R(\cdot - x) < u$  in  $\overline{B(x, R)}$ , for all  $\varepsilon \in [0, \varepsilon_0]$ . Furthermore, for every  $\varepsilon \in [0, \sqrt{1 - \lambda_R}]$ , the function  $\varepsilon\varphi_R(\cdot - x)$  satisfies

$$\Delta(\varepsilon\varphi_R(\cdot - x)) + \varepsilon\varphi_R(\cdot - x) - (\varepsilon\varphi_R(\cdot - x))^3 = \varepsilon\varphi_R(\cdot - x) \times (1 - \lambda_R - (\varepsilon\varphi_R(\cdot - x))^2) \geq 0$$

in  $\overline{B(x, R)}$ , since  $0 \leq \varphi_R \leq 1$  in  $\overline{B_R}$ . As a consequence, for any such  $\varepsilon$ , the function  $\varepsilon\varphi_R(\cdot - x)$  is a subsolution of (1) in the closed ball  $\overline{B(x, R)}$  and it vanishes on  $\partial B(x, R)$ , while  $u > 0$  in  $\overline{B(x, R)}$ . It follows from the strong maximum principle that  $\varepsilon\varphi_R(\cdot - x) < u$  in  $\overline{B(x, R)}$  for every  $\varepsilon \in [0, \sqrt{1 - \lambda_R}]$ . In particular,  $u(x) > \sqrt{1 - \lambda_R}$  since  $\varphi_R(0) = 1$ . Since  $\lambda_R = \lambda_1/R^2 \rightarrow 0$  as  $R \rightarrow +\infty$  and  $-1 < u < 1$  in  $\mathbb{R}^n$ , the conclusion follows.  $\square$

**Remark 9.** Proposition 8 implies that  $\sup_{\mathbb{R}^n} |u| = 1$  if  $\sup_{x \in \mathbb{R}^n} \text{dist}(x, \{u = 0\}) = +\infty$ . However, remember that the property  $\sup_{x \in \mathbb{R}^n} \text{dist}(x, \{u = 0\}) = +\infty$  is not always satisfied, since the Allen-Cahn equation admits non-trivial periodic solutions  $u$  oscillating around the value 0, and for which  $\sup_{\mathbb{R}^n} |u| < 1$ .

We are now in position to prove our Theorem 3.

*Proof of Theorem 3.* Throughout the proof,  $u$  is a bounded non-constant solution of (1) such that

$$u < 0 \text{ in } \mathbb{R}_-^n = \{x_n < 0\}.$$

As recalled in the introduction, we know that  $-1 < u < 1$  in  $\mathbb{R}^n$ .

By Proposition 8, we have that

$$(2) \quad u(x_1, \dots, x_n) \rightarrow -1 \text{ as } x_n \rightarrow -\infty \text{ uniformly in } (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}.$$

since the function  $u$  is negative in  $\mathbb{R}_-^n$ .

Denote  $U(x) = H(x_n) = \tanh(x_n/\sqrt{2})$  and

$$U^\omega(x) = U(x_1, \dots, x_{n-1}, x_n + \omega) = H(x_n + \omega) = \tanh\left(\frac{x_n + \omega}{\sqrt{2}}\right)$$

for  $x \in \mathbb{R}^n$  and  $\omega \in \mathbb{R}$ . We shall now show that  $u \leq U^\omega$  in  $\mathbb{R}^n$  for all  $\omega$  large enough. To do so, let  $A > 0$  be such that

$$(3) \quad u \leq -\frac{1}{\sqrt{3}} \text{ in } \mathbb{R}^{n-1} \times (-\infty, -A] \quad \text{and} \quad U \geq \frac{1}{\sqrt{3}} \text{ in } \mathbb{R}^{n-1} \times [A, +\infty).$$

We claim that

$$u \leq U^\omega \text{ in } \mathbb{R}^n \text{ for all } \omega \geq 2A.$$

To do so, pick any  $\omega \in [2A, +\infty)$ . We shall prove that  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times (-\infty, -A]$ . Assume by way of contradiction that  $M := \sup_{\mathbb{R}^{n-1} \times (-\infty, -A]} (u - U^\omega) > 0$ . Then there is a sequence  $(z_k)_{k \in \mathbb{N}} = (z'_k, z_{k,n})_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times (-\infty, -A]$  such that

$$u(z_k) - U^\omega(z_k) \rightarrow M > 0 \text{ as } k \rightarrow +\infty.$$

By (2), the sequence  $(z_{k,n})_{k \in \mathbb{N}}$  is then bounded. Furthermore, by uniform continuity of  $U$  (or of  $u$ ), property (3) and the assumption  $\omega \geq 2A$  imply that

$$\limsup_{k \rightarrow +\infty} z_{k,n} < -A.$$

Therefore, there is  $\zeta \in (-\infty, -A)$  such that, up to extraction of a subsequence,  $z_{k,n} \rightarrow \zeta$  as  $k \rightarrow +\infty$ . Up to extraction of another subsequence, the functions  $x \mapsto u(x' + z'_k, x_n)$  converge in  $C_{loc}^2(\mathbb{R}^n)$  as  $k \rightarrow +\infty$  to a solution  $w_\infty$  of (1) such that  $w_\infty \leq -1/\sqrt{3}$  and  $w_\infty - U^\omega \leq M$  in  $\mathbb{R}^{n-1} \times (-\infty, -A]$ , while  $w_\infty(0, \zeta) - U^\omega(0, \zeta) = M$ . At the (interior) point  $(0, \zeta) \in \mathbb{R}^{n-1} \times (-\infty, -A)$ , there holds

$$(4) \quad 0 \geq \Delta(w_\infty - U^\omega)(0, \zeta) = -w_\infty(0, \zeta) + w_\infty(0, \zeta)^3 + U^\omega(0, \zeta) - (U^\omega(0, \zeta))^3.$$

But  $-1 < U^\omega(0, \zeta) < U^\omega(0, \zeta) + M = w_\infty(0, \zeta) \leq -1/\sqrt{3}$  and the function  $s \mapsto s - s^3$  is decreasing in  $[-1, -1/\sqrt{3}]$ . Hence the right-hand side of (4) is positive, a contradiction. Therefore,

$$\sup_{\mathbb{R}^{n-1} \times (-\infty, -A]} (u - U^\omega) \leq 0,$$

that is,  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times (-\infty, -A]$ . Similarly, since  $U^\omega \geq 1/\sqrt{3}$  in  $\mathbb{R}^{n-1} \times [-A, +\infty)$  and  $U^\omega \geq u$  on  $\mathbb{R}^{n-1} \times \{-A\}$ , while the function  $s \mapsto s - s^3$  is decreasing in  $[1/\sqrt{3}, 1]$ , one can show that  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times [-A, +\infty)$ . Finally,

$$u \leq U^\omega \text{ in } \mathbb{R}^n \text{ for all } \omega \geq 2A.$$

Define now

$$a = \inf \{ \omega \in \mathbb{R} : u \leq U^\omega \text{ in } \mathbb{R}^n \}.$$

The previous paragraph yields  $a \leq 2A$ . On the other hand, since  $U^\omega \rightarrow -1$  as  $\omega \rightarrow -\infty$  (at least) pointwise in  $\mathbb{R}^n$ , while  $u > -1$  in  $\mathbb{R}^n$ , one infers that  $a \in \mathbb{R}$ . By continuity, there holds  $u \leq U^a$  in  $\mathbb{R}^n$ , that is,

$$(5) \quad u(x) \leq H(x_n + a) \text{ for all } x \in \mathbb{R}^n.$$

This statement corresponds to the first part of the conclusion of Theorem 3.

Let us now show the second part of the conclusion. First of all, if there is a point  $x^* \in \mathbb{R}^n$  such that  $u(x^*) = U^a(x^*) = H(x_n^* + a)$ , then the strong maximum principle implies that  $u \equiv U^a$  in  $\mathbb{R}^n$ , that is,

$$u(x) = H(x_n + a) \text{ for all } x \in \mathbb{R}^n.$$

Let us then assume in the sequel that  $u < U^a$  in  $\mathbb{R}^n$ , that is,

$$(6) \quad u(x) < U^a(x) = H(x_n + a) \text{ for all } x \in \mathbb{R}^n.$$

Let  $B > 0$  be such that  $U^a \geq 2/3 (> 1/\sqrt{3})$  in  $\mathbb{R}^{n-1} \times [B, +\infty)$ . We claim that

$$(7) \quad \sup_{\mathbb{R}^{n-1} \times [-A, B]} (u - U^a) = 0.$$

Indeed, otherwise, one would have  $\sup_{\mathbb{R}^{n-1} \times [-A, B]} (u - U^a) < 0$  and, by uniform continuity of  $U$ , there would exist  $\omega \in (-\infty, a)$  such that  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times [-A, B]$  and  $U^\omega \geq 1/\sqrt{3}$  in  $\mathbb{R}^{n-1} \times [B, +\infty)$ . With the same arguments as in the previous paragraph, one gets that  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times (-\infty, -A]$  and  $u \leq U^\omega$  in  $\mathbb{R}^{n-1} \times [B, +\infty)$ . As a consequence,  $u \leq U^\omega$  in  $\mathbb{R}^n$ , contradicting the minimality of  $a$ . Therefore, (7) holds.

From (7), one infers the existence of a sequence  $(\xi_k)_{k \in \mathbb{N}} = (\xi'_k, \xi_{k,n})_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times [-A, B]$  such that

$$(8) \quad u(\xi_k) - U^a(\xi_k) \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

Up to extraction of a subsequence, one can assume that  $\xi_{k,n} \rightarrow \xi_{\infty,n}$  as  $k \rightarrow +\infty$ , for some  $\xi_{\infty,n} \in [-A, B]$ . Notice that  $|\xi'_k| \rightarrow +\infty$  as  $k \rightarrow +\infty$ , since otherwise there would exist a point  $\xi \in \mathbb{R}^{n-1} \times [-A, B]$  such that  $u(\xi) = U^a(\xi)$ , contradicting (6). Denote

$$y_k = (\xi'_k, 0) \in \mathbb{R}^{n-1} \times \{0\} \text{ and } u_k(x) = u(x + y_k) \text{ for } k \in \mathbb{N} \text{ and } x \in \mathbb{R}^n.$$

To complete the proof of Theorem 3, we just need to show that

$$u_k(x) \rightarrow H(x_n + a)$$

in  $C_{loc}^2(\mathbb{R}^n)$  as  $k \rightarrow +\infty$ . Up to extraction of a subsequence, the functions  $u_k$  converge in  $C_{loc}^2(\mathbb{R}^n)$  to a solution  $u_\infty : \mathbb{R}^n \rightarrow [-1, 1]$  of (1) such that  $u_\infty(x) \leq H(x_n + a)$  in  $\mathbb{R}^n$  from (5) and the definition of  $y_k$ . Furthermore,  $u_\infty(0, \xi_{\infty,n}) = U^a(0, \xi_{\infty,n}) = H(\xi_{\infty,n} + a)$  by (8). It then follows from the strong maximum principle that  $u_\infty(x) = H(x_n + a)$  for all  $x \in \mathbb{R}^n$ . Furthermore, the limit of the functions  $u_k$  being independent of the subsequence, one concludes that the whole sequence  $(u_k)_{k \in \mathbb{N}}$  converges to the function  $x \mapsto H(x_n + a)$  in  $C_{loc}^2(\mathbb{R}^n)$  as  $k \rightarrow +\infty$ . The proof of Theorem 3 is thereby complete.  $\square$

**Remark 10.** In Theorem 3, one has  $-1 < u(x) < H(x_n + a) = \tanh((x_n + a)/\sqrt{2})$  for all  $x \in \mathbb{R}^n$ . Modica's inequality  $|\nabla u|^2 \leq (1 - u^2)^2/2$  (see [25]) then yields

$$(9) \quad |\nabla u(x)| \leq 2\sqrt{2} e^{\sqrt{2}(x_n + a)} \text{ for all } x \in \mathbb{R}^n.$$

By changing  $u$  into  $-u$  in Theorem 3, the following result immediately follows.

**Theorem 11.** *Let  $n \geq 1$  and  $u$  be a bounded non-constant solution of the Allen-Cahn equation (1) in  $\mathbb{R}^n$ . If  $u > 0$  in the half-space  $\{x_n < 0\}$ , then there exists  $a \in \mathbb{R}$  such that*

$$u(x) \geq H(-x_n + a)$$



for all  $x \in \mathbb{R}^n$ , and either  $u(x) = H(-x_n + a)$  for all  $x \in \mathbb{R}^n$ , or  $u(x) > H(-x_n + a)$  for all  $x \in \mathbb{R}^n$  and there exists a sequence  $(y_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times \{0\}$  such that  $|y_k| \rightarrow +\infty$  as  $k \rightarrow +\infty$ , and the functions  $u(\cdot + y_k)$  converge in  $C_{loc}^2(\mathbb{R}^n)$  to the function  $x \mapsto H(-x_n + a)$  as  $k \rightarrow +\infty$ .

Let us now turn to the proof of Corollaries 4, 5 and 6, which follow from Theorems 3 and 11.

*Proof of Corollary 4.* Assume by way of contradiction that  $u$  is a bounded non-constant solution of (1) with  $\{u = 0\}$  contained in a non-degenerate cone. Then, up to changing  $u$  into  $-u$ , and up to translation and rotation of the variables, it follows that

$$(10) \quad u < 0 \quad \text{in} \quad \{(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : x_n < \beta |x'|\},$$

for some  $\beta > 0$ . Theorem 3, together with (10), then yields the existence of a real number  $a$  such that  $u(x) = H(x_n + a)$  for all  $x \in \mathbb{R}^n$ , which leads to a contradiction.  $\square$

*Proof of Corollary 5.* Up to changing  $u$  into  $-u$  and up to translation and rotation of the variables, one can also assume without loss of generality that  $e = (0, \dots, 0, 1)$ , that  $u < 0$  in  $\{x_n \leq 0\}$  and that

$$\{u = 0\} \subset \{0 < x_n < 2A\}.$$

It then follows from Theorem 3 that there exists  $b \in \mathbb{R}$  such that

$$u(x) \leq H(x_n + b)$$

for all  $x \in \mathbb{R}^n$  and either  $u(x) = H(x_n + b)$  for all  $x \in \mathbb{R}^n$ , or  $u(\cdot + y_k) \rightarrow H(x_n + b)$  in  $C_{loc}^2(\mathbb{R}^n)$  as  $k \rightarrow +\infty$ , for some sequence  $(y_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times \{0\}$ . In both cases, one has  $\sup_{\mathbb{R}^{n-1} \times (-\infty, 2A]} u \leq H(2A + b) < 1$  and  $\sup_{\mathbb{R}^n} u = \sup_{\mathbb{R}} H = 1$ . By continuity, one infers that  $u > 0$  in  $\{x_n \geq 2A\}$ . From Theorem 3 applied to the solution

$$x = (x', x_n) \mapsto -u(x', -x_n + 2A),$$

there exists then  $c \in \mathbb{R}$  such that  $-u(x', -x_n + 2A) \leq H(x_n + c)$  for all  $x \in \mathbb{R}^n$ , hence

$$u(x) \geq H(x_n - 2A - c)$$

for all  $x \in \mathbb{R}^n$ . Finally  $H(x_n - 2A - c) \leq u(x) \leq H(x_n + b)$  for all  $x \in \mathbb{R}^n$  and one concludes from [6, Theorem 3.1] or [14, Theorem 2.1] that  $u(x) \equiv H(x_n + a)$  in  $\mathbb{R}^n$ , for some  $a \in \mathbb{R}$ .  $\square$

*Proof of Corollary 6.* Up to changing  $u$  into  $-u$  and up to translation and rotation of the variables, one can also assume without loss of generality that  $e = (0, \dots, 0, 1)$ , that  $u < 0$  in  $\{x_n < 0\}$  and that  $u(x', 0) = 0$  for some  $x' \in \mathbb{R}^{n-1}$ . Theorem 3 then implies that  $u(x) \leq H(x_n + a)$  in  $\mathbb{R}^n$ , for some  $a \in \mathbb{R}$ , and the other parts of the conclusion hold for that real number  $a$ . We claim that  $u(x', 0) = H(a)$ . Indeed, if not, then  $0 = u(x', 0) < H(a)$ , hence  $a > 0$ , while Theorem 3 also yields the existence of a sequence  $(y_k)_{k \in \mathbb{N}} = (y'_k, 0)_{k \in \mathbb{N}}$  in  $\mathbb{R}^{n-1} \times \{0\}$  such that  $u(\cdot + y_k) \rightarrow H(x_n + a)$  in  $C_{loc}^2(\mathbb{R}^n)$  as  $k \rightarrow +\infty$ . In particular,  $u(y'_k, -a/2) \rightarrow H(a/2) > 0$  as  $k \rightarrow +\infty$ , hence  $u(y'_k, -a/2) > 0$  for all  $k$  large enough, which is impossible since  $u < 0$  in  $\{x_n < 0\}$ . Therefore,  $u(x', 0) = H(a)$  (hence,  $a = 0$ ) and  $u(x) \equiv H(x_n)$  in  $\mathbb{R}^n$  from Theorem 3.  $\square$

3. PROOF OF THE HALF-SPACE THEOREM IN DIMENSIONS  $n = 2, 3$

As we mentioned in Section 1, the proof of the half-space theorem for minimal surfaces uses the family of catenoids and the sweeping principle. In the Allen-Cahn case, the solutions are defined in the whole space, and it is not easy to apply this idea.

We remark that the  $n = 2$  case of Theorem 1 can also be proven by applying the method in De Silva and Savin [11]. Our proof uses Pohozaev identity (also called balancing condition, see [12]) and is very different from theirs.

We shall prove Theorem 1 for  $n = 1, 2, 3$ . Notice that if  $n = 1$ , then the solution  $u$  is trivially one-dimensional and since it is not constant, it is then equal to  $H(x_1)$  up to shifts, as follows directly from ODE analysis. The cases of  $n = 2, 3$  are more complicated. Although we can deal with these two cases in a unified way, we choose to first give a simple proof when  $n = 2$ , because this gives us a clear geometric intuition behind the whole proof.

**3.1. The case  $n = 2$ .** Let  $u$  be a solution whose zero level set  $\{u = 0\}$  is contained in the half-space  $\{x_2 > 0\}$ . Up to changing  $u$  into  $-u$  and/or  $x_1$  into  $-x_1$ , and shifting in the direction  $x_2$ , we may assume without loss of generality that  $u < 0$  in  $\{x_2 < 0\}$  and, from Theorem 3, that

$$(11) \quad u(x_1, x_2) \leq H(x_2) \quad \text{for all } (x_1, x_2) \in \mathbb{R}^2$$

and that there exists a sequence  $(t_k^+)_{k \in \mathbb{N}}$  such that  $t_k^+ \rightarrow +\infty$  and  $u(x_1 + t_k^+, x_2) \rightarrow H(x_2)$  in  $C_{loc}^2(\mathbb{R}^2)$  as  $k \rightarrow +\infty$ .

For each  $x_1 \in \mathbb{R}$ , we define

$$g(x_1) = \inf \{x_2 \in \mathbb{R} : u(x_1, x_2) = 0\} \in [0, +\infty].$$

Note that the infimum is a minimum if  $g(x_1)$  is a real number. Note also that  $g(x_1)$  might a priori be  $+\infty$  for some values  $x_1$ , in which case  $u(x_1, x_2) < 0$  for all  $x_2 \in \mathbb{R}$  (nevertheless, the conclusion  $u(x_1, x_2) \equiv H(x_2)$  in  $\mathbb{R}^2$  will show that this case is impossible). We know at this point that  $g$  cannot be equal to  $+\infty$  on  $(-\infty, \xi)$  for some  $\xi \in \mathbb{R}$  since otherwise the zero level set of  $u$  would be included in the quarter-plane  $\{x_1 \geq \xi, x_2 \geq 0\}$ , which is ruled out by Corollary 4. Let us set

$$\alpha = \liminf_{x_1 \rightarrow -\infty} g(x_1) \in [0, +\infty].$$

Let us first consider the case  $0 \leq \alpha < +\infty$ . There exists then a sequence  $(t_k^-, s_k^-)_{k \in \mathbb{N}}$  such that  $u(t_k^-, s_k^-) = 0$ , and  $t_k^- \rightarrow -\infty$  and  $s_k^- \rightarrow \alpha$  as  $k \rightarrow +\infty$ . Up to extraction of a subsequence, the functions  $(x_1, x_2) \mapsto u(x_1 + t_k^-, x_2 + \alpha)$  converge in  $C_{loc}^2(\mathbb{R}^2)$  to a bounded solution  $u_\infty$  of (1) such that  $u_\infty(0, 0) = 0$  and  $u_\infty \leq 0$  in  $\{x_2 \leq 0\}$ , owing to the definition of  $\alpha$ . Furthermore,  $u_\infty(x_1, x_2) \leq H(x_2 + \alpha)$  in  $\mathbb{R}^2$  from (11), hence  $u_\infty(x_1, -\infty) = -1$  for each  $x_1 \in \mathbb{R}$ , and  $u_\infty < 0$  in  $\{x_2 < 0\}$  from the strong maximum principle. Corollary 6 then implies that  $u_\infty(x_1, x_2) \equiv H(x_2 + b)$  in  $\mathbb{R}^2$  for some real number  $b$ . Since  $u_\infty(0, 0) = 0$ , one finally infers that  $b = 0$  and  $u_\infty(x_1, x_2) \equiv H(x_2)$  in  $\mathbb{R}^2$ .

Consider now the semi-infinite vertical strip

$$\Omega_k := \{(x_1, x_2) \in \mathbb{R}^2 : t_k^- < x_1 < t_k^+, -\infty < x_2 < \alpha\}.$$

Let  $X = (0, 1)$ . The balancing condition (see [12, Appendix]) tells us that

$$(12) \quad \int_{\partial\Omega_k} \left[ \left( \frac{1}{2} |\nabla u|^2 + F(u) \right) X \cdot \nu - (\nabla u \cdot X) (\nabla u \cdot \nu) \right] d\sigma = 0,$$

where

$$F(s) = \frac{(1 - s^2)^2}{4}$$

and  $\nu$  is the outward unit normal of the domain  $\Omega_k$  (which is defined everywhere except at the corners  $(t_k^\pm, \alpha)$ ). The previous formula means that

$$\begin{aligned} & \int_{t_k^-}^{t_k^+} \left( \frac{|\nabla u(x_1, \alpha)|^2}{2} + F(u(x_1, \alpha)) - u_{x_2}^2(x_1, \alpha) \right) dx_1 \\ & \quad + \underbrace{\int_{-\infty}^{\alpha} (u_{x_1}(t_k^-, x_2) u_{x_2}(t_k^-, x_2) - u_{x_1}(t_k^+, x_2) u_{x_2}(t_k^+, x_2)) dx_2}_{=: I_k} = 0, \end{aligned}$$

where the second integral  $I_k$  converges absolutely from Remark 10. Since  $u(x_1 + t_k^-, x_2) \rightarrow H(x_2)$  and  $u(x_1 + t_k^-, x_2 + \alpha) \rightarrow H(x_2)$  as  $k \rightarrow +\infty$  in  $C_{loc}^2(\mathbb{R}^2)$ , together with Remark 10, it follows that  $I_k \rightarrow 0$  as  $k \rightarrow +\infty$ . Therefore,

$$\int_{t_k^-}^{t_k^+} \left( \frac{|\nabla u(x_1, \alpha)|^2}{2} + F(u(x_1, \alpha)) - u_{x_2}^2(x_1, \alpha) \right) dx_1 \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

On the other hand, Modica's inequality

$$F(u) \geq \frac{|\nabla u|^2}{2} \left( \geq \frac{u_{x_2}^2}{2} \right)$$

(see [25]) implies that

$$\frac{|\nabla u|^2}{2} + F(u) - u_{x_2}^2 = \frac{u_{x_1}^2}{2} + F(u) - \frac{u_{x_2}^2}{2} \geq \frac{u_{x_1}^2}{2} \geq 0$$

in  $\mathbb{R}^2$ . Since  $t_k^\pm \rightarrow \pm\infty$  as  $k \rightarrow +\infty$ , we finally get that

$$\frac{|\nabla u(x_1, \alpha)|^2}{2} + F(u(x_1, \alpha)) - u_{x_2}^2(x_1, \alpha) = \frac{u_{x_1}^2(x_1, \alpha)}{2} = 0$$

for all  $x_1 \in \mathbb{R}$ , hence

$$F(u(x_1, \alpha)) = \frac{u_{x_2}^2(x_1, \alpha)}{2} = \frac{|\nabla u(x_1, \alpha)|^2}{2}$$

for all  $x_1 \in \mathbb{R}$ . One concludes from [8, 25] that  $u$  is one-dimensional, namely  $u(x_1, x_2) \equiv H(x_2)$  in  $\mathbb{R}^2$  from the first paragraph of the proof.

Let us finally consider the case  $\alpha = +\infty$ . Here, remembering also that  $u < 0$  in  $\{x_2 < 0\}$ , it follows from Proposition 8 that

$$\sup_{x_1 \leq -R, x_2 \leq 0} u(x_1, x_2) \rightarrow -1$$

as  $R \rightarrow +\infty$ , hence  $|\nabla u(x_1, x_2)| \rightarrow 0$  as  $x_1 \rightarrow -\infty$  uniformly with respect to  $x_2 \leq 0$ , from standard elliptic estimates. Together with Remark 10, this implies that

$$\int_{-\infty}^0 u_{x_1}(x_1, x_2) u_{x_2}(x_1, x_2) dx_2 \rightarrow 0 \text{ as } x_1 \rightarrow -\infty.$$

Therefore, by applying (12) in the region

$$\{(x_1, x_2) \in \mathbb{R}^2 : -k < x_1 < t_k^+, -\infty < x_2 < 0\},$$

one gets with the same arguments as before that

$$\int_{-k}^{t_k^+} \left( \frac{|\nabla u(x_1, 0)|^2}{2} + F(u(x_1, 0)) - u_{x_2}^2(x_1, 0) \right) dx_1 \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

This leads to the same conclusion as in the previous paragraph and the proof of Theorem 1 in the case  $n = 2$  is thereby complete.

*Proof of Corollary 2.* Let  $u_1 < u_2$  be two bounded non-constant solutions of (1) in  $\mathbb{R}^2$ . Remember that  $-1 < u_1 < u_2 < 1$  in  $\mathbb{R}^2$ . Using  $u_1$  and  $u_2$  as barriers and applying minimizing arguments, we can construct a stable solution  $u_3$  of (1) with

$$-1 < u_1 \leq u_3 \leq u_2 < 1 \text{ in } \mathbb{R}^2.$$

In dimension two, stable solutions are one-dimensional, as follows from [4, Theorem 1.8]. The function  $u_3$  is then one-dimensional stable and it takes values in  $(-1, 1)$ , hence there exist then a unit vector  $e$  and a real number  $c$  such that  $u_3(x) \equiv H(x \cdot e + c)$  in  $\mathbb{R}^2$ . Therefore, the nodal set of  $u_1$  is contained in the half-space  $\{x \cdot e + c \geq 0\}$  and  $u_1 < 0$  in  $\{x \cdot e + c < 0\}$ . By the weak half-space Theorem 1,  $u_1$  has to be one-dimensional, and more precisely there is  $a \in \mathbb{R}$  such that  $u_1(x) \equiv H(x \cdot e + a)$  in  $\mathbb{R}^2$ . The same is true for  $u_2$ , with  $u_2(x) \equiv H(x \cdot e + b)$  in  $\mathbb{R}^2$ , for some real number  $b$  such that  $b > a$  (since  $u_2 > u_1$ ).  $\square$

**3.2. The case  $n = 3$ .** Next, we shall consider the case of dimension 3. The arguments in this section can also be applied in the two dimensional case, but we preferred to use the more direct proof of the previous section in the case  $n = 2$ .

First of all, as in the case  $n = 2$ , up to changing  $u$  into  $-u$  and/or shifting in the direction  $x_3$ , we may assume without loss of generality that  $u < 0$  in  $\{x_3 < 0\}$  and, from Theorem 3, that

$$(13) \quad -1 < u(x_1, x_2, x_3) \leq H(x_3) \text{ for all } (x_1, x_2, x_3) \in \mathbb{R}^3$$

and there is a sequence  $(y_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^2 \times \{0\}$  such that  $u(\cdot + y_k) \rightarrow H(x_3)$  in  $C_{loc}^2(\mathbb{R}^3)$  as  $k \rightarrow +\infty$ .

Now, let  $A > 0$  be such that

$$\tanh\left(-\frac{A}{\sqrt{2}}\right) \leq -\sqrt{\frac{2}{3}}.$$

For  $s > 0$ , let  $\Omega_s$  be the half-cylinder

$$\Omega_s = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1^2 + x_2^2 < s^2, x_3 < -A\}.$$

Let  $X = (0, 0, 1)$ . Then, for every  $s > 0$ , the following balancing formula (see [12]) holds in  $\Omega_s$ :

$$(14) \quad \int_{\partial\Omega_s} \left[ \left( \frac{1}{2} |\nabla u|^2 + F(u) \right) X \cdot \nu - (\nabla u \cdot X) (\nabla u \cdot \nu) \right] d\sigma = 0,$$

meaning that

$$(15) \quad \begin{aligned} g(s) &:= \int_{\{x_1^2 + x_2^2 < s^2\}} \left( \frac{|\nabla u(x_1, x_2, -A)|^2}{2} + F(u(x_1, x_2, -A)) - u_{x_3}^2(x_1, x_2, -A) \right) dx_1 dx_2 \\ &= s \int_{-\infty}^{-A} \int_0^{2\pi} \left( u_{x_1}(s \cos \theta, s \sin \theta, x_3) \cos \theta + u_{x_2}(s \cos \theta, s \sin \theta, x_3) \sin \theta \right) \\ &\quad \times u_{x_3}(s \cos \theta, s \sin \theta, x_3) d\theta dx_3. \end{aligned}$$

Notice that the integrals converge absolutely from Remark 10. As in the previous section, we infer from Modica's inequality

$$F(u) \geq \frac{|\nabla u|^2}{2}$$

that

$$(16) \quad \frac{|\nabla u|^2}{2} + F(u) - u_{x_3}^2 = \frac{u_{x_1}^2 + u_{x_2}^2}{2} + F(u) - \frac{u_{x_3}^2}{2} \geq \frac{u_{x_1}^2 + u_{x_2}^2}{2} \geq 0$$

in  $\mathbb{R}^3$ . From this, we know that the function  $g$  is nonnegative and nondecreasing in  $(0, +\infty)$ . Hence we can define

$$(17) \quad \alpha = \lim_{s \rightarrow +\infty} g(s) \in [0, +\infty].$$

Let us also define a function  $K : (0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$(18) \quad K(s, x_3) = \int_{\{x_1^2 + x_2^2 < s^2\}} \left( u_{x_1}^2(x_1, x_2, x_3) + u_{x_2}^2(x_1, x_2, x_3) \right) dx_1 dx_2$$

and notice from the definition of  $g$  in (15) and from (16) that

$$(19) \quad K(s, -A) \leq 2g(s) \quad \text{for all } s > 0.$$

Remark 10 yields suitable exponential decay of  $u_{x_1}^2 + u_{x_2}^2$  as  $x_3 \rightarrow -\infty$ , from which we get the following key-property of  $K$ .

**Lemma 12.** *There exist some constants  $C > 0$  and  $\beta > 0$ , such that*

$$K(s, x_3) \leq C(1 + K(s + \beta \ln s, -A)), \quad \text{for all } s \geq 1 \text{ and } x_3 \leq -A.$$

*Proof.* The functions  $u_{x_1}$  and  $u_{x_2}$  satisfy

$$-\Delta u_{x_i} + (3u^2 - 1)u_{x_i} = 0$$

in  $\mathbb{R}^3$ . Thanks to (13), the function  $u^2$  converges to 1 as  $x_3 \rightarrow -\infty$ , hence the operator  $-\Delta + (3u^2 - 1)$  tends to  $-\Delta + 2$ . As a matter of fact, in  $\mathbb{R}^2 \times (-\infty, -A]$ , one has

$$-1 < u \leq H(-A) = \tanh\left(-\frac{A}{\sqrt{2}}\right) \leq -\sqrt{\frac{2}{3}},$$

thus  $3u^2 - 1 \geq 1$  in  $\mathbb{R}^2 \times (-\infty, -A]$ .

We are then going to compare  $u_{x_1}$  (and later  $u_{x_2}$ ) in  $\mathbb{R}^2 \times (-\infty, -A]$  to the bounded solution  $\phi$  of the model problem

$$\begin{cases} -\Delta\phi + \phi = 0, & (x_1, x_2, x_3) \in \mathbb{R}^2 \times (-\infty, -A), \\ \phi(x_1, x_2, -A) = |u_{x_1}(x_1, x_2, -A)|, & (x_1, x_2) \in \mathbb{R}^2. \end{cases}$$

By using the Fourier transform in the  $(x_1, x_2)$  variables, the bounded solution  $\phi$  of the above problem is given in  $\mathbb{R}^2 \times (-\infty, -A)$  by

$$\phi(x_1, x_2, x_3) = \int_{\mathbb{R}^2} \phi(x'_1, x'_2, -A) G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2,$$

where, for  $X_3 < 0$ ,  $G(\cdot, \cdot, X_3)$  is the inverse Fourier transform of the function

$$(\xi_1, \xi_2) \mapsto e^{\sqrt{\xi_1^2 + \xi_2^2 + 1} X_3},$$

that is,

$$G(X_1, X_2, X_3) = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} e^{i(X_1\xi_1 + X_2\xi_2) + \sqrt{\xi_1^2 + \xi_2^2 + 1} X_3} d\xi_1 d\xi_2$$

for  $(X_1, X_2, X_3) \in \mathbb{R}^2 \times (-\infty, 0)$ . Note that we are interested in estimating  $\phi(\cdot, \cdot, x_3)$  in the disks

$$D_s := \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 < s^2\},$$

with  $s \geq 1$  and  $x_3 < -A$ . To do so, for  $s \geq 1$  and  $(x_1, x_2, x_3) \in \mathbb{R}^2 \times (-\infty, -A)$ , we divide  $\phi$  into two parts:

$$\begin{aligned} \phi(x_1, x_2, x_3) &= \underbrace{\int_{\{|(x'_1, x'_2)| \geq s + \beta \ln s\}} \phi(x'_1, x'_2, -A) G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2}_{=:\phi_{1,s}(x_1, x_2, x_3)} \\ &+ \underbrace{\int_{\{|(x'_1, x'_2)| < s + \beta \ln s\}} \phi(x'_1, x'_2, -A) G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2}_{=:\phi_{2,s}(x_1, x_2, x_3)}, \end{aligned}$$

where  $\beta > 0$  will be chosen later.

On the one hand, since

$$|\nabla u| \leq \sqrt{2F(u)} \leq \frac{1}{\sqrt{2}}$$

in  $\mathbb{R}^3$  from Modica's inequality [25], and since  $G \geq 0$  in  $\mathbb{R}^2 \times (-\infty, 0)$  and  $\phi \geq 0$  in  $\mathbb{R}^2 \times (-\infty, -A)$  from the maximum principle, it follows that, for all  $s \geq 1$ ,  $(x_1, x_2) \in D_s$  and  $x_3 < -A$ ,

$$\begin{aligned} 0 \leq \phi_{1,s}(x_1, x_2, x_3) &\leq \frac{1}{\sqrt{2}} \int_{\{|(x'_1, x'_2)| \geq s + \beta \ln s\}} G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2 \\ &\leq \frac{1}{\sqrt{2}} \underbrace{\int_{\{|(x'_1, x'_2)| \geq \beta \ln s\}} G(-x'_1, -x'_2, x_3 + A) dx'_1 dx'_2}_{=:\bar{\phi}_s(0, 0, x_3)} \end{aligned}$$

where  $\bar{\phi}_s$  denotes the bounded solution of

$$(20) \quad \begin{cases} -\Delta \bar{\phi}_s + \bar{\phi}_s = 0, & (x_1, x_2, x_3) \in \mathbb{R}^2 \times (-\infty, -A), \\ \bar{\phi}_s(x_1, x_2, -A) = \begin{cases} \frac{1}{\sqrt{2}} & \text{if } |(x_1, x_2)| \geq \beta \ln s, \\ 0 & \text{otherwise.} \end{cases} \end{cases}$$

It is immediate to check that there is a real number  $\gamma > 0$  small enough so that the function

$$(x_1, x_2, x_3) \mapsto e^{\gamma \sqrt{x_1^2 + x_2^2 + 1}}$$

satisfies

$$-\Delta \bar{\phi} + \bar{\phi} \geq 0$$

in  $\mathbb{R}^3$ . Hence, the function

$$(x_1, x_2, x_3) \mapsto \frac{1}{\sqrt{2}} e^{\gamma \sqrt{x_1^2 + x_2^2 + 1} - \gamma \sqrt{(\beta \ln s)^2 + 1}}$$

is a supersolution of (20), and

$$0 \leq \bar{\phi}_s(x_1, x_2, x_3) \leq \frac{e^{\gamma \sqrt{x_1^2 + x_2^2 + 1} - \gamma \sqrt{(\beta \ln s)^2 + 1}}}{\sqrt{2}} \quad \text{for all } (x_1, x_2, x_3) \in \mathbb{R}^2 \times (-\infty, -A)$$

from the maximum principle. As a consequence,

$$0 \leq \phi_{1,s}(x_1, x_2, x_3) \leq \bar{\phi}_s(0, 0, x_3) \leq \frac{e^{\gamma - \gamma \beta \ln s}}{\sqrt{2}}$$

for all  $s \geq 1$ ,  $(x_1, x_2) \in D_s$  and  $x_3 < -A$ . Therefore, by choosing

$$\beta = \frac{2}{\gamma} > 0,$$

one gets that

$$(21) \quad \int_{D_s} \phi_{1,s}^2(x_1, x_2, x_3) dx_1 dx_2 \leq \frac{\pi e^{2\gamma}}{2s^2} \leq \frac{\pi e^{2\gamma}}{2} \quad \text{for all } s \geq 1 \text{ and } x_3 < -A.$$

On the other hand, remember that  $G \geq 0$  in  $\mathbb{R}^2 \times (-\infty, 0)$ , and notice that  $\|G(\cdot, \cdot, X_3)\|_{L^1(\mathbb{R}^2)} = e^{X_3} \leq 1$  for all  $X_3 < 0$ . Hence, using the Cauchy-Schwarz inequality

and Fubini's theorem, we have, for all  $s \geq 1$  and  $x_3 < -A$ ,

$$\begin{aligned}
& \int_{D_s} \phi_{2,s}^2(x_1, x_2, x_3) dx_1 dx_2 \\
&= \int_{D_s} \left( \int_{\{|(x'_1, x'_2)| < s + \beta \ln s\}} \phi(x'_1, x'_2, -A) G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2 \right)^2 dx_1 dx_2 \\
&\leq \int_{D_s} \left\{ \int_{\{|(x'_1, x'_2)| < s + \beta \ln s\}} \phi^2(x'_1, x'_2, -A) G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2 \right. \\
&\quad \left. \times \int_{\{|(x'_1, x'_2)| < s + \beta \ln s\}} G(x_1 - x'_1, x_2 - x'_2, x_3 + A) dx'_1 dx'_2 \right\} dx_1 dx_2 \\
&\leq \int_{\{|(x'_1, x'_2)| < s + \beta \ln s\}} \phi^2(x'_1, x'_2, -A) dx'_1 dx'_2 = \int_{D_{s+\beta \ln s}} \phi^2(x_1, x_2, -A) dx_1 dx_2.
\end{aligned}$$

Therefore, together with (21), it follows that, for all  $s \geq 1$  and  $x_3 < -A$ ,

$$\int_{D_s} \phi^2(x_1, x_2, x_3) dx_1 dx_2 \leq \pi e^{2\gamma} + 2 \int_{D_{s+\beta \ln s}} \phi^2(x_1, x_2, -A) dx_1 dx_2.$$

Lastly, since  $3u^2 - 1 \geq 1$  in  $\mathbb{R}^2 \times (-\infty, -A]$ , the maximum principle implies that  $|u_{x_1}| \leq \phi$  in  $\mathbb{R}^2 \times (-\infty, -A]$ , hence

$$\begin{aligned}
\int_{D_s} u_{x_1}^2(x_1, x_2, x_3) dx_1 dx_2 &\leq \pi e^{2\gamma} + 2 \int_{D_{s+\beta \ln s}} \phi^2(x_1, x_2, -A) dx_1 dx_2 \\
&= \pi e^{2\gamma} + 2 \int_{D_{s+\beta \ln s}} u_{x_1}^2(x_1, x_2, -A) dx_1 dx_2
\end{aligned}$$

for all  $s \geq 1$  and  $x_3 < -A$ . The same property holds similarly for the function  $u_{x_2}$ . Thus,

$$\begin{aligned}
& \int_{D_s} (u_{x_1}^2(x_1, x_2, x_3) + u_{x_2}^2(x_1, x_2, x_3)) dx_1 dx_2 \\
&\leq 2\pi e^{2\gamma} + 2 \int_{D_{s+\beta \ln s}} (u_{x_1}^2(x_1, x_2, -A) + u_{x_2}^2(x_1, x_2, -A)) dx_1 dx_2
\end{aligned}$$

for all  $s \geq 1$  and  $x_3 < -A$ , that is,

$$K(s, x_3) \leq 2\pi e^{2\gamma} + 2K(s + \beta \ln s, -A),$$

with  $K$  defined in (18). Notice that this last inequality also holds trivially with  $x_3 = -A$  since  $s + \beta \ln s \geq s$ . The proof of Lemma 12 is thereby complete with  $C = 2\pi e^{2\gamma} > 0$ .  $\square$

With a slight abuse of notation, we also write  $u$  in the polar coordinates (in the  $(x_1, x_2)$ -plane) as  $u(r, \theta, x_3)$ . Let us now define, for  $s > 0$ ,

$$(22) \quad f(s) := \int_0^s \int_{-\infty}^{-A} \int_0^{2\pi} u_r(r, \theta, x_3) u_{x_3}(r, \theta, x_3) d\theta dx_3 dr.$$

Note that the above integral converges absolutely for each  $s > 0$ , from Remark 10, and that  $f(s) \rightarrow 0$  as  $s \rightarrow 0$ .



**Lemma 13.** *The quantity  $\alpha$  defined in (17) is such that  $\alpha = 0$ .*

*Proof.* Let us assume by way of contradiction that  $\alpha > 0$ . Observe first that the function  $f$  is of class  $C^1$  in  $(0, +\infty)$  and that, thanks to (15),

$$f'(s) = \frac{g(s)}{s} \quad \text{for all } s > 0.$$

Using the fact that  $g$  is nonnegative and  $g(\tau) \rightarrow \alpha \in (0, +\infty]$  as  $\tau \rightarrow +\infty$ , we deduce that

$$(23) \quad f(s) = \int_0^s f'(\tau) d\tau = \int_0^s \frac{g(\tau)}{\tau} d\tau \geq \alpha' \ln s \quad \text{for all } s \text{ large enough,}$$

with, say,  $\alpha' = \alpha/2 > 0$  if  $0 < \alpha < +\infty$  and  $\alpha' = 1$  if  $\alpha = +\infty$ . On the other hand, one infers Remark 10, with here  $a = 0$  thanks to (13), that

$$(24) \quad |\nabla u| \leq 2\sqrt{2}e^{\sqrt{2}x_3} \quad \text{in } \mathbb{R}^3$$

and from (15) that

$$(25) \quad \frac{g(\tau)}{\tau} = \int_{-\infty}^{-A} \int_0^{2\pi} u_r(\tau, \theta, x_3) u_{x_3}(\tau, \theta, x_3) d\theta dx_3 \leq 4\sqrt{2}\pi \leq 6\pi$$

for all  $\tau > 0$ , hence

$$(26) \quad f(s + \beta \ln s) - f(s) = \int_s^{s+\beta \ln s} \frac{g(\tau)}{\tau} d\tau \leq 6\pi \beta \ln s \quad \text{for all } s \geq 1.$$

Using again (24) and (25), together with the definition (18) of  $K$  and the decomposition of the integral (22) with respect to  $r \in [0, s]$  into two integrals over  $[1, s]$  and  $[0, 1]$ , we get that, for all  $s \geq 1$ ,

$$\begin{aligned} f(s) &\leq \int_{-\infty}^{-A} \left[ \int_1^s \int_0^{2\pi} r u_r^2 d\theta dr \right]^{1/2} \left[ \int_1^s \int_0^{2\pi} \frac{u_{x_3}^2}{r} d\theta dr \right]^{1/2} dx_3 + 6\pi \\ &\leq \sqrt{16\pi \ln s} \int_{-\infty}^{-A} \sqrt{K(s, x_3)} e^{\sqrt{2}x_3} dx_3 + 6\pi. \end{aligned}$$

Applying inequality (19) and Lemma 12, we deduce that, for all  $s \geq 1$ ,

$$\begin{aligned} f(s) &\leq \sqrt{16\pi \ln s} \int_{-\infty}^{-A} \sqrt{C(1 + K(s + \beta \ln s, -A))} e^{\sqrt{2}x_3} dx_3 + 6\pi \\ &\leq \sqrt{8\pi C (\ln s) (1 + 2g(s + \beta \ln s))} + 6\pi \\ &= \sqrt{8\pi C (\ln s) (1 + 2f'(s + \beta \ln s) (s + \beta \ln s))} + 6\pi. \end{aligned}$$

Together with (23), it follows that  $f'(t)t \rightarrow +\infty$  as  $t \rightarrow +\infty$ , and that

$$0 < f(s) \leq \sqrt{17\pi C (s + \beta \ln s) \ln(s + \beta \ln s) f'(s + \beta \ln s)} \quad \text{for all } s \text{ large enough.}$$

Thanks to (23) and (26), we then infer that, for all  $s$  large enough,

$$\begin{aligned} 0 < f(s + \beta \ln s) &\leq f(s) + 6\pi\beta \ln s \leq \left(1 + \frac{6\pi\beta}{\alpha'}\right) f(s) \\ &\leq C_1 \sqrt{(s + \beta \ln s) \ln(s + \beta \ln s) f'(s + \beta \ln s)} \end{aligned}$$

with

$$C_1 = \sqrt{17\pi C} \left(1 + \frac{6\pi\beta}{\alpha'}\right) > 0.$$

In other words, there is  $t_0 > 0$  such that  $f > 0$  on  $[t_0, +\infty)$  and

$$\frac{f'(t)}{f(t)^2} \geq \frac{1}{C_1^2 t \ln t} \text{ for all } t \geq t_0.$$

It follows that the function

$$t \mapsto \frac{1}{f(t)} + \frac{\ln(\ln t)}{C_1^2}$$

is nonincreasing on  $[t_0, +\infty)$ . But since  $f > 0$  on  $[t_0, +\infty)$ , one has

$$\frac{1}{f(t)} + \frac{\ln(\ln t)}{C_1^2} \rightarrow +\infty$$

as  $t \rightarrow +\infty$ . This is a contradiction. Therefore,  $\alpha = 0$  and the proof of Lemma 13 is thereby complete.  $\square$

*End of the proof of Theorem 1 for  $n = 3$ .* As in the case  $n = 2$ , the fact that  $\alpha = 0$  in (15) and (17), together with (16), implies that

$$\begin{aligned} &\frac{|\nabla u(x_1, x_2, -A)|^2}{2} + F(u(x_1, x_2, -A)) - u_{x_3}^2(x_1, x_2, -A) \\ &= \frac{u_{x_1}^2(x_1, x_2, -A) + u_{x_2}^2(x_1, x_2, -A)}{2} = 0 \end{aligned}$$

for all  $(x_1, x_2) \in \mathbb{R}^2$ . Hence

$$F(u(x_1, x_2, -A)) = \frac{u_{x_3}^2(x_1, x_2, -A)}{2} = \frac{|\nabla u(x_1, x_2, -A)|^2}{2}$$

for all  $(x_1, x_2) \in \mathbb{R}^2$  and one concludes from [8, 25] that  $u$  is one-dimensional, namely  $u(x_1, x_2, x_3) \equiv H(x_3)$  in  $\mathbb{R}^3$  from the second paragraph of this subsection.  $\square$

**Remark 14.** In higher dimensions  $n \geq 4$ , one could still apply the balancing condition and define some functions  $g$  and  $f$  with formulas similar to (15) and (22) above. However, in (15), there would be a factor  $s^{n-2}$  instead of  $s$  in the right-hand side. Even if Lemma 12 still extends to that case (with a different value for the constant  $\beta$ ), Lemma 13 does not extend as such. In particular, one would have  $f'(s) = g(s)/s^{n-2}$ , and the integrability of the function  $1/s^{n-2}$  at infinity does not imply that  $f(+\infty) = +\infty$  if  $\alpha := g(+\infty) > 0$ , and then the end of the proof does not work.

## 4. HALF-SPACE THEOREMS FOR FREE BOUNDARY PROBLEMS

In this section, we are interested in half-space properties for free boundary problems. First of all, we consider the following classical one phase free boundary problem:

$$(27) \quad \begin{cases} \Delta u = 0 & \text{in } \Xi := \{u > 0\} \subset \mathbb{R}^n, \\ |\nabla u| = 1 & \text{on } \partial\Xi, \end{cases}$$

where  $u$  is understood in the classical sense in  $\bar{\Xi}$  and  $\partial\Xi$  is globally smooth.

The existence of catenoid type solutions of this problem has been proved using an Allen-Cahn approximation. We refer to [22] and the references therein for more discussion on this problem.

We have the following half-space property:

**Theorem 15.** *Let  $n \leq 3$  and  $u$  be a solution of (27) with  $|\nabla u| \leq 1$ . Suppose that the positive phase  $\Xi$  is contained in the half-space  $\{x_n > 0\}$ . Then  $u$  is one-dimensional, namely there is  $h \geq 0$  such that  $\Xi = \{x_n > h\}$  and  $u$  is the one-dimensional function  $u(x) \equiv x_n - h$  in  $\bar{\Xi}$ .*

*Proof.* The idea of proof is same as that of Section 3.2. We sketch the proof and list the necessary modifications. Let us only consider the case  $n = 3$ .

Up to shift in the  $x_3$ -direction, one can assume without loss of generality that  $\Xi$  is not contained in  $\{x_3 > a\}$  for any  $a > 0$ . From standard elliptic estimates up to the boundary, one can fix  $a > 0$  small enough such that  $u_{x_3} > 0$  in  $\bar{\Xi} \cap \{x_3 \leq a\}$ .

We still adopt the notation of Section 3.2 and, for  $s > 0$ , let  $\Omega_s$  be the half-cylinder

$$\Omega_s := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : r^2 = x_1^2 + x_2^2 < s^2, \quad x_3 < a\}.$$

For  $\varepsilon > 0$ , let us define

$$\bar{\Xi}_\varepsilon := \{Z \in \bar{\Xi} : \text{dist}(Z, \partial\Xi) > \varepsilon\}.$$

Let  $F$  be half the characteristic function of the interval  $(0, +\infty)$ , that is,  $F(\tau) = 1/2$  if  $\tau > 0$  and  $F(\tau) = 0$  if  $\tau \leq 0$ . Then we have the following balancing formula, with  $X = (0, 0, 1)$  and  $\varepsilon \in (0, a)$ :

$$\int_{\partial(\Omega_s \cap \bar{\Xi}_\varepsilon)} \left[ \left( \frac{1}{2} |\nabla u|^2 + F(u) \right) X \cdot \nu - (\nabla u \cdot X) (\nabla u \cdot \nu) \right] d\sigma = 0.$$

Sending  $\varepsilon$  to 0 in this identity and using the free boundary condition, we get

$$\int_{\partial\Omega_s \cap \bar{\Xi}} \left[ \left( \frac{1}{2} |\nabla u|^2 + F(u) \right) X \cdot \nu - (\nabla u \cdot X) (\nabla u \cdot \nu) \right] d\sigma = 0.$$

Now we extend the solution  $u$  to  $\mathbb{R}^3$  such that  $u = 0$  in  $\mathbb{R}^3 \setminus \bar{\Xi}$ . Still denote it as  $u$ . Then we get

$$\int_{\partial\Omega_s} \left[ \left( \frac{1}{2} |\nabla u|^2 + F(u) \right) X \cdot \nu - (\nabla u \cdot X) (\nabla u \cdot \nu) \right] d\sigma = 0.$$

Note that  $u$  is not smooth across the free boundary, but, for any  $Y \in \partial\Xi$ , the quantity  $(|\nabla u(Z)|^2/2 + F(u(Z))) X \cdot \nu(Y) - (\nabla u(Z) \cdot X) (\nabla u(Z) \cdot \nu(Y))$  converges to 0 as  $Z \rightarrow Y$  with  $Z \in \bar{\Xi}$  and it vanishes for all  $Z \in \mathbb{R}^3 \setminus \bar{\Xi}$ .

With the same slight abuse of notation as in Section 3.2, we define, for  $s > 0$ ,

$$f(s) = \int_0^s \int_{-\infty}^a \int_0^{2\pi} u_r(r, \theta, x_3) u_{x_3}(r, \theta, x_3) d\theta dx_3 dr.$$

Since  $|\nabla u| \leq 1$  in  $\Xi$ , one has  $|\nabla u|^2/2 + 1/2 \geq u_{x_3}^2$  in  $\Xi$ , hence the function  $f$  is nonnegative, non-decreasing and differentiable with respect to  $s$ . Similarly to the proof of Section 3.2, we can show that, if

$$\lim_{s \rightarrow +\infty} \int_{\{x_1^2 + x_2^2 < s^2\}} \left( \frac{1}{2} |\nabla u|^2 + F(u) - u_{x_3}^2 \right) dx_1 dx_2 > 0,$$

then there is a positive constant  $C > 0$  such that

$$f'(s) \geq \frac{C f^2(s)}{s \ln s}, \text{ for } s \text{ large.}$$

The previous inequality yields a contradiction as in Section 3.2. This then implies that  $|\nabla u(x_1, x_2, a)| = 1$  and  $u_{x_1}(x_1, x_2, a) = u_{x_2}(x_1, x_2, a) = 0$  for all  $(x_1, x_2) \in \mathbb{R}^2$  such that  $(x_1, x_2, a) \in \Xi$ . Since  $\Delta(|\nabla u|^2) = 2 \sum_{1 \leq i, j \leq 3} u_{x_i x_j}^2 \geq 0$  in  $\Xi$ , we conclude that  $|\nabla u| = 1$  and  $u_{x_1} = u_{x_2} = 0$  in each connected component of  $\Xi$  meeting  $\{x_3 = a\}$ . It finally follows that  $\Xi \supset \{x_3 = a_k\}$ , for a sequence  $(a_k)_{k \in \mathbb{N}}$  with  $a_k \rightarrow 0^+$  (from the normalization made in the second paragraph of the proof) and, remembering that  $u_{x_3} > 0$  in  $\bar{\Xi} \cap \{x_3 \leq a\}$ , we easily conclude that  $\Xi = \{x_3 > 0\}$  and  $u$  is the one-dimensional function  $u(x_1, x_2, x_3) \equiv x_3$ .  $\square$

Similarly, we can consider the following double-well type free boundary problem:

$$(28) \quad \begin{cases} \Delta u = 0 & \text{in } \Xi := \{|u| < 1\} \subset \mathbb{R}^n, \\ |\nabla u| = 1 & \text{on } \partial \Xi. \end{cases}$$

The proof of the following result is essentially same as that of Theorem 15, and we omit the details.

**Theorem 16.** *Let  $n \leq 3$  and  $u$  be a solution of (28) with  $|\nabla u| \leq 1$ . Suppose that  $\{|u| < 1\}$  is contained in the half-space  $\{x_n > 0\}$ . Then  $u$  is one-dimensional, namely there is  $h \geq 1$  such that  $\Xi = \{h - 1 < x_n < h + 1\}$ , and either  $u(x) \equiv x_n - h$  in  $\bar{\Xi}$  or  $u(x) \equiv -(x_n - h)$  in  $\bar{\Xi}$ .*

*Acknowledgements.* F. Hamel is partially supported by: the Excellence Initiative of Aix-Marseille University - A\*MIDEX, a French ‘‘Investissements d’Avenir’’ programme, the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) ERC Grant Agreement n. 321186 - ReaDi - Reaction-Diffusion Equations, Propagation and Modelling, and the ANR NONLOCAL project (ANR-14-CE25-0013). Y. Liu is partially supported by ‘‘The Fundamental Research Funds for the Central Universities WK3470000014’’. P. Sicbaldi is partially supported by the grant ‘‘Ram3n y Cajal 2015’’ RYC-2015-18730 and the grant ‘‘Analisis geom3trico’’ MTM 2017-89677-P. K. Wang is supported by NSFC no. 11871381. J. Wei is partially supported by NSERC of Canada.

Part of the paper was finished while Y. Liu was visiting the University of British Columbia in 2019, and he appreciates the institution for its hospitality and financial support. Part of this work was also completed while F. Hamel and J. Wei were visiting the University of Granada in

2019 in occasion of the conference “Geometry and PDE in front of the Alhambra”, and they also appreciate the institution for the hospitality and the financial support. Finally, part of this work has been carried out in the framework of Archimède Labex of Aix-Marseille University.

#### REFERENCES

- [1] O. Agudelo, M. Del Pino, and J. Wei. Higher-dimensional catenoid, Liouville equation, and Allen-Cahn equation, *Int. Math. Res. Not.* **23** (2016), 7051–7102.
- [2] G. Alberti, L. Ambrosio, and X. Cabré. On a long-standing conjecture of E. De Giorgi: symmetry in 3D for general nonlinearities and a local minimality property, *Acta Appl. Math.* **65** (2001), 9–33.
- [3] L. Ambrosio and X. Cabré. Entire solutions of semilinear elliptic equations in  $\mathbb{R}^3$  and a conjecture of De Giorgi, *J. Amer. Math. Soc.* **13** (2000), 725–739.
- [4] H. Berestycki, L. A. Caffarelli, and L. Nirenberg. Further qualitative properties for elliptic equations in unbounded domains, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **5** (1997), 69–94.
- [5] H. Berestycki, L. A. Caffarelli, and L. Nirenberg. Monotonicity for elliptic equations in unbounded Lipschitz domains, *Comm. Pure Appl. Math.* **50** (1997), 1089–1111.
- [6] H. Berestycki and F. Hamel. Generalized travelling waves for reaction-diffusion equations, In: *Perspectives in Nonlinear Partial Differential Equations. In honor of H. Brezis*, Amer. Math. Soc., Contemp. Math., 2007, 101-123.
- [7] X. Cabré. Uniqueness and stability of saddle-shaped solutions to the Allen-Cahn equation, *J. Math. Pures Appl.* **98** (2012), 239–256.
- [8] L. A. Caffarelli, N. Garofalo, and F. Segala. A gradient bound for entire solutions of quasi-linear equations and its consequences, *Comm. Pure Appl. Math.* **47** (1994), 1457–1473.
- [9] T. H. Colding and W. P. Minicozzi II. The Calabi-Yau conjectures for embedded surfaces, *Ann. Math. (2)* **167** (2008), 211–243.
- [10] E. De Giorgi. Convergence problems for functionals and operators, *Proc. Int. Meeting on Recent Methods in Nonlinear Analysis (Rome, 1978)*, Pitagora, Bologna (1979).
- [11] D. De Silva and O. Savin. Symmetry of global solutions to a class of fully nonlinear elliptic equations in 2D, *Indiana Univ. Math. J.* **58** (2009), 301–315.
- [12] M. Del Pino, M. Kowalczyk, and F. Pacard. Moduli space theory for the Allen-Cahn equation in the plane, *Trans. Amer. Math. Soc.* **365** (2013), 721–766.
- [13] M. Del Pino, M. Kowalczyk, and J. Wei. On De Giorgi’s conjecture in dimension  $N \geq 9$ , *Ann. Math. (2)* **174** (2011), 1485–1569.
- [14] A. Farina. Symmetry for solutions of semilinear elliptic equations in  $\mathbb{R}^n$  and related conjectures, *Ricerche Mat.* **48** (1999), 129–154.
- [15] A. Farina, B. Sciunzi, and E. Valdinoci. Bernstein and De Giorgi type problems: new results via a geometric approach, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5)* **VII** (2008), 741–791
- [16] A. Farina and E. Valdinoci. The state of the art for a conjecture of De Giorgi and related problems, in: *Recent Progress on Reaction–Diffusion Systems and Viscosity Solutions*, World Sci. Publ., Hackensack, NJ, 2009, 74–96.
- [17] A. Farina and E. Valdinoci. Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems, *Arch. Ration. Mech. Anal.* **195** (2010), 1025–1058.
- [18] N. Ghoussoub and C. Gui. On a conjecture of De Giorgi and some related problems, *Math. Ann.* **311** (1998), 481–491.
- [19] B. Gidas, W.-M. Ni, and L. Nirenberg. Symmetry and related properties via the maximum principle, *Comm. Math. Phys.* **68** (1979), 209–243.
- [20] D. Hoffman and W. H. Meeks, The strong halfspace theorem for minimal surfaces, *Invent. Math.* **101** (1990), 373–377.
- [21] Y. Liu, K. Wang, and J. Wei. Global minimizers of Allen-Cahn equation in dimensions  $n \geq 8$ , *J. Math. Pures Appl.* **108** (2017), 818–840.

- [22] Y. Liu, K. Wang, and J. Wei. On smooth solutions to one phase free boundary problem in  $\mathbb{R}^n$ , preprint.
- [23] W. H. Meeks, J. Perez, and A. Ros. Local removable singularity theorems for minimal laminations, *J. Diff. Geom.* 103 (2016), 319–392.
- [24] W. H. Meeks and H. Rosenberg. Maximum principles at infinity, *J. Diff. Geom.* 79 (2008), 141–165.
- [25] L. Modica. A gradient bound and a Liouville theorem for nonlinear Poisson equations, *Comm. Pure Appl. Math.* 38 (1985), 679–684.
- [26] O. Savin. Regularity of flat level sets in phase transitions. *Ann. Math. (2)* 169 (2009), 41–78.
- [27] F. Xavier. Convex hulls of complete minimal surfaces, *Math. Ann.* 269 (1984) 179–182.

F. HAMEL: AIX MARSEILLE UNIV, CNRS, CENTRALE MARSEILLE, I2M, MARSEILLE, FRANCE  
*E-mail address:* francois.hamel@univ-amu.fr

Y. LIU: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, CHINA  
*E-mail address:* yliumath@ustc.edu.cn

P. SICBALDI: UNIVERSIDAD DE GRANADA, DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, CAMPUS FUENTENUEVA, 18071 GRANADA, SPAIN & AIX MARSEILLE UNIV, CNRS, CENTRALE MARSEILLE, I2M, MARSEILLE, FRANCE  
*E-mail address:* pieralberto@ugr.es

K. WANG: SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, HUBEI, CHINA  
*E-mail address:* wangkelei@whu.edu.cn

J. WEI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA  
*E-mail address:* jcwei@math.ubc.ca