



HAL
open science

The Local Last Glacial Maximum of the southern Scandinavian Ice Sheet front: Cosmogenic nuclide dating of erratics in northern Poland

Karol Tylmann, Vincent Rinterknecht, Piotr Woźniak, Didier Bourles, Irene Schimmelpfennig, Valery Guillou, Georges Aumaitre, Karim Keddadouche

► To cite this version:

Karol Tylmann, Vincent Rinterknecht, Piotr Woźniak, Didier Bourles, Irene Schimmelpfennig, et al.. The Local Last Glacial Maximum of the southern Scandinavian Ice Sheet front: Cosmogenic nuclide dating of erratics in northern Poland. *Quaternary Science Reviews*, 2019, 219, pp.36-46. 10.1016/j.quascirev.2019.07.004 . hal-02188748

HAL Id: hal-02188748

<https://hal.science/hal-02188748v1>

Submitted on 13 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The Local Last Glacial Maximum of the southern Scandinavian Ice Sheet front:**
2 **cosmogenic nuclide dating of erratics in northern Poland**

3 Karol Tylmann^{1*}, Vincent R. Rinterknecht², Piotr P. Woźniak¹, Didier Bourlès³, Irene
4 Schimmelpfennig³, Valery Guilou³, ASTER Team^{3a}

5 ¹ University of Gdańsk, Faculty of Oceanography and Geography, Poland
6 (k.tylmann@ug.edu.pl)

7 ² Université Paris 1 Panthéon-Sorbonne, Laboratoire de Géographie Physique, CNRS, UMR
8 8591, France,

9 ³ Aix Marseille Université, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence,
10 France

11 * Corresponding author

12 ^a Georges Aumaître, Karim Keddadouche

13 **Abstract**

14 This paper presents new results of terrestrial cosmogenic nuclide dating of erratics in
15 northern Poland. We report the first exposure ages of erratics located on the pre-Local Last
16 Glacial Maximum and the Local Last Glacial Maximum moraines in Poland. Published
17 radiocarbon ages are calibrated and used as a background indicator of the possible time
18 window for the Local Last Glacial Maximum.

19 The terrestrial cosmogenic nuclide exposure ages presented in this study indicate that:
20 (1) exposure ages of erratics located on the pre-Local Last Glacial Maximum moraines (15.9
21 ± 1.4 to 101.7 ± 8.5 ka) are not clustered around any specific time interval, (2) the age of the
22 ice-sheet retreat from the Local Last Glacial Maximum ice limit in western Poland is $20.7 \pm$
23 0.8 ka and the probable duration of the Local Last Glacial Maximum in western and central
24 Poland is between ~ 25 ka and ~ 21 ka, (3) the age of the ice sheet retreat from the Local Last

25 Glacial Maximum ice limit in eastern Poland is 17.3 ± 0.5 ka and the probable duration of the
26 Local Last Glacial Maximum in eastern Poland is between ~ 22 ka and ~ 18 ka. Our results
27 show that the Local Last Glacial Maximum ice limit in western and central Poland is probably
28 ~ 3 ka older than in eastern Poland. This support a scenario for a complex response of the last
29 Scandinavian Ice Sheet southern margin to the climatic fluctuations and/or to internal
30 dynamics of the ice sheet.

31 **Key words:** Local Last Glacial Maximum, Leszno (Brandenburg) Phase, Poznań (Frankfurt
32 Phase), cosmogenic nuclide dating, Scandinavian Ice Sheet, erratics.

33 1. Introduction

34 The Last Glacial Maximum (LGM) is defined as the most recent period with the
35 maximum global integrated volume of ice sheets and the corresponding minimum global sea
36 level (Mix et al., 2001; Clark et al., 2009; Hughes et al., 2013) and dated to 23–19 cal ka BP
37 or 24–18 cal ka BP (cf. Mix et al., 2001; Hughes and Gibbard, 2015). The termination of the
38 LGM climatic cooling resulted in the progressive shrinking of the inland ice, which for most
39 of the Northern Hemisphere ice sheets is dated at ~ 20 – 17 ka (Hughes and Gibbard, 2015).
40 However, numerical ages (based on calibrated ^{14}C , terrestrial cosmogenic nuclides – TCN or
41 luminescence dating) of terrestrial deposits show significant regional variability in the timing
42 of the ice sheets' maximum extent and the commencement of their retreat. Most of the ice
43 sheets reached their maximum extents in a 7.0 ka time interval between 26.5 ka and 19.0 ka
44 (Clark et al., 2009), but the maximum extents of specific ice sheets (Local Last Glacial
45 Maxima – LLGM) were found to be asynchronous and not always in agreement with the
46 global LGM (Hughes et al., 2013).

47 The Scandinavian Ice Sheet (SIS) was one of the main Northern Hemisphere ice sheet
48 and covered a significant part of northern Europe during the LGM. Timing of the last SIS


49 maximum expansion and retreat is important to understand climatic fluctuations at the end of
50 the Marine Isotope Stage (MIS) 2. The LLGM to the south of the Baltic Basin (Fig. 1A) is
51 dated at ~24 ka in Germany and western Poland, and at ~19 ka in eastern Poland based on
52 calibrated ^{14}C and OSL ages (Marks, 2012 after Stankowska and Stankowski, 1988; Wysota
53 et al., 2009) and at 21–19 ka in Belarus and Lithuania based on calibrated ^{14}C and ^{10}Be ages
54 (Rinterknecht et al., 2007, 2008). Thus, the maximum extent of the of the last SIS in Poland
55 was probably asynchronous with earlier maximum expansion of the ice sheet in western and
56 central Poland and subsequent maximum expansion further to the east (Fig. 1A and B).
57 Nevertheless, there is still a relatively small number of geochronological data constraining the
58 timing of LLGM in north Poland. Specifically, there is a lack of direct dating (TCN) of
59 moraines correlated with the maximum limit of the last ice sheet in this region.

60 Here we present a new set of TCN exposure ages of erratic boulders located within the
61 southern front of the last SIS in northern Poland. It provides the first exposure ages of erratics
62 located on the pre-LLGM moraines and new exposure ages on the LLGM moraines in Poland.
63 Additionally, published ^{14}C ages were calibrated with the most recent radiocarbon calibration
64 curve and used as a background indicator of the possible time window for the LLGM. We
65 discuss our results in a wider chronological context including specifically the growing number
66 of luminescence ages available for the southern fringe of the last SIS (e.g. Houmark-Nielsen
67 and Kjaer, 2003; Wysota et al., 2009; Lüthgens et al., 2011; Bitinas, 2012; Lasberg and Kalm,
68 2013), as well as existing reconstructions of the SIS evolution (e.g. Hughes et al., 2016;
69 Stroeven et al., 2016). The objective of this paper is to date the asynchronous maximum
70 extent of the last SIS in northern Poland based on new exposure ages, and on the extensive
71 dataset of TCN exposure ages already published.

72 2. Study area

73 The study area is located in NW Poland and covers ~20% (~500 km) of the LLGM ice
74 sheet margin south of the Baltic Sea. The investigated region is situated within the eastern
75 part of the extensive, lobe-shaped fringe of the last SIS (Fig. 1A). The region is large enough
76 to find a considerable number of erratics suitable for TCN exposure dating (Tylmann et al.,
77 2018) and to track the last SIS retreat from its maximum limit.

78 The ice sheet limit of the LLGM in northern Poland is mostly based on the spatial
79 distribution of end moraines, outlets of tunnel valleys or proximal parts of outwash plains and
80 fans (Majdanowski, 1947; Galon and Roszkówna, 1961; Marks, 2002). This limit defines the
81 “young glacial” landscape located to the north and the “old glacial” landscape located to the
82 south of it (Fig. 1B). According to recent reconstructions, the LLGM was diachronous and it
83 occurred during the Leszno/Brandenburg Phase in the western part and during the
84 Poznań/Frankfurt Phase in the eastern part of the study area (Marks, 2011, 2012, 2015; Fig.
85 1B). It is supported by lithostratigraphic evidences which suggest asynchronous maxima of
86 the last SIS across the study area. In north-central Poland two till layers separated by ice-
87 wedge casts, are correlated with the glacial phases mentioned above (Wysota et al., 2009;
88 Narloch et al., 2013). However, in the former Vistula Ice Lobe area (Fig. 1B), only one till
89 layer has been identified, and OSL dating of fluvial and glaciofluvial sediments suggests that
90 it is correlated with the Poznań Phase (cf. Wysota et al., 2009; Roman, 2017). In contrast, in
91 the western part of the study area, the till layer correlated with the Leszno Phase occurs close
92 to the maximum extent of the last SIS, and no basal till layer correlated with the Poznań Phase
93 has been documented there. Therefore, the ice marginal belt formed during the Poznań Phase
94 in western Poland is commonly interpreted as a result of the ice margin standstill, which
95 occurred during the last SIS recession after the LLGM of the Leszno Phase (e.g. Kasprzak,
96 1988; Kozarski, 1988, 1995).



97

98 Fig. 1. The main ice margin limits of the last Scandinavian Ice Sheet (SIS) in the area south of
 99 the Baltic Sea and the study area with erratic boulders. (A) Location of the study area within
 100 the southern front of the last SIS. Dark dots indicate location of radiocarbon ages used in the
 101 article. (B) The location of erratic boulders dated with terrestrial cosmogenic nuclides (TCN).
 102 Limits of the last SIS from Marks et al. (2006) and Kozarski (1995).

103 Up until recently, the age of the maximum extent of the last SIS in the study area was
104 interpreted mainly based on available radiocarbon ages (Table 1 and references therein).
105 Published ^{14}C data significant for the interpretation of the last SIS advance and retreat
106 (locations in Fig. 1A) indicate the possible time window for its maximum extent and the
107 beginning of its retreat. Radiocarbon ages from organic deposits resting below the LLGM till
108 layers have been obtained in the vicinity of the last SIS maximum limit in central Poland
109 (Stankowska and Stankowski, 1988; Gogołek and Mańkowska, 1989) as well as in the area of
110 the northern Polish coastline (Rotnicki and Borówka, 1995) and in offshore boreholes drilled
111 into the Baltic Sea bottom (Kramarska, 1998; Krzyszkowski et al., 1999). The youngest
112 calibrated radiocarbon ages of these deposits constrain the maximum reliable age of the
113 LLGM. The most reliable ages for the ice sheet retreat, may be estimated by the oldest
114 calibrated radiocarbon ages of the organic sediments deposited after deglaciation at the
115 bottom of lake basins close to the last SIS maximum extent in central and northern Poland
116 (Niewiarowski, 1995; Stankowski et al., 1999; Gamrat et al., 2017). Because of the risk of
117 organic matter redeposition at the bottom of lake basins and the associated overestimated
118 radiocarbon ages, these ages should be interpreted as a maximum reliable time interval for the
119 deglaciation of north-central Poland after the LLGM.

120 The calibrated radiocarbon ages of organic deposits underlying the LLGM till layers
121 range between 26.5 ± 1.3 and 25.2 ± 3.4 cal ka BP. The oldest minimum age predating the
122 LLGM was found on the northern Polish coastline (Rotnicki and Borówka, 1995) and the
123 youngest one in central Poland (Gogołek and Mańkowska, 1989). The post-LLGM maximum
124 ages of organic sediments deposited after deglaciation range between 21.4 ± 1.9 cal ka BP and
125 19.1 ± 0.2 cal ka BP. The oldest age (21.4 ± 1.9 cal ka BP) from bottom part of the lake
126 deposits resting above the LLGM till in central Poland (Stankowski et al., 1999). A similar
127 age (21.4 ± 0.6 cal ka BP) was obtained from the bottom of the lacustrine deposits overlying

128 the till of the same stratigraphic position, ~60 km further to the north (Niewiarowski, 1995).
129 The youngest maximum post-LLGM age (19.1 ± 0.2 cal ka BP), was found in lacustrine silt
130 filling glacial channel located further to the east (Gamrat et. al., 2017).

131 The distribution of pre-LLGM and post-LLGM calibrated ^{14}C ages in the study area
132 suggests that:


- 133 1) the LLGM ice advance occurred after ~25 cal ka BP,
- 134 2) the beginning of ice sheet retreat after the LLGM in central-western Poland
135 occurred not earlier than ~21.5 cal ka BP and the deglaciation of the regions located
136 further to the east started not earlier than ~19.0 cal ka BP.

137 **3. Methods**

138 *3.1. Sample selection*

139 The largest erratic boulders located in situ on glacial landforms in the vicinity of the
140 maximum extent of the last SIS and to the north of this limit were our targets for sampling for
141 surface exposure dating with TCN (Fig. 1B). The boulders were carefully selected from an
142 extensive dataset of almost 500 erratics, based on their dimensions and geomorphological
143 locations (Tylmann et al., 2018). We selected erratics located on depositional glacial
144 landforms, i.e. terminal moraines, dead-ice moraines, moraine plateaux or proximal zones of
145 outwash fans/plains. Samples were collected with a manual jackhammer from the upper
146 surface of 33 stable and massive boulders of perimeter ranging from 6.0 to 14.3 m and of
147 height ranging from 0.8 to 2.9 m (Table 2; see examples in Fig. 2). Almost all boulders ($n =$
148 32) are characterized by quartz-rich lithologies – granitoid, granite gneiss and gneiss. Because
149 of the availability of quartz, these erratics were sampled for surface exposure dating with in
150 situ-produced ^{10}Be . Out of 33 samples for ^{10}Be analysis (Table 2), one boulder was sampled
151 twice; we took a sample from the top (LGM-02T) and one from the side (LGM-02S). The

152 motivation of this sampling was that from the field observations the boulder looked to have
153 been rotated by 90°, with the original top surface on the side (Fig. 2C). One boulder does not
154 contain quartz (gabbro) and it was sampled for exposure dating with in situ-produced ^{36}Cl
155 (Fig.2D, Tables 3 –5). In total, 34 new samples were analyzed: 33 for ^{10}Be analysis and one
156 for ^{36}Cl analysis.



157
158 Fig. 2. Examples of erratic boulders sampled for TCN exposure dating. (A) Sampling of FRT-
159 01 boulder upper surface with a manual jackhammer. (B) FRT-01 sampling spot with the scar
160 (outlined by a white dashed line) and the collected material. (C) LGM-02 boulder, probably
161 rotated 90° clockwise. Arrows show location of samples LGM-02T (present top of the
162 boulder) and LGM-02S (hypothesized to be the original top). (D) FRT-11 boulder (gabbro);
163 the only one sampled for ^{36}Cl .

164 3.2. ¹⁰Be dating methodology

165 The samples were crushed and sieved. The 0.25–1.0 mm fraction was divided with a
166 Frantz separator into magnetic and non-magnetic subsamples. The non-magnetic fractions
167 were enriched in quartz decontaminated by successive acid leaching (HCl + H₂SiF₆ then
168 dilute HF). The purified quartz was spiked with ~100–300 mg of a 3025 ppm home-made
169 carrier then dissolved in 48% HF. After evaporation and purification with Dowex 1 X 8 and
170 Dowex 50W * 8 (100-200 mesh) ion exchange resins, BeOH was dried and then oxidized to
171 BeO at 800°C. BeO was mixed with 325 mesh Nb-powder prior to measurement at the
172 accelerator mass spectrometer (AMS). Sample preparation was conducted at the Laboratoire
173 National des Nucléides Cosmogéniques (LN₂C) at CEREGE, Aix-en-Provence, France. The
174 French AMS facility ASTER was used for the ¹⁰Be measurements (Arnold et al., 2010). The
175 measured ¹⁰Be/⁹Be ratios were normalized relative to the in-house standard STD-11 using an
176 assigned ¹⁰Be/⁹Be ratio of $(1.191 \pm 0.013) \times 10^{-11}$ (Braucher et al., 2015) and a ¹⁰Be half-life
177 of $(1.387 \pm 0.012) \times 10^6$ years (Chmeleff et al., 2010; Korschinek et al., 2010).

178 ¹⁰Be ages were calculated using the most recent global production rate (Borchers et al.,
179 2016) and the ‘Lm’ time-dependent scaling scheme for spallation according to Lal (1991) and
180 Stone (2000). The modern global ¹⁰Be production rate at sea level and high-latitude is derived
181 from a recalibrated global data set and recent calibration studies (Borchers et al., 2016). This
182 production rate was used as the most adequate, because the local production rate for Poland
183 was not estimated so far. A recently published Swedish ¹⁰Be production rate (Stroeven et al.,
184 2015) would generate slightly younger exposure ages (~5%). The ¹⁰Be production rate was
185 corrected for sample thickness according to an exponential function (Lal, 1991) and assuming
186 an average density of 2.7 g/cm³. A correction for self-shielding (direction and angle of surface
187 dipping) was applied when the surface of a sampling spot was sloping >10°. However, for
188 most of the sampled boulders no correction for surface inclination was necessary. We apply a

189 correction for surface erosion, assuming a maximum erosion rate of 1.3 mm ka^{-1} for granitic
190 surfaces (Gosse et al., 1995; Rinterknecht et al., 2012, 2014). Corrections for snow/vegetation
191 cover were not included due to their minimal impact on ^{10}Be ages in the temperate vegetation
192 zone (e.g. Plug et al., 2007; Rinterknecht et al., 2014). All calculations were performed using
193 the online exposure age calculator formerly known as the CRONUS-Earth online exposure
194 age calculator – version 3 (<http://hess.ess.washington.edu/math/>; accessed: 2019-02-01),
195 which is an updated version of the online calculator described by Balco et al. (2008). ^{10}Be
196 exposure ages corrected and uncorrected for erosion are reported with 1σ uncertainties
197 (including analytical uncertainties and the production rate uncertainty) in ka in Table 2. In the
198 following sections we use ages uncorrected for the erosion and we interpret them as minimum
199 ages for the ice sheet retreat.

200 3.3. ^{36}Cl dating methodology

201 Gabbro sample FRT-11 was crushed and sieved to a grain size fraction of 250–710 μm
202 at CALM (Cosmonucléides Au Laboratoire de Meudon) at the Laboratoire de Géographie
203 Physique (LGP), France, and chemically prepared for in situ ^{36}Cl dating at the LN_2C . The
204 procedure is described in Schimmelpfennig et al. (2011). Sample specific data and
205 compositional results are given in Tables 3, 4 and 5. The ^{36}Cl age of sample FRT-11 was
206 calculated with the Excel® spreadsheet of Schimmelpfennig et al. (2009), using the scaling
207 method of Stone (2000) and employing the following ^{36}Cl production rates, referenced to sea
208 level and high latitude (SLHL): $42.2 \pm 4.8 \text{ atoms } ^{36}\text{Cl} (\text{g Ca})^{-1} \text{ yr}^{-1}$ for spallation of Ca
209 (Schimmelpfennig et al., 2011), $148.1 \pm 7.8 \text{ atoms } ^{36}\text{Cl} (\text{g K})^{-1} \text{ yr}^{-1}$ for spallation of K
210 (Schimmelpfennig et al., 2014), $13 \pm 3 \text{ atoms } ^{36}\text{Cl} (\text{g Ti})^{-1} \text{ yr}^{-1}$ for spallation of Ti (Fink et al.,
211 2000), $1.9 \pm 0.2 \text{ atoms } ^{36}\text{Cl} (\text{g Fe})^{-1} \text{ yr}^{-1}$ for spallation of Fe (Stone et al., 2005), and 696 ± 185
212 neutrons $(\text{g air})^{-1} \text{ yr}^{-1}$ for the production rate of epithermal neutrons from fast neutrons in the
213 atmosphere at the land/atmosphere interface (Marrero et al., 2016b). We used a high-energy

214 neutron attenuation length of 160 g cm^{-2} . The resulting ^{36}Cl age in Table 5 with its 1σ
215 uncertainty including all analytical and production rate errors is shown with and without
216 correction for an erosion rate of 1.3 mm ka^{-1} . The ^{36}Cl age was also calculated using the
217 CRONUScale calculator (Marrero et al., 2016a) for comparison with the results calculated
218 with the Excel® spreadsheet (Schimmelpfennig et al., 2009).


219 **4. Results**

220 *4.1. Erratics located beyond the ice marginal belt of the LLGM*

221 Exposure ages for SAAL samples show significant scatter (16.2 ± 1.4 to 115.2 ± 11.0
222 ka; Table 2). The observed variability (53.2%) is much larger than the average external
223 uncertainty (8.4%). This shows that random uncertainties are dominated by geological
224 uncertainties rather than by analytical ones. Most of the pre-LLGM exposure ages are
225 younger than 100 ka (six samples) with five ages younger than 60 ka (Table 2).

226 *4.2. Erratics of the LLGM ice marginal belt*

227 We sampled 13 erratics located on the ice marginal belt identified geomorphologically
228 as the Leszno (Brandenburg) Phase (Fig. 1B). The exposure ages for samples LGM (01 to 08)
229 and FRT (01 to 04 and 06) range between 19.3 ± 2.1 and 75.3 ± 6.3 ka (Table 2). The
230 distribution of the ages is bimodal with the narrow and high mode containing ages ranging
231 between 19.3 ± 2.1 and 25.1 ± 2.1 ka, and wider and lower mode containing ages between
232 39.5 ± 3.3 and 75.3 ± 6.3 ka (Fig. 3A). The results for the boulder that has been sampled twice
233 (LGM-02) belong to the older distribution, and are: 39.5 ± 3.3 ka for the top sample (LGM-
234 02T) and 58.6 ± 3.9 ka for the side sample (LGM-02S). The older age of the weathered side
235 of the boulder support our hypothesis that this erratic was rotated and the original top surface
236 is on the side (Fig. 2C).



237

238 Fig. 3. Distribution of surface exposure ages for erratic boulders located on the LLGM ice

239 marginal belts. (A) Erratic ^{10}Be ages located on the ice marginal belt of the Leszno Phase. (B)

240 Surface exposure ages (22 ^{10}Be and one ^{36}Cl) of the Poznań Phase ice marginal belt. Yellow

241 areas represent cumulative kernel density estimates of all ^{10}Be ages; blue, red and grey curves

242 represent kernel density estimates for individual ^{10}Be ages: blue – included in the error-

243 weighted mean age calculation, red – outliers, grey – not accepted ages (older mode of ages


244 from ice marginal belt correlated with the Leszno Phase).

245 Ages belonging to the younger distribution are much more clustered around the mode
246 than the ages of the older distribution (Fig. 3A). The reduced chi-squared test for ages of the
247 younger distribution indicates that they represent a single population ($\chi^2_r = 0.8$ and $K = 2.2$),
248 while the ages of the older distribution do not represent a single population ($\chi^2_r = 7.2$ and $K =$
249 2.2). We used the criterion 'K' to test whether the reduced chi-squared value fall within a 2σ
250 envelope (95% confidence). This criterion depends on the degree of freedom, and thus the
251 number of samples (n): $K = 1 + 2\sqrt{2/(n - 1)}$. If $\chi^2_r < K$ then there is a >95% probability that
252 the analyzed measurements represent a single population. We therefore used the younger
253 distribution to interpret the age of the Leszno Phase ice marginal belt. The oldest age of this
254 dataset (25.1 ± 2.1 ka) was identified as an outlier according to the Chauvenet's criterion. A
255 Shapiro-Wilk test, applied after rejecting the outlier, showed that we cannot reject the
256 normality assumption for $p = 0.05$ ($W = 0.97$) for distribution of the remaining six ages. The
257 error-weighted mean and the standard deviation of the mean calculated for these ages is 20.7
258 ± 0.8 ka (Table 2). The spatial distribution of the analyzed erratics indicates that statistically
259 reliable ages (included in the mean age calculation) are located mainly in the western part of
260 the study area (Fig. 4).

261 Twelve boulders located on the Poznań (Frankfurt) ice marginal belt were sampled for
262 ^{10}Be (samples LGM-09 to 12, FRT-05, FRT-07 to 10 and FRT-12 to 14) and one boulder for
263 ^{36}Cl exposure dating (sample FRT-11). We also recalculated and used ^{10}Be ages of ten
264 boulders from published data (Rinterknecht et al., 2005, 2006), located close to the ice sheet
265 limit during the LLGM in the eastern edge of the study area (Table 2; Fig. 1B) using the same
266 procedure as for our samples.

267 ^{10}Be ages range between 5.8 ± 0.8 and 40.6 ± 3.9 ka (Table 2) with narrow modes
268 occurring at $\sim 5\text{--}7$ ka and $\sim 15\text{--}19$ ka, and wide, low mode at $\sim 30\text{--}40$ ka (Fig. 3B). The
269 youngest mode includes two ages 5.8 ± 0.8 and 7.0 ± 1.9 ka, the intermediate mode consists

270 of 14 ages ranging between 15.5 ± 1.6 and 20.2 ± 2.2 ka, and the oldest mode includes four
 271 ages between 30.2 ± 3.2 and 40.6 ± 3.9 ka. The reduced chi-squared test indicates that the 22
 272 ^{10}Be ages represent multiple populations ($\chi^2_r = 18.0$ and $K = 1.6$). The application of
 273 Chauvenet's criterion identifies that the three youngest ages (<13.0 ka) and the five oldest
 274 ages (>24.1 ka) are outliers (Fig. 3B). After rejecting the outliers, the reduced chi-squared test
 275 applied to the remaining 14 ages indicates that they belong to a single population ($\chi^2_r = 0.5$
 276 and $K = 1.8$). Moreover, a Shapiro-Wilk test showed that we cannot reject the normality
 277 assumption for $p = 0.05$ ($W = 0.90$) for the distribution of the 14 ages. The error-weighted
 278 mean and the standard deviation of the error-weighted mean calculated for the ^{10}Be ages is
 279 17.3 ± 0.5 ka (Table 2). The ^{36}Cl age is 20.1 ± 3.1 ka when calculated with the Excel®
 280 spreadsheet (Schimmelpfennig et al., 2009), and 20.0 ± 3.0 ka when calculated with the
 281 CRONUScalc calculator (Marrero et al., 2016b). These two results are very similar and
 282 belong to the same frequency distribution as the 14 ^{10}Be ages (Fig. 3B).



284 Fig. 4. Spatial distribution of ^{10}Be ages and single ^{36}Cl age in the study area. Ages used in the
285 calculation of the error-weighted mean ages for specific ice marginal belts are framed in black
286 frame.

287 5. Discussion

288 5.1. Pre-LLGM erratics

289 The distribution of the exposure ages of erratics located beyond of the ice sheet limit
290 during the LLGM is very large (from 15.9 ± 1.4 to 101.7 ± 8.5 ka; Table 2) and these ages are
291 not clustered around any particular time interval. We have only one exposure age (sample
292 SAAL-05) that may be roughly interpreted as representing the timing of retreat of the MIS 6
293 ice sheet: 101.7 ± 8.5 ka (Table 2). Six out of seven boulders display much too young ages
294 (from 15.9 ± 1.4 to 66.8 ± 5.5 ka) for the retreat of the MIS 6 ice sheet, and probably result
295 from degradation of moraines and erosion of erratics (e.g. Hallet and Putkonen, 1994) during
296 the long time period after deglaciation. Postglacial weathering and erosion of an erratic
297 surface, removes cosmogenic nuclide accumulated in the surface layer of the sample and thus
298 produces too young ages, but we consider the obtained exposure ages most probably represent
299 long-term geomorphic processes acting after deglaciation, and therefore largely underestimate
300 the timing of ice sheet retreat. Similar inconsistencies of ^{10}Be ages for erratics located within
301 Saalian (MIS 6) landscapes were reported in Germany, in the Hoher Fläming area
302 (Rinterknecht et al., 2012), and in Mecklenburg-Vorpommern (Rinterknecht et al., 2014) in
303 front of the ice sheet limit during the Brandenburg Phase. For 14 erratics sampled for ^{10}Be
304 dating in Germany, ten boulders have exposure ages too young (from 27.7 ± 1.3 to 92.8 ± 3.0
305 ka) for the deglaciation at the end of MIS 6 and four erratics have exposure ages ranging from
306 101.7 ± 10.4 to 143 ± 5.0 ka.

307 Two erratics (samples SAAL-04: 22.3 ± 1.9 ka and SAAL-06: 15.9 ± 1.4 ka) have
308 exposure ages close to the expected LLGM age, and they are also located close to the ice
309 sheet limit during the LLGM (Figs. 1 and 4). This may suggest a revision of the last SIS
310 maximum extent in central Poland. However, these erratics are located in the vicinity of the
311 sites of Eemian organic deposits that are not covered by the LLGM till (Tobolski, 1986;
312 Kuszell, 1998; Szałamacha and Skompski, 1999; Balwierz and Roman, 2002; Fig. 1). In
313 addition to geomorphology, this is a strong argument supporting that the ice sheet limit during
314 the LLGM is correct and that these two boulders display underestimated exposure ages
315 resulting from geomorphic processes.

316 *5.2. Age of the LLGM and the first stages of the last SIS retreat*

317 In central-western and northern Poland, deposition of organic sediments resting below
318 the LLGM till is dated to ~ 26.5 – 25.2 cal ka BP (Table 1) indicating that the maximum extent
319 of the last SIS in this region occurred later than ~ 25 cal ka BP. Moreover, the radiocarbon age
320 of the basal part of gyttja (21.4 ± 1.9 cal ka BP) resting on the same till (Stankowski et al.,
321 1999) indicates that the ice sheet retreat from its maximum limit started not earlier than ~ 21.4
322 cal ka BP. Recently, Marks (2012) determined, based on his interpretation of the available
323 ages, the age of the maximum extent of the last SIS in central and western Poland during the
324 Leszno Phase at 24 cal ka BP.

325 Our results for the Leszno (Brandenburg) Phase in western Poland are clustered
326 around two modes at ~ 19 – 22 ka and ~ 40 – 70 ka (Fig. 3A). We used this younger mode to
327 calculate an error-weighted mean surface exposure age of 20.7 ± 0.8 ka (Fig. 5). We interpret
328 this age as a minimum age of the commencement of the ice sheet retreat from its maximum
329 limit at the end of the LLGM in central and western Poland. Ages used in the error-weighted
330 mean age calculation agree, within uncertainty, with the radiocarbon ages coming from
331 deposits resting below and above the till of the ice advance during the Leszno Phase in central

332 Poland (Stankowski et al., 1999). The ages of the older distribution (between 39.5 ± 3.3 and
333 75.3 ± 6.3 ka) probably result from inherited ^{10}Be . This could have happened either due to
334 redeposition of previously exposed boulders by the last SIS with no or limited erosion (cf.
335 Håkansson et al., 2008; Rinterknecht et al., 2012).


336 The new exposure age of the last ice sheet retreat in western Poland is also consistent
337 with most luminescence ages constraining the timing of the Leszno Phase (Fig. 5). OSL
338 dating of fluvial, fluvio-glacial and periglacial sediments underlying the till correlated with the
339 Leszno Phase (Stankowski et al., 1999; Wysota et al., 2009; Molewski and Weckwerth, 2009;
340 Weckwerth et al., 2011) yields ages that mostly range between ~ 29 and ~ 24 ka. Additionally,
341 OSL ages of sediments underlying the LLGM till in the northernmost region of Germany
342 suggest that the last ice sheet reached the southern Baltic coast at Rügen Island after $22.7 \pm$
343 1.9 ka (Pisarska-Jamroży et al., 2018) or 22.0 ± 2.0 ka and 23.0 ± 2.0 ka (Kenzler et al., 2015,
344 2017). OSL ages of fluvial and periglacial deposits overlying the till correlated with the
345 Leszno Phase in Poland range between ~ 21 and 19 ka (Molewski 2007; Wysota et al., 2009;
346 see Fig. 5). We thus suggest that the error-weighted mean exposure age of erratics in western
347 Poland (20.7 ± 0.8 ka) represents the most reliable age for the initiation of the ice sheet retreat
348 after the ice advance during the Leszno Phase.

349 Our results also corroborate TCN data published by Heine et al. (2009), who
350 determined the exposure ^{10}Be age of three boulders located on the Brandenburg (Leszno) and
351 Frankfurt (Poznań) Phase ice marginal belts in north-eastern Germany. Our error-weighted
352 mean exposure age (20.7 ± 0.8 ka) and the results of Heine et al. (2009) (22.0 ± 1.1 ka), are
353 coeval within uncertainties. We suggest that the duration of the LLGM in central and western
354 Poland, constrained with the available radiocarbon ages and our new ^{10}Be exposure ages, is
355 between ~ 25 ka and ~ 21 ka (Fig. 5).

356 Radiocarbon dating from western Belarus (Zimenkov, 1989) indicates an age of the
357 last SIS maximum extent of ~ 22.5 cal ka BP in the eastern part of its southern sector. The
358 deglaciation of this region, in the light of radiocarbon dating in north-eastern Poland (Gamrat
359 et al., 2017), started not earlier than 19.1 ± 0.2 cal ka BP. Our results for erratics located close
360 to the ice sheet limit during the LLGM in the eastern part of the study area also support this
361 view (Fig. 5). TCN exposure ages of boulders located on the ice marginal belt of the Poznań
362 Phase, after rejecting outliers with Chauvenet's criterion, range between 15.5 ± 1.6 and $20.2 \pm$
363 2.2 ka, with an average age of 17.3 ± 0.5 ka (Table 2). We interpret these ages as a minimum
364 age of:

- 365 1) the beginning of the ice sheet retreat during the LLGM in eastern Poland, after the
366 ice advance during the Poznań Phase,
- 367 2) the ice sheet retreat from the Poznań Phase limit in north-central Poland, postdating
368 the LLGM represented by the Leszno Phase.

369 These results are in general agreement with OSL ages (~ 18.7 – 16.7 ka; Fig. 5) of
370 fluvioglacial sand overlying the till correlated with the ice advance during the Poznań Phase
371 in north-central Poland (Stankowski et al., 1999; Wysota et al., 2009; Roman, 2017). ^{10}Be
372 exposure dating of erratics located on moraines correlated with the LLGM in Belarus and
373 Lithuania (Rinterknecht et al., 2007, 2008) yields mean exposure ages of deglaciation at 17.7
374 ± 0.8 and 20.1 ± 1.2 ka respectively. Furthermore, the recent results from Valday Heights in
375 NW Russia show a ^{10}Be age for the LLGM at 18.9 ± 0.5 ka (Rinterknecht et al., 2018). These
376 data suggest an older age for the LLGM further east of eastern Poland. We propose a possible
377 duration for the LLGM in eastern Poland, constrained by radiocarbon ages and our new ^{10}Be
378 exposure ages as well as recalculated data of Rinterknecht et al. (2005, 2006), between ~ 22 ka
379 and ~ 18 ka (Fig. 5).



380

381 Fig. 5. Distribution of TCN, ^{14}C , and OSL ages relevant for the reconstruction of the timing of the LLGM and the initiation of the last ice sheet

382 retreat in the study area. Sources of ^{14}C ages are listed in Table 1.

383 Our results do not agree with OSL data and reinterpretation of ^{10}Be exposure dating of
384 erratics proposed recently by Hardt et al. (2016). Based on OSL dating of fluvioglacial
385 deposits in the vicinity of the Frankfurt (Poznań) ice marginal belts, they assigned the age of
386 $\sim 26.3 \pm 3.7$ ka for the initiation of the ice sheet retreat in north-eastern Germany. This is
387 significantly older than our mean exposure age of 17.3 ± 0.5 ka, which we interpret as a
388 minimum age of the start of deglaciation during the Poznań Phase in northern Poland. The
389 inconsistency may result from the overestimation of the OSL ages obtained by Hardt et al.
390 (2016), due to incomplete bleaching of fluvioglacial sediments. Although these authors
391 applied a single grain method and minimum age model in OSL age calculation, the risk that
392 all quartz grains within the sample are not fully bleached persists, especially when
393 fluvioglacial (outwash) sediments deposited close to the ice margin are dated (e.g. Houmark-
394 Nielsen, 2008; Bickel et al., 2015). However, if the OSL ages were correct for the ice sheet
395 recession, it would suggest a high asynchrony of the ice margin retreat during the Poznań
396 (Frankfurt) Phase: the beginning of the ice sheet retreat in northern Poland occurred at least
397 ~ 5 ka later than in north-eastern Germany. Furthermore, Hardt et al. (2016) interpreted the
398 recalculated surface exposure ages of Heine et al. (2009) as an indicator of landform
399 stabilization after deglaciation rather than a reliable age of the ice-sheet retreat (Houmark-
400 Nielsen et al., 2012). On the other hand, our exposure ages of erratics in the study area
401 correlate strongly with independent geochronological data from Poland, such as OSL dating,
402 and we argue that they represent a reliable age for the initiation of the ice sheet retreat in the
403 study area. They also correlate with the OSL ages cited above and recently obtained on Rügen
404 Island (Kenzler et al., 2015, 2017; Pisarska-Jamroży et al., 2018).

405 *5.3. Asynchrony of LLGM*

406 Although already postulated by other authors (e.g. Marks 2012; Larsen et al., 2016;
407 Stroeven et al., 2016; Patton et al., 2017) the asynchrony of the LLGM ice sheet limit in the

408 southern sector of the SIS was strongly confirmed by our results. TCN dating of the ice front
409 retreat along the last SIS ice margin in northern Poland show clearly a ~3 ka delay of the
410 LLGM in the eastern Poland with respect to the LLGM in western and central Poland. This
411 generally agrees with the recent reconstructions of the diachronous LLGM dated at ~24 ka
412 (Leszno/Brandenburg Phase) in the western part and at ~19 ka (Poznań/Frankfurt Phase) in
413 the eastern part of the study area (Wysota et al., 2009; Marks, 2011, 2012, 2015). This
414 asynchrony of the LLGM may be related to the activation and deactivation of the Vistula and
415 Odra ice streams (Wysota et al., 2009; Woźniak and Czubla, 2015; Spagnolo et al., 2016;
416 Roman, 2017). These ice streams most likely shaped the ice margin (Fig. 5) and to a large
417 extent controlled the timing of the last SIS retreat.

418 Timing of the ice margin retreat within the last SIS have been recently reconstructed
419 based on GIS compilation of available geochronological data and construction of the
420 deglaciation isochrones (Hughes et al., 2016, Stroeve et al., 2016). The reconstruction for the
421 Eurasian ice sheets suggests the maximum ice sheet extent within the southern sector of the
422 last SIS 24–22 ka, with the LLGM in western Poland most likely 24–23 ka and the LLGM in
423 eastern Poland most likely 22 ka (Hughes et al., 2016). Our results of TCN dating indicate
424 larger time offset between the maximum ice sheet expansion in western and eastern Poland.
425 They correspond more closely to the reconstruction of Stroeve et al. (2016), who interpreted
426 the timing of the LLGM to be 22 ka in western and 19 ka in eastern Poland.

427 **6. Conclusions**

- 428 1. A wide distribution of erratic exposure ages located on the pre-LLGM moraine
429 plateaux (15.9 ± 1.4 to 101.7 ± 8.5 ka) is not clustered around any particular time
430 interval. The ^{10}Be exposure ages represent long-term postglacial geomorphic
431 processes.

- 432 2. ^{10}Be exposure dating of erratics resting on the ice marginal belt of the Leszno
433 (Brandenburg) Phase in western Poland show the initiation of the ice sheet retreat 20.7
434 ± 0.8 ka.
- 435 3. The duration of the LLGM in western and central Poland, constrained by our ^{10}Be
436 exposure dating and the available radiocarbon dating is between ~ 25 ka and ~ 21 ka.
- 437 4. ^{10}Be exposure dating of erratics resting on the ice marginal belt of the Poznań Phase
438 show the initiation of the last SIS retreat 17.3 ± 0.5 ka.
- 439 5. The duration of the LLGM in eastern Poland is best constrained by our new ^{10}Be
440 exposure ages, recalculated exposure ages of Rinterknecht et al. (2005, 2006) and
441 calibrated radiocarbon ages is between ~ 22 ka and ~ 18 ka.
- 442 6. Our results confirm that different sectors of the last SIS southern front advanced and
443 retreated asynchronously. The LLGM ice limit in western and central Poland is
444 probably ~ 3 ka older than in eastern Poland. The documented asynchrony supports a
445 complex response of the last SIS southern margin to hemispheric climatic fluctuations
446 and internal dynamics of the ice sheet controlled largely by activation and deactivation
447 of ice streams.

448 **Acknowledgements**

449 We are very grateful to the Regional Directorates of Environmental Protection local
450 communes offices for permissions for sampling large erratics protected by law. We appreciate
451 suggestions of two anonymous reviews and of the editor which help to improve the
452 manuscript.

453 **Research is funded by:** National Science Centre grant no. 2014/15/D/ST10/04113 to Karol
454 Tylmann. The ASTER AMS national facility (CEREGE, Aix en Provence) is supported by

455 the INSU/CNRS, the ANR through the "Projets thématiques d'excellence" program for the
456 "Equipements d'excellence" ASTER-CEREGE action and IRD.

457 7. References

- 458 1. Arnold, M., Merchel, S., Bourlés, D.L., Braucher, R., Benedetti, L., Finkel, R.C.,
459 Aumaître, G., Gott dang, A., Klein, M., 2010. The French accelerator mass
460 spectrometry facility ASTER: Improved performance and developments. *Nuclear
461 Instruments and Methods in Physics Research B* 268, 1954–1959.
- 462 2. Balco, G., Stone, J.O., Lifton, N.A., Dunai, T.J., 2008. A complete and easily
463 accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and
464 ^{26}Al measurements. *Quaternary Geochronology* 3, 174–195.
- 465 3. Balwierz, Z., Roman, M., 2002. A new Eemian Interglacial to Early Vistulian site at
466 Łanięta, central Poland. *Geological Quarterly* 46, 207–217.
- 467 4. Bickel, L., Lüthgens, C., Lomax, J., Fiebig, M., 2015. Luminescence dating of
468 glaciofluvial deposits linked to the penultimate glaciation in the Eastern Alps
469 *Quaternary International* 357, 110–124.
- 470 5. Bitinas, A., 2012. New sights into the last deglaciation of the southeastern flank of the
471 Scandinavian Ice Sheet. *Quaternary Science Reviews* 44, 69–80.
- 472 6. Borchers, B., Marrero, S., Balco, G., Caffee, M., Goehring, B., Lifton, N., Nishiizumi,
473 K., Philips, F., Schaefer, J., Stone, J., 2016. Geological calibration of spallation
474 production rates in the CRONUS-Earth project. *Quaternary Geochronology* 31, 188–
475 198.

- 476 7. Braucher, R., Guillou, V., Bourlès, D. L., Arnold, M., Aumaître, G., Keddadouche, K.,
477 Nottoli, E., 2015. Preparation of ASTER in-house $^{10}\text{Be}/^9\text{Be}$ standard solutions.
478 Nuclear Instruments and Methods in Physics Research B 361, 335–340.
- 479 8. Chmeleff, J., von Blanckenburg, F., Kossert, K., Jakob, D., 2010. Determination of the
480 ^{10}Be half-life by multicollector ICP-MS and liquid scintillation counting. Nuclear
481 Instruments and Methods in Physics Research Section B: Beam Interactions with
482 Materials and Atoms 268, 192–199.
- 483 9. Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B.,
484 Mitrovica, J.X., Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial Maximum.
485 Science 325, 710–714.
- 486 10. Fink, D., Vogt, S., Hotchkis, M., 2000. Cross-sections for ^{36}Cl from Ti at $E_p=35\text{--}150$
487 MeV: Applications to in-situ exposure dating. Nuclear Instruments and Methods in
488 Physics Research Section B: Beam Interactions with Materials and Atoms 172, 861–
489 866.
- 490 11. Galon, R., Roszkówna, L., 1961. Extents of the Scandinavian glaciations and of their
491 recession stages on the territory of Poland in the light of analysis of the marginal
492 forms of inland ice. *Przegląd Geograficzny* 33, 347–364.
- 493 12. Gamrat, W.W., Błaszkiwicz, M., Andrzejewski, L., Krześlak, I., 2017.
494 Asynchronous development of two Late Glacial lake basins near the Drwęca ice-
495 marginal valley (N Poland). *Geological Quarterly* 61, 450–464.
- 496 13. Gogołek, W., Mańkowska, A. 1989. Vistulian of the Turek Upland in the light of new
497 data. *Kwartalnik Geologiczny* 33, 573–586.

- 498 14. Gosse, J.C., Evenson, E.B., Klein, J., Lawn, B., Middleton, R., 1995. Precise
499 cosmogenic ^{10}Be measurements in western North America: support for a global
500 Younger Dryas cooling event. *Geology* 23, 877–880.
- 501 15. Hallet, B., Putkonen, J., 1994. Surface exposure dating of dynamic landforms: young
502 boulders on aging moraines. *Science* 265, 937–940.
- 503 16. Hardt, J., Lüthgens, C., Hebenstreit, R., Böse, M., 2016. Geochronological (OSL) and
504 geomorphological investigations at the presumed Frankfurt ice marginal position in
505 northeast Germany. *Quaternary Science Reviews* 154, 85–99.
- 506 17. Håkansson, L., Alexanderson, H., Hjort, C., Möller, P., Briner, J.P., Aldahan, A.,
507 Possnert, G., 2008. Late Pleistocene glacial history of Jameson land, central east
508 Greenland, derived from cosmogenic ^{10}Be and ^{26}Al exposure dating. *Boreas* 38, 244–
509 260.
- 510 18. Heine, K., Reuther, A.U., Thieke, H.U., Schulz, R., Schlaak, N., Kubik, P.W., 2009.
511 Timing of Weichselian ice marginal positions in Brandenburg (northeastern Germany)
512 using cosmogenic in situ ^{10}Be . *Zeitschrift für Geomorphologie NF* 53, 433–454.
- 513 19. Houmark-Nielsen, M., 2008. Testing OSL failures against a regional Weichselian
514 glaciations chronology from southern Scandinavia. *Boreas* 37, 660–677.
- 515 20. Houmark-Nielsen, M., Kjaer, K.H., 2003. Southwest Scandinavia, 40–15 kyr BP:
516 palaeogeography and environmental change. *Journal of Quaternary Science* 18, 769–
517 786.
- 518 21. Houmark-Nielsen, M., Linge, H., Fabel, D., Schnabel, C., Xu, S., Wilcken, K.M.,
519 Binnie, S., 2012. Cosmogenic surface exposure dating the last deglaciation in

- 520 Denmark: Discrepancies with independent age constraints suggest delayed periglacial
521 landform stabilisation. *Quaternary Geochronology* 13, 1–17.
- 522 22. Hughes, A.L.C., Gyllencreutz, R., Lohne, Ø.S., Mangerud, J., Svendsen, J.I., 2016.
523 The last Eurasian ice sheets e a chronological database and time-slice reconstruction,
524 DATED-1. *Boreas* 45, 1–45.
- 525 23. Hughes, P.D., Gibbard, P.L., 2015. A stratigraphical basis for the Last Glacial
526 Maximum (LGM). *Quaternary International* 383, 174–15.
- 527 24. Hughes, P.D., Gibbard, P.L., Ehlers, J., 2013. Timing of glaciation during the last
528 glacial cycle: evaluating the concept of a global ‘Last Glacial Maximum’ (LGM).
529 *Earth Science Reviews* 125, 171–198.
- 530 25. Kasprzak, L., 1988. Differentiation mechanism in the formation of marginal zones of
531 Leszno and Poznań Phases of the Last Glaciation, Great Poland Lowland.
532 *Dokumentacja Geograficzna* 5–6, 159 pp (in Polish with English summary).
- 533 26. Kenzler, M., Tsukamoto, S., Meng, S., Frechen, M., Hüneke, H., 2017. New age
534 constraints from the SW Baltic Sea area – implications for Scandinavian Ice Sheet
535 dynamics and palaeo-environmental conditions during MIS 3 and early MIS 2. *Boreas*
536 46, 34–52.
- 537 27. Kenzler, M., Tsukamoto, S., Meng, S., Thiel, C., Frechen, M., Hüneke, H., 2015.
538 Luminescence dating of Weichselian interstadial sediments from the German Baltic
539 Sea coast. *Quaternary Geochronology* 30, 215–256.
- 540 28. Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U.C., Knie, K., Rugel,
541 G., Wallner, A., Dillmann, I., Dollinger, G., von Gostomski, C.L., 2010. A new value

- 542 for the half-life of ^{10}Be by Heavy-Ion Elastic Recoil Detection and liquid scintillation
543 counting. *Nuclear Instruments and Methods in Physics Research Section B: Beam*
544 *Interactions with Materials and Atoms* 268, 187–191.
- 545 29. Kozarski, S., 1988. Time and dynamics of the last Scandinavian ice-sheet retreat from
546 northwestern Poland. *Geographia Polonica* 55, 91–101.
- 547 30. Kozarski, S., 1995. Deglaciation of northwestern Poland: environmental conditions
548 and geosystem transformation ~20 ka → 10 ka BP. *Dokumentacja Geograficzna* 1, 82
549 pp. (in Polish).
- 550 31. Kramarska, R., 1998. Origin and development of the Odra bank in the light of the
551 geologic structure and radiocarbon dating. *Geological Quarterly* 42, 277–288.
- 552 32. Krzyszkowski, D., Dobracka, E., Dobracki, R., Czerwonka, J., Kuszell, T., 1999.
553 Stratigraphy of Weichselian deposits in the cliff sections between Łukęcin and
554 Niechorze, Baltic coast, Northwestern Poland. *Quaternary Studies in Poland* 16, 27–
555 45.
- 556 33. Kuszell, T., 1998. New sites of the interglacial deposits in south-western Poland.
557 *Biuletyn Państwowego Instytutu Geologicznego* 385, 127–142 (in Polish).
- 558 34. Lal, D., 1991. Cosmic ray labeling of erosion surfaces: In situ nuclide production rates
559 and erosion models. *Earth Planetary Science Letters* 104, 424–439.
- 560 35. Larsen, E., Fredin, O., Lyså, A., Amantov, A., Fjeldskaar, W., Ottesen, D., 2016.
561 Causes of time-transgressive glacial maxima positions of the last Scandinavian Ice
562 Sheet. *Norwegian Journal of Geology* 96, 159–170.

- 563 36. Lasberg, K., Kalm, V., 2013. Chronology of Late Weichselian glaciation in the
564 western part of the East European Plain. *Boreas* 42, 995–1007.
- 565 37. Lüthgens, C., Böse, M., Preusser, F., 2011. Age of the Pomeranian ice-marginal
566 position in northeastern Germany determined by Optically Stimulated Luminescence
567 (OSL) dating of glaciofluvial sediments. *Boreas* 40, 598–615.
- 568 38. Majdanowski, S., 1947. Rozmieszczenie, gęstość i kierunki rynien jeziornych na Niziu
569 Polskim. *Przegląd Geograficzny* 21, 37–69.
- 570 39. Marks, L., 2002. Last Glacial Maximum in Poland. *Quaternary Science Reviews* 21,
571 103–110.
- 572 40. Marks, L., 2011. Quaternary glaciations in Poland. In: Ehlers, J., Gibbard, P.L.,
573 Hughes, P.D. (eds.), *Quaternary Glaciations – Extent and Chronology: A Closer Look*.
574 *Developments in Quaternary Science*, 15. Elsevier, Amsterdam, pp. 299–304.
- 575 41. Marks, L., 2012. Timing of the Late Vistulian (Weichselian) glacial phases in Poland.
576 *Quaternary Science Reviews* 44, 81–88.
- 577 42. Marks, L., 2015. Last deglaciation of northern continental Europe. *Cuadernos de*
578 *Investigación Geográfica* 41(2), 279–293.
- 579 43. Marks, L., Ber A., Gogolek W., Piotrowska, K. (eds.), 2006. *Mapa Geologiczna Polski*
580 *1:500 000*. Ministerstwo Środowiska, PIG-PIB, Warszawa.
- 581 44. Marrero, S.M., Philips, F.M., Borchers, B., Lifton, N., Aumer, R., Balco, G., 2016a.
582 *Cosmogenic nuclide systematics and the CRONUScale program*. *Quaternary*
583 *Geochronology* 31, 160–187.

- 584 45. Marrero, S.M., Phillips, F.M., Caffee, M.W., Gosse, J.C., 2016b. CRONUS-Earth
585 cosmogenic ³⁶Cl calibration. *Quaternary Geochronology* 31, 199–219.
- 586 46. Mix, A.C., Bard, E., Schneider, R., 2001. Environmental processes of the ice age:
587 land, oceans, glaciers (EPILOG). *Quaternary Science Reviews* 20, 627–657.
- 588 47. Molewski P, 2007. Neotectonic and glaciodynamic conditions for the formation of the
589 Pleistocene of the Kujawy Moraine Plateau). *Wydawnictwo Naukowe Uniwersytetu*
590 *Mikołaja Kopernika, Toruń*, 140 pp. (in Polish).
- 591 48. Molewski, P., Weckwerth, P., 2009. Explanatory text to the Detailed geological map
592 of Poland in the scale 1 : 50 000, ark. Toruń. *Centr. Arch. Geol. PIG-PIB, Warszawa*
593 (in Polish).
- 594 49. Narloch, W., Wysota, W., Piotrowski, J.A., 2013. Sedimentological record of
595 subglacial conditions and ice sheet dynamics of the Vistula Ice Stream (north-central
596 Poland) during the Last Glaciation. *Sedimentary Geology* 293, 30–44.
- 597 50. Niewiarowski, W., 1995. Fluctuations of the water level in the Biskupin lake and their
598 reasons. In: Niewiarowski W. (ed.) *Outline of geographical environment`s changes*
599 *under the influence of natural and anthropogenic factors in Late Glacial and Holocene:*
600 *215–234. Wydawnictwo Turpress, Toruń* (in Polish).
- 601 51. Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P.L, Stroeven, A.P.,
602 Shackleton, C., Winsborrow, M., Heyman, J., Hall, A.M., 2017. Deglaciation of the
603 Eurasian ice sheet complex. *Quaternary Science Reviews* 169, 148–172.
- 604 52. Pisarska-Jamroży, M., Belżyty, Sz., Börner, A., Hoffmann, G., Hüneke, H., Kenzler M.,
605 Obst, K., Rother, H., van Loon, A.J. (Tom), 2018. Evidence from seismites for glacio-

- 606 isostatically induced crustal faulting in front of an advancing land-ice mass (Rügen
607 Island, SW Baltic Sea). *Tectonophysics* 745, 338–348.
- 608 53. Plug, L.J., Gosse, J.C., McIntosh, J.J., Bigley, R., 2007. Attenuation of cosmic ray flux
609 in temperate forests. *Journal of Geophysical Research* 112,
610 <http://dx.doi.org/10.1029/2006JF000668>.
- 611 54. Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck,
612 C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P.,
613 Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G.,
614 Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W.,
615 Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht,
616 J., 2013. INTCAL13 AND MARINE13 radiocarbon age calibration curves 0-50,000
617 years cal BP. *Radiocarbon* 55, 1869–1887.
- 618 55. Rinterknecht V.R., Bitinas, A., Clark, P.U., Raisbeck, G.M., Yiou, F. & Brook, E.J.,
619 2008, Timing of the last deglaciation in Lithuania, *Boreas* 37, 426–433.
- 620 56. Rinterknecht, V.R., Börner, A., Bourlès, D., Braucher, R., 2014. Cosmogenic ^{10}Be
621 dating of ice sheet marginal belts in Mecklenburg-Vorpommern, Western Pomerania
622 (northeast Germany). *Quaternary Geochronology* 19, 42–51.
- 623 57. Rinterknecht, V., Braucher, R., Böse, M., Bourlès, D., Mercier, J.L., 2012. Late
624 Quaternary ice sheet extents in northeastern Germany inferred from surface exposure
625 dating. *Quaternary Science Reviews* 44, 89–95.
- 626 58. Rinterknecht, V.R., Marks, L., Piotrowski, J.A., Raisbeck, G.M., Yiou, F., Brook, E.J.,
627 Clark, P.U., 2005. Cosmogenic ^{10}Be ages on the Pomeranian moraine, Poland. *Boreas*
628 34, 186–191.

- 629 59. Rinterknecht, V.R., Clark, P.U., Raisbeck, G.M., Yiou, F., Bitinas, A., Brook, E.J.,
630 Marks, L., Zelčs, V., Lunkka, J.-P., Pavlovskaya, I.E., Piotrowski, J.A., Raukas, A.,
631 2006. The last deglaciation of the southeastern sector of the Scandinavian ice sheet.
632 Science 311, 1449–1452.
- 633 60. Rinterknecht, V., Hang, T., Gorlach, A., Kohv M., Kalla, K., Kalm, V., Subetto, D.,
634 Bournès, D., Léanni, L., Guillou, V., ASTER Team, 2018. The Last Glacial Maximum
635 extent of the Scandinavian Ice Sheet in the Valdai Heights, western Russia: Evidence
636 from cosmogenic surface exposure dating using ^{10}Be . Quaternary Science Reviews
637 200, 106–113.
- 638 61. Rinterknecht V.R., I.E. Pavlovskaya, P.U. Clark, G.M. Raisbeck, F. Yiou, E.J. Brook,
639 2007, Timing of the Last Deglaciation in Belarus, Boreas 36, 307–313.
- 640 62. Roman, M., 2017. Ice-flow directions of the last Scandinavian Ice Sheet in central
641 Poland, Quaternary International, <https://doi.org/10.1016/j.quaint.2017.11.035>.
- 642 63. Rotnicki, K., Borówka, R.K., 1995. The Last Cold Period in the Gardno-Łeba Coastal
643 Plain. Journal of Coastal Research 22, 225–229.
- 644 64. Schimmelpfennig, I., Benedetti, L., Finkel, R., Pik, R., Blard, P.H., Bournès, D.,
645 Burnard, P., Williams, A., 2009. Sources of in-situ ^{36}Cl in basaltic rocks. Implications
646 for calibration of production rates. Quaternary Geochronology 4, 441–461.
- 647 65. Schimmelpfennig, I., Benedetti, L., Garreta, V., Pik, R., Blard, P.H., Burnard, P.,
648 Bournès, D., Finkel, R., Ammon, K., Dunai, T., 2011. Calibration of cosmogenic ^{36}Cl
649 production rates from Ca and K spallation in lava flows from Mt. Etna (38°N, Italy)
650 and Payun Matru (36°S, Argentina). Geochimica Cosmochimica Acta 75, 2611–2632.

- 651 66. Schimmelpfennig, I., Schaefer, J.M., Putnam, A.E., Koffman, T., Benedetti, L., Ivy-
652 Ochs, S., Team, A., Schlüchter, C., 2014. ^{36}Cl production rate from K-spallation in the
653 European Alps (Chironico landslide, Switzerland). *Journal of Quaternary Science* 29,
654 407–413.
- 655 67. Spagnolo, M., Philips, E., Piotrowski, J.A., Rea, B.R., Clark, C.D., Stokes, C.R., Carr,
656 S.J., Ely, J.C., Ribolini, A., Wysota, W., Szuman, I., 2016. Ice stream motion
657 facilitated by a shallow-deforming and accreting bed. *Nature Communications* 7,
658 10723, doi:10.1038/ncomms10723.
- 659 68. Stankowska, A., Stankowski, W., 1988. Maximum extent of the Vistulian ice sheet in
660 the vicinity of Konin, Poland: a geomorphological, sedimentological and radiometric
661 evidence. *Geographia Polonica* 55, 141–150.
- 662 69. Stankowski, W., Bluszcz, A., Nita, M., 1999. Upper Quaternary deposits sites
663 Mikorzyn and Sławoszewek in the light of geological investigation, radiocarbon and
664 thermoluminescence dating, and palynological analysis. In: Pazur, A., Bluszcz, A.,
665 Stankowski, W., Starkel, L. (eds.), *Geochronology of Upper Quaternary of Poland in*
666 *the light of radiocarbon and luminescence dating*. Institute of Physics Silesian
667 University of Technology Press, Gliwice, 87–111 (in Polish).
- 668 70. Stone, J., 2000. Air pressure and cosmogenic isotope production. *Journal of*
669 *Geophysical Research* 105, 23753–23760.
- 670 71. Stone, J.O., Fifield, K., Vasconcelos, P., 2005. Terrestrial chlorine-36 production
671 from spallation of iron, in: *Abstract of 10th International Conference on Accelerator*
672 *Mass Spectrometry*. Berkeley, CA.

- 673 72. Stroeven, A.P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O.,
674 Goodfellow, B.W., Harbor, J.M., Jansen, J.D., Olsen, L., Caffee, M.W., Fink, D.,
675 Lundqvist, J., Rosqvist, G.C., Strömberg, B., Jansson, K.N., 2016. Deglaciation of
676 Fennoscandia. *Quaternary Science Reviews* 147, 91–121.
- 677 73. Stroeven, A.P., Heyman, J., Fabel, D., Björck, S., Caffee, M.W., Fredin, O., Harbore,
678 J.M. 2015. A new Scandinavian reference ^{10}Be production rate. *Quaternary*
679 *Geochronology* 29, 104–115.
- 680 74. Stuiver, M., Reimer, P.J., Reimer, R.W., 2017. CALIB 7.1 [WWW program] at
681 <http://calib.org>, accessed 2017-1-19.
- 682 75. Szałamacha, G., Skompski, S., 1999. Biogenic sediments of the Eemian Interglacial at
683 Krzyżówki near Koło, central Poland. *Geological Quarterly* 43, 99–106.
- 684 76. Tobolski, K., 1986. Paleobotanical studies of the Eemian interglacial and early
685 Vistulian, Władysławów in the vicinity of Turek (preliminary report). *Quaternary*
686 *Studies in Poland* 7, 1–101.
- 687 77. Tylmann, K., Woźniak, P.P., Rinterknecht V., 2018. Erratics selection for cosmogenic
688 nuclide exposure dating – an optimization approach. *Baltica* 31(2), 100–114.
- 689 78. Weckwerth, P., Przegiętka, K., Chruścińska, A., Woronko, B., Oczkowski, H.L., 2011.
690 Age and sedimentological features of fluvial series in the Toruń basin and the Drwęca
691 valley (Poland). *Geochronometria* 38, 397–412.
- 692 79. Woźniak, P.P., Czubla P., 2015. The Late Weichselian glacial record in northern
693 Poland – towards a wider perspective: a new look at debris transport routes by the FIS.
694 *Quaternary International* 386, 3–17.

- 695 80. Wysota, W., Molewski, P., Sokołowski, R.J., 2009. Record of the Vistula ice lobe
696 advances in the Late Weichselian glacial sequence in north-central Poland. *Quaternary*
697 *International* 207, 26–41.
- 698 81. Zimenkov, O., 1989. Age of the maximum extent of the Poozerian Glaciation in
699 Belarus. New data about Cenozoic geology of Belarus and adjacent areas. In.
700 Matveyev, A. (ed.) *Nauka i tekhnika*, 30–44, Minsk, Belarus (in Russian).

Table 1. Published radiocarbon ages significant for the timing of the last SIS advance and retreat along the LLGM limit in Poland.

Laboratory symbol	¹⁴ C age (ka BP)	Calibrated age (cal ka BP)*	Dated material	Locality	Reference
Post-LLGM ages					
Poz-46881	15.9 ± 0.1	19.1 ± 0.2	dispersed organic matter in laminated lacustrine silt	Piotrowo (Poland)	Gamrat et al. (2017)
Gd-5777	17.7 ± 0.2	21.4 ± 0.6	dispersed organic matter in lacustrine silt	Biskupińskie Lake (Poland)	Niewiarowski (1995)
Gd-9604	17.7 ± 0.8	21.4 ± 1.9	peat at the bottom of gytja	Sławoszewek (Poland)	Stankowski et al. (1999)
Pre-LLGM ages					
Gd-786	21.2 ± 1.6	25.2 ± 3.4	dispersed organic matter in silt below till	Żdzary (Poland)	Gogolek and Mańkowska (1989)
Gd-1038	21.5 ± 0.4	25.6 ± 1.1	dispersed organic matter in silt below till	Odra Bank (Poland)	Kramarska (1998)
Gd-9686	21.6 ± 1.1	25.6 ± 2.2	organic matter below till	Niechorze (Poland)	Krzyszowski et al. (1999)
Gd-645	22.1 ± 0.5	26.4 ± 0.9	peat below till	Maliniec (Poland)	Stankowska and Stankowski (1988)
Gd-4581	22.3 ± 0.7	26.5 ± 1.3	dispersed organic matter in silt	Łeba Barrier (Poland)	Rotnicki and Borówka (1995)

* Calibrated using IntCal13 calibration curve (Reimer et al., 2013) with the Calib 7.0.4 software (Stuiver et al., 2017).

Table 2. Surface exposure ^{10}Be ages of pre-LLGM and LLGM erratic boulders in northern Poland. The list consists of 33 new exposure ages and 10 ^{10}Be ages recalculated from the original data of Rinterknecht et al. (2005, 2006). Densities of all samples (granitoids, granite gneisses and gneisses) were assumed at 2.7 g/cm^3 . All ^{10}Be exposure ages are calculated with ‘Lm’ time-dependent scaling scheme for spallation according to Lal (1991) and Stone (2000) and global production rate according to Borchers et al. (2016).

Sample ID	Latitude DD	Longitude DD	Elevation (m a.s.l.)	Boulder perimeter/height (m)	Boulder lithology	Sample thickness (cm)	Shielding factor ¹	Quartz (g)	Spike Be (g)	[^{10}Be] (10^4 at g^{-1})	Age (ka) $\varepsilon = 1.3$ (mm ka^{-1})	Age (ka) $\varepsilon = 0$ (mm ka^{-1})
Pre-LLGM erratics (Late Saalian)												
SAAL-01	51.9411	17.2008	122	9.7/1.3	granitoid	1.6	1.0000	18.38	0.1015	25.35 ± 0.81	57.9 ± 5.1	54.5 ± 4.5
SAAL-02	51.9410	17.2008	122	11.1/1.7	granitoid	1.2	0.9938	17.38	0.1018	27.76 ± 0.88	64.1 ± 5.7	59.8 ± 4.9
SAAL-03	51.9690	17.4716	120	9.5/1.1	granitoid	1.0	0.9975	16.00	0.1017	31.04 ± 1.00	72.2 ± 6.5	66.8 ± 5.5
SAAL-04	52.0883	18.3700	116	7.0/1.2	granitoid	2.5	1.0000	19.11	0.1005	10.30 ± 0.40	22.9 ± 2.0	22.3 ± 1.9
SAAL-05	52.1714	19.2166	138	8.8/1.4	granitoid	1.0	1.0000	17.09	0.1032	47.95 ± 1.51	115.2 ± 11.0	101.7 ± 8.5
SAAL-06	52.7110	19.9693	131	13.0/1.2	granitoid	1.0	1.0000	17.50	0.1019	7.58 ± 0.32	16.2 ± 1.4	15.9 ± 1.4
SAAL-07	53.2182	20.0510	157	9.5/1.9	granitoid	1.5	1.0000	15.23	0.1016	17.93 ± 0.79	38.3 ± 3.5	36.8 ± 3.2
LLGM ice marginal belts												
Leszno (Brandenburg) Phase												
											Mean ^{10}Be age (n = 6): 20.7 ± 0.8	
LGM-01	52.3703	16.2724	99	8.7/0.8	gneiss	2.5	1.0000	15.53	0.1017	9.14 ± 0.50	20.5 ± 2.0	$20.1 \pm 1.9^\wedge$
LGM-02T	52.1739	15.6817	133	6.0/2.0	granite gneiss	0.9	1.0000	15.01	0.1004	18.82 ± 0.60	41.3 ± 3.6	39.5 ± 3.3
LGM-02S	52.1739	15.6817	133	6.0/2.0	granite gneiss	1.1	1.0000	15.87	0.1019	27.74 ± 0.93	62.7 ± 5.6	58.6 ± 4.9
LGM-03	52.1185	15.6663	111	10.5/2.3	granite gneiss	2.2	1.0000	5.83	0.1021	8.91 ± 0.70	19.7 ± 2.2	$19.3 \pm 2.1^\wedge$
LGM-04	51.9690	17.1330	120	13.5/1.2	granitoid	1.1	1.0000	18.32	0.1007	19.80 ± 0.63	44.4 ± 3.8	42.3 ± 3.5
LGM-05	51.9609	17.1492	111	10.4/1.3	granitoid	1.6	1.0000	17.98	0.1036	10.08 ± 0.50	22.3 ± 2.0	$21.8 \pm 1.9^\wedge$
LGM-06	52.1457	18.3311	159	6.8/1.0	granitoid	1.4	1.0000	18.20	0.1024	25.08 ± 0.80	54.9 ± 4.8	51.7 ± 4.3
LGM-07	52.1303	18.3199	156	7.8/1.5	granitoid	1.3	1.0000	14.67	0.1024	21.31 ± 0.76	46.2 ± 4.1	43.9 ± 3.7
LGM-08	52.1511	18.3218	160	6.3/1.2	granitoid	1.6	0.9980	16.96	0.1019	30.68 ± 0.97	68.4 ± 6.1	63.6 ± 5.3
FRT-01	52.4411	15.0488	140	6.5/1.3	granitoid	0.8	1.0000	15.17	0.1024	10.13 ± 0.54	21.5 ± 2.0	$21.0 \pm 1.9^\wedge$
FRT-02	52.3949	15.0943	133	7.0/0.8	granitoid	1.0	0.9981	14.91	0.1024	35.51 ± 1.22	82.2 ± 7.6	75.3 ± 6.3
FRT-03	52.4471	15.1529	161	10.1/1.8	granitoid	0.4	1.0000	15.95	0.1023	10.19 ± 0.72	21.1 ± 2.2	$20.6 \pm 2.1^\wedge$
FRT-04	52.4489	15.1606	145	12.7/2.5	granitoid	2.5	1.0000	16.47	0.1036	11.96 ± 0.42	25.8 ± 2.2	25.1 ± 2.1
FRT-06	52.4426	15.2819	177	8.2/1.6	granitoid	1.2	0.9988	16.11	0.1032	10.45 ± 0.53	21.5 ± 2.0	$21.0 \pm 1.9^\wedge$

Poznań (Frankfurt) Phase											Mean ¹⁰ Be age (n = 14): 17.3 ± 0.5		
LGM-09	52.7390	19.6783	143	14.3/1.4	granitoid	1.0	1.0000	15.21	0.1036	15.93 ± 0.77	34.3 ± 3.2	33.0 ± 3.0	
LGM-10	52.8749	19.6290	114	9.0/0.8	granitoid	1.0	1.0000	17.31	0.1021	7.95 ± 0.38	17.3 ± 1.6	17.0 ± 1.5 [^]	
LGM-11	53.4187	19.9013	166	9.5/1.2	granitoid	1.1	1.0000	14.68	0.1032	8.67 ± 0.47	17.8 ± 1.7	17.5 ± 1.6 [^]	
LGM-12	53.3874	19.5752	130	8.0/1.0	granitoid	1.2	1.0000	15.07	0.1026	11.50 ± 0.53	24.8 ± 2.3	24.1 ± 2.1	
FRT-05	52.4897	15.1843	113	8.7/1.6	granitoid	1.1	1.0000	15.60	0.1023	14.13 ± 1.06	31.3 ± 3.5	30.2 ± 3.2	
FRT-07	53.2823	17.5074	116	8.9/1.2	granitoid	1.0	1.0000	9.71	0.3080	8.91 ± 1.07	19.3 ± 2.8	18.9 ± 2.7 [^]	
FRT-08	52.9537	17.1255	89	17.5/2.2	gneiss	2.5	1.0000	15.03	0.1023	7.74 ± 0.43	17.5 ± 1.7	17.1 ± 1.6 [^]	
FRT-09	52.9525	17.3370	94	13.5/1.3	granitoid	1.2	1.0000	15.65	0.1025	7.66 ± 0.38	17.0 ± 1.6	16.7 ± 1.5 [^]	
FRT-10	52.7259	17.3231	105	20.5/1.1	granitoid	0.7	1.0000	16.46	0.1025	7.89 ± 0.65	17.3 ± 2.0	16.9 ± 1.9 [^]	
FRT-12	53.5238	16.2029	150	19.0/3.4	gneiss	1.6	0.9996	22.03	0.3049	9.81 ± 0.76	20.6 ± 2.3	20.2 ± 2.2 [^]	
FRT-13	53.6903	17.1372	171	20.5/1.9	gneiss	2.0	1.0000	9.29	0.3055	9.69 ± 0.57	20.0 ± 2.0	19.5 ± 1.9 [^]	
FRT-14	53.7271	17.1342	187	14.6/2.1	granitoid	2.5	0.9994	20.20	0.3052	8.55 ± 0.54	17.4 ± 1.7	17.0 ± 1.7 [^]	
LES-3	53.4806	20.2222	212	n.a./1.6	granitoid	2.0	1.0000	39.99	0.2500	3.45 ± 0.35	7.1 ± 0.9	7.0 ± 0.9	
LES-4	53.4944	20.2389	172	n.a./1.8	granitoid	2.0	1.0000	40.00	0.2500	7.40 ± 0.54	15.8 ± 1.7	15.5 ± 1.6 ^{^*}	
LES-5	53.5792	20.0944	180	11.7/2.5	granitoid	2.0	1.0000	40.00	0.2500	19.24 ± 1.16	42.5 ± 4.3	40.6 ± 3.9 [*]	
LES-6	53.6006	20.0611	151	17.1/2.2	gneiss	2.0	1.0000	40.00	0.2500	8.08 ± 0.58	17.7 ± 1.9	17.4 ± 1.8 ^{^*}	
LES-7	53.6250	20.0417	132	13.0/2.8	granitoid	2.0	1.0000	60.51	0.2500	2.64 ± 0.33	5.8 ± 0.9	5.8 ± 0.8 [*]	
LES-8	53.5111	19.9000	255	4.5/1.1	granitoid	2.0	1.0000	40.00	0.2500	10.14 ± 1.10	20.1 ± 2.7	19.7 ± 2.6 ^{^*}	
LES-10	53.5764	19.9417	270	8.7/1.7	granitoide	2.0	1.0000	40.01	0.2500	6.78 ± 0.57	13.2 ± 1.5	13.0 ± 1.5 [*]	
LES-11	53.5222	19.8375	218	10.5/1.5	gneiss	2.0	1.0000	39.99	0.2500	7.94 ± 0.77	16.3 ± 2.0	16.0 ± 2.0 ^{^*}	
LES-12	53.5625	19.9528	275	10.1/1.6	granitoid	2.0	1.0000	40.01	0.2500	8.46 ± 0.70	16.4 ± 1.9	16.1 ± 1.8 ^{^*}	
LES-13	53.5528	19.9250	302	11.0/1.2	granitoid	2.0	1.0000	40.01	0.2500	19.15 ± 1.33	37.0 ± 4.0	35.5 ± 3.7 [*]	

All samples measured at the ASTER facility. AMS results are standardized to STD-11 (samples LGM and FRT) and SRM 4325 (samples LES). ¹⁰Be/⁹Be ratios were corrected for a process blank value of 6.08 x 10⁻¹² (samples LGM and FRT-01 to FRT-10), 5.97 x 10⁻¹² (samples FRT-11 to FRT-14), and 26.8 ± 1.4 x 10⁻¹² (samples LES).

¹ Corresponding to self-shielding (direction and angle of surface dipping).

* Surface exposure ages recalculated from the original data of Rinterknecht et al. (2005, 2006).

[^] Ages used in the error-weighted mean age calculation for the LLGM moraines.

n.a. – not available

Table 3. Bulk composition of sample FRT-11 before chemical treatment, analysed at the SARM-CRPG (Nancy, France) by ICP-OES (major elements), ICP-MS (trace element), atomic absorption (Li), colorimetry (B) and spectrophotometry (Cl).

Sample Name	SiO ₂ %	Al ₂ O ₃ %	Fe ₂ O ₃ %	MnO %	MgO %	CaO %	Na ₂ O %	K ₂ O %	TiO ₂ %	P ₂ O ₅ %	LOI %	Total	Cl (ppm)	Li (ppm)	B (ppm)	Sm (ppm)	Gd (ppm)	Th (ppm)	U (ppm)
FRT-11	50.46	23.48	6.17	0.08	1.38	10.38	3.90	0.77	1.12	0.44	1.10	99.26	13	6.9	3.4	3.4	3.4	1.17	0.25

Table 4. Major element concentrations in sample splits after acid etching, which removed 14% of the sample, analysed at the SARM-CRPG (Nancy, France) by ICP-OES. Ca, K, Ti and Fe are target elements for the spallogenic/muogenic production of ³⁶Cl.

Sample name	SiO ₂ %	Al ₂ O ₃ %	Fe ₂ O ₃ %	MnO %	MgO %	CaO %	Na ₂ O %	K ₂ O %	TiO ₂ %	P ₂ O ₅ %	LOI %	Total
FRT-11	51.52 ± 0.51	23.61 ± 0.23	4.56 ± 0.09	0.069 ± 0.007	0.99 ± 0.01	11.10 ± 0.22	4.04 ± 0.61	0.71 ± 0.04	1.43 ± 0.07	0.88 ± 0.18	0.59	99.51

Table 5. Surface exposure ³⁶Cl age of LLGM gabbro erratic with characteristics of FRT-11 sample. Sample and blank were spiked with a solution enriched in ³⁵Cl (99.9%). AMS measurements were normalized to in-house standard SM-CL-12, using an assigned value of 1.428 (± 0.021) x 10¹² (Merchel et al., 2011), and assuming a natural ratio of 3.127 for the stable ratio ³⁵Cl/³⁷Cl. The chemistry blank correction was performed by subtracting the number of atoms of Cl and ³⁶Cl in the blank from those in the sample.

Sample ID	Latitude DD	Longitude DD	Elevation (m a.s.l.)	Boulder perimeter/height (m)	Boulder lithology	Shielding factor	Sample thickness (cm)	Sample density (g cm ⁻³)	Sample weight (g)	[Cl] (ppm)
FRT-11	52.8035	17.9141	107	13.1/2.9	gabbro	1.0000	1.9	3.0	48.75	101.7 ± 8.6
Sample ID	Mass of Cl in spike (mg)	^[35Cl/37Cl]	^[36Cl/35Cl] (10 ⁻¹⁴)	Number of atoms Cl (10 ¹⁷)*	^[36Cl] (10 ⁴ atoms)*	^[36Cl] (10 ⁴ at g ⁻¹)	Excel® spreadsheet		CRONUScale	
							Age (ka) ε = 1.3 (mm yr ⁻¹)	Age (ka) ε = 0 (mm yr ⁻¹)	Age (ka) ε = 1.3 (mm yr ⁻¹)	Age (ka) ε = 0 (mm yr ⁻¹)
FRT-11	1.601	4.470 ± 0.093	10.27 ± 0.64	842 ± 71	939 ± 81	19.0 ± 1.6	19.2 ± 3.0	20.1 ± 3.1	18.7 ± 2.9	20.0 ± 3.0
Blank	1.599	590 ± 16	0.447 ± 0.081	0.691 ± 0.063	12.3 ± 2.2	-	-	-	-	-

* Sample values are not yet corrected for blank.